
A
ATM alterations by histology
LUAD vs. LUSC (Fisher's exact test):
any p = 0.002; truncated p = 0.0003
Samples with ATM alteration (%)
ATM-altered (any)
ATM-truncated
7.7%
(n = 311)
3.0%
(n = 120)
4.1%
(n = 22)
0.6%
(n = 3)
Lung adeno-
carcinoma
(n = 4,064)
Squamous cell
carcinoma
(n = 537)
B
Overall survival by ATM status (LUAD, n = 4,041)
ATM wild-type (n = 3,733)
ATM-altered, missense/in-frame only (n = 189)
ATM-truncated (n = 119)
Overall survival probability
log-rank vs. wild-type: any alteration p = 0.10; truncated p = 0.13
median OS: wild-type 40.8, missense/in-frame 38.1, truncated 29.5 months
Time from sequencing (months)
C
ATM-truncated (nonsense, frameshift, splice; n = 120) vs. ATM wild-type (n = 3,753)
ATM-altered (any non-synonymous variant; n = 311) vs. ATM wild-type
KRAS mutation
STK11 mutation
KEAP1 mutation
TP53 mutation
EGFR mutation
BRAF mutation
MET exon 14
PIK3CA mutation
MET amplification
CDKN2A deletion
group vs. WT OR Fisher p
71.7% vs 30.7% 5.71 2e-19
57.9% vs 30.7% 3.10 3e-21
39.2% vs 14.8% 3.71 2e-10
27.0% vs 14.8% 2.13 1e-7
15.8% vs 13.6% 1.19 0.50
15.8% vs 13.6% 1.18 0.30
25.8% vs 48.4% 0.37 9e-7
34.4% vs 48.4% 0.56 2e-6
5.8% vs 27.8% 0.16 2e-9
8.4% vs 27.8% 0.24 2e-16
2.5% vs 3.6% 0.68 0.80
1.9% vs 3.6% 0.52 0.15
0.8% vs 3.4% 0.24 0.19
1.0% vs 3.4% 0.28 0.02
2.5% vs 4.2% 0.58 0.49
2.6% vs 4.2% 0.60 0.18
1.7% vs 2.2% 0.75 1.00
1.9% vs 2.2% 0.87 1.00
8.3% vs 11.1% 0.73 0.46
8.7% vs 11.1% 0.76 0.22
Odds ratio vs. ATM wild-type LUAD (95% CI)