

Online learning in repeated matrix games

Yoav Freund

February 24, 2020

Based on “Adaptive Game Playing Using Multiplicative Weights” Freund and Schapire.

Outline

Repeated Matrix Games

Outline

Repeated Matrix Games

Specific games

Outline

Repeated Matrix Games

Specific games

Minmax vs. Regret

Outline

Repeated Matrix Games

Specific games

Minmax vs. Regret

Fictitious play

Outline

Repeated Matrix Games

Specific games

Minmax vs. Regret

Fictitious play

Strategy using Hedge

Outline

Repeated Matrix Games

Specific games

Minmax vs. Regret

Fictitious play

Strategy using Hedge

The basic analysis

Outline

Repeated Matrix Games

Specific games

Minmax vs. Regret

Fictitious play

Strategy using Hedge

The basic analysis

Proof of minmax theorem

Outline

- Repeated Matrix Games

- Specific games

- Minmax vs. Regret

- Fictitious play

- Strategy using Hedge

- The basic analysis

- Proof of minmax theorem

- Approximately solving games

 - Fixed Learning rate

 - Variable learning rate

Zero sum games in matrix form

- ▶ Game between two players.

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ **Row** player chooses $i \in \{1, \dots, n\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix **M**
- ▶ **Row** player chooses $i \in \{1, \dots, n\}$
- ▶ **Column** player chooses $j \in \{1, \dots, m\}$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$

Zero sum games in matrix form

- ▶ Game between two players.
- ▶ Defined by $n \times m$ matrix \mathbf{M}
- ▶ Row player chooses $i \in \{1, \dots, n\}$
- ▶ Column player chooses $j \in \{1, \dots, m\}$
- ▶ Row player gains $\mathbf{M}(i, j) \in [0, 1]$
- ▶ Column player loses $\mathbf{M}(i, j)$
- ▶ Game repeated many times.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**


Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .

Pure vs. mixed strategies

- ▶ Choosing a **single** action = **pure** strategy.
- ▶ Choosing a **Distribution** over actions = **mixed** strategy.
- ▶ **Row** player chooses dist. over rows **P**
- ▶ **Column** player chooses dist. over columns **Q**
- ▶ **Row** player gains **$M(P, Q)$** .
- ▶ **Column** player loses **$M(P, Q)$** .


Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

- **Q** is a **column** vector. **P^T** is a row vector.

Mixed strategies in matrix notation



$$(A \times B)_{12} = \sum_{r=1}^4 a_{1r} b_{r2} = a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} + a_{14}b_{42}$$

- ▶ **Q** is a **column** vector. **P^T** is a row vector.
- ▶ **M(P, Q) = P^TMQ = $\sum_{i=1}^n \sum_{j=1}^m \mathbf{P}(i)\mathbf{M}(i,j)\mathbf{Q}(j)$**

The minmax Theorem

When using pure strategies, second player has an advantage.

The minmax Theorem

When using pure strategies, second player has an advantage.

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

In words:

- ▶ for **pure** strategies, choosing second can be better.

The minmax Theorem

When using pure strategies, second player has an advantage.

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

In words:

- ▶ for **pure** strategies, choosing second can be better.
- ▶ for **mixed** strategies, choosing second gives no advantage.

The minmax Theorem

When using pure strategies, second player has an advantage.

John von Neumann, 1928.

$$\min_P \max_Q M(P, Q) = \max_Q \min_P M(P, Q)$$

In words:

- ▶ for **pure** strategies, choosing second can be better.
- ▶ for **mixed** strategies, choosing second gives no advantage.
- ▶ There are min-max optimal mixed Strategies: **P^*, Q^***

The minmax Theorem

When using pure strategies, second player has an advantage.

John von Neumann, 1928.

$$\min_P \max_Q M(P, Q) = \max_Q \min_P M(P, Q)$$

In words:

- ▶ for **pure** strategies, choosing second can be better.
- ▶ for **mixed** strategies, choosing second gives no advantage.
- ▶ There are min-max optimal mixed Strategies: **P^*, Q^***
- ▶ **$M(P^*, Q^*)$** is the **value** of the game.

Online Learning as matrix game

- ▶ Row = action

	$t = 1$	$t = 2$...
<i>expert1</i>	0	1	...
<i>expert2</i>	0.2	0.1	...
<i>expert3</i>	0.5	0.2	...
...
<i>Master</i>	0.35	0.13	...

Online Learning as matrix game

- ▶ Row = action
- ▶ Column = iteration.

	$t = 1$	$t = 2$...
<i>expert1</i>	0	1	...
<i>expert2</i>	0.2	0.1	...
<i>expert3</i>	0.5	0.2	...
...
<i>Master</i>	0.35	0.13	...

Online Learning as matrix game

- ▶ Row = action
- ▶ Column = iteration.
- ▶ Player chooses mixed strategy \mathbf{P}_t

	$t = 1$	$t = 2$...
<i>expert1</i>	0	1	...
<i>expert2</i>	0.2	0.1	...
<i>expert3</i>	0.5	0.2	...
...
<i>Master</i>	0.35	0.13	...

Online Learning as matrix game

- ▶ Row = action
- ▶ Column = iteration.
- ▶ Player chooses mixed strategy \mathbf{P}_t
- ▶ adversary chooses pure strategy
 $\mathbf{Q}_t = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ the 1 is at position t

	$t = 1$	$t = 2$...
<i>expert1</i>	0	1	...
<i>expert2</i>	0.2	0.1	...
<i>expert3</i>	0.5	0.2	...
...
<i>Master</i>	0.35	0.13	...

Online Learning as matrix game

- ▶ Row = action
- ▶ Column = iteration.
- ▶ Player chooses mixed strategy \mathbf{P}_t
- ▶ adversary chooses pure strategy
 $\mathbf{Q}_t = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ the 1 is at position t
- ▶ Goal - minimize regret: $\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) - \sum_{t=1}^T \mathbf{M}(\mathbf{P}^*, \mathbf{Q}_t)$

	$t = 1$	$t = 2$...
<i>expert1</i>	0	1	...
<i>expert2</i>	0.2	0.1	...
<i>expert3</i>	0.5	0.2	...
...
<i>Master</i>	0.35	0.13	...

Boosting as a matrix game (1)

- ▶ Row = example (x, y)

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Boosting as a matrix game (1)

- ▶ Row = example (x, y)
- ▶ Column = Weak Rule h_t

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Boosting as a matrix game (1)

- ▶ Row = example (x, y)
- ▶ Column = Weak Rule h_t
- ▶ Matrix entry for $(x, y), h_t$ is 0 if $h_t(x) = y$, 1 $h_t(x) \neq y$

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Boosting as a matrix game (2)

- ▶ Boosting assumption: for any distribution over examples, there exists a weak rule with weighted error $< 1/2$

Boosting as a matrix game (2)

- ▶ Boosting assumption: for any distribution over examples, there exists a weak rule with weighted error $< 1/2$
- ▶ In game terms: For any **mixed strategy** of the row player **P**, there is a **pure strategy** for column player **Q** = $\langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ such that $M(\mathbf{P}, \mathbf{Q}) < 1/2$

Boosting as a matrix game (2)

- ▶ Boosting assumption: for any distribution over examples, there exists a weak rule with weighted error $< 1/2$
- ▶ In game terms: For any mixed strategy of the row player \mathbf{P} , there is a pure strategy for column player $\mathbf{Q} = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ such that $M(\mathbf{P}, \mathbf{Q}) < 1/2$
- ▶ From Min-Max theorem: There exists a column mixed strategy (a distribution over weak rules), that has expected value larger than zero for any row pure strategy (= any example).

Boosting as a matrix game (2)

- ▶ Boosting assumption: for any distribution over examples, there exists a weak rule with weighted error $< 1/2$
- ▶ In game terms: For any **mixed strategy** of the row player **P**, there is a **pure strategy** for column player **Q** = $\langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ such that $M(\mathbf{P}, \mathbf{Q}) < 1/2$
- ▶ From Min-Max theorem: There exists a column mixed strategy (a distribution over weak rules), that has expected value larger than zero for any row pure strategy (= any example).
- ▶ The weighted majority vote over the weak rule is **always** correct.

Adaboost as a repeated matrix game

- ▶ Booster chooses distribution over examples = mixed strategy over rows \mathbf{P}_t

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Adaboost as a repeated matrix game

- ▶ Booster chooses distribution over examples = mixed strategy over rows \mathbf{P}_t
- ▶ adversary chooses weak rule $\mathbf{Q}_t = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$
the 1 is at position t

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Adaboost as a repeated matrix game

- ▶ Booster chooses distribution over examples = mixed strategy over rows \mathbf{P}_t
- ▶ adversary chooses weak rule $\mathbf{Q}_t = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ the 1 is at position t
- ▶ **Goal 1:** produce a weighted majority rule that is highly accurate.

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Adaboost as a repeated matrix game

- ▶ Booster chooses distribution over examples = mixed strategy over rows \mathbf{P}_t
- ▶ adversary chooses weak rule $\mathbf{Q}_t = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle$ the 1 is at position t
- ▶ **Goal 1:** produce a weighted majority rule that is highly accurate.
- ▶ **Goal 2:** Find a “hard” distribution over the training examples.

	h_1	h_2	...
<i>example1</i>	0	1	...
<i>example2</i>	1	0	...
<i>example3</i>	0	0	...
...

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.

Minmax is weaker than diminishing regret

- ▶ The minmax theorem proves the existence of an **Equilibrium**.
- ▶ Learning guarantees no regret with respect to the past.
- ▶ If all sides use learning, then game will converge to minmax equilibrium.
- ▶ If opponent is not optimally adversarial (limited by knowledge, computational power...) then learning gives **better** performance than min-max.
- ▶ Our goal is to minimize regret.

Fictitious play

- ▶ also called “Follow the leader”

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.
- ▶ Might not converge to optimal mixed strategy.

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.
- ▶ Might not converge to optimal mixed strategy.
- ▶ Consider playing the matching coins game against an adversary that alternates **HTHTHTHT**

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.
- ▶ Might not converge to optimal mixed strategy.
- ▶ Consider playing the matching coins game against an adversary that alternates **HTHTHTHT**
- ▶ If $\#H > \#T$ the next element is T

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.
- ▶ Might not converge to optimal mixed strategy.
- ▶ Consider playing the matching coins game against an adversary that alternates **HTHTHTHT**
- ▶ If $\#H > \#T$ the next element is T
- ▶ If $\#T > \#H$ the next element is H

Fictitious play

- ▶ also called “Follow the leader”
- ▶ Choose the best action with respect to the sum of past loss vectors.
- ▶ Might not converge to optimal mixed strategy.
- ▶ Consider playing the matching coins game against an adversary that alternates **HTHTHTHT**
- ▶ If $\#H > \#T$ the next element is T
- ▶ If $\#T > \#H$ the next element is H
- ▶ follow the leader makes an error on each iteration.

Randomized Fictitious play

- ▶ Also called 'Follow the perturbed leader'

Randomized Fictitious play

- ▶ Also called 'Follow the perturbed leader'
- ▶ Choose the best action with respect to the sum of past loss vectors **plus noise**.

Randomized Fictitious play

- ▶ Also called 'Follow the perturbed leader'
- ▶ Choose the best action with respect to the sum of past loss vectors **plus noise**.
- ▶ Adding noise allows us to choose responses that are slightly worse than best response.

Randomized Fictitious play

- ▶ Also called 'Follow the perturbed leader'
- ▶ Choose the best action with respect to the sum of past loss vectors **plus noise**.
- ▶ Adding noise allows us to choose responses that are slightly worse than best response.
- ▶ **Hannan 1957** Randomized fictitious play converges to regret minimizing strategy.

Randomized Fictitious play

- ▶ Also called 'Follow the perturbed leader'
- ▶ Choose the best action with respect to the sum of past loss vectors **plus noise**.
- ▶ Adding noise allows us to choose responses that are slightly worse than best response.
- ▶ **Hannan 1957** Randomized fictitious play converges to regret minimizing strategy.
- ▶ regret is $O(1/\sqrt{n})$ where n is number of actions.

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$

The basic algorithm

- ▶ Choose an initial distribution \mathbf{P}_1



$$\mathbf{P}_{t+1}(i) = \mathbf{P}_t(i) \frac{e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}}{Z_t}$$

- ▶ Where $Z_t = \sum_{i=1}^n \mathbf{P}_t(i) e^{-\eta \mathbf{M}(i, \mathbf{Q}_t)}$
- ▶ $\eta > 0$ is the learning rate.

Generalized regret bound

- Regret relative to the best *pure strategy* i

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_i \left[\eta \sum_{t=1}^T \mathbf{M}(i, \mathbf{Q}_t) - \ln \mathbf{P}_1(i) \right]$$

Generalized regret bound

- ▶ Regret relative to the best *pure strategy* i

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_i \left[\eta \sum_{t=1}^T \mathbf{M}(i, \mathbf{Q}_t) - \ln \mathbf{P}_1(i) \right]$$

- ▶ regret with respect the the best *mixed strategy* \mathbf{P} :

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_{\mathbf{P}} \left[\eta \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \text{RE}(\mathbf{P} \parallel \mathbf{P}_1) \right]$$

Generalized regret bound

- ▶ Regret relative to the best *pure strategy* i

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_i \left[\eta \sum_{t=1}^T \mathbf{M}(i, \mathbf{Q}_t) - \ln \mathbf{P}_1(i) \right]$$

- ▶ regret with respect the the best *mixed strategy* \mathbf{P} :

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_{\mathbf{P}} \left[\eta \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \text{RE}(\mathbf{P} \parallel \mathbf{P}_1) \right]$$

- ▶ Where

$$\text{RE}(\mathbf{P} \parallel \mathbf{Q}) \doteq \sum_{i=1}^n \mathbf{P}(i) \ln \frac{\mathbf{P}(i)}{\mathbf{Q}(i)}$$

Main Theorem

- ▶ For **any** game matrix **M**.

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. $\mathbf{Q}_1, \dots, \mathbf{Q}_T$

Main Theorem

- ▶ For **any** game matrix **M**.
- ▶ Any sequence of mixed strat. **$\mathbf{Q}_1, \dots, \mathbf{Q}_T$**
- ▶ The sequence **$\mathbf{P}_1, \dots, \mathbf{P}_T$** produced by basic alg using **$\eta > 0$** satisfies

$$\sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \left(\frac{1}{1 - e^{-\eta}} \right) \min_{\mathbf{P}} \left[\eta \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \text{RE}(\mathbf{P} \parallel \mathbf{P}_1) \right]$$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is

$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

Corollary

- ▶ Setting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ the average per-trial loss is


$$\frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{M}(\mathbf{P}, \mathbf{Q}_t) + \Delta_{T,n}$$

- ▶ Where

$$\Delta_{T,n} = \sqrt{\frac{2 \ln n}{T}} + \frac{\ln n}{T} = O\left(\sqrt{\frac{\ln n}{T}}\right).$$


Visual intuition

- **Hedge(η)** : If $M(\mathbf{P}_t, \mathbf{Q}_t) \gg M(\tilde{\mathbf{P}}, \mathbf{Q}_t)$ then:
distance between \mathbf{P}_{t+1} and $\tilde{\mathbf{P}}$ smaller than
distance between \mathbf{P}_t and $\tilde{\mathbf{P}}$



Visual intuition

- ▶ **Hedge(η)** : If $M(\mathbf{P}_t, \mathbf{Q}_t) \gg M(\tilde{\mathbf{P}}, \mathbf{Q}_t)$ then:
distance between \mathbf{P}_{t+1} and $\tilde{\mathbf{P}}$ smaller than
distance between \mathbf{P}_t and $\tilde{\mathbf{P}}$
- ▶ $RE(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) - RE(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) \leq$
 $\eta M(\tilde{\mathbf{P}}, \mathbf{Q}_t) - (1 - e^{-\eta})M(\mathbf{P}_t, \mathbf{Q}_t)$



The minmax Theorem

John von Neumann, 1928.

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

The minmax Theorem

John von Neumann, 1928.

$$\min_P \max_Q \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_Q \min_P \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

In words: for **mixed** strategies, choosing second gives no advantage.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

Let $\bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t$ and $\bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} \leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q}$$

$$= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}}$$

Proving minmax Theorem using online learning (1)

Row player chooses \mathbf{P}_t using learning alg.

Column player chooses \mathbf{Q}_t after row player so that

$$\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$$

$$\text{Let } \bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t \text{ and } \bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$$

$$\begin{aligned} \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{P}^T \mathbf{M} \mathbf{Q} &\leq \max_{\mathbf{Q}} \bar{\mathbf{P}}^T \mathbf{M} \mathbf{Q} \\ &= \max_{\mathbf{Q}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \quad \text{by definition of } \bar{\mathbf{P}} \\ &\leq \frac{1}{T} \sum_{t=1}^T \max_{\mathbf{Q}} \mathbf{P}_t^T \mathbf{M} \mathbf{Q} \end{aligned}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t$$

by definition of \mathbf{Q}_t

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \bar{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \bar{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

Proving minmax Theorem using online learning (2)

$$= \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t^T \mathbf{M} \mathbf{Q}_t \quad \text{by definition of } \mathbf{Q}_t$$

$$\leq \min_{\mathbf{P}} \frac{1}{T} \sum_{t=1}^T \mathbf{P}^T \mathbf{M} \mathbf{Q}_t + \Delta_{T,n} \quad \text{by the Corollary}$$

$$= \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \overline{\mathbf{Q}} + \Delta_{T,n} \quad \text{by definition of } \overline{\mathbf{Q}}$$

$$\leq \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{P}^T \mathbf{M} \mathbf{Q} + \Delta_{T,n}.$$

but $\Delta_{T,n}$ can be set arbitrarily small.

Solving a game

- ▶ to **solve** a game is to find the min-max mixed strategies **P, Q**

Solving a game

- ▶ to **solve** a game is to find the min-max mixed strategies \mathbf{P}, \mathbf{Q}
- ▶ Suppose that **Hedge**(η) is playing $\mathbf{P}_1, \mathbf{P}_2$, against a worst case adversary that plays second: adversary that plays $\mathbf{Q}_1, \mathbf{Q}_2, \dots$ such that $\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$.

Solving a game

- ▶ to **solve** a game is to find the min-max mixed strategies \mathbf{P}, \mathbf{Q}
- ▶ Suppose that **Hedge**(η) is playing $\mathbf{P}_1, \mathbf{P}_2$, against a worst case adversary that plays second: adversary that plays $\mathbf{Q}_1, \mathbf{Q}_2, \dots$ such that $\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$.
- ▶ Without loss of generality \mathbf{Q}_t is a pure strategy (prob. 1 on a single action).

Solving a game

- ▶ to **solve** a game is to find the min-max mixed strategies \mathbf{P}, \mathbf{Q}
- ▶ Suppose that **Hedge**(η) is playing $\mathbf{P}_1, \mathbf{P}_2$, against a worst case adversary that plays second: adversary that plays $\mathbf{Q}_1, \mathbf{Q}_2, \dots$ such that $\mathbf{Q}_t = \arg \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}_t, \mathbf{Q})$.
- ▶ Without loss of generality \mathbf{Q}_t is a pure strategy (prob. 1 on a single action).
- ▶ Let $\bar{\mathbf{P}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{P}_t$, $\bar{\mathbf{Q}} \doteq \frac{1}{T} \sum_{t=1}^T \mathbf{Q}_t$

Using average distributions

- ▶ Von Neumann Min/Max Thm:

$$v \doteq \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

Using average distributions

- ▶ Von Neumann Min/Max Thm:

$$v \doteq \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

- ▶ Fixing T and letting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$

Using average distributions

- ▶ Von Neumann Min/Max Thm:

$$v \doteq \min_{\mathbf{P}} \max_{\mathbf{Q}} \mathbf{M}(\mathbf{P}, \mathbf{Q}) = \max_{\mathbf{Q}} \min_{\mathbf{P}} \mathbf{M}(\mathbf{P}, \mathbf{Q})$$

- ▶ Fixing T and letting $\eta = \ln \left(1 + \sqrt{\frac{2 \ln n}{T}} \right)$
- ▶ Two immediate corollaries of the proof of the min/max Thm:

$$\max_{\mathbf{Q}} \mathbf{M}(\bar{\mathbf{P}}, \mathbf{Q}) \leq v + \Delta_{T,n} \cdot \min_{\mathbf{P}} \mathbf{M}(\mathbf{P}, \bar{\mathbf{Q}}) \geq v - \Delta_{T,n}$$

- └ Approximately solving games
- └ Variable learning rate

Using the final row distribution v^{MW}

- Can we make the row distribution converge?

Using the final row distribution vMW

- ▶ Can we make the row distribution converge?
- ▶ Suppose we have an upper bound on the value of the game $u \geq v$

Using the final row distribution v MW

- ▶ Can we make the row distribution converge?
- ▶ Suppose we have an upper bound on the value of the game $u \geq v$
- ▶ **Good Enough:** If $\mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq u$ the row player does nothing $\mathbf{P}_{t+1} = \mathbf{P}_t$

Using the final row distribution v MW

- ▶ Can we make the row distribution converge?
- ▶ Suppose we have an upper bound on the value of the game $u \geq v$
- ▶ **Good Enough:** If $\mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \leq u$ the row player does nothing $\mathbf{P}_{t+1} = \mathbf{P}_t$
- ▶ **Learn:** If $\mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) > u$ set

$$\eta = \ln \frac{(1 - u)\mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)}{u(1 - \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t))} .$$

- └ Approximately solving games
- └ Variable learning rate

Bound for v MW

- ▶ Let $\tilde{\mathbf{P}}$ be any mixed strategy for the rows such that $\max_{\mathbf{Q}} \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}) \leq u$

Bound for vMW

- ▶ Let $\tilde{\mathbf{P}}$ be any mixed strategy for the rows such that $\max_{\mathbf{Q}} \mathbf{M}(\tilde{\mathbf{P}}, \mathbf{Q}) \leq u$
- ▶ Then on any iteration of algorithm vMW in which $\mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t) \geq u$ the relative entropy between $\tilde{\mathbf{P}}$ and \mathbf{P}_{t+1} satisfies

$$\text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_{t+1}) \leq \text{RE}(\tilde{\mathbf{P}} \parallel \mathbf{P}_t) - \text{RE}(u \parallel \mathbf{M}(\mathbf{P}_t, \mathbf{Q}_t)) .$$