

# Vovk's aggregating algorithm

## Mixable and unmixable loss functions

Yoav Freund

January 28, 2020

Section 3.5 in “Prediction, Learning and Games”

# Outline

Log Loss and Absolute loss

The general prediction game

Analysis for specific loss functions

Vovk's algorithm

mixable loss functions

The convexity condition

Log loss

Square loss

    Square loss using simple averaging

Summary table

## Binary log-loss

- ▶ Prediction:  $p \in [0, 1]$  outcome  $x \in \{0, 1\}$
- ▶ Loss:  $\lambda(p, x) = -x \log p - (1 - x) \log(1 - p)$
- ▶  $N$  experts, expert  $i$  at time  $t$  outputs  $q_i^t \in [0, 1]$
- ▶ Cumulative loss of expert  $i$  at time  $t$ :  $L_i^t = \sum_{s=1}^t \lambda(q_i^s, x^s)$
- ▶ Experts algorithm (Bayes Algorithm):
  - ▶ Assign weights:  $w_i^t = \frac{1}{N} \exp(-L_i^{t-1})$
  - ▶ Master prediction:  $q_M^t = \frac{\sum_{i=1}^N q_i^t w_i^t}{\sum_{i=1}^N w_i^t}$
- ▶ Regret Bound:

$$L_A^T \leq \min_i L_i^T + \ln N$$

# Absolute loss

- ▶ Prediction:  $p \in [0, 1]$  outcome  $x \in \{0, 1\}$
- ▶ Loss:  $\lambda(p, x) = |x - p|$
- ▶  $N$  experts, expert  $i$  at time  $t$  outputs  $q_i^t \in [0, 1]$
- ▶ Cumulative loss of expert  $i$  at time  $t$ :  $L_i^t = \sum_{s=1}^t \lambda(q_i^s, x^s)$
- ▶ Experts algorithm (Hedge):
  - ▶ Assign weights:  $w_i^t = \frac{1}{N} \exp(-\eta L_i^{t-1})$
  - ▶ Master prediction:  $q_M^t = \frac{\sum_{i=1}^N q_i^t w_i^t}{\sum_{i=1}^N w_i^t}$
- ▶ Regret Bound for known horizon.
  - ▶ Set  $\eta$  according to  $T$ :  $\eta \approx \sqrt{\frac{2 \ln N}{T}}$
  - ▶ Regret bound:

$$L_A \leq \min_i L_i + \sqrt{2T \ln N} + \ln N$$

## Two other loss functions over $[0, 1]$

- Prediction:  $p \in [0, 1]$  outcome  $x \in [0, 1]$  (note  $x$  is not restricted to  $\{0, 1\}$ )
- Square loss (Breier Loss):  $\lambda(p, x) = (p - x)^2$
- Hellinger Loss

$$\lambda(p, x) = \frac{1}{2} \left( (\sqrt{p} + \sqrt{x})^2 + (\sqrt{1-p} + \sqrt{1-x})^2 \right)$$

## Vovk's general prediction game

$\Gamma$  - prediction space.  $\Omega$  - outcome space.

On each trial  $t = 1, 2, \dots$

1. Each expert  $i \in \{1 \dots N\}$  makes a prediction  $\gamma_i^t \in \Gamma$
2. The learner, after observing  $\langle \gamma_1^t \dots \gamma_N^t \rangle$ , makes its own prediction  $\gamma^t$
3. Nature chooses an outcome  $\omega^t \in \Omega$
4. Each expert incurs loss  $\ell_t(i) = \lambda(\omega^t, \gamma_i^t)$   
The learner incurs loss  $\ell_t(\mathbf{A}) = \lambda(\omega^t, \gamma^t)$

## Achievable loss bounds

- ▶  $L_A \doteq \sum_{t=1}^T \ell_t(A)$  - total loss of algorithm
- ▶  $L_i \doteq \sum_{t=1}^T \ell_t(i)$  - total loss of expert  $i$
- ▶ **Goal:** find an algorithm which guarantees that

$$(a, c) \in [0, \infty), \quad L_A \leq aL_{\min} + c \ln N$$


For any sequence of events.

- ▶ We say that the pair  $(a, c)$  is **achievable**.

## The set of achievable bounds

- ▶ Fix loss function  $\lambda : \Omega \times \Gamma \rightarrow [0, \infty)$
- ▶ The pair  $(a, c)$  is *achievable* if there exists *some* prediction algorithm such that for *any*  $N > 0$ , *any* set of  $N$  prediction sequences and *any* sequence of outcomes

$$L_A \leq aL_{\min} + c \ln N$$





## Analysis for specific loss functions

- ▶ Outcomes:  $\omega^1, \omega^2, \dots, \omega^t \in [0, 1]$
- ▶ Predictions:  $\gamma^1, \gamma^2, \dots, \gamma^t \in [0, 1]$

## Log loss (Entropy loss)



$$\lambda_{\text{ent}}(\omega, \gamma) = \omega \ln \frac{\omega}{\gamma} + (1 - \omega) \ln \frac{1 - \omega}{1 - \gamma}$$

- ▶ When  $q_t \in \{0, 1\}$  Cumulative log loss = coding length  $\pm 1$
- ▶ If  $P[\omega_t = 1] = q$ , optimal prediction  $\gamma^t = q$
- ▶ Unbounded loss.
- ▶ Not symmetric  $\exists p, q \lambda(p, q) \neq \lambda(q, p)$ .
- ▶ No triangle inequality  
 $\exists p_1, p_2, p_3 \lambda(p_1, p_3) > \lambda(p_1, p_2) + \lambda(p_2, p_3)$

## Square loss (Breier Loss)



$$\lambda_{\text{sq}}(\omega, \gamma) = (\omega - \gamma)^2$$

- ▶  $P[\omega^t = 1] = q, P[\omega^t = 0] = 1 - q,$   
optimal prediction  $\gamma^t = q$
- ▶ Bounded loss.
- ▶ Defines a metric (symmetric and triangle ineq.)
- ▶ Corresponds to regression.

## Hellinger Loss



$$\lambda_{\text{hel}}(\omega, \gamma) = \frac{1}{2} \left( (\sqrt{\omega} + \sqrt{\gamma})^2 + (\sqrt{1-\omega} + \sqrt{1-\gamma})^2 \right)$$

- ▶ If  $P[\omega^t = 1] = q$ ,  $P[\omega^t = 0] = 1 - q$ ,  
optimal prediction  $\gamma^t = q$
- ▶ Loss is bounded.
- ▶ Defines a metric.
- ▶  $\lambda_{\text{hel}}(p, q) \approx \lambda_{\text{ent}}(p, q)$  when  $p \approx q$  and  $p, q \in (0, 1)$

## Absolute loss



$$\lambda(\omega, \gamma) = |\omega - \gamma|$$

- ▶ Probability of making a mistake if predicting 0 or 1 using a biased coin
- ▶ If  $P[\omega^t = 1] = q$ ,  $P[\omega^t = 0] = 1 - q$ , then the optimal prediction is

$$\gamma^t = \begin{cases} 1 & \text{if } q > 1/2, \\ 0 & \text{otherwise} \end{cases}$$

## Structureless bounded loss

- ▶ Prediction is a distribution  $\gamma = \langle p_1, \dots, p_N \rangle$ ,  $p_i \geq 0$ ,  
 $\sum_{i=1}^N p_i = 1$
- ▶ Outcome is a loss vector  $\omega = \langle \omega_1, \dots, \omega_N \rangle$ ,  $0 \leq \omega_i \leq 1$
- ▶ Loss is the dot product:  $\lambda_{\text{dot}}(\omega, \gamma) = \gamma \cdot \omega$
- ▶ Corresponds to the hedging game.
- ▶ For hedge loss the regret is  $\Omega(\sqrt{T \log N})$ .
- ▶ For the log loss the regret is  $O(\log N)$
- ▶ Which losses behave like **entropy loss** and which behave like **hedge loss**?

## Some technical requirements

- ▶ There should be a **topology** on the prediction set  $\Gamma$  such that
- ▶  $\Gamma$  is compact.
- ▶  $\forall \omega \in \Omega$ , the function  $\gamma \rightarrow \lambda(\omega, \gamma)$  is **continuous**
- ▶ There is a **universally reasonable prediction**  
 $\exists \gamma \in \Gamma, \forall \omega \in \Omega, \lambda(\omega, \gamma) < \infty$
- ▶ There is **no universally optimal prediction**  
 $\neg \exists \gamma \in \Gamma, \forall \omega \in \Omega, \lambda(\omega, \gamma) = 0$

## Vovk's meta-algorithm

- Fix an **achievable** pair  $(a, c)$  and set  $\eta = a/c$
- 1.

$$w_i^t = \frac{1}{N} e^{-\eta L_i^t}$$

- 2. Choose  $\gamma_t$  so that, for all  $\omega^t \in \Omega$ :

$$\lambda(\omega^t, \gamma^t) - c \ln \sum_i w_i^t \leq -c \ln \left( \sum_i w_i^t e^{-\eta \lambda(\omega^t, \gamma_i^t)} \right)$$

- If choice of  $\gamma_t$  always exists, then the total loss satisfies:

$$\sum_t \lambda(\omega^t, \gamma^t) \leq -c \ln \sum_i w_i^{T+1} \leq a L_{\min} + c \ln N$$


- Vovk's result: **yes!** a good choice for  $\gamma_t$  always exists!



# Vovk's algorithm is the the highest achiever [Vovk95]

The pair  $(a, c)$  is achieved by some algorithm if and only if it is achieved by Vovk's algorithm.

The separation curve is  $\left\{ \left( a(\eta), \frac{a(\eta)}{\eta} \right) \mid \eta \in [0, \infty] \right\}$



## Mixable Loss Functions

- ▶ A Loss function is **mixable** if a pair of the form  $(1, c)$ ,  $c < \infty$  is achievable.

$$L_A \leq L_{\min} + c \ln N$$

- ▶ Vovk's algorithm with  $\eta = 1/c$  achieves this bound.
- ▶  $\lambda_{\text{ent}}, \lambda_{\text{sq}}, \lambda_{\text{hel}}$  are **mixable**
- ▶  $\lambda_{\text{abs}}, \lambda_{\text{dot}}$  are **not mixable**

## The convexity condition

- ▶ requirement for loss to be  $(1, 1/\eta)$  mixable
- ▶  $\forall \langle (\gamma_1, W_1), \dots, (\gamma_N, W_N) \rangle$   
 $\exists \gamma \in \Gamma$   
 $\forall \omega \in \Omega$ :

$$\lambda(\omega, \gamma) - \frac{1}{\eta} \ln \sum_i W_i \leq -\frac{1}{\eta} \ln \left( \sum_i W_i e^{-\eta \lambda(\omega, \gamma_i)} \right)$$

- ▶ Can be re-written as:


$$e^{-\eta \lambda(\omega, \gamma)} \geq \sum_i \left( \frac{W_i}{\sum_j W_j} \right) e^{-\eta \lambda(\omega, \gamma_i)}$$

- ▶ Equivalently - the image of the set  $\Gamma$  under the mapping  $F(\gamma) = \langle e^{-\eta \lambda(\omega, \gamma)} \rangle_{\omega \in \Omega}$  is concave.

## convexity condition: Pictorially

- **Example:** Suppose  $\Omega = \{0, 1\}$ ,  $\Gamma = [0, 1]$ . then

$$F(\gamma) = \left\langle e^{-\eta\lambda(0,\gamma)}, e^{-\eta\lambda(1,\gamma)} \right\rangle$$



## Vovk Algorithm for log loss

- ▶ The log loss is mixable with  $\eta = 1$
- ▶ The image of  $[0, 1]$  through  $F(\gamma) = \langle e^{-\eta\lambda(0,\gamma)}, e^{-\eta\lambda(1,\gamma)} \rangle$  is a straight line segment.
- ▶ The **only** satisfactory prediction is

$$\gamma = \frac{\sum_i w_i \gamma_i}{\sum_i w_i}$$

- ▶ We are back to the online Bayes algorithm.

## Vovk algorithm for square loss

- ▶ The square loss is mixable with  $\eta = 2$ .
- ▶ Prediction must satisfy

$$1 - \sqrt{-\frac{1}{2} \ln \sum_i V_i^t e^{-2(1-p_i^t)^2}} \leq p^t \leq \sqrt{-\frac{1}{2} \ln \sum_i V_i^t e^{-2(p_i^t)^2}}$$

where  $V_i^t = \frac{W_i^t}{\sum_s W_i^s}$ .

▶

$$L_A \leq L_{\min} + \frac{1}{2} \ln N$$

## Simple prediction for square loss

- ▶ We can use the prediction

$$\gamma = \frac{\sum_i W_i \gamma_i}{\sum_i W_i}$$

- ▶ But in that case we must use  $\eta = 1/2$  when updating the weights.
- ▶ Which yields the bound

$$L_A \leq L_{\min} + 2 \ln N$$

## Summary of bounds for mixable losses

| Loss Functions:        | $c$ values: ( $\eta = 1/c$ )       |                                   |
|------------------------|------------------------------------|-----------------------------------|
|                        | $\text{pred}_{\text{wmean}}(v, x)$ | $\text{pred}_{\text{Vovk}}(v, x)$ |
| $L_{\text{sq}}(p, q)$  | 2                                  | $1/2$                             |
| $L_{\text{ent}}(p, q)$ | 1                                  | 1                                 |
| $L_{\text{hel}}(p, q)$ | 1                                  | $1/\sqrt{2}$                      |

Figure 2.  $(c, 1/c)$ -realizability:  $c$  values for loss and prediction