

Synthetic calculation · wb-loading

Quality: not assessed · Template: valid

计算快照 / Calculation snapshot

wb-loading · default · wb-loading-v3 · valid

Operations: liquid-operations-v2 · Pipetting: below-minimum

WB-A · 样品量 (μL) / Sample (μL): 10 μL
样品量 (μL) / Sample (μL) → WB-A
WB-A · 缓冲液 (μL) / Buffer (μL): 5 μL
缓冲液 (μL) / Buffer (μL) → WB-A
WB-A · Synthetic stock: 1 μL
Synthetic stock → WB-A
WB-A · 稀释液 (μL) / Diluent (μL): 4 μL
稀释液 (μL) / Diluent (μL) → WB-A

有效样本 / Valid samples1

配液明细 / Preparation row 1	
样本ID	WB-A
状态	有效 / Valid
总量 (μL)	20
缓冲液 (μL)	5
样品量 (μL)	10
稀释液 (μL)	4
原始体积单位	μL
可用体积 (μL)	
添加方式	volume-fraction
还原剂原液	Synthetic stock
还原剂 (μL)	1
目标蛋白量 (μg)	20
输入浓度单位	μg/μL
原液与目标定义	最终体积的 5% v/v 原液；不代表有效成分终浓度 / Fraction of final volume, not active concentration
原浓度 (μg/μL)	2

WB-A · Synthetic stock (theoretical): 1 μL，低于所设 2 μL 下限 / below configured minimum. 调整制备规模，或评估中间液方案 / Adjust preparation scale or assess an intermediate dilution.

最终体积的 5% v/v 原液；不代表有效成分终浓度 / Fraction of final volume, not active concentration

Inputs / Units: {
 "samples": [
 {
 "id": "WB-A",
 "available": "",
 "concentration": "2" }
]
}

```
    }  
  ],  
  "__context": {  
    "experimentId": "calculator-acceptance-experiment",  
    "experimentStepId": "calculator-acceptance-step"  
  },  
  "bufferFold": 4,  
  "reducingMode": "volume-fraction",  
  "__displayUnits": {  
    "table:sampleUL": "µL"  
  },  
  "targetProteinUg": 20,  
  "pipetteMinimumUL": 2,  
  "reducingAgentName": "Synthetic stock",  
  "finalLoadingVolumeUL": 20,  
  "reducingAgentPercent": 5,  
  "bufferContainsReducingAgent": "no"  
}
```