

CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL

MASTER THESIS

Thesis Title

Author:
Anne Fischer

Supervisor:
Prof. Dr.
Wimmer-Schweingruber

Research Group Name
Department or School Name

October 13, 2019

Abstract

The Thesis Abstract is written here (and usually kept to just this page). The page is kept centered vertically so can expand into the blank space above the title too...

Contents

Abstract	iii
1 Motivation	1
2 Pickup Ions	3
2.1 The Heliosphere	3
2.2 Pickup Ions	3
2.3 Interstellar Pickup Ions	4
2.4 Inner-source Pickup Ions	4
2.5 VDF	5
3 Instrumentation	7
3.1 ULYSSES	7
3.2 SWICS	7
A Frequently Asked Questions	9
A.1 How do I change the colors of links?	9
Bibliography	11
Declaration of Authorship	13
Acknowledgements	15

Chapter 1

Motivation

Chapter 2

Pickup Ions

Pickup ions are created when neutral atoms inside the heliosphere become ionised and are subsequently swept away with the heliospheric magnetic field that is embedded within the solar wind.

2.1 The Heliosphere

Oder Überkapitel Solar Physics?

Heliosphere: Grenze zu LISM

Solar Wind: Zusammensetzung, schneller und langsamer

B-Feldgleichung

2.2 Pickup Ions

A neutral atom inside the heliosphere is only subjected to the gravitational force and radiation pressure of the sun. It is not sensitive to any electromagnetic forces until it becomes ionised by solar ultra-violet radiation, charge exchange with solar wind protons or electron impact (Q?). After ionisation the particle starts interacting with the solar wind plasma. In particular it is forced onto a gyro orbit about the heliospheric magnetic field that is embedded within the solar wind. As the freshly created ion is swept away with the magnetic field line it is “picked up” from its location of ionisation – a new pickup ion (PUI) has been created.

PUIs were first observed by Möbius et al. (1985) with the SULEICA Instrument on the AMPTE spacecraft. The particles measured at 1 AU were He⁺ ions of interstellar origin.

Once the particle is ionised, its probability to become ionised another time decreases (Quelle). This characteristic of being only singly charged can help to discriminate PUIs from solar wind ions, that are mostly more often charged (Q?).

PUIs are mostly only single charged. This characteristic can help to distinguish them from solar wind ions of coronal origin which often have been ionized multiple times, if not completely. (Q?)

VDF non-maxwellian, spatial density pattern

There have been observed several species of PUIs:

2.3 Interstellar Pickup Ions

Heliosheath, relative motion

The neutral part of the LISM can enter the heliosphere as it is not affected by the heliosheath (Todo). Inside the heliosphere the neutrals are guided only by the gravitational force and radiation pressure of the sun. The neutral particle's species determines how deep it can travel into the heliosphere before it becomes ionized. Species with a higher First Ionization Potential will be able to approach the sun much closer without being ionized. This results in He⁺ being the dominant PUI species at a solar distance of 1 AU even if in the LISM the abundance of hydrogen is about 10 times the one of helium.

- ionisation process is also dependent on the species
- radiation pressure only important for H (and He?). Kepler orbit...
- Spatial distribution:
gravitational force and radiation pressure lead to two regions of enhanced density of neutrals (in the ecliptic): Focusing cone and crescent. Focusing cone: For species with high FIP (as the others are ionized before and do not reach the downwind side of the sun)
- variation of He⁺ with the solar cycle: Rucinski 2003
- H, O and N are depleted in the filtration region (Baranov Malama 1995), Wimmer Skript: even before ionization: density is determined by ratio of gravitational force and photon pressure
- neutral density determines PUI production rate

2.4 Inner-source Pickup Ions

The idea of an additional source for the PUI's neutral seed population was born when Geiss, Gloeckler, Fisk, et al. (1995) measured a global distribution of C⁺ PUIs with the SWICS instrument on Ulysses. Interstellar carbon exists almost exclusively in a single charged state (Frisch et al., 2011) in the LISM. As only neutral atoms can enter the heliosphere it was not expected to find a distinct signature of C⁺ pickup ions. However, pickup carbon was observed with about the same ratio as oxygen, of which, in contrast, 80% is in a neutral charge state in the interstellar medium. These findings suggested that there must be another source for neutrals that has its origin somewhere inside the heliosphere.

In following studies (e.g. Geiss, Gloeckler, and Steiger, 1995) there were found also other species like O⁺ and Ne⁺ of these, so called, inner-source PUIs.

Inner-source PUIs show a composition that is similar to the one of the solar wind (Gloeckler et al., 2000; Allegrini et al., 2005) as well as a velocity distribution function that is centered around $w_{SC} \approx 1$ (Schwadron et al., 2000) and seems to have thermalized with the solar wind.

Beneath those two characteristics there are other aspects concerning inner-source PUIs, that are still under debate. In particular that is the production mechanism of their neutral seed population. Allegrini et al. (2005) has summarized current candidates for possible scenarios. Two of those give an explanation for the ion's composition as they directly incorporate solar wind ions in the process:

- Solar wind recycling (Gloeckler et al., 2000; Schwadron et al., 2000): Absorption of solar wind ions by heliospheric grains and subsequent reemission of neutral atoms
- Solar wind neutralization (Wimmer-Schweingruber and Bochsler, 2002): Solar wind ions penetrate sub-micron-sized dust grains and undergo (partial) neutralization by charge exchange

2.5 VDF

After the particle has been ionised it is forced onto a gyro motion about the local field line of the heliospheric magnetic field due to the Lorentz force.

No velocity compared to v_{sw}

B field is convected by the SW and moves radially outwards from the sun

Form of the torus depends on several factors: velocity of the particle before the PU process, orientation of the magnetic field

Chapter 3

Instrumentation

3.1 ULYSSES

sfsf

3.2 SWICS

sffsf

Appendix A

Frequently Asked Questions

A.1 How do I change the colors of links?

The color of links can be changed to your liking using:

```
\hypersetup{urlcolor=red}, or  
\hypersetup{citecolor=green}, or  
\hypersetup{allcolor=blue}.
```

If you want to completely hide the links, you can use:

```
\hypersetup{allcolors=.}, or even better:  
\hypersetup{hidelinks}.
```

If you want to have obvious links in the PDF but not the printed text, use:

```
\hypersetup{colorlinks=false}.
```


Bibliography

- Allegrini, F., N. A. Schwadron, D. J. McComas, G. Gloeckler, and J. Geiss (May 2005). "Stability of the inner source pickup ions over the solar cycle". In: *Journal of Geophysical Research* 110.A5, A05105, A05105. DOI: [10.1029/2004JA010847](https://doi.org/10.1029/2004JA010847).
- Frisch, Priscilla C., Seth Redfield, and Jonathan D. Slavin (Sept. 2011). "The Interstellar Medium Surrounding the Sun". In: *Annual Review of Astronomy and Astrophysics* 49.1, pp. 237–279. DOI: [10.1146/annurev-astro-081710-102613](https://doi.org/10.1146/annurev-astro-081710-102613).
- Geiss, J., G. Gloeckler, L. A. Fisk, and R. von Steiger (1995). "C+ Pickup ions in the heliosphere and their origin". In: *Journal of Geophysical Research: Space Physics* 100.A12, pp. 23373–23377. DOI: [10.1029/95JA03051](https://doi.org/10.1029/95JA03051). eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/95JA03051>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/95JA03051>.
- Geiss, J., G. Gloeckler, and R. von Steiger (Apr. 1995). "Origin of the solar wind from composition data". In: *Space Science Reviews* 72.1, pp. 49–60. ISSN: 1572-9672. DOI: [10.1007/BF00768753](https://doi.org/10.1007/BF00768753). URL: <https://doi.org/10.1007/BF00768753>.
- Gloeckler, G., L. A. Fisk, J. Geiss, N. A. Schwadron, and T. H. Zurbuchen (Apr. 2000). "Elemental composition of the inner source pickup ions". In: *Journal of Geophysical Research* 105.A4, pp. 7459–7464. DOI: [10.1029/1999JA000224](https://doi.org/10.1029/1999JA000224).
- Möbius, E., D. Hovestadt, B. Klecker, G. Scholer, and G. Gloeckler (1985). "Direct observation of He+ pick-up ions of interstellar origin in the solar wind". In: *nature* 318 (6045), pp. 426–429. DOI: [10.1038/318426A0](https://doi.org/10.1038/318426A0). URL: <https://www.nature.com/articles/318426a0>.
- Schwadron, N. A., J. Geiss, L. A. Fisk, G. Gloeckler, T. H. Zurbuchen, and R. von Steiger (Apr. 2000). "Inner source distributions: Theoretical interpretation, implications, and evidence for inner source protons". In: *Journal of Geophysical Research* 105.A4, pp. 7465–7472. DOI: [10.1029/1999JA000225](https://doi.org/10.1029/1999JA000225).
- Wimmer-Schweingruber, R. F. and P. Bochsler (May 2002). "On the Origin of Inner-Source Pickup Ions". In: *Geophysical Research Letters* 30.2, pp. 49-1 - 49-4.

Declaration of Authorship

I, Anne Fischer, declare that this thesis titled, and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed:

Date:

Acknowledgements

The acknowledgments and the people to thank go here, don't forget to include your project advisor...