

Completeness in Generalized Ultrametric Spaces

Nathanael Leedom Ackerman
presented by Anne Johnson

August 29, 2018

"The goal of this paper is to characterize the relationship between four important notions of completeness for Γ -ultrametric spaces: spherical completeness, Cauchy completeness, strong Cauchy completeness and injectivity"

Instead, we'll talk about:

- ▶ equivalence of categories between the category of flabby, separated presheaves on Γ^{op} and the category of Γ -ultrametric spaces and non-expanding maps.
- ▶ Injective hulls

Some background

Definition

A partial order (Γ, \leq_Γ) is a **complete lattice** if every subset of Γ has a supremum and infimum. A complete lattice (Γ, \leq_Γ) is an **op-frame** if it satisfies

$$(\forall x \in \Gamma, A \subseteq \Gamma) x \vee_\Gamma (\bigwedge_{\alpha \in A} \alpha) = \bigwedge_{\alpha \in A} (x \vee_\Gamma \alpha)$$

Some background

Notation:

- ▶ (Γ, \leq_Γ) or Γ – complete lattice, always an op-frame
- ▶ \perp_Γ – least element, ie $\bigwedge^\Gamma \{\gamma : \gamma \in \Gamma\}$
- ▶ \top_Γ – greatest element, ie $\bigvee^\Gamma \{\gamma : \gamma \in \Gamma\}$
- ▶ Γ^{op} – Γ but with the order reversed

Some background

Definition

A Γ -**ultrametric space** is a pair (M, d_M) , where M is a set and $d_M : M \times M \rightarrow \Gamma$ is a function that satisfies:

- ▶ $(\forall x, y \in M) \ d_M(x, y) = \perp \iff x = y$
- ▶ $(\forall x, y \in M) \ d_M(x, y) = d_M(y, x)$
- ▶ $(\forall x, y, z \in M) \ d_M(x, y) \vee d_M(y, z) \geq d_M(x, z)$

Definition

A **non-expanding map** between Γ -ultrametric spaces (M, d_M) and (N, d_N) is a function $f : M \rightarrow N$ such that $(\forall a, b \in M)$
 $d_N(f(a), f(b)) \leq d_M(a, b)$

Some background

A few things to note

- ▶ Γ –ultrametric spaces and non-expanding maps form a category, denoted $UMet(\Gamma)$
- ▶ Given a Γ –ultrametric space (M, d_M) , the balls of fixed radius γ for $\gamma \in \Gamma$ also form a generalized ultrametric space, with the distance function taking values in $\Gamma_\gamma = \{\zeta \in \Gamma : \zeta \geq_\Gamma \gamma\}$

Some background: closed ball functor

Denote by (M_γ, d_{M_γ}) the generalized ultrametric space having points

$$M_\gamma = \{B^M(x; \gamma) : x \in M\}$$

and distance function

$$d_{M_\gamma}(B^M(x; \gamma), B^M(y; \gamma)) = d_M(x, y) \vee \gamma$$

Lastly if $f : (M, d_M) \rightarrow (N, d_N)$ is a map of Γ -ultrametric spaces, let $f_\gamma : (M_\gamma, d_{M_\gamma}) \rightarrow (N_\gamma, d_{N_\gamma})$ be such that

$$f(B^M(x; \gamma)) = B^N(f(x), \gamma)$$

Then f_γ is non-expanding and well-defined.

Some background: closed ball functor

Lemma 2.7 $(\cdot)_\gamma$ is a functor from $UMet(\Gamma)$ to $UMet(\Gamma_\gamma)$

Let's see an example

Motivation

"In order to show that each Γ -ultrametric space has a minimal, strongly Cauchy complete extension, we first show that the category of Γ -ultrametric spaces is equivalent to the category of flabby separated presheaves on Γ^{op} ."

- ▶ Theorem 2.25: a Γ -ultrametric space is injective if and only if is strongly Cauchy complete.
- ▶ Then an injective hull of a Γ -ultrametric space is a minimal extension that is strongly Cauchy complete
- ▶ Idea: show every object in $Flab(\Gamma^{op})$ has an injective hull and use the equivalence.

More notation:

- ▶ $Flab(\Gamma^{op})$: the category of flabby separated presheaves on Γ^{op}
- ▶ $Sep(\Gamma^{op})$: the category of separated presheaves on Γ^{op}
- ▶ $i_\Gamma : Flab(\Gamma^{op}) \rightarrow Sep(\Gamma^{op})$: the inclusion functor

First we show an equivalence of categories between $UMet(\Gamma)$ and $Flab(\Gamma^{op})$ via two functors, F_Γ and G_Γ .

the functor F_Γ

We define $F_\Gamma : \mathbf{UMet}(\Gamma) \rightarrow \mathbf{Flab}(\Gamma^{op})$ as follows:

Definition

Let $(M, d_M) \in \mathbf{UMet}(\Gamma)$ and define $F_\Gamma(M, d_M)(\gamma) = (M_\gamma, d_{M_\gamma})$.
Let the restriction maps be given by: $B(a, \gamma) \restriction_{\gamma^*} = B(a, \gamma^*)$ for $\gamma^* \geq \gamma$.

Check that:

- ▶ The restriction maps are well-defined
- ▶ $F_\Gamma(M, d_M)$ is flabby, i.e. if $B(a, \gamma) \in F_\Gamma(M, d_M)(\gamma)$, then there exists $\tilde{\gamma} \leq \gamma \in \Gamma$ such that $B(a, \gamma) \restriction_{\tilde{\gamma}} = B(a, \tilde{\gamma})$
- ▶ $F_\Gamma(M, d_M)$ is separated
- ▶ If $f : (M, d_M) \rightarrow (N, d_N)$ is a non-expanding map, then $F_\Gamma(f) : F_\Gamma(M, d_M) \rightarrow F_\Gamma(N, d_N)$ is a map of flabby separated presheaves.

the functor G_Γ

We define $G_\Gamma : Flab(\Gamma^{op}) \rightarrow UMet(\Gamma)$ as follows:

Definition

Let $A \in Flab(\Gamma^{op})$ and define $G_\Gamma(A) = (A(\perp), d_{A\perp})$, where $d_\perp(a, b) = \bigwedge \{ \gamma : a \upharpoonright_\gamma = b \upharpoonright_\gamma \}$.

Check that:

- ▶ The distance function satisfies the strong triangle identity
- ▶ $(\forall a, b \in A(\perp)) \ d_\perp(a, b) = \perp \iff a = b$
- ▶ If $A, C \in Flab(\Gamma^{op})$ and $f : A \rightarrow C$ is a map of presheaves then $G_\Gamma(f) : G_\Gamma(A) \rightarrow G_\Gamma(C)$ is a non-expanding map.

equivalence of categories

Claim 3.19

There is a natural isomorphism $\eta : F_{\Gamma} \circ G_{\Gamma} \rightarrow id_{Flab(\Gamma^{op})}$ such that $(\forall x \in A(\perp)) \eta_A(\{x\}) = x$ when $A \in Flab(\Gamma^{op})$.

Check that:

- ▶ η is well-defined and injective (since A being flabby will imply that is surjective)
- ▶ η is natural

equivalence of categories

Claim 3.20

There is a natural isomorphism $\epsilon : id_{UMet(\Gamma)} \rightarrow G_\Gamma \circ F_\Gamma$.

- ▶ Suppose $(M', d_{M'}) = G_\gamma(F_\gamma(M, d_M))$, so that $M' = \{\{a\} : a \in M\}$.
- ▶ Check directly that the map given by $\epsilon_{M, d_M}(a) = \{a\}$ is an isomorphism. Naturality is immediate.

Theorem 3.12

There is an equivalence of categories between $Flab(\Gamma^{op})$ and $UMet(\Gamma)$

injective hull

Definition

Suppose C is a category and $A \in C$. An **injective hull** for A is a monic $e : A \rightarrow I$ where

1. I is injective
2. If $k : A \rightarrowtail I'$ is a monomorphism with I' injective, there is a monomorphism $k' : I \rightarrowtail I'$ such that $k' \circ e = k$.

We show every object of $Flab(\Gamma^{op})$ has an injective hull.

injective hull

Claim 3.23 For every object \mathcal{F} of $Flab(\Gamma^{op})$, the following are equivalent:

1. \mathcal{F} is injective in $Flab(\Gamma^{op})$.
2. $i_{\Gamma}(\mathcal{F})$ is injective in $Sep(\Gamma^{op})$

In fact, this is enough, since we also have [2],

Claim 3.24 The injective objects of $Sep(\Gamma^{op})$ are exactly the flabby sheaves and every object of $Sep(\Gamma^{op})$ has an injective hull.

injective hull

$\mathcal{F} \in \text{Flab}(\Gamma^{op})$ is injective $\Rightarrow i_{\Gamma}(\mathcal{F})$ is injective in $\text{Sep}(\Gamma^{op})$

Proof.

Let A be a separated presheaf on Γ . Define A^* as follows:

- ▶ $A^*(V) = \cup_{U \leq V} A(U) \times \{U\} \times \{V\}$
- ▶ If $U \leq V$ and $\langle x, W, V \rangle \in A^*(V)$ then
 $\langle x, W, V \rangle|_U = \langle x|_{U \wedge W}, U \wedge W, U \rangle \in A^*(U)$.

Define m, m^*, e_A



references



N.L. Ackerman, "Completeness in Generalized Ultrametric Spaces," *p-Adic Numbers, Ultrametric Analysis and Applications*, **5**(2),89-105 (2013).



J. Coulon and J-L.Coulon, "Injectivity in the quasi-topos of separated presheaves on a complete Heyting algebra," *Portugal. Math.* **47**(3),227-241 (1990).