

[Login](#)[Register](#)[News](#)[Journals](#)[Events](#)[Compendia](#)[Media](#)[Continuing Education](#)[Formulary](#)[X](#)

Currently Viewing:

American Thoracic Society 2018

Public Health

May 22, 2018

How IMPACT Design Allowed Trelegy Ellipta to Expand Indication

May 22, 2018

Currently Reading **Smartphones, Apps, Smart Inhalers: Using Tech to Personalize COPD Care**

May 22, 2018

Medicaid Expansion Linked to Decline in ICU Stays, Study at ATS Finds

May 22, 2018

Screen More Former Smokers for Lung Cancer, Study Suggests

May 23, 2018

Welcome the the new and improved AJMC.com, the premier managed mail.
Tell us about yourself so that we can serve you better.



Conferences > American Thoracic Society 2018 – Published on: May 22, 2018



Smartphones, Apps, Smart Inhalers: Using Tech to Personalize COPD Care



Mary Caffrey



From smartphones to smart rescue inhalers, researchers offered ideas to make chronic obstructive pulmonary disease care more data-driven and personalized.

A randomized clinical trial in telehealth was a disappointment. But a host of small studies using smartphone apps, pulse oximeters, and a digital rescue inhaler showed promise during Tuesday's technology-focused poster session at the American Thoracic Society 2018 International Conference in San Diego, California.

As one researcher put it, right now it's hard to predict which patients with chronic obstructive pulmonary disease (COPD) will end up in the emergency department (ED), or which ones will have a second exacerbation within days of the first. So, studies presented at the session put forth a host of ideas for tracking patient data with the goal of tackling this problem. There were also ideas for using technology for better remote management without adding significant staff time or costs.

"Smartphone ownership has grown over the last several years in the target demographic," said Jennifer T. Krall, MD, the lead author on one of the abstracts. Thus, asking patients with COPD to use personal digital tools to collect data or manage their disease is rarely the challenge it might have been just a few years ago.

might have been just a few years ago.

Personalization is a priority, said Andrew Miller, MD, another study author, due to the Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017 mandates, which call for greater emphasis on individualized care and were cited frequently throughout ATS 2018.

Smartphone management. Krall, Cone Health in Greensboro, North Carolina, presented results of a feasibility study using a smartphone application to promote patient engagement. Of

Welcome the the new and improved AJMC.com, the premier managed mail ^x |

Tell us about yourself so that we can serve you better.

77% to 97%. It took staff less than 1 person/hour a day to manage the patient portal.¹

Pulse oximeters. Home- or smartphone-based tools offer a noninvasive way to measure oxygen saturation, but commercial apps can have varying levels of accuracy. A team from the University of Illinois tested 3 “wellness” pulse oximeter apps—from Walgreens C20, iCare, and SHealth—against a reference app that received FDA clearance, the Nonin Onyx II. The 19 patients had a mean peripheral capillary oxygen saturation (SpO₂) of 92.7 (7.2)% as measured by the reference standard. The mean percent error for the other pulse oximeters ranged from -1.8% to +5.7%; the Walgreens C20 was the most accurate; but none of these apps could identify severe hypoxemia, which is SpO₂ ≤ 88%. Authors found a substantial price difference; the most expensive of the 3 non-FDA cleared apps was \$39.99, compared with \$350 for the reference app.²

A “smart” rescue inhaler. Miller, of the University of Colorado, captured the audience’s attention with his zeal for his pilot study, which involves equipping rescue inhalers with passive sensors to capture the exact time and date when patients are using them. Not only does this eliminate recall bias, but as Miller explained, it also opens up a host of possibilities for “machine learning.” There’s a difference between “frequent use,” and “patterned use,” he said; in his home state of Colorado, for example, running patient use data against known weather patterns would likely show how rescue inhaler use increase when there is a forest fire. In time, the data would give clinicians the ability to meet the GOLD 2017 mandate for more personalized approaches to COPD management.³ While the pilot is small, so far, the patients who have infrequent rescue inhaler use without a pattern have a body mass index (BMI) of less than 28 kg/m².

Negative results, but patients liked telehealth. PROMETE II was a multicenter, randomized clinical trial that randomized 229 patients to either telehealth or routine clinical practice, with the primary endpoint of reducing exacerbations that led to a visit to the emergency department (ED) and hospital admission. Participants had an average age of 71 years and 80% were men. After 12 months, the number of COPD-related ED visits and all-cause deaths was comparable between the 2 groups. The number of mean exacerbations was 1.0 in the

**Welcome the the new and improved AJMC.com, the premier managed market
Tell us about yourself so that we can serve you better.**

average). On the plus side, doctors and patients both liked telehealth, despite the results.⁴

Reference

1. Krall JT, Alva R. Feasibility of using a novel smartphone application to improve compliance and engagement in chronic obstructive pulmonary disease (COPD). Presented at the American Thoracic Society 2018 International Conference; San Diego, California; May 18-23, 2018; Abstract A4528.
2. Brilliante C, Binder A, Luo J, Prieto-Centurion V. Can smartphone-based pulse oximeters be used for monitoring patients with chronic obstructive pulmonary disease? Presented at the American Thoracic Society 2018 International Conference; San Diego, California; May 18-23, 2018; Abstract A4554.
3. Miller AL, Bowler RP, Allinder M, et al. Real-time data collection of rescue inhaler use in COPD patients: an important step to achieving the new GOLD standard. Presented at the American Thoracic Society 2018 International Conference; San Diego, California; May 18-23, 2018; Abstract A4559.
4. Soriano JB, Garcia-Rio F, Vazquez-Espinosa E, et al. Efficacy and costs of telehealth for the management of COPD: a multicenter, randomized controlled trial. Presented at the American Thoracic Society 2018 International Conference; San Diego, California; May 18-23, 2018; Abstract A4546.

Related Articles

[Out With the Stethoscope, In With the Smartphone: How Genomics and Digital Advances Will](#)

[Reshape Practice](#)

[Study Investigates New COPD Categories With GOLD 2017](#)

[Claims-Based Risk Model for First Severe COPD Exacerbation](#)

Welcome to the new and improved AJMC.com, the premier managed market
Tell us about yourself so that we can serve you better.



Copyright AJMC 2006-2019 Clinical Care Targeted Co