

# Sections and Chapters

John Paul Guzman

$\langle \frac{2}{3} \rangle$

## 1 Basic Logic

=====

### 1.1 (Remark) Modern Logic

- Proof theory - properties of proofs
  - Model theory - properties of interpretations
  - Recursion theory - properties of algorithms
  - Set theory - structureMaster
- =====

### 1.2 (Remark) Bootstrapping

- Informal math contains all the real ideas, while formal math contains encoded symbols
  - Formal math is mere toy-replica of the informal math
  - Properties of the toy can tell us about properties of the real thing
  - Formal reasoning (syntax) is much safer than informal reasoning (syntax + semantics)
  - The informal theory about a formal theory is called a metatheory
- =====

### 1.3 (Remark) Notions in the metatheory are informal

- Metamathematics exists outside and independently of our effort to build this or that formal system
  - - All its constructs are available to us for use in the analysis of the behaviour of a formal system
  - Properties about objects that they must satisfy can only be recognized in the metatheory
  - - Formal theories are pre-built systems that are designed to run not knowing about these properties
  - The formal theory is just a generator of theorems, and not a parser
  - - It cannot remember what it generates nor state the properties of some given string
- =====

### 1.4 (Definition) Pre-requisite informal notions

- Subset  $X \subseteq Y$  for sets  $X$  and  $Y$  iff for all  $x \in X$ ,  $x \in Y$
- Union  $X \cup Y$  for sets  $X$  and  $Y$  iff for all  $x \in X$  and  $y \in Y$ ,  $x \in X \cup Y$  and  $y \in X \cup Y$
- Kleene star  $S^*$  is the set of all sequences that can be made from the elements of set  $S$
- The string  $S$  from a set  $V$  is defined by  $S \in V^*$
- String concatenation  $A * B$  for strings  $A$  and  $B$  is defined by  $A$  appended with  $B$
- The empty string  $\lambda$  is defined by for all strings  $S$ ,  $S = \lambda * S = S * \lambda$
- String  $A$  occurs in string  $B$  iff there exists strings  $C$  and  $D$ ,  $B = C * A * D$
- Variadic notation  $[a_i]_{i=1}^n$  is an abbreviation for  $a_1, a_2, a_3, \dots, a_n$
- TODO: rules + rule metatheorems, schema substitution =====

### 1.5 (Definition) Rule

- The rule  $Q$  is a metatheoretical function that takes in a sequence of strings and outputs strings
- The rule with  $n$  arguments and an output is called  $(n + 1)$ -ary
- The rule has an associated function *Arity* which outputs the number of arguments for a given function or relation
- The function *ArityR* outputs the arity of given a given rule - Rules must be algorithmic and can be executed within a

finite number of steps - The immediate predecessors of a string  $d$  on the  $(n + 1)$ -ary rule  $Q$  are  $[s_i]_{i=1}^n$  iff  $Q([s_i]_{i=1}^n, d)$  holds  
=====

## 1.6 (Definition) Closed set under a rule

- TODO: use/meaning/intent - wat - The set  $S$  is closed under the  $(n + 1)$ -ary rule  $Q$  iff for all  $d$  where  $Q([s_i]_{i=1}^n, d)$ , if  $\{[s_i]_{i=1}^n\} \subseteq S$ , then  $d \in S$   
=====

## 1.7 (Definition) Rule-defined set

- The rule-defined set  $Cl(J, R)$  consists of:
  - - The set of base objects  $J$
  - -  $J$  can be treated like a set of rules that takes no arguments
  - - The set of rules for generating inductive objects  $R$
  - Closure  $Cl(J, R)$  is the smallest set that satisfies all of the following:
  - -  $J \subseteq Cl(J, R)$
  - - for all  $Q \in R$ ,  $closed(Cl(J, R), Q)$
  - Sets built via rules satisfy a smallest qualifier to remove erroneous structures and elements
- 
- =====

## 1.8 (Metatheorem) Induction on a rule-defined set

- TODO: use/meaning/intent - If a property  $P$  holds for all in  $J$ , and propagates (true for input (Induction Hypothesis) -> true for output (Inductive Step)) through for all in  $R$ , then it holds for the entire closure ??? - For any property  $P$  and  $w \in Cl(J, R)$ ,  $P(w)$  iff - For all  $j \in J$ ,  $P(j)$  - For all  $r \in R$ ,  $P$  closed blablablablabl TODO fuck this shit - If  $J \subseteq T$  (Basis Step) and for any  $Q \in R$ ,  $T$  is closed under  $Q$  (Inductive Step), then  $Cl(J, R) \subseteq T$   
=====

## 1.9 (Definition) Ambiguous pair

- TODO: use/meaning/intent - ??? - The pair  $(J, R)$  is ambiguous if there exists  $d \in Cl(J, R)$  satisfies any of the following:
  - - There exists  $Q \in R$  where  $Q([x_i]_{i=1}^{ArityR(Q)})$  and  $Q([y_i]_{i=1}^{ArityR(Q)})$  and  $\langle [x_i]_{i=1}^{ArityR(Q)} \rangle \neq \langle [y_i]_{i=1}^{ArityR(Q)} \rangle$
  - - There exists  $P \in R$  and  $Q \in R$  where  $P([s_i]_{i=1}^n, d)$  and  $Q([s_i]_{i=1}^n, d)$  and  $P \neq Q$
  - - There exists  $Q \in R$  where  $Q([s_i]_{i=1}^n, d)$  and  $d \in J$
- 
- =====

## 1.10 (Metatheorem) Definition by recursion

TOFIX - TODO: use/meaning/intent -  $Cl(J, R)$  from some unambiguous  $(J, R)$  has some nice unique mapping - TODO: [ABSTRACTED] - All inductive or recursive definitions must be unambiguous to be well-defined  
=====

## 1.11 (Remark) The state of the metatheory

- The metatheory houses assumptions or axioms about behavior of theories
  - Is it necessary to prove the consistency of the metatheory?
  - - No because otherwise, it would require a meta-metatheory which would require a meta-meta-metatheory, and so on which would never end
  - - The metatheory should be small and simple and close to intuition such that it would not require formalized verification of its consistency
  - Is it okay to use infinite sets and induction in the metatheory?
  - - This is mostly political, but we should avoid using suspicious inconsistent tools such as full-blown naive set theory
  - - There are ways we could simulate infinite sets using safer finite notions like using finite calculations then arbitrarily large finite can be a stand-in for infinite, but it is not worth it
- 
- =====

## 2 First Order Languages

  
=====

## 2.1 (Definition) Formal theory

- TODO: use/meaning/intent - notionmaster - The formal theory  $T$  is defined by  $T = (V, Wff, Thm)$
  - The alphabet  $V$  is defined by the set of all symbols allowed in the theory
  - The set of all strings  $String$  from an alphabet  $V$
  - The set of all well-formed formulas  $Wff$  is defined by  $Wff \subseteq String$
  - The set of all theorems  $Thm$  is defined by  $Thm \subseteq Wff$
- =====

## 2.2 (Definition) Formal language

- The formal language encodes the notions of a theory
  - The formal language  $L$  is defined by  $L = (V, Term, Wff)$
  - $Term$  encodes the objects of a theory
  - $Wff$  encodes the statements of a theory
  - $L$  is a (one-sorted) language with only one type of variable in  $V$ , and then predicates are used to simulate typing
  - $Term, Wff, Thm$  can be explicitly provided or they can be rule-defined
- =====

## 2.3 (Definition) Alphabet

- TODO: use/meaning/intent - writeables - An alphabet  $V$  consists of elements from  $LS$  and  $NLS$ :
  - - The set of all logical symbols  $LS$  consists of:
    - - - Elements from the set of all variables  $Var$
    - - - The boolean connectives  $\neg, \vee$
    - - - The existential quantifier  $\exists$
    - - - The equality predicate  $\equiv$
  - - The set of all nonlogical symbols  $NLS$  consists of:
    - - - Elements from the set of all constants  $Const$
    - - - Elements from the set of all predicates  $Pred$
    - - - Elements from the set of all functions  $Func$
  - $NLS$  can be extended to contain additional symbols
- =====

## 2.4 (Definition) Set of all terms

- TODO: use/meaning/intent - talkables - The set of all terms  $Term$  is defined by  $Term = Cl(J_{Term}, R_{Term})$
  - The set of all base terms  $J_{Term}$  consists of:
    - - Variables from  $Var$
    - - Constants from  $Const$
  - The set of term generating rules  $R_{Term}$  is defined by the rules:
    - - If  $f \in Func$  and  $\{[t_i]_{i=1}^{Ar(f)}\} \subseteq Term$ , then  $f[t_i]_{i=1}^{Ar(f)} \in Term$
- =====

## 2.5 (Definition) Set of all wffs

- TODO: use/meaning/intent - truthables - The set of all wffs  $Wff$  is defined by  $Wff = Cl(J_{Wff}, R_{Wff})$
  - The set of all base (or atomic) wffs  $J_{Wff}$  is the set of all predicates that take only terms as arguments and it is defined by the rules:
    - - If  $\{t, s\} \subseteq Term$ , then  $(t \equiv s) \in J_{Wff}$
    - - If  $p \in Pred$  and  $\{[t_i]_{i=1}^{Ar(p)}\} \subseteq Term$ , then  $p[t_i]_{i=1}^{Ar(p)} \in J_{Wff}$
  - The set of wff generating rules  $R_{Wff}$  is defined by the rules:
    - - If  $A \in Wff$ , then  $\neg A \in Wff$
    - - If  $\{A, B\} \subseteq Wff$ , then  $\vee AB \in Wff$
    - - If  $x \in Var$  and  $A \in Wff$ , then  $\exists_x A \in Wff$
  - Allowing only variables to be quantified is what makes the language “first-order”
  - First-order languages ...
- =====

## 2.6 (Definition) Set of all thms

- TODO: use/meaning/intent - truths - The set of all thms  $Thm$  is defined by  $Thm = Cl(J_{Thm}, R_{Thm})$
- The set of all base (or axiomatic) thms  $J_{Thm}$  is defined by  $J_{Thm} \subseteq Wff$

- The set of thm generating rules (or rules of inference)  $R_{Thm}$  is defined a set of rules where each rule takes in wffs and outputs a wff

### 3 Constructs for Thm

#### 3.1 (Definition) Wff abbreviations

- $(A \vee B)$  is an abbreviation for  $\vee AB$
- $\forall_x A$  is an abbreviation for  $\neg \exists_x \neg A$
- $(A \wedge B)$  is an abbreviaton for  $\neg(\neg A \vee \neg B)$
- $(A \implies B)$  is an abbreviation for  $\neg A \vee B$
- $(A \equiv B)$  is an abbreviation for  $((A \implies B) \wedge (B \implies A))$
- Precedence:
  - -  $\neg, \wedge, \vee, \implies, \equiv$
  - - Equal precedence invokes right associativity

#### 3.2 (Definition) Free variable in a term

- TODO: use/meaning/intent - ??? - The variable  $x$  is free in a term  $t$  iff it  $x$  occurs in  $t$

#### 3.3 (Definition) Closed term

- TODO: use/meaning/intent - ??? - The term  $t$  is closed iff it contains no free variable

#### 3.4 (Definition) Free and bound variable in a wff

- TODO: use/meaning/intent - ??? - The variable  $x$  is free in a wff  $B$  if it satisfies any of the following:
  - - Iff  $B$  is an atomic wff and  $x$  occurs in  $B$
  - - Iff  $B$  is  $\neg C$  and  $x$  is free in  $C$
  - - Iff  $B$  is  $C \vee D$  and ( $x$  is free in  $C$  or  $x$  is free in  $D$ )
  - - Iff  $B$  is  $\exists_y(C)$  and  $x \neq y$  and  $x$  is free in  $C$
- The variable  $x$  is bounded in a wff  $B$  iff  $x$  occurs in  $B$  and  $x$  is not free in  $B$

#### 3.5 (Definition) Closed and open wff

- TODO: use/meaning/intent - ??? - The wff  $C$  is closed iff it contains no free variable
- The wff  $C$  is open iff it contains no quantifier
- Open wffs are also closed

#### 3.6 (Definition) Variable occurrence notation

- $A[[v_i]_{i=1}^n]$  iff the variables  $[v_i]_{i=1}^n$  occur in the wff  $A$
- $A([v_i]_{i=1}^n)$  iff the variables  $[v_i]_{i=1}^n$  occur in the wff  $A$  and no other variables occur in  $A$

#### 3.7 (Definition) The set all prime wffs

- TODO: use/meaning/intent - atom truth assignable - The set of all base terms *Prime* consists of:
  - Atomic wffs from  $J_{Wff}$
  - Inductive wffs under from the  $\exists_x A$  rule in  $R_{Wff}$

### 3.8 (Definition) Propositional valuation

- TODO: use/meaning/intent - ??? - The valuation  $v$  is a function defined by  $v : Prime \rightarrow \{\perp, \top\}$  - The propositional valuation  $p_v$  from the valuation  $v$  is defined by  $p_v : Wff \rightarrow \{\perp, \top\}$  and  $p_v$  is an extension of  $v$  from  $Prime$  to  $Wff$
- If wff  $A$  is a prime wff, then  $p_v(A) = v(A)$
- If wff  $A$  is not a prime wff, then  $p_v$  is defined as follows:
  - $p_v(\neg A) = F_{\neg}(A)$
  - $F_{\neg}(\top) = \perp$
  - $F_{\neg}(\perp) = \top$
  - $p_v(A \vee B) = F_{\vee}(A, B)$
  - $F_{\vee}(\perp, \perp) = \perp$
  - $F_{\vee}(\perp, \top) = \top$
  - $F_{\vee}(\top, \perp) = \top$
  - $F_{\vee}(\top, \top) = \top$
- This definition relies on the definition of prime wff and parsing to be unambiguous =====

### 3.9 (Definition) Tautology

- TODO: use/meaning/intent - ??? - The wff  $T$  is a tautology ( $\models_{Taut} T$ ) iff for any valuation  $v$ , the propositional valuation  $p_v(T) = \top$
- =====

### 3.10 (Definition) Satisfiable wff

- TODO: use/meaning/intent - ??? - The wff  $U$  is wffsatisfiable iff there exists a valuation  $v$  where,  $p_v(U) = \top$
- $v$  wffsatisfies  $U$
- =====

### 3.11 (Definition) Satisfiable set

- TODO: use/meaning/intent - ??? - The set of wffs  $\Gamma$  is setsatisfiable iff there exists a valuation  $v$  where, for any  $A \in \Gamma$ ,  $p_v(A) = \top$
- $v$  setsatisfies  $\Gamma$
- =====

### 3.12 (Definition) Unsatisfiable wff

- Inverted version of wffsatisfiable
- The wff  $U$  is wffunsatisfiable iff for any valuation  $v$ ,  $p_v(U) = \perp$  =====

### 3.13 (Definition) Unsatisfiable set

- Inverted version of setsatisfiable
- An set of wffs  $\Gamma$  is setunsatisfiable iff for any valuation  $v$ , there exists  $A \in \Gamma$  where,  $p_v(A) = \perp$
- =====

### 3.14 (Definition) Tautological implication

- TODO: use/meaning/intent - ????? - The set of wffs  $\Gamma$  tautologically implies a wff  $A$  ( $\Gamma \models_{Taut} A$ ) iff for any valuation  $v$ , if  $v$  setsatisfies  $\Gamma$ , then  $v$  wffsatisfies  $A$
- This can also be checked via truth tables =====

### 3.15 (Remark) Satisfiable and unsatisfiable

- """"Satisfiable" and "unsatisfiable" are terms introduced here in the propositional or Boolean sense. These terms have a more complicated meaning when we decide to "see" the object variables and quantifiers that occur in formulas."""" TODO
- =====

### 3.16 (Definition) Substitution of a term for a variable in term

- TODO: use/meaning/intent - nifty rule dependency - The substitution ( $s[x \leftarrow t]$ ) of the variable  $x$  by the term  $t$  in the term  $s$  is defined as following rule:
  - If  $s$  is  $x$ , then the substitution outputs  $t$
  - If  $s$  is a constant or a variable that is not  $x$ , then the substitution outputs  $s$
  - If  $s$  is a function  $f$  applied to the terms  $[t_i]_{i=1}^n$ , then the substitution outputs  $f[t_i[x \leftarrow t]]_{i=1}^{Ariety(f)}$
- =====

### 3.17 (Metatheorem) Substituted terms are terms

- By induction, this is true for B.S. terms and joining terms I.H. implies joined is also a term I.S.
- =====

### 3.18 (Definition) Substitution of a term for a variable in wff

- TODO: use/meaning/intent - nifty rule - The substitution ( $A[x \leftarrow t]$ ) of the variable  $x$  by the term  $t$  in the wff  $A$  is defined as following rule:
  - If  $A$  is  $s \equiv r$  on the terms  $s$  and  $r$ , then the substitution outputs  $s[x \leftarrow t] \equiv r[x \leftarrow t]$
  - If  $A$  is a predicate  $p$  applied to the terms  $[t_i]_{i=1}^{Ariety(p)}$ , then the substitution outputs  $p[t_i[x \leftarrow t]]_{i=1}^{Ariety(p)}$
  - If  $A$  is  $\neg B$  on the wff  $B$ , then the substitution outputs  $\neg B[x \leftarrow t]$
  - If  $A$  is  $(B \vee C)$  on the wffs  $B$  and  $C$ , then the substitution outputs  $(B[x \leftarrow t] \vee C[x \leftarrow t])$
  - If  $A$  is  $\exists_x(B)$  on the wff  $B$ , then the substitution outputs  $A$
  - If  $A$  is  $\exists_y(B)$  on the variable  $y$  and the wff  $B$  and  $y$  is not  $x$  and  $y$  does not occur in  $t$ , then the substitution outputs  $\exists_y(B[x \leftarrow t])$
  - If  $x$  is not quantified over, then all variables in the term  $t$  must not be quantified over to allow the wff to preserve its intended meaning
  - The substitution is defined iff it satisfies a defined condition in the rule
  - Using the substitution notation immediately implies that the substitution output is defined
  - The symbols  $[ , ] , \leftarrow$  are symbols in the metatheory and they have the higher precedence compared to formal symbols
  - The variable  $x$  is substitutable by the term  $t$  in the wff  $A$  iff  $A[x \leftarrow t]$  is defined
- =====

### 3.19 (Metatheorem) Substituted wffs are wffs

- By induction, this true for B.S (terms) and joining wffs I.H. implies joined is also a wff I.S.
- =====

### 3.20 (Definition) Simultaneous substitution

- The simultaneous substitution  $A[[y_i]_{i=1}^n \leftarrow [t_i]_{i=1}^n]$  of the variables  $[y_i]_{i=1}^n$  by the terms  $[t_i]_{i=1}^n$  in the wff  $A$  is an abbreviation for  $A[[y_i \leftarrow t_i]]_{i=1}^n$
- =====

### 3.21 (Definition) Schemata

- The schemata is a rule written down as a metaformula that consists of metasymbols
  - These metasymbols can be replaced by formal symbols to output a wff called an instance of the schema
  - It is a rule that takes in formal symbols, and substitutes them in place of metavariables, then outputs the instance wff
- =====

### 3.22 (Definition) The set of all axioms

- TODO: use/meaning/intent - assumptions - The set of all axioms  $J_{Thm}$  consists of:
- Elements from the set of logical axioms  $\Lambda$
- Elements from the set of nonlogical axioms  $\Gamma$

### 3.23 (Definition) The set of all logical axioms

- TODO: use/meaning/intent - assumed logic behavior - The set of all logical axioms  $\Lambda$  is defined by  $\Lambda = Cl(J_\Lambda, R_\Lambda)$
- The set of all base logical axioms  $J_\Lambda$  is the set of all tautological wffs
- The set of logical axiom generating rules  $R_\Lambda$  is defined by the rules:
  - Substitution Axiom: If  $A \in \Lambda$ , then for any term  $t$  and variable  $x$  substitutable for  $t$ ,  $(A[x \leftarrow t] \implies \exists_x(A)) \in \Lambda$
  - Self-equivalence Axiom: For any variable  $x$ ,  $(x \equiv x) \in \Lambda$
  - Leibniz Axiom: For any (countable) wff  $A$ , variable  $x$ , terms  $t$  and  $s$ ,  $(t \equiv s \implies (A[x \leftarrow t] \equiv A[x \leftarrow s])) \in \Lambda$
- The rules / axioms are what endow the symbols with the intended behavior
- The Leibniz Axiom is still first-order because ??? =====

### 3.24 (Definition) The set of all nonlogical axioms

- TODO: use/meaning/intent - assumed notions behavior - The set of all nonlogical axioms  $\Gamma$  are the wffs that are assumed to be true or the hypotheses =====

### 3.25 (Definition) Rules of inference

- TODO: use/meaning/intent - truth preserving - The rules of inference  $R_{Thm}$  is defined by the rules:
    - Primary rules of inference:
      - Modus Ponens Rule: Given  $A, A \implies B$ , the output is  $B$
      - E-Introduction Rule: Given  $A \implies B$  and  $x$  is not free in  $B$ , the output is  $\exists_x(A) \implies B$
    - Derived rules of inference:
      - Rules of inference derived from the proofs made by other rules of inference
      - The validity of these rules are only provable within the metatheory
- =====

### 3.26 (Definition) The set of all Gamma-Theorems

- TODO: use/meaning/intent - Provable wffs - The set of all Gamma-Theorems  $Thm_\Gamma$  is defined by  $Thm_\Gamma = Cl(J_{Thm}, R_{Thm})$
  - The wff  $A$  is a Gamma-theorem or  $\Gamma \vdash A$  is an abbreviation for  $A \in Thm_\Gamma$
  - $\Gamma \vdash \Lambda$  is an abbreviation for for all  $L \in \Lambda, L \in Thm_\Gamma$
- =====

### 3.27 (Definition) Abstract formal theory

- The formal theory  $T_L$  appropriate for the language  $L$  is the set of wffs  $T_L \subseteq Wff$  that are considered to be correct within a theory
  - The formal theory  $T$  is defined by  $T = Cl(J_{Thm}, R_{Thm})$
  - The formal theory  $T$  is inconsistent if  $T = Wff$
  - We often like to find the smallest set of set of axioms  $\Gamma$  for axiomatizing  $T$  such that  $T = Thm_\Gamma$
  - The set of axioms  $\Gamma$  is recognizable if there exists an algorithmic process to decide if  $A \in \Gamma$
  - The theory  $T$  is recursively axiomatized if  $\Gamma$  is recognizable
- =====

## 4 Constructs and Metatheorems for Provability

=====

### 4.1 (Definition) Derivation and stage

- The  $(J, R)$ -derivation is the finite sequence of wffs from the construction of  $Cl(J, R)$
  - The stage  $X_i$  is some collection of all wffs from the  $(J, R)$ -derivation until step  $i$
- =====

### 4.2 (Metatheorem) Equivalent ways of generating closures

- $Cl(J, R) = x : x \text{ occurs in } (J, R) - \text{derivation} = \cup X_i$
- TODO: [ABSTRACTED] duh =====

### 4.3 (Definition) Gamma-proof

- The Gamma-proof of the wff  $A$  is some  $(J, R)$ -derivation of a  $\Gamma$ -Theorem =====

### 4.4 (Metatheorem) Transitivity of $\vdash$

- If  $\Gamma \vdash \Delta$  and  $\Delta \vdash B$ , then  $\Gamma \vdash B$
- The existing Gamma-proofs can be concatenated to form the Gamma-proof for  $B$
- This allows us to re-use collections of previously established theorems =====

### 4.5 (Metatheorem) Hypothesis strengthening

- If  $\Gamma \subseteq \Delta$  and  $\Gamma \vdash A$ , then  $\Delta \vdash A$
  - The Gamma-proof for  $A$  is also a valid Delta-proof for  $A$  since it contains all the required wffs
- =====

### 4.6 (Metatheorem) Post's Tautology Theorem

- If  $\{[A_i]_{i=1}^n\} \models_{Taut} B$ , then  $\{[A_i]_{i=1}^n\} \vdash B$
  - $\models_{Taut} [A_i \implies ]_{i=1}^n B$  i(1) from Hypothesis 1 and truth table
  - $\vdash [A_i \implies ]_{i=1}^n B$  i(2) from tautology 1 in  $J_{Thm}$
  - $\{[A_i]_{i=1}^n\} \vdash [A_i \implies ]_{i=1}^n B$  i(3) from Hypothesis strengthening on (2)
  - $\{[A_i]_{i=1}^n\} \vdash B$  i(4) from Modus Ponens Rule on (3)  $n$  times
- =====

### 4.7 (Definition) Provably equivalent wffs in a theory

- The wff  $A$  and wff  $B$  are provably equivalent in the theory  $T$  iff  $\Gamma \vdash A \equiv B$  =====

### 4.8 (Metatheorem) Theorems are provably equivalent

- If  $\Gamma \vdash \{A, B\}$ , then  $\Gamma \vdash (A \equiv B)$
  - $\Gamma \vdash A$  i(1) from Hypothesis 1
  - $A \models_{Taut} B \implies A$  i(2) from tautological implication
  - $A \vdash B \implies A$  i(3) from Post's Tautology Theorem on (2)
  - $\Gamma \vdash B \implies A$  i(4) from Transitivity of  $\vdash$  on (1, 3)
  - $\Gamma \vdash A \implies B$  i(5) Ditto of (1-4) but utilizing Hypothesis 2
  - $\Gamma \vdash \{B \implies A, A \implies B\}$  i(6) Collection of established theorems on (4, 5)
  - $\{A \implies B, B \implies A\} \models_{Taut} A \equiv B$  i(7) from tautological implication
  - $\{A \implies B, B \implies A\} \vdash A \equiv B$  i(8) from Post's Tautology Theorem on (7)
  - $\Gamma \vdash A \equiv B$  i(9) Transitivity of  $\vdash$  on (6, 8)
- =====

### 4.9 (Metatheorem) A-Introduction

- If the variable  $x$  is not free in the wff  $\neg A$ , then  $A \implies B \vdash A \implies \forall_x(B)$
  - $A \implies B \models_{Taut} \neg B \implies \neg A$  i(1) from tautological implication
  - $A \implies B \vdash \neg B \implies \neg A$  i(2) from Post's Tautology Theorem on (1)
  - $\neg B \implies \neg A \vdash \exists_x(\neg B) \implies \neg A$  i(3) from E-Introduction WITH Hypothesis 1
  - $\exists_x(\neg B) \implies \neg A \models_{Taut} A \implies \neg(\exists_x(\neg B))$  i(4) from tautological implication
  - $\exists_x(\neg B) \implies \neg A \vdash A \implies \neg(\exists_x(\neg B))$  i(5) from Post's Tautology Theorem on (4)
  - $A \implies \neg(\exists_x(\neg B)) \vdash A \implies \forall_x(B)$  i(6) from abbreviation of forall on (5)
  - $A \implies B \vdash A \implies \forall_x(B)$  i(8) from Transitivity of  $\vdash$  on (2, 3, 5, 6)
- =====

### 4.10 (Metatheorem) Specialization

- The succeeding proofs will be less based less on metatheory and based more on derivation
  - For any wff  $A$  and term  $t$ ,  $\vdash \forall_x(A) \implies A[x \leftarrow t]$
  - $(\neg A)[x \leftarrow t] \implies \exists_x(\neg A)$  i(1) from Substitution Axiom
  - $\neg \exists_x(\neg A) \implies A[x \leftarrow t]$  i(2) from tautological implication and Post's Tautology Theorem on (1)
  - $\forall_x(A) \implies A[x \leftarrow t]$  i(3) from abbreviation of forall on (2)
- =====



#### 4.11 (Metatheorem) Specialization corollary

- For any wff  $A$ ,  $\vdash \forall_x(A) \implies A$
  - $\forall_x(A) \implies A[x \leftarrow x]$  ;(1) from Specialization<sub>i</sub> -  $\forall_x(A) \implies A$  ;(2) from definition of substitution on (1)<sub>i</sub>
- =====

#### 4.12 (Metatheorem) Generalization

- For any wff  $A$ ,  $\vdash A \implies \forall_x(A)$  -  $A \implies A$  ;(1) from tautological implication and Post's Tautology Theorem<sub>i</sub>
  - $A \implies \forall_x(A)$  ;(2) A-Introduction on (1) WITH  $x$  is not free in  $A$ <sub>i</sub>
- =====

#### 4.13 (Metatheorem) Generalization corollary

- For any wff  $A$ ,  $\vdash A \equiv \forall_x(A)$
  - $\vdash A \implies \forall_x(A)$  ;(1) from Generalization<sub>i</sub>
  - $\vdash \forall_x(A) \implies \vdash A$  ;(2) from Specialization corollary<sub>i</sub>
  - $\vdash A \equiv \forall_x(A)$  ;(3) from tautological implication on (1, 2) and Post's Tautology Theorem<sub>i</sub>
- =====

#### 4.14 (Metatheorem) CorollaryMaster

- For any wff  $A$ ,  $A \vdash \forall_x(A)$  and  $\forall_x(A) \vdash A$
  - $A \vdash \forall_x(A)$  ;(1) from Modus Ponens Rule on  $(A \vdash A, \text{Generalization corollary})$ <sub>i</sub>
  - $\forall_x(A) \vdash A$  ;(2) from Modus Ponens Rule on  $(\forall_x(A) \vdash \forall_x(A), \text{Generalization corollary})$ <sub>i</sub>
  - $\vdash A \equiv \forall_x(A)$  ;(3) from tautological implication on (1, 2) and Post's Tautology Theorem<sub>i</sub>
- =====

#### 4.15 (Definition) Universal closure

- The universal closure of a wff  $A$  with free variables  $[y_i]_{i=1}^n$  is defined to be  $[\forall_{y_i}]_{i=1}^n(A)$
  - Any formula deduces and is deduced by its universal closure ;from CorollaryMaster<sub>i</sub>
- =====

#### 4.16 (Metatheorem) Substitution of terms

- For any terms  $[t_i]_{i=1}^n$ ,  $A[[x_i]_{i=1}^n] \vdash A[[t_i]_{i=1}^n]$  -  $[\forall_{x_i}]_{i=1}^n(A)$  ;(1) Generalization on (Hypothesis 1)  $n$  times<sub>i</sub> -  $A[[t_i]_{i=1}^n]$  ;(2) Specialization on (1)  $n$  times<sub>i</sub>
- =====

#### 4.17 (Metatheorem) Renaming

- For any wff  $A$  and variable  $z$ , if  $z$  does not occur in  $A$ , then  $\vdash (\exists_x(A)) \equiv (\exists_z(A[x \leftarrow z]))$  -  $A[x \leftarrow z] \implies \exists_x(A)$  ;(1) from Substitution Axiom WITH Hypothesis 1 (the variables in  $z$  does not occur in  $A$ )<sub>i</sub> -  $\exists_z(A[x \leftarrow z]) \implies \exists_x(A)$  ;(2) from E-Introduction on (1) WITH Hypothesis 1 ( $z$  is not free in  $\exists_x(A)$ )<sub>i</sub> -  $A[x \leftarrow z][z \leftarrow x] \implies \exists_z(A[x \leftarrow z])$  ;(3) from Substitution Axiom WITH Hypothesis 1 (the variables in  $x$  does not occur in  $A[x \leftarrow z]$  and the variables in  $z$  does not occur in  $A$ )<sub>i</sub>
  - $\exists_x(A) \implies \exists_z(A[x \leftarrow z])$  ;(4) from simplification and E-introduction on (3) WITH Hypothesis 1 ( $x$  is not free in  $\exists_z(A[x \leftarrow z])$ )<sub>i</sub>
  - $\exists_x(A) \equiv \exists_z(A[x \leftarrow z])$  ;(5) from tautological implication on (2, 4) and Post's Tautology Theorem<sub>i</sub>
  - Since the supply of variables are effectively infinite and Renaming states that a wff is provably equivalent to another wff with a dummy variable, then substitutability can be guaranteed by using dummy variables for variables in the substituted term
- =====

#### 4.18 (Definition) Theory extension

- The theory  $T'$  is an extension of the theory  $T$  iff  $T \subseteq T'$
  - The theory  $T'$  is aware of the theory  $T$  iff  $V \subseteq V'$
  - Given multiple theories  $T$  and  $T'$ ,  $\vdash_T A$  is an abbreviation for  $A \in T$  and  $\vdash_{T'} A$  is an abbreviation for  $A \in T'$
  - The extension  $T'$  of the theory  $T$  is conservative iff for any  $A \in L$ , if  $\vdash_{T'} A$ , then  $\vdash_T A$
- =====

#### 4.19 (Metatheorem) On constants

- Given that  $V' = V \cup \{[e_i]_{i=1}^n\}$  and  $\Gamma' = \Gamma, \vdash_{L'} A[[e_i]_{i=1}^n]$  iff  $\vdash_L A[[e_i]_{i=1}^n]$
- TODO: [ABSTRACTED] from dummy renaming =====

#### 4.20 (Metatheorem) Important corollary

- Given  $\{[e_i]_{i=1}^n\} \cap \Gamma = \emptyset$ , if  $\Gamma \vdash A[[e_i]_{i=1}^n]$ , then  $\Gamma \vdash [x_i]_{i=1}^n$  - TODO: [ABSTRACTED] from dummy renaming, try after deduction theorem =====

#### 4.21 (Metatheorem) Deduction theorem

- For any closed wff  $A$  and wff  $B$  and set of wffs  $\Gamma$ , if  $\Gamma \cup \{A\} \vdash B$ , then  $\Gamma \vdash A \implies B$
- The following is a proof by induction on  $\vdash$  ?? - Basis cases: - - When  $B = A$  - -  $A \implies B$  is tautological Q.E.D - - When  $B \neq A$  - -  $B < (1) \text{ must be } \Gamma \text{ axiomatic} > - - \vdash A \implies B$  (2) from (1) and Post's Tautology Theorem (3) from Hypothesis Strengthening on (2) and (1) - Inductive cases: - - Closed under the Modus Ponens Rule: let the inputs be  $\Gamma \cup \{A\} \vdash C$  and  $\Gamma \cup \{A\} \vdash C \implies B$ :
  - - - Inductive hypothesis:  $\Gamma \vdash A \implies C$  and  $\Gamma \vdash A \implies (C \implies B)$
  - - - Inductive step:
    - - - -  $\{A \implies C, A \implies (C \implies B)\} \vdash_{Taut} A \implies B$  (1) tautological implication
    - - - -  $\Gamma \vdash A \implies B$  (2) Post's Tautology Theorem on (1) and Transitivity of  $\vdash$
  - - Closed under the E-Introduction Rule: let the inputs be  $\Gamma \cup \{A\} \vdash (C \implies D)$  where  $x$  is not free in  $D$
  - - - Inductive hypothesis:  $\Gamma \vdash A \implies (C \implies D)$
  - - - Inductive step: - - - -  $A \implies (C \implies D) \vdash_{Taut} C \implies (A \implies D)$  (1) tautological implication
  - - - -  $\exists_x(C) \implies (A \implies D)$  (2) E-Introduction on (1) WITH Hypothesis ( $A$  is closed) and  $x$  is not free in  $D$
  - - - -  $\exists_x(C) \implies (A \implies D) \vdash_{Taut} A \implies (\exists_x(C) \implies D)$
  - - - -  $\Gamma \vdash A \implies (\exists_x(C) \implies D)$  (3) from Post's Tautology theorem
- The condition on  $A$  must be closed in important to enable one to preserve semantic meaning - - Suppose  $A$  is not closed like  $x \iff y$ , then  $x \iff y \vdash \forall_x(x \iff y)$  from Generalization, but this is a contradiction since there exists  $y$  where  $x \neq y$  =====

#### 4.22 (Metatheorem) Proof by contradiction

- For any closed wff  $A$ ,  $\Gamma \vdash A$  iff  $Thm_{\Gamma \cup \{\neg A\}} = Wff$
- When  $\Gamma \vdash A$ 
  - -  $\Gamma \cup \{\neg A\} \vdash A$  (1) Hypothesis Strengthening
  - -  $\Gamma \cup \{\neg A\} \vdash \neg A$  (2) nonlogical axiomed
  - -  $A, \neg A \vdash_{Taut} B < () \text{ tautological implication} >$
  - - When  $Thm_{\Gamma \cup \{\neg A\}} = Wff$
  - -  $\Gamma \cup \{\neg A\} \vdash A$  (3) hypothesis
  - -  $\Gamma \vdash \neg A \implies A$  (4) Deduction theorem
  - -  $\neg A \implies A \vdash_{Taut} A$  (5) tautological implication
  - -  $\Gamma \vdash A$  (6) Post's Tautology Theorem and Deduction theorem
- $A$  is closed necessary for Deduction theorem, add WITH in annotation - TODO =====

#### 4.23 (Metatheorem) Principle of Explosion

- TODO: same tautological implication gimmick =====

#### 4.24 (Metatheorem) Distributivity of exists and forall

- For any variable  $x$  and wff  $A$  and wff  $B$ ,  $A \implies B \vdash \exists_x(A) \implies \exists_x(B)$
- For any variable  $x$  and wff  $A$  and wff  $B$ ,  $A \implies B \vdash \forall_x(A) \implies \forall_x(B)$  =====

#### 4.25 (Metatheorem) Leibniz rule

- If  $A \equiv B$ , then  $C[A] \equiv C[B]$  or maybe  $C \equiv C'$  ?? =====

## 4.26 (Metatheorem) Proof by cases

- If  $\Gamma \vdash [A_i \vee]_{i=1}^n$  and ????

## 4.27 (Metatheorem) Proof by auxiliary constant

- ??? WIP lazy atm

# 5 Semantics

## 5.1 (Definition) Structure

- TODO: meaning ?? - The structure  $S$  appropriate for a language  $L$  is defined by  $S_L = (M, I)$
- The model  $M$  is defined by a non-empty set of concrete objects or the universe
- The interpretation  $I$  is a function defined by  $I : V + OTHERSTUFF??? \rightarrow M$  that satisfies the following:
- For any  $a \in Const$ ,  $I(a) \in M$
- For any  $f \in Func$ ,  $I(f) : M^{Arity(f)} \rightarrow M$  and  $I$  is total (defined everywhere)
- For any  $p \in Pred$ ,  $I(p) : M^{Arity(p)} \rightarrow M$  and  $I(p) \subseteq M^{Arity(p)}$

## 5.2 (Definition) Extension and restriction of a language

- The language  $L'$  is the expansion of the language  $L$  iff  $LS' = LS$  and  $NLS \subseteq NLS'$
- $L$  is the restriction of  $L'$

## 5.3 (Definition) Expansion and reduction of a structure

- The structure  $S'$  is an expansion of the structure  $S$  iff  $L'$  is an extension of  $L$  and for all  $v \in V$ ,  $I'(v) = I(v)$
- $S$  is the reduct of  $S'$
- The restriction/reduct symbol can be used as  $I = I' \upharpoonright L$  or  $S = S' \upharpoonright L$

## 5.4 (Definition) Model imported to the language

- TODO: meaning ?? - The imported language  $L(S)$  is defined by the language  $L$  and the appropriate structure  $S_L$
  - The imported alphabet  $V(S)$  or S-Alphabet is defined by  $V(S) = V \cup M$  - The imported set of all terms  $Term(S)$  or S-Term is defined by  $V(S)$
  - The imported set of all wffs  $Wff(S)$  or S-Wff is defined by  $Term(S)$  and  $V(S)$
  - This allows formal substitutions on model objects
  - Imported symbols are denoted by  $\bar{m}$  and  $I(\bar{m}) = I(m)$
- TODO  $I$  will now accept new shit

## 5.5 (Definition) Interpretation on closed S-terms

- TODO: use/meaning/intent - talkables - The interpretation  $I$  on closed S-terms is defined by the following:
- If the term is  $a \in Const$ , then  $I(a)$  and  $I(\bar{a}) = I(a)$
- If the term is  $f \in Func$  applied to the closed S-terms  $[t_i]_{i=1}^{Arity(f)}$ , then  $I(f([t_i]_{i=1}^{Arity(f)})) = I(f) \left( [I(t_i)]_{i=1}^{Arity(f)} \right)$