

Supplementary Material for: Methodological Flexibility in the Iowa Gambling Task

Table S2.

Interrater Reliability for Procedural Variables

Variable	Cohen's Kappa (κ)	SE (Bootstrap)	95% CI (low)	95% CI (high)
sample size	0.76	0.04	0.68	0.84
variant	0.88	0.06	0.76	1.00
protocol	0.55	0.08	0.40	0.71
goal	0.69	0.08	0.53	0.82
instructions	0.44	0.09	0.26	0.61
feedback	0.57	0.08	0.41	0.71
incentives	0.53	0.07	0.40	0.66
currency	0.75	0.05	0.65	0.86
starting Capital	0.80	0.06	0.69	0.91
reward per card (advantageous decks)	0.65	0.06	0.52	0.77
reward per card (disadvantageous decks)	0.63	0.06	0.51	0.75
net gain advantageous deck 1	0.70	0.06	0.57	0.80
net gain advantageous deck 2	0.71	0.06	0.58	0.82
net loss disadvantageous deck 1	0.69	0.06	0.57	0.81
net loss disadvantageous deck 2	0.68	0.06	0.55	0.79
loss interval advantageous deck 1	0.73	0.06	0.60	0.85
loss interval advantageous deck 2	0.73	0.07	0.60	0.86
loss interval disadvantageous deck 1	0.73	0.07	0.60	0.87

Variable	Cohen's Kappa (κ)	SE (Bootstrap)	95% CI (low)	95% CI (high)
loss interval disadvantageous deck 2	0.73	0.07	0.59	0.85
loss frequency advantageous deck 1	0.70	0.07	0.56	0.82
loss frequency advantageous deck 2	0.72	0.07	0.59	0.85
loss frequency disadvantageous deck 1	0.69	0.07	0.56	0.82
loss frequency disadvantageous deck 2	0.72	0.06	0.60	0.84
number of blocks	0.80	0.05	0.69	0.90
number of trials	0.90	0.06	0.76	1.00

Note. SE = standard error; CI = confidence interval. The standard error was calculated using a bootstrapping method with N = 1,000 iterations.