

Задача 5.2

In [121]:

```
import numpy as np
import math as mth
import scipy.optimize as opt
import scipy.stats as sps
import matplotlib.pyplot as plt
from sklearn.datasets import load_iris
%pylab inline
```

Populating the interactive namespace from numpy and matplotlib

Загрузка данных и разбиение на компоненты

In [122]:

```
data = load_iris()
comp = [[], [], []]
for i in np.arange(150):
    comp[data.target[i]].append(data.data[i])
```

Оценка матожиданий и матриц ковариаций

Я сразу написала функции для любых размерностей, тк это понадобится далее

In [141]:

```
def expect(sample, k):
    return np.array([sample[:,j].mean() for j in np.arange(k)])

a = [expect(np.array(comp[0]), 4), expect(np.array(comp[1]), 4), expect(np.
array(comp[2]), 4)]
```

In [142]:

```
def covar(sample, k):
    return np.array([([(sample[:,i] * sample[:,j]).mean() - sample[:,i].mean() * sample[:,j].mean())\
                        for j in np.arange(k)] for i in np.arange(k))).reshape(k,k)

sigma = [covar(np.array(comp[0]), 4), covar(np.array(comp[1]), 4), covar(np.array(comp[2]), 4)]
```

Значение вектора матожиданий и матрицы ковариаций по для всех компонент

1 компонента

Матожидание

In [143]:

```
print(a[0])
```

```
[ 5.006  3.418  1.464  0.244]
```

Матрица ковариаций

In [125]:

```
print(sigma[0])
```

```
[[ 0.121764  0.098292  0.015816  0.010336]
 [ 0.098292  0.142276  0.011448  0.011208]
 [ 0.015816  0.011448  0.029504  0.005584]
 [ 0.010336  0.011208  0.005584  0.011264]]
```

2 компонента

Матожидание

In [145]:

```
print(a[1])
```

```
[ 5.936  2.77  4.26  1.326]
```

Матрица ковариаций

In [126]:

```
print(sigma[1])
```

```
[[ 0.261104  0.08348  0.17924  0.054664]
 [ 0.08348  0.0965  0.081  0.04038 ]
 [ 0.17924  0.081  0.2164  0.07164 ]
 [ 0.054664  0.04038  0.07164  0.038324]]
```

3 компонента

Матожидание

In [146]:

```
print(a[2])
```

```
[ 6.588  2.974  5.552  2.026]
```

Матрица ковариаций

In [127]:

```
print(sigma[2])  
  
[[ 0.396256  0.091888  0.297224  0.048112]  
 [ 0.091888  0.101924  0.069952  0.046676]  
 [ 0.297224  0.069952  0.298496  0.047848]  
 [ 0.048112  0.046676  0.047848  0.073924]]
```


Построение графиков плотностей для различных проекций различных компонент смеси

In [175]:

```
def showdens(pair, com, x1=0, x2=9, y1=-1, y2=6):  
    sample = np.array([[comp[com-1][i][pair[0]-1], comp[com-1][i][pair[1]-1]] for i in np.arange(50)])  
    X,Y = np.mgrid[0:8:0.01, -1:8:0.01]  
    pos = np.empty(X.shape + (2,))  
    pos[:, :, 0] = X  
    pos[:, :, 1] = Y  
    plt.figure(figsize = (10, 10))  
    plt.contour(X, Y, sps.multivariate_normal.pdf(pos, mean=expect(sample, 2), cov=covar(sample, 2)), cmap = 'BrBG')  
    plt.title(r'Coordinates - '$' + str(pair) + r'$, Component # '$' + str(com) + r'$')  
    plt.scatter(sample[:, 0], sample[:, 1], alpha=0.2, s=40, label='sample')  
    plt.xlim(x1, x2)  
    plt.ylim(y1, y2)  
    plt.legend()  
    plt.xlabel(r'$x$')  
    plt.ylabel(r'$y$')  
    plt.show()
```


In [166]:

```
showdens((1, 2), 1, 4, 6.5, 2, 4.5)
```




In [168]:

```
showdens((1, 2), 2, 4.7, 7.3, 1.7, 3.7)
```




In [169]:

```
showdens((1, 2), 3, 4.3, 8.2, 2, 4)
```




In [170]:

```
showdens((2, 3), 1, 2, 5, 0.7, 2.1)
```




In [155]:

```
showdens((2, 3), 2, 1.5, 4, 2.5, 5.5)
```




In [176]:

```
showdens((2, 3), 3, 2, 4, 4.1, 7)
```




In [177]:

```
showdens((1, 3), 1, 4, 6.1, 0.8, 2.1)
```




In [178]:

```
showdens((1, 3), 2, 4.6, 8, 2.8, 5.3)
```



In [181]:

```
showdens((1, 3), 3, 4.8, 8.1, 4, 7.1)
```



Графики условной плотности

In [187]:

```
def print_cond_density(pair, i):
    K=[1,2,3]
    K.remove(i)

    X,Y = np.mgrid[0:8:0.01, -1:6:0.01]
    pos = np.empty(X.shape + (2,))
    pos[:, :, 0] = X
    pos[:, :, 1] = Y


    sample = [np.array([[x[pair[0]], x[pair[1]]] for x in data.data[50*(k-
1):50*k]]) for k in K ]
    exp=[expect(sample[0],2),expect(sample[1],2)]
    cov=[covar(sample[0],2),covar(sample[1],2)]
    density=(mlab.bivariate_normal(X, Y, sigmax=numpy.sqrt(cov[0][0][0]), s
igmay=numpy.sqrt(cov[0][1][1]),
                                mux=exp[0][0], muy=exp[0][1], sigmaxy=cov[0]
[0][1]) \
            + mlab.bivariate_normal(X, Y, sigmax=numpy.sqrt(cov[1][0][0]), sigm
ay=numpy.sqrt(cov[1][1][1]),
                                mux=exp[1][0], muy=exp[1][1], sigmaxy=cov
[1][0][1])) \
            / 2

    plt.figure(figsize = (10, 10))
    plt.contour(X,Y,density, cmap = 'spectral')
    plt.title(r'Coordinates - '$' + str(pair) + r'$, i - '$' + str(i) + r'$')
    plt.xlabel(r'$x$')
    plt.ylabel(r'$y$')
    plt.show()
```


In [188]:

```
for i in np.arange(1,4):  
    for pair in ((0,1), (1,3), (2,3)):  
        print_cond_density(pair,i)
```


Coordinates - $(0, 1)$, $i - 1$




Coordinates - (1, 3), i - 1




Coordinates - (2, 3), i - 1




Coordinates - $(0, 1)$, $i - 2$




Coordinates - $(1, 3)$, $i - 2$




Coordinates - $(2, 3)$, $i - 2$



Coordinates - $(0, 1)$, $i - 3$



Coordinates - $(1, 3)$, $i - 3$



Coordinates - $(2, 3)$, $i - 3$

