

# Lesson 26: Practice Problems

## Converting Higher-Order to First-Order Systems

### Part A: Basic Conversions (6 problems)

1. Convert to a first-order system:  $y'' + 4y = 0$
2. Convert with initial conditions:  $y'' - 3y' + 2y = 0$ ,  $y(0) = 1$ ,  $y'(0) = -1$
3. Write the companion matrix for:  $y''' + 2y'' - y' + 3y = 0$
4. Convert the fourth-order equation:  $y^{(4)} - y = 0$
5. Convert with forcing:  $y'' + \omega^2 y = \cos(\omega t)$
6. Convert the Airy equation:  $y'' - ty = 0$

### Part B: Companion Matrix Construction (5 problems)

7. Find the companion matrix for:  $y^{(5)} + y''' - 2y' = 0$
8. Given  $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -6 & -11 & -6 \end{bmatrix}$ , find the corresponding scalar ODE.
9. Construct the companion matrix for:  $2y''' + 3y'' - y' + 4y = 0$  (note the leading coefficient)
10. Find the companion matrix eigenvalues for:  $y'' + 2y' + 2y = 0$
11. Write the companion matrix for the Bessel equation:  $t^2 y'' + ty' + (t^2 - n^2)y = 0$

### Part C: Reverse Conversion (5 problems)

12. Convert the system  $\begin{cases} x' = y \\ y' = -4x - 4y \end{cases}$  to a single second-order equation.
13. Given  $\mathbf{x}' = \begin{bmatrix} 0 & 1 \\ -\omega^2 & -2\zeta\omega \end{bmatrix} \mathbf{x}$ , find the scalar equation.

14. Convert  $\begin{cases} x' = y \\ y' = z \\ z' = 6x - 11y + 6z \end{cases}$  to a third-order equation.
15. The system  $\mathbf{x}' = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \mathbf{x}$  corresponds to which scalar ODE?
16. Convert the coupled system  $\begin{cases} u' = v \\ v' = -u + 2v \end{cases}$  to find equations for both  $u$  and  $v$ .

## Part D: Initial Condition Conversion (5 problems)

17. Convert IVP:  $y''' + y' = 0$ ,  $y(0) = 1$ ,  $y'(0) = 0$ ,  $y''(0) = -1$
18. Transform:  $y'' + 3y' + 2y = e^{-t}$ ,  $y(0) = 0$ ,  $y'(0) = 1$
19. Convert the IVP:  $t^2 y'' - 2ty' + 2y = 0$ ,  $y(1) = 1$ ,  $y'(1) = 2$  (for  $t > 0$ )
20. Set up the system for:  $y^{(4)} + 4y'' + 4y = 0$  with  $y(0) = y'(0) = y''(0) = 0$ ,  $y'''(0) = 1$
21. Convert:  $y'' + (1 - \frac{1}{4t^2})y = 0$ ,  $y(\pi) = 0$ ,  $y'(\pi) = 1$

## Part E: Theoretical Problems (4 problems)

22. Prove that the characteristic polynomial of a companion matrix equals the characteristic equation of the original ODE.
23. Show that if  $y(t)$  solves  $y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_0y = 0$ , then  $\mathbf{x}(t) = [y, y', \dots, y^{(n-1)}]^T$  solves  $\mathbf{x}' = A\mathbf{x}$  where  $A$  is the companion matrix.
24. Prove that the companion matrix for a constant coefficient equation has  $n$  linearly independent eigenvectors if and only if all characteristic roots are distinct.
25. Show that the determinant of a companion matrix equals  $(-1)^{n+1}a_0$ .

## Part F: Exam-Style Problems (5 problems)

26. (Prof. Ditkowski style) Given  $y''' - 3y'' + 3y' - y = 0$ :
- Write the companion matrix
  - Find its eigenvalues
  - Explain the connection to the general solution

27. Convert the system and solve for eigenvalues:

$$y^{(4)} + 5y'' + 4y = 0$$

28. A mass-spring-damper system satisfies  $m\ddot{x} + c\dot{x} + kx = F(t)$ .

- (a) Convert to first-order form with  $m = 1$ ,  $c = 2$ ,  $k = 5$
- (b) Write the companion matrix
- (c) Find conditions on  $c$  for complex eigenvalues

29. Given the companion matrix  $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -\alpha & -\beta & -\gamma \end{bmatrix}$ :

- (a) Write the scalar ODE
- (b) Find  $\alpha, \beta, \gamma$  so that  $e^t, e^{2t}, e^{3t}$  are solutions
- (c) Verify using eigenvalues

30. (Comprehensive) The equation  $y''' + py' + qy = 0$  has solutions  $e^t, e^{-t}, e^{2t}$ .

- (a) Find  $p$  and  $q$
- (b) Write the companion matrix
- (c) Convert the IVP with  $y(0) = 1$ ,  $y'(0) = 0$ ,  $y''(0) = 3$
- (d) Express the solution using the state vector

## Solutions and Hints

### Selected Solutions:

**Problem 1:**  $\mathbf{x} = \begin{bmatrix} y \\ y' \end{bmatrix}$ ,  $A = \begin{bmatrix} 0 & 1 \\ -4 & 0 \end{bmatrix}$

**Problem 8:** The scalar ODE is  $y''' + 6y'' + 11y' + 6y = 0$

**Problem 13:**  $x'' + 4x' + 4x = 0$  (repeated roots at  $\lambda = -2$ )

**Problem 21:** Characteristic polynomial:  $\det(A - \lambda I) = (-\lambda)^{n-1}(-\lambda - a_{n-1}\lambda^{n-1} - \dots - a_0)$

**Problem 26:** Eigenvalues are 1, 1, 1 (triple root), indicating  $y = (c_1 + c_2t + c_3t^2)e^t$

### Key Insights:

- Always check dimensions when setting up state vectors
- The companion matrix structure is universal - memorize it!
- Initial conditions transform directly:  $\mathbf{x}(t_0) = [y(t_0), y'(t_0), \dots, y^{(n-1)}(t_0)]^T$  *Eigenvalues of companion matrix are roots of characteristic equation*