

Lesson 44: Practice Problems

Repeated Roots and t^k Terms

ODE 1 with Prof. Adi Ditkowski

Tel Aviv University

Part A: Identifying Multiplicities (5 problems)

1. Given the characteristic polynomial $(r-3)^2(r+1) = 0$, *identify all roots and their multiplicities.*
2. Given the characteristic polynomial $r^4 - 8r^3 + 24r^2 - 32r + 16 = 0$, *show that it equals $(r-2)^4$ and identify the multiplicity.*
3. The characteristic equation $(r^2 - 2r + 5)^2 = 0$ *has complex roots. Find them and their multiplicities.*
4. Factor completely: $r^3 + 3r^2 + 3r + 1 = 0$ *and identify multiplicities.*
5. Given $(r^2 + 4)^3 = 0$, *identify all roots and multiplicities.*

Part B: Solutions from Multiplicities (5 problems)

1. If $r = 3$ is a triple root, write all three linearly independent solutions.
2. If $r = -2$ has multiplicity 4, write the form of the general solution.
3. Complex roots $1 \pm 2i$ each have multiplicity 2. Write all four real solutions.
4. The characteristic polynomial is $(r + 1)^2(r-2)^3$. *Write the general solution.*
5. If $\pm 3i$ are both double roots, write the general real solution.

Part C: Complete ODEs with Repeated Roots (5 problems)

6. Solve: $y'' - 6y' + 9y = 0$
7. Solve: $y''' + 3y'' + 3y' + y = 0$
8. Solve: $y^{(4)} - 4y''' + 6y'' - 4y' + y = 0$

9. Solve: $y'' + 4y' + 4y = 0$ with $y(0) = 2, y'(0) = -3$
10. Solve: $y^{(4)} + 4y'' + 4y = 0$

Part D: Reduction of Order Applications (5 problems)

11. Given that $y_1 = e^{-t}$ solves $(D + 1)^2[y] = 0$, use reduction of order to find y_2 .
12. Verify that te^{3t} solves $y'' - 6y' + 9y = 0$ by direct substitution.
13. Show that if $y_1 = e^{rt}$ solves $L[y] = 0$ where r is a double root, then $y_2 = te^{rt}$ also solves it.
14. Given $y'' - 4y' + 4y = 0$ has solution e^{2t} , find the second solution using $y_2 = v(t)e^{2t}$.
15. For the equation with characteristic polynomial $(r-a)^3 = 0$, verify the Wronskian of $e^{at}, te^{at}, t^2e^{at}$ is nonzero.

Part E: Complex Repeated Roots (5 problems)

16. Solve: $y^{(4)} + 8y'' + 16y = 0$
17. Find the general real solution when the characteristic polynomial is $(r^2 + 2r + 2)^2 = 0$.
18. Solve: $y^{(4)} + 4y''' + 8y'' + 8y' + 4y = 0$ (Hint: $(r^2 + 2r + 2)^2$)
19. If $2 \pm 3i$ are both triple roots, how many linearly independent real solutions are there? List them.
20. Solve the IVP: $y'' + 2y' + 5y = 0$ with $y(0) = 0, y'(0) = 4$, given that $-1 \pm 2i$ are roots.

Part F: Theoretical and Exam-Style Problems (5 problems)

21. Prove that if r_0 is a root of multiplicity m of the characteristic polynomial, then $p(r_0) = p'(r_0) = \dots = p^{(m-1)}(r_0) = 0$ but $p^{(m)}(r_0) \neq 0$.
22. Show that the dimension of the solution space equals the degree of the differential equation, even with repeated roots.
23. A fourth-order equation has characteristic polynomial with roots: $r = 2$ (mult. 2), $r = -1$ (mult. 1), $r = 3$ (mult. 1). If a solution satisfies $y(0) = 1, y'(0) = 0, y''(0) = -1, y'''(0) = 2$, find it.
24. Explain why the functions $\{t^k e^{rt} : k = 0, 1, \dots, m-1\}$ span the solution space for a root of multiplicity m .

25. (Prof. Ditkowski style) Consider the family of equations $y'' - 2ay' + a^2y = 0$ parameterized by $a \in \mathbb{R}$.
- (a) Show that for all a , the characteristic equation has a repeated root.
 - (b) Find the general solution in terms of a .
 - (c) For which values of a do all solutions remain bounded as $t \rightarrow \infty$?
 - (d) Find the solution with $y(0) = 1, y'(0) = 0$.

Solutions

Part A: Identifying Multiplicities

1. Roots: $r = 3$ (multiplicity 2), $r = -1$ (multiplicity 1)
2. Expand $(r - 2)^4 = r^4 - 8r^3 + 24r^2 - 32r + 16$
Root :r = 2(multiplicity4)
3. $(r^2 - 2r + 5)^2 = 0 \Rightarrow r^2 - 2r + 5 = 0$
 $r = 2 \pm \sqrt{4 - 20} = 2 \pm 2\sqrt{-1} = 2 \pm 2i$
Roots: $1 + 2i$ (mult. 2), $1 - 2i$ (mult. 2)
4. $r^3 + 3r^2 + 3r + 1 = (r+1)^3 = 0$
Root :r = -1(multiplicity3)
5. $(r^2 + 4)^3 = 0 \Rightarrow r^2 = -4 \Rightarrow r = \pm 2i$
Roots: $2i$ (mult. 3), $-2i$ (mult. 3)

Part B: Solutions from Multiplicities

6. $e^{3t}, te^{3t}, t^2e^{3t}$
7. $y(t) = (c_1 + c_2t + c_3t^2 + c_4t^3)e^{-2t}$
8. Real solutions: $e^t \cos(2t), te^t \cos(2t), e^t \sin(2t), te^t \sin(2t)$
9. $y(t) = (c_1 + c_2t)e^{-t} + (c_3 + c_4t + c_5t^2)e^{2t}$
10. $y(t) = (c_1 + c_2t)\cos(3t) + (c_3 + c_4t)\sin(3t)$

Part C: Complete ODEs with Repeated Roots

11. Char. eq.: $r^2 - 6r + 9 = (r-3)^2 = 0$
Root :r = 3(mult.2)
Generalsolution :y(t) = (c_1 + c_2t)e^{3t}
12. Char. eq.: $r^3 + 3r^2 + 3r + 1 = (r+1)^3 = 0$
Root :r = -1(mult.3)
Generalsolution :y(t) = (c_1 + c_2t + c_3t^2)e^{-t}
13. Char. eq.: $(r - 1)^4 = 0$
Root :r = 1(mult.4)
Generalsolution :y(t) = (c_1 + c_2t + c_3t^2 + c_4t^3)e^t
14. $(r + 2)^2 = 0$, so $y = (c_1 + c_2t)e^{-2t}$
 $y(0) = c_1 = 2$
 $y'(t) = c_2e^{-2t} - 2(c_1 + c_2t)e^{-2t}$
 $y'(0) = c_2 - 2c_1 = -3$

$$c_2 = 1$$

$$\text{Solution : } y(t) = (2 + t)e^{-2t}$$

$$15. r^4 + 4r^2 + 4 = (r^2 + 2)^2 = 0$$

$$\text{Roots : } \pm i\sqrt{2} \text{ (each mult. 2)}$$

$$y(t) = (c_1 + c_{2t})\cos(\sqrt{2}t) + (c_3 + c_{4t})\sin(\sqrt{2}t)$$

Part D: Reduction of Order Applications

$$16. \text{ Let } y_2 = v(t)e^{-t}$$

$$\text{Substituting into } (D + 1)^2[y] = y'' + 2y' + y = 0:$$

$$\text{Aftersimplification : } v'' = 0$$

$$\text{Thus } v = c_{1t} + c_2$$

$$\text{Secondsolution : } y_2 = te^{-t}$$

$$17. y = te^{3t}$$

$$y' = e^{3t} + 3te^{3t} = (1 + 3t)e^{3t}$$

$$y'' = 3e^{3t} + 3(1 + 3t)e^{3t} = (6 + 9t)e^{3t}$$

$$\text{Substitute: } (6 + 9t)e^{3t} - 6(1 + 3t)e^{3t} + 9te^{3t}$$

$$= e^{3t}[6 + 9t - 6 - 18t + 9t] = 0 \checkmark$$

$$18. \text{ For double root } r: (D - r)^2[y] = 0$$

$$\text{If } y = te^{rt}: y' = e^{rt} + rte^{rt}$$

$$y'' = 2re^{rt} + r^2te^{rt}$$

$$(D - r)^2[y] = D[(D - r)[te^{rt}]] = D[e^{rt}] = re^{rt} - re^{rt} = 0 \checkmark$$

$$19. y_2 = v(t)e^{2t}$$

$$y_2' = (v' + 2v)e^{2t}$$

$$y_2'' = (v'' + 4v' + 4v)e^{2t}$$

$$\text{Substituting: } e^{2t}[v'' + 4v' + 4v - 4v' - 8v + 4v] = e^{2t} \cdot v'' = 0$$

$$v'' = 0 \Rightarrow v = c_{1t} + c_2$$

$$\text{Secondsolution : } y_2 = te^{2t}$$

$$20. W = \begin{vmatrix} e^{at} & te^{at} & t^2e^{at} \\ ae^{at} & (1 + at)e^{at} & (2t + at^2)e^{at} \\ a^2e^{at} & (2a + a^2t)e^{at} & (2 + 4at + a^2t^2)e^{at} \end{vmatrix}$$

$$\text{After calculation: } W = 2e^{3at} \neq 0$$

Part E: Complex Repeated Roots

$$21. (r^2 + 4)^2 = 0 \Rightarrow r = \pm 2i \text{ (each mult. 2)}$$

$$y(t) = (c_1 + c_{2t})\cos(2t) + (c_3 + c_{4t})\sin(2t)$$

$$22. r^2 + 2r + 2 = 0 \Rightarrow r = -1 \pm i$$

$$\text{Each root has mult. 2}$$

$$y(t) = e^{-t}[(c_1 + c_{2t})\cos(t) + (c_3 + c_{4t})\sin(t)]$$

23. Roots: $-1 \pm i$ (each mult. 2)
 $y(t) = e^{-t}[(c_1 + c_{2t})\cos(t) + (c_3 + c_{4t})\sin(t)]$
24. 6 linearly independent real solutions:
 $e^{2t} \cos(3t), te^{2t} \cos(3t), t^2 e^{2t} \cos(3t)$
 $e^{2t} \sin(3t), te^{2t} \sin(3t), t^2 e^{2t} \sin(3t)$
25. Simple roots (not repeated): $r = -1 \pm 2i$
 $y(t) = e^{-t}[c_1 \cos(2t) + c_2 \sin(2t)]$
 $y(0) = c_1 = 0$
 $y'(t) = e^{-t}[-c_1 \cos(2t) - c_2 \sin(2t) - 2c_1 \sin(2t) + 2c_2 \cos(2t)]$
 $y'(0) = -c_1 + 2c_2 = 4$
 $c_2 = 2$
Solution : $y(t) = 2e^{-t} \sin(2t)$

Part F: Theoretical and Exam-Style Problems

26. If $p(r) = (r - r_0)^m q(r)$ where $q(r_0) \neq 0$:
 $p'(r) = m(r - r_0)^{m-1} q(r) + (r - r_0)^m q'(r)$
 $\text{At } r = r_0: p'(r_0) = 0 \text{ if } m \geq 1$
Continued if differentiating to show $p^{(k)}(r_0) = 0 \text{ for } k \leq m$
But $p^{(m)}(r_0) = m! \cdot q(r_0) \neq 0$
27. Total number of solutions = sum of all multiplicities = degree of polynomial = order of ODE
28. Char. poly: $(r - 2)^2(r + 1)(r - 3) = 0$
 $y = (c_1 + c_2 t)e^{2t} + c_3 e^{-t} + c_4 e^{3t}$
 Apply initial conditions (solve 4×4 system) :
Final solution : $y(t) = (1 - 2\frac{3t}{3t}e^{2t} + \frac{1}{4}e^{-t} - \frac{1}{4}e^{3t})$
29. The operator $(D - r)^m$ has an m -dimensional kernel. The functions $t^k e^{rt}$ for $k = 0, \dots, m - 1$ are linearly independent (nonzero Wronskian) and all satisfy $(D - r)^m[y] = 0$.
30. (a) Char. eq.: $r^2 - 2ar + a^2 = (r - a)^2 = 0$
Always a double root at $r = a$
- (b) $y(t) = (c_1 + c_2 t)e^{at}$
- (c) Bounded as $t \rightarrow \infty$ only if $a \leq 0$
- (d) $y(0) = c_1 = 1$
 $y'(t) = c_2 e^{at} + a(c_1 + c_2 t)e^{at}$
 $y'(0) = c_2 + ac_1 = 0$
 $c_2 = -a$
Solution : $y(t) = (1 - at)e^{at}$