

ODE Lesson 37: 2D Linear Systems: Complete Classification

ODE 1 - Prof. Adi Ditkowski

1 Classification Framework

Consider the 2D linear system:

$$\dot{\mathbf{x}} = A\mathbf{x}, \quad \text{where } A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Master Classification Parameters:

$$\text{Trace: } \tau = \text{tr}(A) = a + d \quad (1)$$

$$\text{Determinant: } \Delta = \det(A) = ad - bc \quad (2)$$

$$\text{Discriminant: } D = \tau^2 - 4\Delta \quad (3)$$

These three quantities completely determine the phase portrait!

2 Eigenvalue Analysis

The characteristic equation is:

$$\lambda^2 - \tau\lambda + \Delta = 0$$

with eigenvalues:

$$\lambda_{1,2} = \frac{\tau \pm \sqrt{\tau^2 - 4\Delta}}{2} = \frac{\tau \pm \sqrt{D}}{2}$$

Theorem 1 (Eigenvalue Classification). *The nature of eigenvalues depends on the discriminant D :*

- $D > 0$: Two distinct real eigenvalues
- $D = 0$: One repeated real eigenvalue
- $D < 0$: Two complex conjugate eigenvalues $\alpha \pm i\beta$

3 Complete Classification Table

Type	Condition	Eigenvalues	Stability	Behavior
Saddle	$\Delta < 0$	$\lambda_1 < 0 < \lambda_2$	Unstable	Hyperbolic
Stable Node	$\Delta > 0, D > 0, \tau < 0$	$\lambda_1, \lambda_2 < 0$	Asymp. Stable	Approach
Unstable Node	$\Delta > 0, D > 0, \tau > 0$	$\lambda_1, \lambda_2 > 0$	Unstable	Repelling
Stable Spiral	$\Delta > 0, D < 0, \tau < 0$	$\alpha \pm i\beta, \alpha < 0$	Asymp. Stable	Spiral in
Unstable Spiral	$\Delta > 0, D < 0, \tau > 0$	$\alpha \pm i\beta, \alpha > 0$	Unstable	Spiral out
Center	$\Delta > 0, \tau = 0$	$\pm i\beta$	Neutrally Stable	Ellipses

4 Detailed Portrait Descriptions

4.1 Saddle Point

Classification 1 (Saddle). **Condition:** $\det(A) < 0$

Eigenvalues: Real with opposite signs

Key Features:

- Stable manifold: $E^s = \text{span}\{v_1\}$ where $\lambda_1 < 0$
- Unstable manifold: $E^u = \text{span}\{v_2\}$ where $\lambda_2 > 0$
- Trajectories approach along E^s , depart along E^u Four hyperbolic sectors

• Drawing a Saddle:

1. Draw stable eigenvector (arrows pointing in)
2. Draw unstable eigenvector (arrows pointing out)
3. Add hyperbolic trajectories in each sector
4. These are the separatrices - label them!

4.2 Nodes

Classification 2 (Proper Node). **Condition:** $\Delta > 0, D > 0, \lambda_1 \neq \lambda_2$

Behavior:

- All trajectories tangent to slow eigenvector at origin
- Fast eigenvector: $|\lambda_1| > |\lambda_2|$ determines initial direction
- Slow eigenvector: determines final approach direction

Classification 3 (Improper Node). **Condition:** $D = 0$, single eigenvector

Special Case: $A = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$ (Jordan form)

Behavior: Trajectories curve, all tangent to eigenvector

Classification 4 (Star Node). **Condition:** $A = \lambda I$ (scalar multiple of identity)
Behavior: Straight-line trajectories in all directions

4.3 Spirals and Centers

Classification 5 (Spiral/Focus). **Condition:** $D < 0$ (complex eigenvalues)
Eigenvalues: $\lambda = \alpha \pm i\beta$
Behavior:

- Rotation frequency: $\omega = \beta$
- Decay/growth rate: α
- Clockwise if $\det(A) > 0$ and $b < 0$ (or $c > 0$)
- Counterclockwise if $\det(A) > 0$ and $b > 0$ (or $c < 0$)

Quick Spiral Direction: Look at the off-diagonal elements of A :


- Upper-right positive ($b > 0$) \rightarrow counterclockwise
 Upper-right negative ($b < 0$) \rightarrow clockwise

Classification 6 (Center). **Condition:** $\tau = 0$, $\Delta > 0$
Eigenvalues: Pure imaginary $\pm i\beta$
Behavior: Closed elliptical orbits, period $T = 2\pi/\beta$

5 Degenerate Cases

- When $\det(A) = 0$:
 - One zero eigenvalue: Line of equilibria
 - Two zero eigenvalues: Every point is an equilibrium
 - Parallel trajectories if one eigenvalue is nonzero

6 The Trace-Determinant Diagram



7 Classification Algorithm

Prof. Ditkowski's Quick Classification:

1. Compute $\det(A)$
 - If $\det(A) < 0 \rightarrow$ **SADDLE**(done!) If $\det(A) = 0 \rightarrow$ *Degenerate case*
 - If $\det(A) > 0 \rightarrow$ *Continue...*
2. Compute $\tau = \text{tr}(A)$
 - If $\tau = 0 \rightarrow$ **CENTER** If $\tau \neq 0 \rightarrow$ *Continue...*
3. Compute $D = \tau^2 - 4\det(A)$
 - If $D > 0 \rightarrow$ **NODE**(stable if $\tau < 0$)
 - If $D < 0 \rightarrow$ **SPIRAL**(stable if $\tau < 0$)
 - If $D = 0 \rightarrow$ **IMPROPER/STAR NODE**

8 Examples Gallery

Example 1 (Complete Classification). Classify $A = \begin{pmatrix} 2 & 3 \\ 3 & 2 \end{pmatrix}$:

$$\det(A) = 4 - 9 = -5 < 0 \quad (4)$$

Therefore: **SADDLE** (unstable)

Verification: $\lambda = \frac{4 \pm \sqrt{16+20}}{2} = \frac{4 \pm 6}{2} = 5, -1 \checkmark$

Example 2 (Using Trace-Det Only). *Classify* $A = \begin{pmatrix} -3 & 2 \\ -2 & -3 \end{pmatrix}$:

$$\tau = -6 < 0 \quad (\text{stable}) \tag{5}$$

$$\Delta = 9 + 4 = 13 > 0 \quad (\text{not a saddle}) \tag{6}$$

$$D = 36 - 52 = -16 < 0 \quad (\text{complex eigenvalues}) \tag{7}$$

Therefore: **STABLE SPIRAL**