

SAMPLE-1 trial results: a synthetic study drug versus placebo in adults with type 2 diabetes.

Table 2. Baseline characteristics of the study population

Characteristic	Placebo (n=120)	Study drug (n=118)	P value
Age, years, mean (SD)	61.2 (9.8)	60.7 (10.1)	0.71
Male sex, n (%)	72 (60.0)	69 (58.5)	0.81
Hypertension, n (%) ^a	80 (66.7)	71 (60.2)	0.30
Diabetes duration, y, median (IQR)	8 (5-12)	9 (5-13)	0.55
Prior MI, n/N (%)	11/120 (9.2)	14/118 (11.9)	0.52
HbA1c, %	7.9	8.0	0.64

^a Systolic blood pressure of at least 140 mmHg or treatment with an antihypertensive agent.

Table 3. Primary and secondary outcomes at week 24

Outcome	Study drug	Placebo	Hazard ratio (95% CI)
Primary composite, n/N (%)	22/118 (18.6)	38/120 (31.7)	0.52 (0.31–0.88)
Death from any cause, n/N (%)	6/118 (5.1)	9/120 (7.5)	0.66 (0.23–1.86)
Hospitalisation, n/N (%)	15/118 (12.7)	27/120 (22.5)	0.50 (0.27–0.94)