■■■■■■■■■■

■■■■

SGLT2 inhibitors versus placebo for chronic kidney disease progression in adults

■■■■

■■■■■■■■ evidence_gap■■■■ blocked■■■■■■■■■■■ Meta ■■■■■■■■■■■■■■■■■■■■■■■■■■■

PICO

- ■■■Adults (age ≥18 years) with chronic kidney disease of any stage, with or without diabetes mellitus - ■■■Sodium-glucose co-transporter-2 (SGLT2) inhibitors (e.g., canagliflozin, dapagliflozin, empagliflozin, ertugliflozin, sotagliflozin) at any dose, as monotherapy or as part of combination therapy - ■■■Placebo, no pharmacological treatment, lifestyle intervention alone, active pharmacological comparator (other glucose-lowering drugs or other CKD treatments), or standard of care without an SGLT2 inhibitor - ■■■■■Composite of kidney disease progression: sustained decline in estimated glomerular filtration rate (eGFR) of ≥50%, progression to end-stage kidney disease (requiring dialysis or transplantation), or death due to kidney disease, as defined by the trial authors

■■■■■■■■

- ■■/■■■PubMed, curated literature index, OpenAlex, ClinicalTrials.gov - ■■■■■576 - ■■■■■■20 - ■■/■■■■■20 - ■■■■■3 - ■■■■■■/■■■2 - ■■■■■■■■■2

■■■■■■■■■■■■■

■■■■■■■■ 2 ■■■■■RR 0.7992881557729065 (95% CI 0.7272984471350266 ■ 0.8784035748673147)■■■■■■■■■■■■■■■■■■■■■■■■■

■■■■

- primary_counts_not_source_verified: Primary-effect row 39453837:1 does not source-verify all arm-level event/total counts used for pooling: total_intervention=3304, total_control=3305.

■■■■■

- limited_text_sources_present: 2 retrieved/screened record(s) use limited source text (2 abstract-only).

■■■■■■■■

■	■■	■■	■■	■■■■	■■■

Retrieval and Processing Notes

1 run-level warning(s) were recorded during retrieval, parsing, analysis, or output generation. Review the project run-warning log before external use. - retrieval/clinicaltrials_fallback_failed: ClinicalTrials.gov fallback had 1 failed request(s); registry-first trials may be missing.

■■■■■■■■

1. ■■■■■■■■■■■■ abstract-only ■■■■ PDF■ 2. ■ extraction review ■■■■■■■■■■■■■■■■■■■■■■■■■■■■ 3. ■■■■■■■■■■■■■■■■■■■■■■■■■■■■ narrative-only■ 4. ■■ blockers ■■■■■■■■■■ manuscript■

Evidence readiness warning: This run is classified as evidence_gap and is not cleared for quantitative synthesis because unresolved evidence blockers remain (primary_counts_not_source_verified). See manuscript_facts.json, extraction/extraction_audit.json, and analysis/effect_selection_audit.json before external use.