

INTERPOLATION OF POWER MAPPINGS

JACK BURKART AND KIRILL LAZEBNIK

ABSTRACT. Let $(M_j)_{j=1}^\infty \in \mathbb{N}$ and $(r_j)_{j=1}^\infty \in \mathbb{R}^+$ be increasing sequences satisfying some mild rate of growth conditions. We prove that there is an entire function $f : \mathbb{C} \rightarrow \mathbb{C}$ whose behavior in the large annuli $\{z \in \mathbb{C} : r_j \cdot \exp(\pi/M_j) \leq |z| \leq r_{j+1}\}$ is given by a perturbed rescaling of $z \mapsto z^{M_j}$, such that the only singular values of f are rescalings of $\pm r_j^{M_j}$. We describe several applications to the dynamics of entire functions.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Interpolation of Power Maps	6
4. Entire Functions	14
5. A Dynamical Application	17
References	20

1. INTRODUCTION

The problem of constructing analytic functions with prescribed geometry arises in areas across function theory. One general approach consists of defining a convergent infinite product, which has the advantage, for instance, of yielding an explicit expression for the zeros of the function. We mention several dynamical applications of this approach: the first construction of an entire function with a wandering domain [Bak76], constructions of entire functions of slow growth with Julia set $= \mathbb{C}$ [BE00], in the study of escaping sets of entire functions [RS13], the construction of transcendental entire functions with 1-dimensional Julia sets [Bis18], and in the dynamics of sine maps [ERS20].

A general principle used in the above works is that the behavior of an infinite product is dominated within certain regions by certain factors in the product. Our main result similarly proves the existence of an entire function whose behavior is dominated within certain annuli by certain prescribable monomials. One advantage of our approach, however, is a precise description of the singular values of the entire function. This information is in general difficult to glean from an infinite product, and is often crucial in applications. Before describing further motivation and applications, we state our main result. First we will need

the following definition, which will serve as an assumption on the growth rate of the degree of the aforementioned monomials.

Definition 1.1. Let $(M_j)_{j=1}^\infty \in \mathbb{N}$ be increasing, and $(r_j)_{j=1}^\infty \in \mathbb{R}^+$. We say that $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are *permissible* if

$$(1.1) \quad r_{j+1} \geq \exp(\pi/M_j) \cdot r_j \text{ for all } j \in \mathbb{N}, r_j \xrightarrow{j \rightarrow \infty} \infty, \text{ and } \sup_j \frac{M_{j+1}}{M_j} < \infty.$$

With Definition 1.1, we can state our main result:

Theorem A. Let $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ be permissible, $r_0 := 0$ and $c \in \mathbb{C}^* := \mathbb{C} \setminus \{0\}$. Set

$$(1.2) \quad c_1 := c, \text{ and } c_j := c_{j-1} \cdot r_{j-1}^{M_{j-1}-M_j} \text{ for } j \geq 2.$$

Then there exists an entire function $f : \mathbb{C} \rightarrow \mathbb{C}$ and a homeomorphism $\phi : \mathbb{C} \rightarrow \mathbb{C}$ such that

$$(1.3) \quad f \circ \phi(z) = c_j z^{M_j} \text{ for } r_{j-1} \cdot \exp(\pi/M_{j-1}) \leq |z| \leq r_j, j \in \mathbb{N}.$$

Moreover, if $\sum_{j=1}^\infty M_j^{-1} < \infty$, then $|\phi(z)/z - 1| \rightarrow 0$ as $z \rightarrow \infty$. The only singular values of f are the critical values $(\pm c_j r_j^{M_j})_{j=1}^\infty$.

Remark 1.2. The homeomorphism ϕ is a K -quasiconformal homeomorphism (see Definition 2.1), where K depends only on $\sup_j (M_{j+1}/M_j)$. The conclusion $|\phi(z)/z - 1| \rightarrow 0$ is deduced from the Teichmüller-Wittich-Belinskii Theorem (see Theorem 2.6), and in many applications, we can even deduce uniform estimates $\|\phi(z)/z - 1\|_{L^\infty(\mathbb{C})} < \varepsilon$ (see Section 5). We further remark that a precise description of the critical points and zeros of f are also given (see Proposition 4.9 and Remark 4.10), up to the perturbation ϕ . The “scaling” constants $(c_j)_{j=1}^\infty$ ensure that $z \mapsto c_j z^{M_j}$ and $z \mapsto c_{j+1} z^{M_{j+1}}$ both map $|z| = r_j$ to the same scale. Lastly, we comment that if we replace the assumption $r_j \rightarrow \infty$ of Definition 1.1 to $r_j \rightarrow r_\infty < \infty$, we obtain a result similar to Theorem A, but with the domain of f equal to $r_\infty \cdot \mathbb{D}$ rather than \mathbb{C} (see Theorem B in Section 4).

As indicated in Remark 1.2, our methods rely on *quasiconformal surgery*, a large collection of techniques to which we refer to [FH09] for a survey. Among these techniques, there are at least two distinct approaches both termed *quasiconformal surgery*. The first consists in the construction of a quasiregular function g possessing a g -invariant Beltrami coefficient μ . The integrating map for μ then conjugates g to a holomorphic function. This approach appears first in [Sul85] (to the best of the authors’ knowledge), and is the most common use of the Measurable Riemann Mapping Theorem in complex dynamics, as it is inherently dynamical.

A different approach also termed *quasiconformal surgery* consists of constructing a quasiregular function g which in turn yields a holomorphic function $g \circ \phi^{-1}$, where ϕ is the integrating map for the Beltrami coefficient $g_{\bar{z}}/g_z$. This is the approach used in the present work, and has long found fundamental application in the type problem and value distribution theory (see, for instance, [Dra86], [Ere86]), and, surprisingly, has found recent application in complex




FIGURE 1. Illustrated is an interpolation, defined on $1 \leq |z| \leq 2$, between the identity on \mathbb{T} and the angle-doubling map on $2\mathbb{T}$. In (A), the white (respectively, black) vertices on the outer circle represent preimages under -2 (respectively, $+2$) of the angle doubling map, and on the inner circle we represent -1 (respectively, $+1$) by a white (respectively, black) vertex.

dynamics despite the lack of a conjugacy between g and $g \circ \phi^{-1}$. Indeed, this approach was used in [Bis15] in settling a long-standing question about the existence of wandering domains for functions with bounded singular set (see also [FGJ15], [FJL19], [Laz19], [MPS20]).

A general difficulty in this approach lies in proving that ϕ is only a small perturbation of the identity, which may be deduced, for instance, from showing that g is holomorphic (and hence $g_{\bar{z}}/g_z = 0$) except on a very small set. To this end, the work [Bis15] introduced a technique termed *quasiconformal folding*, which has found many recent applications in complex dynamics and in function theory more broadly (see, for instance, [Bis14], [BP15], [OS16], [BRS17], [Laz17], [BL19], [AB20]). Our main methods of proof for Theorem A draw on this technique, but we emphasize that the present work does not rely directly on the results of [Bis15], and indeed may be read and understood independently of the aforementioned works.

To conclude the Introduction, we remark on several applications of Theorem A and present a sketch of the proof. In the present manuscript, we briefly present in Section 5 how Theorem A yields a robust approach to the construction of entire functions with multiply connected wandering domains (first constructed in [Bak76] - see also [Bak85], [Hin94], [KS08], [BZ11], [Ber11], [BRS13]). In a companion manuscript [BL], we show how Theorem A gives a different approach to the result of [Bis18] on existence of transcendental entire functions with 1-dimensional Julia set (see also [Bur19]).

The proof of Theorem A consists of defining a quasiregular function g by $z \mapsto c_j z^{M_j}$ in the large annuli $r_{j-1} \cdot \exp(\pi/M_{j-1}) \leq |z| \leq r_j$, and finding an efficient interpolation between these power mappings in the thin annuli $r_j \leq |z| \leq r_j \cdot \exp(\pi/M_j)$. We sketch the main idea in the simple case of interpolation in $1 \leq |z| \leq 2$ between $z \mapsto z$ on $\mathbb{T} := \{z : |z| = 1\}$ and $z \mapsto z^2$ on $2\mathbb{T}$. We will refer to Figure 1. It will suffice to interpolate between the identity




FIGURE 2. Illustrated is the main interpolation used in Theorem A in a slightly more elaborate case than in Figure 1: interpolation between $z \mapsto z^2$ and $z \mapsto z^6$. The map $(B) \mapsto (C)$ is given by $\theta \rightarrow 6\theta$.

on \mathbb{T} and the angle doubling map on $2\mathbb{T}$. The map $(A) \mapsto (B)$ is the identity on $2\mathbb{T}$ and piecewise-linear in logarithmic coordinates, except on the red segment where it is 2-valued as illustrated (the inverse $(B) \mapsto (A)$ is called the folding map in [Bis15]). The map $(B) \mapsto (C)$ is given by angle-doubling. The composition $(A) \mapsto (C)$ is still 2-valued on the red segment, whence the map $(C) \mapsto (D)$ identifies these 2 values on the real-axis. Although useful for a sketch of the proof, we remark that we do not actually use a separate map $(C) \mapsto (D)$ to identify these 2 values, but rather this procedure is built into the definition of $(B) \mapsto (C)$ (see Definition 3.11).

The more general case of interpolation between $(z \mapsto z^n)|_{\mathbb{T}}$ and $(z \mapsto z^m)|_{r\mathbb{T}}$, $r > 1$, for general $n < m$ is similar (see Figure 2): we attach radial segments (illustrated in red) to each preimage of -1 under $z \mapsto z^n$ on \mathbb{T} which play a similar role as above, and use in general a more elaborate placement of vertices along these radial segments. A careful analysis shows that the desired interpolation can be achieved in an annulus of modulus $1/2n$, with distortion constants depending only on the ratio m/n and not otherwise on n, m . This essentially yields the first conclusion in Theorem A, and the statement about critical values follows by an inspection of which points the above quasiregular function branches at.

We briefly outline the paper. After collecting a couple preliminary results we will need in Section 2, we detail the specifics of the main interpolation in Section 3 where the primary contributions of the present work are contained. Section 4 applies the results of Section 3 to build the entire function of Theorem A. In Section 5, we consider dynamical applications of Theorem A.

2. PRELIMINARIES

Definition 2.1. An orientation-preserving homeomorphism $\phi : \mathbb{C} \rightarrow \mathbb{C}$ is said to be a *quasiconformal mapping* if $\phi \in W_{\text{loc}}^{1,2}(\mathbb{C})$ and $\|\phi_{\bar{z}}/\phi_z\|_{L^\infty(\mathbb{C})} \leq k$ for some $0 \leq k < 1$.

Remark 2.2. We refer the reader to [LV73] for a detailed study of quasiconformal mappings. We will assume a familiarity with the basic theory in what follows. We remark that a *quasiregular* mapping is one which may be represented by a composition $f \circ \phi$, where f is holomorphic, and ϕ is quasiconformal (see [LV73], Chapter VI).

Notation 2.3. We abbreviate piecewise linear by PWL. Given a quasiregular map f , we will denote the dilatation constant of f by $K(f)$, where $1 \leq K(f) < \infty$. We denote $k(f) := (K(f) - 1)/(K(f) + 1)$, and occasionally we will use the notation $A(r_1, r_2) := \{z \in \mathbb{C} : r_1 < |z| < r_2\}$.

We record a Theorem due to Teichmüller, Wittich, and Belinskii (Theorem 2.6 below). As already mentioned in the Introduction, we will use this result to deduce the conclusion $|\phi(z)/z - 1| \rightarrow 0$ of Theorem A. The statement of the result is taken from Theorem 6.1 of [LV73], to which we refer for the relevant bibliography. We note that in Theorem 2.6, $dA(z)$ refers to area measure. Before stating the result, we recall two definitions which will appear in the Theorem.

Definition 2.4. Let $\psi : \mathbb{C} \rightarrow \mathbb{C}$ be a quasiconformal mapping. The *dilatation quotient* of ψ is defined by

$$(2.1) \quad D(z) := \frac{|\psi_z(z)| + |\psi_{\bar{z}}(z)|}{|\psi_z(z)| - |\psi_{\bar{z}}(z)|}.$$

The quantity $D(z)$ is defined for a.e. z for a quasiconformal mapping ψ (see [LV73], Section IV.1.5).

Definition 2.5. Let ψ be a quasiconformal mapping defined in a neighborhood of a point z_0 . The map ψ is said to be *conformal* at z_0 if the limit

$$(2.2) \quad \lim_{z \rightarrow z_0} \frac{\psi(z) - \psi(z_0)}{z - z_0}$$

exists, in which case we denote the limit by $\psi_z(z_0)$.

Theorem 2.6. *Let ψ be a K -quasiconformal mapping of the finite plane onto itself with $\psi(0) = 0$ and*

$$(2.3) \quad I(r) := \frac{1}{2\pi} \int_{|z| < r} \frac{D(z) - 1}{|z|^2} dA(z) < \infty \text{ for } r < \infty.$$

Then ψ is conformal at $z = 0$ and

$$(2.4) \quad \left| \frac{\psi(z)}{z} - \psi_z(0) \right| < |\psi_z(0)| \varepsilon(|z|), \quad \varepsilon(|z|) \xrightarrow{|z| \rightarrow 0} 0,$$

where the function ε depends only on K and I and not otherwise on the mapping ψ .

3. INTERPOLATION OF POWER MAPS

This Section contains the primary technical contributions of the present work. We will describe in detail the interpolation procedure sketched in the Introduction and prove the relevant estimates. We begin by describing the first step of the interpolation in logarithmic coordinates.

Definition 3.1. We define a region

$$W := \{z \in \mathbb{C} : 0 < \operatorname{Re}(z) < 1\} \setminus \{z \in \mathbb{C} : 0 \leq \operatorname{Re}(z) \leq 1/2 \text{ and } \operatorname{Im}(z) \in 2\mathbb{Z} + 1\}.$$

Given $m \in \mathbb{N}$, we also define a triangulation T_m of W as follows (see Figure 3). Place vertices at

$$(3.1) \quad 0, \left(1 + \frac{j}{m} \cdot i\right)_{j=0}^m, \text{ and } \left(i + \frac{j}{2m-2}\right)_{j=0}^{m-1}.$$

Label the vertices black or white as follows: 0 and 1 are black, i is white, and the other vertices are colored so that adjacent vertices on $\operatorname{Re}(z) = 1$ or $(\operatorname{Im}(z) = 1) \cap (\operatorname{Re}(z) \leq 1/2)$ have different colors. There is a triangulation of $W \cap [0, 1]^2$ formed by connecting each vertex with real part $\leq 1/2$ to $1 + i \cdot (m-1)/m$. Iteratively reflecting this triangulation of $W \cap [0, 1]^2$ along a subset of horizontal lines $\operatorname{Im}(z) = k$, $k \in \mathbb{Z}$ defines the triangulation T_m .

Remark 3.2. The coloring of the vertices in Definition 3.1 is not essential, however it is a useful convention: as already discussed in the Introduction, the colored vertices represent preimages of ± 1 under a map defined below in Definition 3.11, with the two colors corresponding to the two choices ± 1 .

Definition 3.3. Given $m \in \mathbb{N}$, we define a triangulation T'_m of $\{z \in \mathbb{C} : 0 < \operatorname{Im}(z) < 1\}$ as follows (see Figure 3). First place vertices at

$$(3.2) \quad \left(\frac{j}{m} \cdot i\right)_{j=0}^m \text{ and } \left(1 + \frac{j}{m} \cdot i\right)_{j=0}^m.$$

We color 0 and 1 black, and require adjacent vertices with the same real part to have different colors. There is a triangulation of $[0, 1]^2$ defined by connecting each vertex on $\operatorname{Re}(z) = 0$ to $1 + i \cdot (m-1)/m$. Iteratively reflecting this triangulation of $[0, 1]^2$ along a subset of the horizontal lines $\operatorname{Im}(z) = k$, $k \in \mathbb{Z}$ defines the triangulation T'_m .

Definition 3.4. We define a PWL-homeomorphism

$$(3.3) \quad \psi_m : W \rightarrow \{z \in \mathbb{C} : 0 < \operatorname{Im}(z) < 1\}$$




FIGURE 3. Illustrated is the definition of ψ_m in $W \cap [0, 1]^2$ in the cases $m = 2, 3$.

as follows. We specify that

$$(3.4) \quad \psi_m(0) := 0 \text{ and } \psi_m(1 + \frac{j}{m} \cdot i) := 1 + \frac{j}{m} \cdot i \text{ for } 1 \leq j \leq m, \text{ and}$$

$$(3.5) \quad \psi_m\left(i + \frac{j}{2m-2}\right) := \frac{j+1}{m} \cdot i \text{ for } 1 \leq j \leq m-1.$$

As the triangulations $T_m \cap [0, 1]^2$ and $T'_m \cap [0, 1]^2$ are compatible, (3.4) and (3.5) uniquely determine a triangulation-preserving PWL-homeomorphism

$$(3.6) \quad \psi_m : W \cap [0, 1]^2 \rightarrow [0, 1]^2.$$

We extend the map ψ_m to a PWL-homeomorphism $\psi_m : W \rightarrow \{z \in \mathbb{C} : 0 < \text{Im}(z) < 1\}$ by repeated applications of the Schwarz-reflection principle.

Proposition 3.5. *For any $m \in \mathbb{N}$, the map*

$$(3.7) \quad \psi_m : W \rightarrow \{z \in \mathbb{C} : 0 < \text{Im}(z) < 1\}$$

of Definition 3.4 is quasiconformal.

Proof. The map

$$(3.8) \quad \psi_m : W \cap [0, 1]^2 \rightarrow [0, 1]^2$$

is defined as a PWL map, and hence the dilatation constant of (3.8) is the supremum of the dilatations of the $m+2$ \mathbb{R} -linear maps in the definition of (3.8). Thus (3.8) is quasiconformal. As the definition of ψ_m in W is obtained by the Schwarz-reflection principle, it follows that ψ_m is also quasiconformal with the same constant. \square

Remark 3.6. $K(\psi_m) \rightarrow \infty$ as $m \rightarrow \infty$.

The above essentially defines the first step in our interpolation in logarithmic coordinates. We now revert back to the z -plane (see also Figure 5).

Definition 3.7. We define a planar region

$$E_n := \exp\left(\frac{\pi}{n} \cdot W\right).$$

Given $n, m \in \mathbb{N}$, we also define a map

$$(3.9) \quad \eta_{n,m}(z) := \left(z \mapsto \exp\left(\frac{\pi}{n} \cdot z\right)\right) \circ \psi_m \circ \left(z \mapsto \frac{n}{\pi} \log z\right) \text{ for } z \in E_n.$$

Proposition 3.8. *For any $m \in \mathbb{N}$, the map*

$$(3.10) \quad \eta_{n,m} : E_n \rightarrow \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(\pi/n)\}$$

is a quasiconformal homeomorphism. Moreover, $K(\eta_{n,m})$ depends only on m .

Proof. This follows from (3.9) and Proposition 3.5. \square

Having transferred the first step of our interpolation back to the z -plane, we now have a function $\eta_{n,m}$ which is the desired map (A) \mapsto (B) of the Introduction. The composition (A) \mapsto (C) is given by $(z \mapsto z^{nm}) \circ \eta_{n,m}$. As discussed in the Introduction, $(z \mapsto z^{nm}) \circ \eta_{n,m}$ is 2-valued across the radial arcs on the boundary of E_n , and we will need a procedure to “identify” these 2 values. This is given in Definition 3.11 below, for which we will first need the following:

Definition 3.9. Following [BL19], we define a 3-quasiconformal map

$$(3.11) \quad \sigma : \{z \in \mathbb{C} : |z| > 1\} \rightarrow \mathbb{C} \setminus [-1, 1]$$

as follows (see Figure 4). Denote by $\mu(z) := (z+1)/(z-1)$ the Möbius transformation mapping $|z| > 1$ conformally to the right-half plane. Let ν denote the 3-quasiconformal map sending the right-half plane to $\mathbb{C} \setminus (-\infty, 0]$ that is the identity on $|\arg(z)| \leq \pi/4$ and triples angles in the remaining sector. Then

$$(3.12) \quad \sigma := \mu \circ \nu \circ \mu.$$




FIGURE 4. Illustrated is the map σ of Definition 3.9. This Figure is taken from [BL19].

Remark 3.10. We will denote $X := \text{supp}(\sigma_{\bar{z}}/\sigma_z)$ (illustrated as the dark gray region in the left-most copy of \mathbb{C} in Figure 4). Note that σ is the identity on $\{z \in \mathbb{C} : |z| > 1\} \setminus X$.

Definition 3.11. Given $n, m \in \mathbb{N}$, we define a quasiregular map $g_{n,m}$ on the region E_n as follows. Consider first the map

$$(3.13) \quad (z \mapsto z^{mn}) \circ \eta_{n,m} : E_n \rightarrow \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(m\pi)\}.$$

Note that

$$X \subset \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(m\pi)\} \text{ for any } m \in \mathbb{N}.$$

There is a partition of the boundary of E_n into the two circular arcs $|z| = 1$, $|z| = \exp(\pi/n)$ and n radial arcs perpendicular to $|z| = 1$ (see Figure 5 for the case $n = 2$ where the radial arcs are colored red). The preimage of X under (3.13) consists of $2mn$ components: let U denote the union of those components which neighbor one of the n radial arcs on the boundary of E_n . We define

$$(3.14) \quad g_{n,m}(z) := \begin{cases} \sigma \circ (z \mapsto z^{mn}) \circ \eta_{n,m}(z) & z \in U \\ (z \mapsto z^{mn}) \circ \eta_{n,m}(z) & z \in E_n \setminus U. \end{cases}$$

Remark 3.12. Note that the two formulas in (3.14) agree on ∂U as $\sigma(z) = z$ for $z \in \partial X$.

Proposition 3.13. For any $n, m \in \mathbb{N}$, The map $g_{n,m}$ of Definition 3.11 is quasiregular on $1 \leq |z| \leq \exp(\pi/n)$. Moreover, $K(g_{n,m})$ depends only on m , and:

- (1) $g_{n,m}(z) = z^n$ if $|z| = 1$, and
- (2) $g_{n,m}(z) = z^{mn}$ if $|z| = \exp(\pi/n)$.

Proof. It is evident from Definition 3.11 that $g_{n,m}$ is quasiregular in E_n as it is a composition of quasiregular maps in E_n . Thus to show that $g_{n,m}$ is quasiregular on $1 \leq |z| \leq \exp(\pi/n)$, it suffices (by a standard removability result) to show that $g_{n,m}$ extends continuously across the radial arcs on the boundary of E_n . For z on such a radial arc, $(z \mapsto z^{mn}) \circ \eta_{n,m}(z)$ is




FIGURE 5. The two dark-gray regions in (C) represent the set X of Remark 3.10, and the dark-gray regions in (B) and (A) represent the pullback of X under the maps $z \mapsto z^6$ and $\eta_{2,3} \circ (z \mapsto z^6)$, respectively. The definition of $g_{2,3}$ differs from that of $(z \mapsto z^6) \circ \eta_{2,3}$ only in the four dark-gray regions in (A) neighboring the radial arcs colored red. The map $(z \mapsto z^6) \circ \eta_{2,3}$ is 2-valued on these arcs, whereas $g_{2,3}$ is single-valued and indeed continuous across these arcs.

2-valued with both values lying at complex-conjugate points on \mathbb{T} , whence σ identifies these two points. Thus $g_{n,m}$ extends continuously across the radial arcs on the boundary of E_n .

It remains to show (1) and (2). (2) is evident since $|z| = \exp(\pi/n)$ is disjoint from U , and $\eta_{n,m}(z) = z$ for $|z| = \exp(\pi/n)$ (see Definition 3.4 of ψ_m). The verification of (1) is straightforward: it follows from the definition of $\eta_{n,m}$ and fact that for $|z| = 1$, $g_{n,m}(z) = (z \mapsto z^{mn}) \circ \eta_{n,m}(z)$. We leave the details to the reader. That $K(g_{n,m})$ depends only on m follows from the Formula (3.14) and Proposition 3.8. \square

In order to understand the singular values of the function f of Theorem A, we will need to keep track of those points at which our interpolating function is locally $n : 1$ for $n > 1$. To this end, we introduce the following definition:

Definition 3.14. Let g be a quasiregular function, defined in a neighborhood of a point $z \in \mathbb{C}$. We say that z is a *branched point* of g if for any neighborhood U of z , the map $g|_U$ is $n : 1$ onto its image for $n > 1$. If, further, $n = 2$, we say that z is a *simple branched point*. We say $w \in \mathbb{C}$ is a *branched value* of g if $w = g(z)$ for a branched point z of g .

Proposition 3.15. Let g be as in Definition 3.11. Define $g(z) := z^n$ for $|z| < 1$, and $g(z) := z^{mn}$ for $|z| > \exp(\pi/n)$. Then the only non-zero branched points of g are:

$$(3.15) \quad \exp\left(\frac{\pi}{n} \cdot \frac{l}{(2m-2)} + i \cdot \frac{\pi(2k-1)}{n}\right) \quad \text{for } 0 \leq l \leq m-2, \text{ and } 1 \leq k \leq n.$$

Moreover, all non-zero branched points of g are simple, and the only non-zero branched values of g are ± 1 .

Remark 3.16. The expression (3.15) is simply a formula for the black and white vertices located on the radial arcs colored red in, for instance, Figure 5(A), but excluding those vertices at the outer-most tips of the radial arcs.

Proof. We first note that the extended formula defines a quasiregular function on \mathbb{C} by removability of analytic arcs for quasiregular mappings. It is readily verified then from the definition of g that any neighborhood of each of the points (3.15) is mapped 2 : 1 onto its image by g . The points (3.15) are sent to ± 1 by g , where we note that the sign may be determined by the coloring of the vertex as described in Remark 3.2. Lastly, again from the definition of g , one verifies directly that there are no remaining non-zero branched points of g . □

We will also need to record the zeros of our interpolating function for later application. The formulas are listed below in Proposition 3.17, but they are readily seen to just be the midpoint between each adjacent black/white vertex on a radial segment pictured in, for instance, Figure 5(A).

Proposition 3.17. *Let g be as in Proposition 3.15. Then the zeros of g are given by:*

$$(3.16) \quad 0 \text{ and } \exp\left(\frac{\pi}{n} \cdot \frac{l + \frac{1}{2}}{2m - 2} + i \cdot \frac{\pi(2k - 1)}{n}\right) \text{ for } 0 \leq l \leq m - 2 \text{ and } 1 \leq k \leq n,$$

all of which are simple except for 0 which is of multiplicity n .

Proof. Recalling Definition 3.9 of σ , we see that $\sigma^{-1}(0) = \{\pm i\}$. Let $S := (z \mapsto z^{mn})^{-1}(\pm i)$. Then $g^{-1}(0)$ is readily seen to be those points x such that $\eta_{n,m}(x) = \{\pm i\}$ and $x \in U$ for U as in Definition 3.11. Such x are listed in (3.16). □

We now generalize our interpolating function slightly to allow for interpolation between $z \mapsto z^n$ and $z \mapsto z^M$ for $M > n$, where M is not necessarily a multiple of n . While there is a necessary level of complication in the formulas in the following proof, the idea is simple. Denoting $m := \lfloor M/n \rfloor$, we will use $\eta_{n,m}$ in some parts of the interpolating annulus, and $\eta_{n,m+1}$ in others, so that we add the necessary amount $M - n$ of new vertices (branched points). Next, we adjust by a homeomorphism τ of the circle which sends the n old vertices and $M - n$ new vertices on \mathbb{T} together to span the M^{th} roots of unity, and finally post-compose with an adjustment of the map $z \mapsto z^M$ as in Definition 3.11.

Proposition 3.18. *Let $n, M \in \mathbb{N}$ with $M > n$. Then there exists a quasiregular function*

$$(3.17) \quad g : \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(\pi/n)\} \rightarrow \{z \in \mathbb{C} : |z| \leq \exp(M\pi/n)\},$$

such that:

- (1) $g(z) = z^n$ if $|z| = 1$,
- (2) $g(z) = z^M$ if $|z| = \exp(\pi/n)$, and
- (3) $K(g)$ depends only on M/n , and not otherwise on M or n .

Proof. Consider the region E_n as in Definition 3.7. Let $m := \lfloor M/n \rfloor$. Let

$$(3.18) \quad E_{n,j} := E_n \cap \left\{ z \in \mathbb{C} \setminus \{0\} : \frac{2(j-1)\pi}{n} \leq \arg(z) \leq \frac{2j\pi}{n} \right\} \text{ for } 1 \leq j \leq n.$$

Let $p := n - M + nm$, and recall the mapping $\eta_{n,m}$ as in Definition 3.7, where we use the convention $\eta_{n,0}(z) \equiv z$. We define the mapping

$$(3.19) \quad \eta(z) := \begin{cases} \eta_{n,m}(z) & z \in \cup_{j=1}^p E_{n,j} \\ \eta_{n,m+1}(z) & z \in \cup_{j=p+1}^n E_{n,j}. \end{cases}$$

It is readily verified that $\eta : E_n \rightarrow \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(\pi/n)\}$ is a homeomorphism, since ψ_m, ψ_{m+1} are both the identity on the lines $y = 2\mathbb{Z}$. Next, we define a homeomorphism $\tau : \mathbb{T} \rightarrow \mathbb{T}$ by

$$(3.20) \quad \tau(\theta) := \begin{cases} \frac{nm}{M} \cdot \theta & \theta \in [0, 2p\pi/n] \\ \frac{(m+1)n}{M} \cdot \theta & \theta \in [-2\pi(n-p)/n, 0]. \end{cases}$$

Extend τ to a self-homeomorphism of $A(1, \exp(\pi/n))$ by linearly interpolating $\tau|_{\mathbb{T}}$ with the identity on $|z| = \exp(\pi/n)$. Since $n - p = (M/n)/(M/n - m)$, it is readily seen from (3.20) that $K(\tau)$ depends only on M/n , and not otherwise on M or n .

Finally, we define g by appropriately adjusting Definition 3.11. Namely, let $X := \text{supp}(\sigma_{\bar{z}}/\sigma_z)$ as in Remark 3.10. Then the pullback of X under

$$(3.21) \quad (z \mapsto z^M) \circ \tau \circ \eta : E_n \rightarrow A(1, \exp(M\pi/n))$$

consists of $2M$ components. Denoting by U the union of those $2(M-n)$ components which neighbor a radial arc on the boundary of E_n , we again define

$$(3.22) \quad g(z) := \begin{cases} \sigma \circ (z \mapsto z^M) \circ \eta(z) & z \in U \\ (z \mapsto z^M) \circ \eta(z) & z \in E_n \setminus U. \end{cases}$$

The proof that g satisfies (1)-(3) then is the same as in Proposition 3.13. □

Following Proposition 3.15, we will also list the branched points, branched values, and zeros of the map g of Proposition 3.18. But first, we describe a simple rescaling which allows our annular region of interpolation to have an inner boundary lying on a circle of variable radius.

Proposition 3.19. *Let $r \geq 1$, $c \in \mathbb{C}^*$ and $n, M \in \mathbb{N}$ with $M > n$. There exists a quasiregular function*

$$(3.23) \quad g : \{z \in \mathbb{C} : r \leq |z| \leq r \exp(\pi/n)\} \rightarrow \{z \in \mathbb{C} : |z| \leq cr^n \exp(M\pi/n)\},$$

such that:

- (1) $g(z) = c \cdot z^n$ if $|z| = r$,
- (2) $g(z) = c \cdot z^M / r^{M-n}$ if $|z| = r \exp(\pi/n)$, and
- (3) $K(g)$ depends only on M/n , and not otherwise on M, n, r or c .

Proof. We first consider the case when $M = m \cdot n$ for $m \in \mathbb{N}$. Let

$$(3.24) \quad E_{n,r} := \exp\left(\frac{\pi}{n} \cdot W + \log r\right).$$

By Proposition 3.8, the map

$$(3.25) \quad \eta_{n,m,r} := \eta_{n,m} \circ (z \mapsto z/r) : E_{n,r} \rightarrow \{z \in \mathbb{C} : 1 \leq |z| \leq \exp(\pi/n)\}$$

is a quasiconformal homeomorphism where $K(\eta_{n,m,r})$ depends only on m . Following Definition 3.11, we can define a map \tilde{g} in $E_{n,r}$ by adjusting the values of $(z \mapsto z^{mn}) \circ \eta_{n,m,r}$ in a neighborhood of the radial arcs on the boundary of $E_{n,r}$. The proof that \tilde{g} is quasiregular on $\{z \in \mathbb{C} : r \leq |z| \leq r \exp(\pi/n)\}$ is then the same as in Proposition 3.13. Set

$$(3.26) \quad g(z) := c \cdot r^n \cdot \tilde{g}(z).$$

Then the proof that g satisfies (1)-(3) similarly follows as in Proposition 3.13. Lastly, the case when $M > n$ is not necessarily a multiple of n follows from the same adjustment of the above interpolation as in Proposition 3.18. \square

Remark 3.20. When we wish to emphasize the dependence of g on the parameters n, M, r, c , we will write $g_{n,M,r,c}$ (in that order).

Lastly, we record the branched points, branched values, and zeros of our interpolating function. The same proofs as for Propositions 3.15 and 3.17 prove the following:

Proposition 3.21. *Let g , and notation be as in Proposition 3.19. Set $m := \lfloor M/n \rfloor$, and $p := n - M + nm$. Define $g(z) := cz^n$ for $|z| < r$, and $g(z) := cz^M / r^{M-n}$ for $|z| > r \exp(\pi/n)$. Then the only non-zero branched points of g are:*

$$(3.27) \quad \begin{aligned} & r \cdot \exp\left(\frac{\pi}{n} \cdot \frac{l}{(2m-2)} + i \cdot \frac{\pi(2k-1)}{n}\right) \quad \text{for } 0 \leq l \leq m-2 \text{ and } 1 \leq k \leq p, \text{ and} \\ & r \cdot \exp\left(\frac{\pi}{n} \cdot \frac{l}{2m} + i \cdot \frac{\pi(2k-1)}{n}\right) \quad \text{for } 0 \leq l \leq m-1, \text{ and } p+1 \leq k \leq n. \end{aligned}$$

Moreover, all non-zero branched points of g are simple, and the only non-zero branched values of g are $\pm cr^n$. The zeros of g are given by the multiplicity n zero at 0, and the simple zeros described by replacing l with $l + 1/2$ in the expressions (3.27).

4. ENTIRE FUNCTIONS

With the technical work of Section 3 behind us, we will now apply our interpolation repeatedly between power maps of increasing degree in “increasing” annuli in the plane. As long as the degree of the power maps increases by at most a fixed constant factor in each consecutive annulus, this procedure gives a function quasiregular in \mathbb{C} . We formalize this below.

Definition 4.1. Let $c \in \mathbb{C}^*$, $(M_j)_{j=1}^\infty \in \mathbb{N}$ be increasing, and $(r_j)_{j=1}^\infty \in \mathbb{R}^+$. We set $r_0 := 0$, $M_0 := 1$. Suppose that

$$(4.1) \quad r_{j+1} \geq \exp(\pi/M_j) \cdot r_j \text{ for all } j \in \mathbb{N}, \text{ and } r_j \xrightarrow{j \rightarrow \infty} \infty.$$

Set

$$(4.2) \quad c_1 := c, \text{ and } c_j := c_{j-1} \cdot r_{j-1}^{M_{j-1}-M_j} = c \cdot \prod_{k=2}^j r_{k-1}^{M_{k-1}-M_k} \text{ for } j \geq 2.$$

We then define:

$$(4.3) \quad h(z) := \begin{cases} c_j \cdot z^{M_j} & \text{if } r_{j-1} \cdot \exp(\pi/M_{j-1}) \leq |z| \leq r_j \\ g_{M_j, M_{j+1}, r_j, c_j}(z) & \text{if } r_j \leq |z| \leq r_j \cdot \exp(\pi/M_j). \end{cases}$$

over all $j \in \mathbb{N}$.

Remark 4.2. By Proposition 3.19, the two definitions in (4.3) agree on $|z| = r_j$. By Proposition 3.19 and the definition (4.2) of c_j , the two definitions in (4.3) agree on $|z| = r_j \cdot \exp(\pi/M_j)$. Thus the formula (4.3) determines a well-defined function $h : \mathbb{C} \rightarrow \mathbb{C}$.

Definition 4.3. We will say that $(M_j)_{j=1}^\infty \in \mathbb{N}$, $(r_j)_{j=1}^\infty \in \mathbb{R}^+$ are *weakly permissible* if (4.1) is satisfied.

Remark 4.4. Suppose $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are weakly permissible. Then, according to Definition 1.1, we have that $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are permissible if and only if the sequence $(M_j/M_{j-1})_{j=1}^\infty$ is bounded.

Remark 4.5. The map h of Definition 4.1 is determined by a choice of $c \in \mathbb{C}^*$ and weakly permissible $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$.

Proposition 4.6. Suppose $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are weakly permissible, and $c \in \mathbb{C}^*$. Then the function h as defined in Definition 4.1 is quasiregular on compact subsets of \mathbb{C} . If, moreover, $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are permissible, then h is quasiregular on \mathbb{C} and $K(h)$ depends only on $\sup_j (M_j/M_{j-1})$ and not otherwise on $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$.

Proof. In each annular region in the Definition (4.3), the map h is either analytic, or quasiregular by Proposition 3.19. By removability of analytic arcs, the map h is therefore quasiregular across the boundaries of the annular regions. As any compact subset of \mathbb{C} meets only

finitely many such annular regions, it follows that h is locally quasiregular. If the sequence $(M_j/M_{j-1})_{j=1}^\infty$ is bounded (or equivalently $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are permissible), then each of the maps $g_{M_j, M_{j+1}, r_j, c_j}(z)$ have a dilatation bounded uniformly over j (with a bound depending only on $\sup_j (M_j/M_{j-1})$) by (3) of Proposition 3.19, and so the last statement of the Proposition follows. \square

The definition of *permissible* thus ensures that the formula (4.3) defines a quasiregular function, in which case we may integrate the Beltrami coefficient $h_{\bar{z}}/h_z$ to obtain a quasiconformal map ϕ such that $h \circ \phi^{-1}$ is holomorphic. If, moreover, the M_j increase sufficiently quickly (see (4.4)), the total region of interpolation is sufficiently small to guarantee conformality of ϕ at ∞ . This is the content of Theorem 4.8 below, which will be presented after the following definition:

Definition 4.7. Let $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ be permissible. We say that $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ are *strongly permissible* if also

$$(4.4) \quad \sum_j M_j^{-1} < \infty.$$

Theorem 4.8. Let $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ be strongly permissible, and $c \in \mathbb{C}^*$. Then there exists a quasiconformal mapping $\phi : \mathbb{C} \rightarrow \mathbb{C}$ such that $f := h \circ \phi^{-1}$ as in Theorem A is holomorphic, and

$$(4.5) \quad \left| \frac{\phi(z)}{z} - 1 \right| \xrightarrow{z \rightarrow \infty} 0.$$

Proof. The proof is an application of Theorem 2.6. The existence of a quasiconformal $\phi : \mathbb{C} \rightarrow \mathbb{C}$ such that $h \circ \phi^{-1}$ is holomorphic and $\phi(0) = 0$ follows from applying the Measurable Riemann Mapping Theorem to $h_{\bar{z}}/h_z$. Let

$$(4.6) \quad \psi(z) := 1/\phi(1/z).$$

Then ψ is a quasiconformal self-mapping of \mathbb{C} satisfying $\psi(0) = 0$. Let K denote the quasiconformal constant of ψ , and take $r < \infty$. We calculate

$$(4.7) \quad I(r) := \frac{1}{2\pi} \int_{|z| < r} \frac{D(z) - 1}{|z|^2} dA(z) < \frac{1}{2\pi} \int_{|z| < 1} \frac{K - 1}{|z|^2} dA(z) = \frac{1}{2\pi} \sum_{j=1}^\infty \int_{B_j} \frac{K - 1}{|z|^2} dA(z),$$

where

$$(4.8) \quad B_j := \{z \in \mathbb{C} : r_j^{-1} \exp(-\pi/M_j) \leq |z| \leq r_j^{-1}\}.$$

We continue our calculation:

$$(4.9) \quad \begin{aligned} \frac{1}{2\pi} \sum_{j=1}^{\infty} \int_{B_j} \frac{K-1}{|z|^2} dA(z) &\leq \frac{K-1}{2\pi} \sum_{j=1}^{\infty} \int_{B_j} \frac{1}{r_j^{-2} \exp(-2\pi/M_j)} dA(z) = \\ \frac{K-1}{2\pi} \sum_{j=1}^{\infty} \frac{\pi(r_j^{-2} - r_j^{-2} \exp(-2\pi/M_j))}{r_j^{-2} \exp(-2\pi/M_j)} &= \frac{K-1}{2} \sum_{j=1}^{\infty} (\exp(2\pi/M_j) - 1). \end{aligned}$$

The infinite sum on the right-hand side of (4.9) converges if and only if

$$(4.10) \quad \prod_{j=1}^{\infty} \exp(2\pi/M_j)$$

converges, which is readily seen to be the case by assumption of (4.4). Thus, by Theorem 2.6, the limit

$$(4.11) \quad \psi_z(0) := \lim_{z \rightarrow 0} \frac{\psi(z)}{z}$$

exists, and we have

$$(4.12) \quad \left| \frac{\psi(z)}{z} - \psi_z(0) \right| \xrightarrow{|z| \rightarrow 0} 0.$$

By multiplying ψ by $1/\psi_z(0)$ if necessary, we can further assume that $\psi_z(0) = 1$. Since $\psi(z) = 1/\phi(1/z)$, (4.5) follows. \square

Next we list the singularities of the entire function f :

Proposition 4.9. *Let $(M_j)_{j=1}^{\infty}$, $(r_j)_{j=1}^{\infty}$ be permissible, $c \in \mathbb{C}^*$, and f, ϕ as in Theorem 4.8. Set $m_j := \lfloor M_j/M_{j-1} \rfloor$, and $p_j := M_{j-1} - M_j + M_{j-1}m_j$. Then the only critical points of f are 0 and the simple critical points given by*

$$(4.13) \quad \begin{aligned} &\phi \left(r_j \cdot \exp \left(\frac{\pi}{M_{j-1}} \cdot \frac{l_j}{2m_j - 2} + i \frac{(2k_j - 1)\pi}{M_{j-1}} \right) \right), \text{ and} \\ &\phi \left(r_j \cdot \exp \left(\frac{\pi}{M_{j-1}} \cdot \frac{l'_j}{2m_j} + i \frac{(2k'_j - 1)\pi}{M_{j-1}} \right) \right), \end{aligned}$$

where $j \in \mathbb{N}$, and $1 \leq k_j \leq p_j$, $0 \leq l_j \leq m_j - 2$, and $p_j + 1 \leq k'_j \leq M_{j-1}$, $0 \leq l'_j \leq m_j - 1$. The only singular values of f are the critical values $(\pm c_j r_j^{M_j})_{j=0}^{\infty}$.

Proof. By Proposition 3.21, the only branched points of h are 0 and the points given by (4.13) without the ϕ factor. Thus, as $f := h \circ \phi^{-1}$, it follows the only critical points of f are $\phi(0) = 0$ and those given in (4.13). Again, as $f := h \circ \phi^{-1}$, by Proposition 3.21 each of the points in (4.13) is mapped to $\pm c_j r_j^{M_j}$ as j ranges over \mathbb{N} . There are no asymptotic values of f : if $\gamma \rightarrow \infty$ is a curve, then $f(\gamma)$ is unbounded. \square

Remark 4.10. The same proof as for Proposition 4.9 shows that the zeros of f are given by 0 (multiplicity M_1) and the simple zeros whose formulae is given by replacing l_j and l'_j in the expressions in (4.13) by $l_j + 1/2$, $l'_j + 1/2$, respectively.

Theorem A now follows:

Proof of Theorem A: We take $f := h \circ \phi^{-1}$ as in Theorem 4.8. The conclusions of Theorem A are then included in the statements of Theorem 4.8 and Proposition 4.9.

The applications of the present manuscript of which the authors are aware are in regards to entire functions with an essential singularity at ∞ , and so it is natural to impose the assumption $r_j \rightarrow \infty$ in Definition 1.1 of “permissibility”. However, this is inessential to most of our arguments. Indeed, if we assume instead that $r_j \rightarrow r_\infty < \infty$, we can define h exactly as in (4.3), and then the argument of Proposition 4.6 together with an application of the Measurable Riemann Mapping theorem yields the following:

Theorem B. *Let $c \in \mathbb{C}^*$, $0 < r_\infty < \infty$, $(M_j)_{j=1}^\infty \in \mathbb{N}$ be increasing, $(r_j)_{j=1}^\infty \in \mathbb{R}^+$ such that*

$$(4.14) \quad r_{j+1} \geq \exp(\pi/M_j) \cdot r_j \text{ for all } j \in \mathbb{N}, r_j \xrightarrow{j \rightarrow \infty} r_\infty, \text{ and } \sup_j \frac{M_{j+1}}{M_j} < \infty.$$

Set $r_0 := 0$ and

$$(4.15) \quad c_1 := c, \text{ and } c_j := c_{j-1} \cdot r_{j-1}^{M_{j-1}-M_j} \text{ for } j \geq 2.$$

Then there exists a holomorphic function $f : r_\infty \mathbb{D} \rightarrow \mathbb{C}$ and a K -quasiconformal homeomorphism $\phi : r_\infty \mathbb{D} \rightarrow r_\infty \mathbb{D}$ with K depending only on $\sup_j (M_{j+1}/M_j)$ such that:

$$(4.16) \quad f \circ \phi(z) = c_j z^{M_j} \text{ for } r_{j-1} \cdot \exp(\pi/M_{j-1}) \leq |z| \leq r_j, j \in \mathbb{N}.$$

Moreover, the only singular values of f are the critical values $(\pm c_j r_j^{M_j})_{j=1}^\infty$.

Missing from the statement of Theorem B is an application of criteria for conformality at a point (in Theorem A the criteria is $\sum_j M_j^{-1} < \infty$, and the point is ∞). Although we do not record them here, one can produce analogous criteria in Theorem B for conformality of ϕ at points on $|z| = r_\infty$. Indeed, we may extend ϕ to a self-map of $\hat{\mathbb{C}}$ by the Schwarz reflection principle, and apply Theorem 2.6 at a point lying on $|z| = r_\infty$.

5. A DYNAMICAL APPLICATION

In this Section, we discuss some brief dynamical applications of Theorem A. We will discuss an approach to constructing entire functions with multiply-connected wandering domains using Theorem A. But first we will show how, in certain applications, we can conclude much more than conformality at ∞ of the map ϕ in Theorem A. Namely we can conclude a uniform estimate $\|\phi(z)/z - 1\|_{L^\infty(\mathbb{C})} < \varepsilon$ in the following situation: given a function $f = h \circ \phi^{-1}$ as in Theorem A, if we define h_n by replacing h in $|z| \leq r_n$ with

$z \mapsto c_n z^{M_n}$, we obtain a sequence of entire functions $f_n = h_n \circ \phi_n^{-1}$ with $f_n \approx f$ in $|z| \geq r_n$. As $n \rightarrow \infty$, the maps ϕ_n converge uniformly (in the spherical metric on $\hat{\mathbb{C}}$) to the identity so that $\|\phi_n(z)/z - 1\|_{L^\infty(\mathbb{C})} < \varepsilon$ for large n . This argument is very useful in dynamical applications: it will be used in the companion paper [BL], and a related argument is used in [Bis15], [FJL19], [Laz19]. We formalize the above discussion below.

Definition 5.1. Let

$$(5.1) \quad (M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty \text{ be strongly permissible, and } c \in \mathbb{C}^\star.$$

Denote by $f = h \circ \phi^{-1}$ the entire function obtained by applying Theorem A to (5.1), where h is quasiregular and ϕ quasiconformal. We define a sequence $(f_n)_{n=0}^\infty$ of entire functions as follows. Let $h_0 := h$ and for $n \geq 1$:

$$h_n(z) = \begin{cases} c_n z^{M_n} & \text{if } |z| \leq r_n \\ h(z) & \text{if } |z| \geq r_n. \end{cases}$$

Define $f_n := h_n \circ \phi_n^{-1}$, where $\phi_n : \mathbb{C} \rightarrow \mathbb{C}$ is the unique quasiconformal mapping such that:

- (1) f_n is holomorphic,
- (2) $\phi_n(0) = 0$, and
- (3) $|\phi_n(z)/z - 1| \rightarrow 0$ as $z \rightarrow \infty$.

Remark 5.2. That we may normalize ϕ_n in Definition 5.1 so as to satisfy (3) requires justification. To this end, let

$$(5.2) \quad I_n(r) := \frac{1}{2\pi} \int_{|z| < r} \frac{D_n(z) - 1}{|z|^2} dA(z) \text{ for } r > 0,$$

where D_n is the dilatation quotient (see Definition 2.4) of $\psi_n(z) := 1/\phi_n(1/z)$. Then

$$(5.3) \quad D_n(z) = 0 \text{ if } |z|^{-1} \leq r_n, \text{ and } D_n(z) \stackrel{a.e.}{=} D_0(z) \text{ if } |z|^{-1} \geq r_n.$$

In particular, $I_n(r) \leq I_0(r)$. Thus by Theorem 2.6, ψ_n is conformal at 0, and hence ϕ_n is conformal at ∞ . Thus we may normalize ϕ_n as claimed.

Theorem 5.3. *Let $(M_j)_{j=1}^\infty, (r_j)_{j=1}^\infty$ be strongly permissible, $c \in \mathbb{C}^\star$, and notation as in Definition 5.1. Then for any $\varepsilon > 0$, there exists $N_\varepsilon \in \mathbb{N}$ such that for $n \geq N_\varepsilon$,*

$$(5.4) \quad \|\phi_n(z)/z - 1\|_{L^\infty(\mathbb{C})} < \varepsilon.$$

Proof. Briefly, this is a consequence of the error term in the conclusion (2.4) of Theorem 2.6 depending only on K and I , and not otherwise on the quasiconformal mapping under consideration. Let us explain further. We follow the notation of Remark 5.2. For I_n as in (5.2), we have already noted that $I_n(r) \leq I_0(r)$. Moreover, $K(\psi_n) \leq K(\psi_0)$ by (5.3). Thus, by Theorem 2.6, there is a function $\iota : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with

$$(5.5) \quad \left| \frac{\psi_n(z)}{z} - 1 \right| < \iota(|z|)$$

where $\iota(r) \rightarrow 0$ as $r \rightarrow 0$, and ι does not depend on n . Thus, given $\varepsilon > 0$, there exists $R > 0$ such that:

$$(5.6) \quad \left\| \frac{\phi_n(z)}{z} - 1 \right\|_{L^\infty(|z| \geq R)} < \varepsilon \text{ for all } n \in \mathbb{N}.$$

On the other hand, for all $z \in \mathbb{C}$ we have $D_n(z) \rightarrow 0$ as $n \rightarrow \infty$. Thus, by a standard argument, we have

$$(5.7) \quad \|\phi_n(z) - z\|_{L^\infty(|z| \leq R)} \xrightarrow{n \rightarrow \infty} 0.$$

The result now follows from (5.6), (5.7), and a K  be distortion estimate for $\phi'_n(0)$. \square

We now turn to an application: we will show that as long as our parameters satisfy certain simple relations, the resulting entire function of Theorem A has a multiply connected wandering domain.

Notation 5.4. For $0 < \alpha < 1$, we let $A_j^\alpha := \{z \in \mathbb{C} : \alpha^{-1}r_{j-1} \cdot \exp(\pi/M_{j-1}) < |z| < \alpha r_j\}$.

Theorem 5.5. *Suppose $(M_j)_{j=1}^\infty$, $(r_j)_{j=1}^\infty$ are strongly permissible, $c \in \mathbb{C}^*$, and*

$$(5.8) \quad r_{j+1} = c_j r_j^{M_j}.$$

Then, for any $\alpha > 1$, there exists N_α such that, for $j \geq N_\alpha$, A_j^α is contained in a wandering domain for f as in Theorem A.

Remark 5.6. The condition (5.8) is sufficient to guarantee a wandering domain for the function f of Theorem A, but is far from necessary.

Proof. As usual, we denote $f = h \circ \phi^{-1}$. By Theorem A, there exists $N = N_\alpha \in \mathbb{N}$ such that

$$(5.9) \quad |\phi(z)/z - 1| < |\sqrt{\alpha} - 1| \text{ for } |z| \geq r_N.$$

It follows that

$$(5.10) \quad \phi(A_j^\alpha) \subset A_j^{\sqrt{\alpha}} \text{ for } j \geq N.$$

By (5.8), we have $h(r_j) = r_{j+1}$ for all j . Together with expansivity of h , this implies:

$$(5.11) \quad f(A_j^\alpha) \subset A_{j+1}^\alpha \text{ for } j \geq N,$$

perhaps after increasing N . Thus the iterates of f converge uniformly to ∞ on A_j^α for $j \geq N$, and so each such A_j^α is contained in a Fatou component for f , which we will call Ω_j .

If we suppose by way of contradiction that $\Omega_j = \Omega_k$ for some $j \neq k$, then $\Omega_j = \Omega_{j+1}$, and this would imply that Ω_j is unbounded. But Ω_j is multiply connected since $f(0) = 0$, and this is a contradiction since all multiply connected Fatou components of f must be bounded by [Bak75, Theorem 1]. Thus, we may conclude that the $(\Omega_j)_{j=N}^\infty$ form a distinct sequence of Fatou components. Since $f(\Omega_j) \subset \Omega_{j+1}$, it follows that each such Ω_j is a wandering component for f . \square

Remark 5.7. Let notation be as in Theorem 5.5, and consider the family f_n of Definition 5.1. It is not difficult to show using Theorem 5.3 that if $0 < \alpha < 1$, then for sufficiently large n we have that *each* A_j^α for $j \in \mathbb{N}$ is contained in a wandering component Ω_j of f_n .

REFERENCES

- [AB20] Simon Albrecht and Christopher J. Bishop. Speiser class Julia sets with dimension near one. *J. Anal. Math.*, 141(1):49–98, 2020.
- [Bak75] I. N. Baker. The domains of normality of an entire function. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 1(2):277–283, 1975.
- [Bak76] Irvine N. Baker. An entire function which has wandering domains. *J. Austral. Math. Soc. Ser. A*, 22(2):173–176, 1976.
- [Bak85] I. N. Baker. Some entire functions with multiply-connected wandering domains. *Ergodic Theory Dynam. Systems*, 5(2):163–169, 1985.
- [BE00] Walter Bergweiler and Alexandre Eremenko. Entire functions of slow growth whose Julia set coincides with the plane. *Ergodic Theory Dynam. Systems*, 20(6):1577–1582, 2000.
- [Ber11] Walter Bergweiler. An entire function with simply and multiply connected wandering domains. *Pure Appl. Math. Q.*, 7(1):107–120, 2011.
- [Bis14] Christopher J. Bishop. True trees are dense. *Invent. Math.*, 197(2):433–452, 2014.
- [Bis15] Christopher J. Bishop. Constructing entire functions by quasiconformal folding. *Acta Math.*, 214(1):1–60, 2015.
- [Bis18] Christopher J. Bishop. A transcendental Julia set of dimension 1. *Invent. Math.*, 212(2):407–460, 2018.
- [BL] Jack Burkart and Kirill Lazebnik. Interpolation of Power Maps II: Transcendental Julia Sets of Dimension 1 (Working title). *in preparation*.
- [BL19] Christopher J. Bishop and Kirill Lazebnik. Prescribing the postsingular dynamics of meromorphic functions. *Math. Ann.*, 375(3-4):1761–1782, 2019.
- [BP15] Christopher J. Bishop and Kevin M. Pilgrim. Dynamical dessins are dense. *Rev. Mat. Iberoam.*, 31(3):1033–1040, 2015.
- [BRS13] Walter Bergweiler, Philip J. Rippon, and Gwyneth M. Stallard. Multiply connected wandering domains of entire functions. *Proc. Lond. Math. Soc. (3)*, 107(6):1261–1301, 2013.
- [BRS17] Karl F. Barth, Philip J. Rippon, and David J. Sixsmith. The MacLane class and the Eremenko-Lyubich class. *Ann. Acad. Sci. Fenn. Math.*, 42(2):859–873, 2017.
- [Bur19] Jack Burkart. Transcendental Julia Sets with Fractional Packing Dimension. *arXiv e-prints*, page arXiv:1905.07811, May 2019.
- [BZ11] Walter Bergweiler and Jian-Hua Zheng. On the uniform perfectness of the boundary of multiply connected wandering domains. *J. Aust. Math. Soc.*, 91(3):289–311, 2011.
- [Dra86] D. Drasin. On the Teichmüller-Wittich-Belinskii theorem. *Results Math.*, 10(1-2):54–65, 1986.
- [Ere86] A. È. Eremenko. Inverse problem of the theory of distribution of values for finite-order meromorphic functions. *Sibirsk. Mat. Zh.*, 27(3):87–102, 223, 1986.
- [ERS20] Vasiliki Evdoridou, Lasse Rempe, and David J. Sixsmith. Fatou’s Associates. *Arnold Math. J.*, 6(3-4):459–493, 2020.
- [FGJ15] Núria Fagella, Sébastien Godillon, and Xavier Jarque. Wandering domains for composition of entire functions. *J. Math. Anal. Appl.*, 429(1):478–496, 2015.
- [FH09] Núria Fagella and Christian Henriksen. The Teichmüller space of an entire function. In *Complex dynamics*, pages 297–330. A K Peters, Wellesley, MA, 2009.

- [FJL19] Núria Fagella, Xavier Jarque, and Kirill Lazebnik. Univalent wandering domains in the Eremenko-Lyubich class. *J. Anal. Math.*, 139(1):369–395, 2019.
- [Hin94] A. Hinkkanen. On the size of Julia sets. In *Complex analysis and its applications (Hong Kong, 1993)*, volume 305 of *Pitman Res. Notes Math. Ser.*, pages 178–189. Longman Sci. Tech., Harlow, 1994.
- [KS08] Masashi Kisaka and Mitsuhiro Shishikura. On multiply connected wandering domains of entire functions. In *Transcendental dynamics and complex analysis*, volume 348 of *London Math. Soc. Lecture Note Ser.*, pages 217–250. Cambridge Univ. Press, Cambridge, 2008.
- [Laz17] Kirill Lazebnik. Several constructions in the Eremenko-Lyubich class. *J. Math. Anal. Appl.*, 448(1):611–632, 2017.
- [Laz19] Kirill Lazebnik. Oscillating Wandering Domains for Functions with Escaping Singular Values. *arXiv e-prints*, page arXiv:1904.10086, April 2019.
- [LV73] O. Lehto and K. I. Virtanen. *Quasiconformal mappings in the plane*. Springer-Verlag, New York-Heidelberg, second edition, 1973. Translated from the German by K. W. Lucas, Die Grundlehren der mathematischen Wissenschaften, Band 126.
- [MPS20] David Martí-Pete and Mitsuhiro Shishikura. Wandering domains for entire functions of finite order in the Eremenko-Lyubich class. *Proc. Lond. Math. Soc. (3)*, 120(2):155–191, 2020.
- [OS16] John W. Osborne and David J. Sixsmith. On the set where the iterates of an entire function are neither escaping nor bounded. *Ann. Acad. Sci. Fenn. Math.*, 41(2):561–578, 2016.
- [RS13] P. J. Rippon and G. M. Stallard. Baker’s conjecture and Eremenko’s conjecture for functions with negative zeros. *J. Anal. Math.*, 120:291–309, 2013.
- [Sul85] Dennis Sullivan. Quasiconformal homeomorphisms and dynamics. I. Solution of the Fatou-Julia problem on wandering domains. *Ann. of Math. (2)*, 122(3):401–418, 1985.