

Constructing Entire Functions (a summary)

Kirill Lazebnik

SUNY Stony Brook

Kirill.Lazebnik@stonybrook.edu

July 29, 2015

$$p(z) = \frac{z^3}{2} - \frac{3z}{2}$$

$$p(z) = \frac{z^3}{2} - \frac{3z}{2}$$

$$p'(z) = (z - 1)(z + 1)$$

$$p(z) = \frac{z^3}{2} - \frac{3z}{2} \text{ (two critical values } \pm 1)$$

$$p'(z) = (z - 1)(z + 1)$$

$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z$$

$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z$$

$$p'(z) = (z - 1)^2(z + 1)$$

$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z \text{ (two critical values } 5/12, -11/12)$$

$$p'(z) = (z - 1)^2(z + 1)$$

Shabat polynomial -

Shabat polynomial - only has two critical values ± 1

Shabat polynomial - only has two critical values ± 1

Proposition: For any *Shabat* polynomial $p(z)$, it is true that $p^{-1}[-1, 1]$ is a tree.

Shabat polynomial - only has two critical values ± 1

Proposition: For any *Shabat* polynomial $p(z)$, it is true that $p^{-1}[-1, 1]$ is a tree, with $\deg(p)$ edges.

Shabat polynomial - only has two critical values ± 1

Proposition: For any *Shabat* polynomial $p(z)$, it is true that $p^{-1}[-1, 1]$ is a tree.

Theorem (Grothendieck): ALL combinatorial trees occur as $p^{-1}[-1, 1]$ for some *Shabat* polynomial $p(z)$.

Shabat polynomial - only has two critical values ± 1

Proposition: For any *Shabat* polynomial $p(z)$, it is true that $p^{-1}[-1, 1]$ is a tree.

Theorem (Grothendieck): ALL combinatorial trees occur as $p^{-1}[-1, 1]$ for some *Shabat* polynomial $p(z)$.

Theorem (Bishop): Any continua can be ϵ -approximated in the Hausdorff metric by some $p^{-1}[-1, 1]$.

trees \iff *Shabat* polynomials

trees \iff *Shabat* polynomials

infinite trees \iff Transcendental Functions

trees \iff *Shabat* polynomials

infinite trees \iff Subclass of Transcendental Functions

trees \iff *Shabat* polynomials

infinite trees \iff Subclass of Transcendental Functions

$\mathcal{S}_{2,0}$ - transcendental functions with two critical values ± 1 and no asymptotic values

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh'(z) = \frac{e^z - e^{-z}}{2}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$
$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$
$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z} \text{ (critical points)}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z} \text{ (critical points)}$$

critical values: ± 1

Put in somewhere notion of **conformally balanced**
are inverse images in $\mathcal{S}_{2,0}$ always trees?
I am adding a new line to see if GitHub sees the change.

References



Chris Bishop (2014)

Constructing Entire Functions By Quasiconformal Folding

Acta Mathematica



Nuria Fagella, Sebastien Godillion, and Xavier Jarque (2014)

Wandering domains for composition of entire functions

Journal of Mathematical Analysis and Applications