

# Constructing Entire Functions (a summary)

Kirill Lazebnik

SUNY Stony Brook

*Kirill.Lazebnik@stonybrook.edu*

August 4, 2015

# Constructing Entire Functions By Quasiconformal Folding (a summary)

Kirill Lazebnik

SUNY Stony Brook

*Kirill.Lazebnik@stonybrook.edu*

August 4, 2015

$$p(z) = \frac{z^3}{2} - \frac{3z}{2}$$

$$p(z) = \frac{z^3}{2} - \frac{3z}{2}$$


$$p'(z) = \frac{3}{2}(z-1)(z+1)$$

$$p(z) = \frac{z^3}{2} - \frac{3z}{2} \text{ (two critical values } \pm 1)$$

$$p'(z) = \frac{3}{2}(z - 1)(z + 1)$$

$$p(z) = \frac{z^3}{2} - \frac{3z}{2} \text{ (two critical values } \pm 1)$$

$$p'(z) = \frac{3}{2}(z-1)(z+1)$$



$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z$$

$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z$$

$$p'(z) = (z - 1)^2(z + 1)$$




$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z \text{ (two critical values } 5/12, -11/12)$$

$$p'(z) = (z - 1)^2(z + 1)$$

$$p(z) = \frac{z^4}{4} - \frac{z^3}{3} - \frac{z^2}{2} + z \text{ (two critical values } 5/12, -11/12\text{)}$$

$$p'(z) = (z - 1)^2(z + 1)$$



*Shabat* polynomial -

*Shabat* polynomial - only has two critical values  $\pm 1$

*Shabat* polynomial - only has two critical values  $\pm 1$

**Proposition:** For any *Shabat* polynomial  $p(z)$ , it is true that  $p^{-1}[-1, 1]$  is a tree.

*Shabat* polynomial - only has two critical values  $\pm 1$

**Proposition:** For any *Shabat* polynomial  $p(z)$ , it is true that  $p^{-1}[-1, 1]$  is a tree, with  $\deg(p)$  edges.

*Shabat* polynomial - only has two critical values  $\pm 1$

**Proposition:** For any *Shabat* polynomial  $p(z)$ , it is true that  $p^{-1}[-1, 1]$  is a tree, with  $\deg(p)$  edges.

**Theorem (Grothendieck):** ALL combinatorial trees occur as  $p^{-1}[-1, 1]$  for some *Shabat* polynomial  $p(z)$ .

*Shabat* polynomial - only has two critical values  $\pm 1$

**Proposition:** For any *Shabat* polynomial  $p(z)$ , it is true that  $p^{-1}[-1, 1]$  is a tree, with  $\deg(p)$  edges.

**Theorem (Grothendieck):** ALL combinatorial trees occur as  $p^{-1}[-1, 1]$  for some *Shabat* polynomial  $p(z)$ .

**Theorem (Bishop):** Any continua can be  $\epsilon$ -approximated in the Hausdorff metric by some  $p^{-1}[-1, 1]$ .



trees  $\iff$  *Shabat* polynomials

trees  $\iff$  *Shabat* polynomials

infinite trees  $\iff$  Transcendental Functions

trees  $\iff$  *Shabat* polynomials

infinite trees  $\iff$  Subclass of Transcendental Functions

trees  $\iff$  *Shabat* polynomials

infinite trees  $\iff$  Subclass of Transcendental Functions

$\mathcal{S}_{2,0}$  - transcendental functions with two critical values  $\pm 1$  and no asymptotic values

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh'(z) = \frac{e^z - e^{-z}}{2}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$
$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z}$$

$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$
$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z} \text{ (critical points)}$$



$$\cosh(z) := \frac{e^z + e^{-z}}{2}$$

$$\cosh'(z) = \frac{e^z - e^{-z}}{2} = 0 \implies z = \pi i n : n \in \mathbb{Z} \text{ (critical points)}$$

critical values:  $\pm 1$

$T$  - unbounded, locally finite tree

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$

$V$  - the vertices of  $T$ .

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$



$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

(1) The edges of  $T$  are  $C^2$  with uniform bounds.

(2) The angles between adjacent edges are bounded uniformly from zero

(3) Adjacent edges have uniformly comparable lengths

(4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau: \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

## Theorem:

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry*



$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry and every edge has  $\tau$ -size  $\geq \pi$ .*

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry and every edge has  $\tau$ -size  $\geq \pi$ . Then there is an  $r_0 > 0$ , an entire  $f$ , and a  $K$ -quasiconformal  $\phi$*

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry and every edge has  $\tau$ -size  $\geq \pi$ . Then there is an  $r_0 > 0$ , an entire  $f$ , and a  $K$ -quasiconformal  $\phi$  so that  $f \circ \phi = \cosh \circ \tau$  off  $T(r_0)$ .*

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry and every edge has  $\tau$ -size  $\geq \pi$ . Then there is an  $r_0 > 0$ , an entire  $f$ , and a  $K$ -quasiconformal  $\phi$  so that  $f \circ \phi = \cosh \circ \tau$  off  $T(r_0)$ .  $K$  depends only on the bounded geometry constants of  $T$ .*

$T$  - unbounded, locally finite tree, with a bipartite labeling of vertices.

$\Omega_j$  - components of  $\mathbb{C} - T$ .

$\tau : \cup \Omega_j \rightarrow \mathbb{C}$  - the map conformal on each  $\Omega_j$  to  $\mathbb{H}_r$ .

$V$  - the vertices of  $T$ .

$V_j$  - the image of  $V$  under  $\tau$  restricted to  $\Omega_j$ .

For  $r > 0$ , define  $T(r) = \cup_{e \in T} \{z : \text{dist}(z, e) < r \text{diam}(e)\}$

The  $\tau$ -size of edge  $e$  is the minimum length of the two images  $\tau(e)$

$T$  has uniformly bounded geometry if:

- (1) The edges of  $T$  are  $C^2$  with uniform bounds.
- (2) The angles between adjacent edges are bounded uniformly from zero
- (3) Adjacent edges have uniformly comparable lengths
- (4) For non-adjacent edges  $e, f$ , we have  $\text{diam}(e)/\text{dist}(e, f)$  uniformly bounded.

**Theorem:** *Suppose  $T$  has bounded geometry and every edge has  $\tau$ -size  $\geq \pi$ . Then there is an  $r_0 > 0$ , an entire  $f$ , and a  $K$ -quasiconformal  $\phi$  so that  $f \circ \phi = \cosh \circ \tau$  off  $T(r_0)$ .  $K$  depends only on the bounded geometry constants of  $T$ . The only critical values of  $f$  are  $\pm 1$  and  $f$  has no asymptotic values.*

$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f^{\circ n}$  is normal in an open set  $U$

$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f^{\circ n}$  is normal in an open set  $U$  if every sequence of  $f^{\circ k}$  contains a further subsequence converging locally uniformly to a holomorphic function  $g : U \rightarrow \overline{\mathbb{C}}$



$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f^{\circ n}$  is normal in an open set  $U$  if every sequence of  $f^{\circ k}$  contains a further subsequence converging locally uniformly to a holomorphic function  $g : U \rightarrow \overline{\mathbb{C}}$

The Fatou set of  $f$  is the set of points  $z \in \mathbb{C}$  for which  $f$  is normal in some neighborhood of  $z$ .

$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f^{\circ n}$  is normal in an open set  $U$  if every sequence of  $f^{\circ k}$  contains a further subsequence converging locally uniformly to a holomorphic function  $g : U \rightarrow \overline{\mathbb{C}}$

The Fatou set of  $f$  is the set of points  $z \in \mathbb{C}$  for which  $f$  is normal in some neighborhood of  $z$ . The components of the Fatou set are called Fatou components.

$f : \mathbb{C} \rightarrow \mathbb{C}$  entire function

$f^{\circ n}$  is normal in an open set  $U$  if every sequence of  $f^{\circ k}$  contains a further subsequence converging locally uniformly to a holomorphic function  $g : U \rightarrow \overline{\mathbb{C}}$

The Fatou set of  $f$  is the set of points  $z \in \mathbb{C}$  for which  $f$  is normal in some neighborhood of  $z$ . The components of the Fatou set are called Fatou components.

A Fatou component  $U$  is called *wandering* if  $f^n(U) \cap f^m(U) = \emptyset$  for all  $n, m \in \mathbb{N}, n \neq m$

**Theorem:** *(Sullivan) Rational maps don't have wandering domains.*

**Theorem:** *(Sullivan) Rational maps don't have wandering domains.*

For  $f : \mathbb{C} \rightarrow \mathbb{C}$ , the singular set  $S(f)$  consists of the critical values and asymptotic values of  $f$ .

**Theorem:** *(Sullivan) Rational maps don't have wandering domains.*

For  $f : \mathbb{C} \rightarrow \mathbb{C}$ , the singular set  $S(f)$  consists of the critical values and asymptotic values of  $f$ .

The *Speiser* class  $\mathcal{S}$  consists of those transcendental functions for which  $S(f)$  is finite.

**Theorem:** *(Sullivan) Rational maps don't have wandering domains.*

For  $f : \mathbb{C} \rightarrow \mathbb{C}$ , the singular set  $S(f)$  consists of the critical values and asymptotic values of  $f$ .

The *Speiser* class  $\mathcal{S}$  consists of those transcendental functions for which  $S(f)$  is finite.

**Theorem:** *(Golberg and Keen, Eremenko and Lyubich) Functions in the Speiser Class don't have wandering domains.*

**Theorem:** *(Sullivan) Rational maps don't have wandering domains.*

For  $f : \mathbb{C} \rightarrow \mathbb{C}$ , the singular set  $S(f)$  consists of the critical values and asymptotic values of  $f$ .

The *Speiser* class  $\mathcal{S}$  consists of those transcendental functions for which  $S(f)$  is finite.

**Theorem:** *(Golberg and Keen, Eremenko and Lyubich) Functions in the Speiser Class don't have wandering domains.*

The Eremenko-Lyubich class  $\mathcal{B}$  consists of those transcendental functions with bounded singular set.



$$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ .

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$$z_n = a_n + i\pi$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$$z_n = a_n + i\pi, D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ .

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho$  of the disc so that:



$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

$$(1) \quad \rho_n(z) = z \text{ for } z \in \partial\mathbb{D}$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:**

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$ , there exists a transcendental  $f$  and a quasiconformal  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  so that:

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$ , there exists a transcendental  $f$  and a quasiconformal  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  so that:

$$(1) f(\bar{z}) = \overline{f(z)}, f(-z) = f(z)$$



$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left[ \frac{\lambda}{\pi} \cosh(n\pi) \right] \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$ , there exists a transcendental  $f$  and a quasiconformal  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  so that:

- (1)  $f(\bar{z}) = \overline{f(z)}, f(-z) = f(z)$
- (2)

$$f(z) = \begin{cases} \cosh(\lambda \sinh(\phi(z))) & \text{if } \phi(z) \in S^+ \\ \rho_n((\phi(z) - z_n)^{d_n}) & \text{if } \phi(z) \in D_n \end{cases}$$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$ , there exists a transcendental  $f$  and a quasiconformal  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  so that:

- (1)  $f(\bar{z}) = \overline{f(z)}, f(-z) = f(z)$
- (2)

$$f(z) = \begin{cases} \cosh(\lambda \sinh(\phi(z))) & \text{if } \phi(z) \in S^+ \\ \rho_n((\phi(z) - z_n)^{d_n}) & \text{if } \phi(z) \in D_n \end{cases}$$

(3)  $f$  has no asymptotic values and its set of critical values is  $\pm 1$  and  $\overline{\{w_n : n \geq 1\}}$

$S^+ = \{x + iy : x > 0, |y| < \pi/2\}$  is mapped conformally to  $\mathbb{H}_r$  by  $\lambda \cdot \sinh$ . Then holomorphically to  $\mathbb{C} - [-1, 1]$  by  $\cosh$ .

$$a_n = \cosh^{-1} \left( \frac{\pi}{\lambda} \left\lfloor \frac{\lambda}{\pi} \cosh(n\pi) \right\rfloor \right)$$

$z_n = a_n + i\pi$ ,  $D_n = \{z \in \mathbb{C} : |z - z_n| < 1\}$  is mapped holomorphically to  $|z| < 1$  by  $z \rightarrow (z - z_n)^{d_n}$ . Then by a quasiconformal map  $\rho_n$  of the disc so that:

- (1)  $\rho_n(z) = z$  for  $z \in \partial\mathbb{D}$
- (2)  $\rho_n(0) = w_n$  where  $w_n$  is a point near  $1/2$ .
- (3)  $\rho_n$  is conformal on  $\frac{3}{4}\mathbb{D}$
- (4)  $\rho_n$  is quasiconformal on  $\mathbb{D}$ .

**Theorem:** For every choice of parameters  $\lambda, (d_n), (w_n)$  with  $\lambda \in \pi\mathbb{N}, d_n \in 2\mathbb{N}$ , there exists a transcendental  $f$  and a quasiconformal  $\phi : \mathbb{C} \rightarrow \mathbb{C}$  so that:

- (1)  $f(\bar{z}) = \overline{f(z)}, f(-z) = f(z)$
- (2)

$$f(z) = \begin{cases} \cosh(\lambda \sinh(\phi(z))) & \text{if } \phi(z) \in S^+ \\ \rho_n((\phi(z) - z_n)^{d_n}) & \text{if } \phi(z) \in D_n \end{cases}$$

- (3)  $f$  has no asymptotic values and its set of critical values is  $\pm 1$  and  $\{w_n : n \geq 1\}$
- (4)  $\phi(0) = 0, \phi(\mathbb{R}) = \mathbb{R}$  and  $\phi$  is conformal in  $S^+$ .

# References



Chris Bishop (2014)

Constructing Entire Functions By Quasiconformal Folding

*Acta Mathematica*



Nuria Fagella, Sebastien Godillion, and Xavier Jarque (2014)

Wandering domains for composition of entire functions

*Journal of Mathematical Analysis and Applications*