Please check the examination details below	ow before ente	ring your candidate	information
Candidate surname		Other names	
Centre Number Candidate Nu	ımber		
Pearson Edexcel International Advanced Level			
Monday 16 October 2023			
Morning (Time: 1 hour 30 minutes)	Paper reference	WME	01/01
Mathematics			
International Advanced Subsidiary/Advanced Level			
Mechanics M1			
You must have:			Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \,\mathrm{m \, s^{-2}}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶



P74321A
©2023 Pearson Education Ltd.
Z:1/1/1/



1.




Figure 1

Figure 1 shows a beam AB with weight 24N and length 6m.

The beam is suspended by two light vertical ropes. The ropes are attached to the points C and D on the beam where AC = x metres and CD = 2 m.

The tension in the rope attached to the beam at C is double the tension in the rope attached to the beam at D.

The beam is modelled as a uniform rod, resting horizontally in equilibrium.

Find

- (i) the tension in the rope attached to the beam at D.
- (ii) the value of x.

(5)

(Total for Ouestian 1 is 5 marks)	Question 1 continued		
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion Lis 5 marks)			
(Total for Onestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Ouestion 1 is 5 marks)			
(Total for Question 1 is 5 marks)			
(2000 101 Question 1 10 0 marks)	(Total for Question 1 is 5 marks)		



2.




Figure 2

Two fixed points, A and B, are on a straight horizontal road.

The **acceleration-time** graph in Figure 2 represents the motion of a car travelling along the road as it moves from A to B.

At time t = 0, the car passes through A with speed $8 \,\mathrm{m \, s}^{-1}$

At time t = 20 s, the car passes through B with speed $v \text{ m s}^{-1}$

(a) Show that v = 18

(3)

(b) Sketch a speed-time graph for the motion of the car from A to B.

(3)

(c) Find the distance AB.

(4)

Question 2 continued		



Question 2 continued		

Question 2 continued		
	(Total for Question 2 is 10 marks)	



3.	A hammer is used to hit a tent peg into soft ground.	
	The hammer has mass 1.8 kg and the tent peg has mass 0.2 kg.	
The hammer and tent peg are both modelled as particles and the impact is modelled direct collision.		
	Immediately before the impact, the tent peg is stationary and the hammer is moving vertically downwards with speed $10\mathrm{ms}^{-1}$	
	Immediately after the impact, the hammer and tent peg move together, vertically downwards, with the same speed v m s ⁻¹	
	(a) Find the value of v	(2)
		(2)
	(b) Find the magnitude of the impulse exerted on the tent peg by the hammer, stating the units of your answer.	
		(3)
	The ground exerts a constant vertical resistive force of magnitude R newtons, bringing the hammer and tent peg to rest after they travel a distance of 12cm .	
	(c) Find the value of R.	(5)
		(5)



Question 3 continued		



Question 3 continued		

Question 3 continued		
(Total for Question 3 is 10 mar	ks)	



4. [*In this question* **i** *and* **j** *are horizontal unit vectors directed due east and due north respectively.*]

A particle *P* moves with constant acceleration $(-\lambda \mathbf{i} + 2\lambda \mathbf{j}) \,\mathrm{m\,s}^{-2}$, where λ is a positive constant.

At time t = 0, the velocity of P is $(5\mathbf{i} - 8\mathbf{j}) \,\mathrm{m \, s}^{-1}$

(a) Find the velocity of P when t = 5 s, giving your answer in terms of **i**, **j** and λ .

(2)

The speed of *P* when $t = 5 \,\mathrm{s}$ is $13 \,\mathrm{m \, s}^{-1}$

(b) Show that

$$25\lambda^2 - 42\lambda - 16 = 0$$

(3)

(c) Find the direction of motion of P when t = 4 s, giving your answer as a bearing to the nearest degree.

(5)



Question 4 continued		



Question 4 continued		

Question 4 continued		
	Total for Organism Alic 10 - 1	
	Total for Question 4 is 10 marks)	






Figure 3

A small ring of mass 0.2 kg is attached to one end of a light inextensible string.

The ring is **threaded** onto a fixed rough vertical rod.

The string is taut and makes an angle θ with the rod, as shown in Figure 3, where $\tan \theta = \frac{12}{5}$

Given that the ring is in equilibrium and that the tension in the string is 10 N,

(a) find the magnitude of the frictional force acting on the ring,

(3)

(b) state the direction of the frictional force acting on the ring.

(1)

The coefficient of friction between the ring and the rod is $\frac{1}{4}$

Given that the ring is in equilibrium, and that the tension in the string, T newtons, can now vary,

- (c) (i) find the minimum value of T
 - (ii) find the maximum value of T

(8)

Question 5 continued



Question 5 continued

Question 5 continued	
	Total for Question 5 is 12 marks)



6. [In this question **i** and **j** are horizontal unit vectors directed due east and due north respectively and position vectors are given relative to a fixed origin O.]

At 12:00, a ship P sets sail from a harbour with position vector (15i + 36j)km.

At 12:30, P is at the point with position vector $(20\mathbf{i} + 34\mathbf{j})$ km.

Given that P moves with constant velocity,

(a) show that the velocity of P is $(10\mathbf{i} - 4\mathbf{j}) \,\mathrm{km} \,\mathrm{h}^{-1}$

(2)

At time t hours after 12:00, the position vector of P is \mathbf{p} km.

(b) Find an expression for \mathbf{p} in terms of \mathbf{i} , \mathbf{j} and t.

(2)

A second ship Q is also travelling at a constant velocity.

At time t hours after 12:00, the position vector of Q is given by \mathbf{q} km, where

$$\mathbf{q} = (42 - 8t)\mathbf{i} + (9 + 14t)\mathbf{j}$$

Ships P and Q are modelled as particles.

If both ships maintained their course,

- (c) (i) verify that they would collide at 13:30
 - (ii) find the position vector of the point at which the collision would occur.

(4)

At 12:30 Q changes speed and direction to avoid the collision.

Ship Q now travels due north with a constant speed of 15 km h⁻¹

Ship *P* maintains the same constant velocity throughout.

(d) Find the exact distance between P and Q at 14:30

(7)

Question 6 continued	



Question 6 continued

Question 6 continued	
(Total	for Question 6 is 15 marks)






Figure 4

Figure 4 shows a block A of mass m held at rest on a rough plane.

The plane is inclined at an angle α to the horizontal and the coefficient of friction between the block and the plane is μ .

One end of a light inextensible string is now attached to A. The string passes over a small smooth pulley which is fixed at the top of the plane.

The other end of the string is attached to a block *B* of mass *km*.

Block *B* hangs vertically below the pulley, with the string taut.

The string from A to the pulley lies along a line of greatest slope of the plane.

Both A and B are modelled as particles.

When the system is released from rest, A moves up the plane and the tension in the string is $\frac{4mg}{3}$

Given that $\mu = \frac{1}{3}$ and $\tan \alpha = \frac{7}{24}$

- (a) (i) find the magnitude of the acceleration of A, giving your answer in terms of g,
 - (ii) find the value of k.

(9)

(b) Find the magnitude of the resultant force exerted on the pulley by the string, giving your answer in terms of m and g.

(4)

Question 7 continued



Question 7 continued

Question 7 continued



Question 7 continued	
	(Total for Question 7 is 13 marks)
7	TOTAL FOR PAPER IS 75 MARKS

