



Mark Scheme (Results)

January 2022

Pearson Edexcel International Advanced
Level in Physics (WPH14) Paper 01 Physics
Further Mechanics, Fields and Particles



Question Number	Answer	Mark
1	The only correct answer is D <i>A is not correct because 8 is the number of protons</i> <i>B is not correct because 8 is the number of protons</i> <i>C is not correct because 8 is the number of protons</i>	1
2	The only correct answer is D <i>A is not correct because this is the mass of an electron</i> <i>B is not correct because this is the mass of an electron</i> <i>C is not correct because this is the charge of a proton</i>	1
3	The only correct answer is A <i>B is not correct because $p^2/2m = 0.5 \text{ J}$</i> <i>C is not correct because $p^2/2m = 0.5 \text{ J}$</i> <i>D is not correct because $p^2/2m = 0.5 \text{ J}$</i>	1
4	The only correct answer is B <i>A is not correct because momentum is conserved and E_k decreases</i> <i>C is not correct because momentum is conserved and E_k decreases</i> <i>D is not correct because momentum is conserved and E_k decreases</i>	1
5	The only correct answer is D <i>A is not correct because the process described is thermionic emission</i> <i>B is not correct because the process described is thermionic emission</i> <i>C is not correct because the process described is thermionic emission</i>	1
6	The only correct answer is B <i>A is not correct because this would have no effect on the deflection</i> <i>C is not correct because this would increase the deflection</i> <i>D is not correct because this would increase the deflection</i>	1
7	The only correct answer is A <i>B is not correct because force is out of page on QR</i> <i>C is not correct because force is zero on RS</i> <i>D is not correct because force is into page on SP</i>	1
8	The only correct answer is C <i>A is not correct because this was a correct conclusion</i> <i>B is not correct because this was a correct conclusion</i> <i>D is not correct because this was a correct conclusion</i>	1
9	The only correct answer is D <i>A is not correct because this would not change the energy</i> <i>B is not correct because this would not change the energy for one revolution</i> <i>C is not correct because this would not change the energy</i>	1
10	The only correct answer is B <i>A is not correct because if the time of orbit stays constant and the radius of orbit increases then the speed must increase</i> <i>C is not correct because if the time of orbit stays constant and the radius of orbit increases then the speed must increase</i> <i>D is not correct because the angular velocity stays constant</i>	1

Question Number	Answer	Mark
11	at least 4 radial straight lines (1)	3
	distributed equally (1)	
	arrow on at least one line pointing towards centre (1)	
	Total for question 11	3

Question Number	Answer	Mark
12	The blades exert a downward force on the air (1)	4
	The air exerts an equal upwards force on the blades/helicopter Or By Newton's 3 rd law there is an equal upwards force (1)	
	This upwards force equals the weight of helicopter (1)	
	The resultant force is zero, so (by Newton's 1 st or 2 nd law) there is no acceleration (and the helicopter maintains a constant height) (1)	
	Total for question 12	4

Question Number	Answer	Mark
13ai	<p>Use of trigonometry appropriate for determination of angle (1)</p> <p>Use of $W = mg$ (1)</p> <p>Resolves tension in thread vertically or horizontally Or draw triangle of forces (1)</p> <p>Force of repulsion = 1.2×10^{-3} (N) (1)</p> <p><u>Example of calculation</u></p> <p>$\sin \theta = 13/122$ Angle of thread to vertical $\theta = 6.12^\circ$</p> <p>$T \cos 6.12^\circ = 1.1 \times 10^{-3} \text{ kg} \times 9.81 \text{ N kg}^{-1}$</p> <p>Tension in thread = 0.0109 N</p> <p>Force of repulsion = $0.0109 \sin 6.12^\circ = 1.16 \times 10^{-3}$ N</p>	4
13aii	<p>Use of $F = \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2}$ (accept use of $F = \frac{k Q_1 Q_2}{r^2}$) (1)</p> <p>$Q = 1.7 \times 10^{-7}$ (C) (allow ecf from ai) (1)</p> <p><u>Example of calculation</u></p> <p>$1.16 \times 10^{-3} \text{ N} = 8.99 \times 10^9 \text{ Nm}^2 \text{C}^{-2} \frac{Q^2}{0.47^2 \text{ m}^2}$</p> <p>$Q = 1.69 \times 10^{-7} \text{ C}$</p>	2
13b	<p>Use of $V = \frac{Q}{4\pi\epsilon_0 r}$ (accept use of $V = \frac{kQ}{r}$) (1)</p> <p>$V = 5100 \text{ V}$ (allow ecf from aii) (1)</p> <p><u>Example of calculation</u></p> <p>$V = 8.99 \times 10^9 \text{ Nm}^2 \text{C}^{-2} \frac{(-)1.7 \times 10^{-7} \text{ C}}{0.30 \text{ m}} = (-)5094 \text{ V}$</p>	2
Total for question 13		8

Question Number	Answer	Mark
14a	Use of $I = V / R$ (1)	2
	$I = 0.15 \text{ mA}$ which is consistent (with the value on the graph) (1)	
	<u>Example of calculation</u>	
	$I = 5.0 \text{ V} / 33 \text{ k}\Omega = 1.5 \times 10^{-4} \text{ A} = 0.15 \text{ mA}$	
14b	The current would vary with time in the same way as on ammeter A_1 (1)	2
	Because (current is same everywhere) in a series circuit (1)	
14c	Either	3
	Takes two corresponding values of I and t from graph (1)	
	Use of $\ln I = \ln I_0 - t/RC$ (1)	
	$C = 2.27 \times 10^{-4} \text{ F}$ ($2.0 \times 10^{-4} \text{ F} - 2.3 \times 10^{-4} \text{ F}$) (1)	
	Or	
	Draws initial tangent to curve and determines t intercept: T (0.65-0.75 s) (1)	
	Use of $T = RC$ (1)	
	$C = 2.2 \times 10^{-4} \text{ F}$ ($2.0 \times 10^{-4} \text{ F} - 2.3 \times 10^{-4} \text{ F}$) (1)	
	Or	
	Read value of t at which $I = I_0 / e$ (0.56 A, 0.7 s) (1)	
	Use of $T = RC$ (1)	
	$C = 2.1 \times 10^{-4} \text{ F}$ ($2.0 \times 10^{-4} \text{ F} - 2.3 \times 10^{-4} \text{ F}$) (1)	
	<u>Example of calculation</u> eg $I = 0.04 \text{ mA}$ and $t = 10 \text{ s}$	
	$\ln 0.04 = \ln 0.152 - \frac{10\text{s}}{C \times 33\text{k}}$	
	$C = 2.27 \times 10^{-4} \text{ F}$ range: $2.0 \times 10^{-4} \text{ F} - 2.3 \times 10^{-4} \text{ F}$	
14d	Attempt to determine an area under the curve (1)	2
	$Q = 1.1 \times 10^{-3} \text{ C}$ ($1.0 \times 10^{-3} \text{ C}$ to $1.2 \times 10^{-3} \text{ C}$) (1)	
	Or	
	Use of $Q = CV$ with 5.0 V (1)	
14e	$Q = 1.1 \times 10^{-3} \text{ C}$ (allow ecf from (c)) (1)	2
	Use of $W = \frac{QV}{2}$ or $W = \frac{1}{2} CV^2$ or $W = Q^2/2C$ (1)	
	$W = 2.8 \times 10^{-3} \text{ J}$ (allow ecf from 14c and 14d) (1)	
	<u>Example of calculation</u> $W = 1.1 \times 10^{-3} \text{ C} \times 5 \text{ V} / 2 = 2.8 \times 10^{-3} \text{ J}$	
Total for question 14		11

Question Number	Answer	Mark	
15a	<p>Correct vector diagram showing velocity change</p> <p>(Small angle, so) $\delta\theta = \delta v / v$</p> <p>Use of $\delta\theta / \delta t = \omega$ and $v = r\omega$</p> <p>Use of $\delta v / \delta t = a$</p> <p>Algebra to show $a = v^2 / r$</p> <p><u>Example of derivation</u></p> <div></div> <div></div> <p>Small angle, so $\delta\theta = \delta v / v$ $\delta\theta = \omega\delta t$ $\delta\theta = v\delta t / r$ $v\delta t / r = \delta v / v$ $\delta v / \delta t = v^2 / r$</p>	(1) (1) (1) (1) (1) (1)	5
15b	<p>Use of velocity $= f \times 2\pi r$</p> <p>Use of $a = v^2 / r$</p> <p>$a = 39 \text{ m s}^{-2}$</p> <p>Or</p> <p>Use of $\omega = 2\pi f$</p> <p>Use of $a = \omega^2 / r$</p> <p>$a = 39 \text{ m s}^{-2}$</p> <p><u>Example of calculation</u></p> <p>$v = 1.3 \text{ s}^{-1} \times 2\pi \times 0.58 \text{ m} = 4.74 \text{ m s}^{-1}$</p> <p>$a = 4.74^2 (\text{m s}^{-1})^2 / 0.58 \text{ m} = 38.7 \text{ m s}^{-2}$</p>	(1) (1) (1)	3
15c	<p>Tension in cord is force on hand</p> <p>Centripetal force is constant</p> <p>Weight of ball is added to tension at top</p> <p>Weight is subtracted from tension at bottom so force on hand varies and child correct</p>	(1) (1) (1) (1)	4

	<p>OR</p> <p>Tension in cord is force on hand (1)</p> <p>Centripetal force is constant (1)</p> <p>Weight of ball is subtracted from tension at bottom (1)</p> <p>Weight is added to tension at top so force on hand varies and child correct (1)</p> <p><u>Example of discussion</u></p> <p>At top of motion $W + T = \text{centripetal force}$</p> <p>Or at bottom of motion $T - W = \text{centripetal force}$</p> <p>So T varies from (centripetal force $- W$) to (centripetal force $+ W$)</p>	
	Total for question 15	12

Question Number	Answer	Mark																																
16a	(the particle is) ionising Or it knocks electrons out of atoms in its path (1) A track is formed by the ionised particles produced (1)	2																																
16bi	A baryon is three quarks (or three antiquarks) (1) A meson is an anti-quark and a quark (1)	2																																
16bii	into the page/paper (1)	1																																
16biii	charge: applies charge conservation (1) omega baryon charge −1 (1) baryon number: before = 1 identified as the proton so the omega particle = 1 (1)	3																																
*16c	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table><tr><th>Number of indicative marking points seen in answer</th><th>Number of marks awarded for indicative marking points</th><th>Max linkage mark available</th><th>Max final mark</th></tr><tr><td>6</td><td>4</td><td>2</td><td>6</td></tr><tr><td>5</td><td>3</td><td>2</td><td>5</td></tr><tr><td>4</td><td>3</td><td>1</td><td>4</td></tr><tr><td>3</td><td>2</td><td>1</td><td>3</td></tr><tr><td>2</td><td>2</td><td>0</td><td>2</td></tr><tr><td>1</td><td>1</td><td>0</td><td>1</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td></tr></table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	Max linkage mark available	Max final mark	6	4	2	6	5	3	2	5	4	3	1	4	3	2	1	3	2	2	0	2	1	1	0	1	0	0	0	0	6
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	Max linkage mark available	Max final mark																															
6	4	2	6																															
5	3	2	5																															
4	3	1	4																															
3	2	1	3																															
2	2	0	2																															
1	1	0	1																															
0	0	0	0																															

The following table shows how the marks should be awarded for structure and lines of reasoning.

	Number of marks awarded for structure of answer and sustained line of reasoning
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout	2
Answer is partially structured with some linkages and lines of reasoning	1
Answer has no linkages between points and is unstructured	0

Guidance on how the mark scheme should be applied: The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points which is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning). If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).

Indicative content:

Energy:

IC1 As (Rest) mass-energy of proton and kaon + Initial $E_k =$ (rest) mass-energy of omega and kaon + kinetic energies of both particles

Or (Total) mass-energy conserved

IC2 Incoming K^- had **high** kinetic energy

IC3 some of this initial kinetic energy converted to mass of the omega particle (– mass of proton)

IC4 $\Delta E = \Delta mc^2$


momentum:

IC5 momentum of $K^- =$ sum of x components of $K^+ + \Omega^-$

Or vector sum of momentum of $K^+ + \Omega^- =$ momentum of K^-

Or an attempt to sketch a triangle of vectors

eg



IC6 y component of K^+ equals y component Ω^-

Or

all vectors correctly labelled

Total for question 16

14

Question Number	Answer	Mark
17a	It is a lepton (1)	2
	It is a fundamental particle Or second generation (1)	
17b	negative pion (1)	2
	muon anti-neutrino (1)	
17c	Use of $\Delta E = c^2 \Delta m$ (1)	3
	Divide mass of muon by mass of electron Or multiplies the mass of an electron by 200 (1)	
	Mass of muon is 207 times that of an electron so true statement Or 200 times electron mass is $1.82 \times 10^{-28} \text{ kg}$ which is just under $1.88 \times 10^{-28} \text{ kg}$ so it is correct (1)	
	<u>Example of calculation</u>	
	$\text{mass} = \frac{106 \text{ MeV}/c^2 \times 10^6 \times 1.6 \times 10^{-19} \text{ J eV}^{-1}}{(3 \times 10^8)^2 (\text{m s}^{-1})^2} = 1.88 \times 10^{-28} \text{ kg}$ $\text{mass} = \frac{1.88 \times 10^{-28} \text{ kg}}{9.11 \times 10^{-31} \text{ kg}}$ <p>mass = 207 times that of an electron</p>	
17di	A unit of energy is GeV Or a unit of mass is GeV/c^2 (1)	2
	The unit of momentum is the same as the unit of energy/velocity so GeV divided by $c = \text{GeV}/c$ Or The unit of momentum is the same as the unit of mass \times velocity so $\text{GeV}/c^2 \times c = \text{GeV}/c$ (1)	
17dii	Use of Circumference = $2\pi r$ (1)	3
	Use of $r = p / BQ$ (1)	
	Show that a momentum of $1.65 \times 10^{-18} \text{ N s}$ is consistent with the correct radius (7.11 m) by determination of p , r , B or Q and statement that it is correct (1)	
	<u>Example of calculation</u>	
	$r = 44.7 \text{ m} / 2\pi = 7.11 \text{ m}$ $r = \frac{1.65 \times 10^{-18} \text{ N s}}{1.45 \text{ T} \times 1.6 \times 10^{-19} \text{ C}} = 7.11 \text{ m}$	
17diii	muons travelling close to speed of light (1)	2
	relativistic effect on particle lifetime (1)	
Total for question 17		14

Question Number	Answer	Mark
18a	Flux linkage (1) weber / Wb (1)	2
18b	Evidence of attempt to determine maximum gradient of graph (1) Use of $\Delta B / \Delta t$ (1) use of area of coil = πr^2 (1) use of $\Phi = BA$ (1) Use of $\mathcal{E} = \frac{d(N\Phi)}{dt}$ (1) $\mathcal{E} = 1.4 \text{ V}$ range 1.2 to 1.6 V (1) <u>Example of calculation</u> gradient = $0.6 \text{ T} / 0.006 \text{ s} = 100 \text{ T s}^{-1}$ Area of coil = $\pi(0.003 \text{ m})^2 = 2.83 \times 10^{-5} \text{ m}^2$ $\mathcal{E} = 500 \times 100 \text{ T s}^{-1} \times 2.83 \times 10^{-5} \text{ m}^2$ $\mathcal{E} = 1.4 \text{ V}$	6
18c	There is a change in the magnetic flux (linkage with aluminium disc) (1) Or disc is cutting magnetic field/flux So an <u>e.m.f.</u> is <u>induced</u> (1) Leads to a current (in the disc) (accept eddy current) (1) Force acts on the disc, as there is a current in a magnetic field (accept reference to motor effect, FLHR or $F = BIl$ if current in disc has been mentioned) Or field due to current in disc interacting with field due to magnet to cause force on disc (1) According to Lenz's law Or the direction of e.m.f./current is such to oppose (the cause of) the change in flux (1) The disc moves to reduce this change (the same direction as the magnet) so correct suggestion (1)	6
Total for question 18		14