Please check the examination details belo	ow before ente	ering your candidate information
Candidate surname		Other names
Centre Number Candidate Nu	ımber	
Pearson Edexcel Inter	nation	al Advanced Level
Tuesday 24 October	2023	
Afternoon (Time: 1 hour 45 minutes)	Paper reference	WPH15/01
Physics		• •
International Advanced Le	evel	
UNIT 5: Thermodynamics	, Radiat	ion, Oscillations
and Cosmology	•	
and cosmology		
(V. 1)		
You must have:		Total Marks
Scientific calculator, ruler		

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Show all your working out in calculations and include units where appropriate.

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- The list of data, formulae and relationships is printed at the end of this booklet.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over



P75625A ©2023 Pearson Education Ltd. 2:1/1/



SECTION A

Answer ALL questions.

For questions 1–10, in Section A, select one answer from A to D and put a cross in the box \boxtimes . If you change your mind, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

1 Fusion reactions are the primary energy source of stars.

Which row of the table gives the conditions necessary for fusion in stars?

		Density	Temperature
X	A	high	high
X	В	high	low
X	C	low	high
X	D	low	low

(Total for Question 1 = 1 mark)

2 Astronomers use stellar parallax to determine the distances to nearby stars.

Which of the following must be known to use stellar parallax?

- A distance between the Earth and the Sun
- **B** intensity of radiation received from the star
- C luminosity of the Sun
- **D** value of the Hubble constant

(Total for Question 2 = 1 mark)

- Which of the following statements about nuclear binding energy (B.E.) is correct?
 - A B.E. per nucleon increases steadily as proton number increases.
 - B B.E. per nucleon decreases steadily as proton number increases.
 - B.E. per nucleon is approximately constant for all elements.
 - **D** B.E. per nucleon increases to a maximum and then decreases as proton number increases.

(Total for Question 3 = 1 mark)

4 A mass suspended from a spring is oscillating vertically. The oscillations are damped.

Which of the following statements is correct?

- A The damping force is always in the opposite direction to the acceleration.
- **B** The damping force is always in the opposite direction to the velocity.
- C The damping force is always in the same direction as the acceleration.
- **D** The damping force is always in the same direction as the velocity.

(Total for Question 4 = 1 mark)

5 A star is moving away from the Earth.

The wavelengths of spectral lines in the light received from the star are measured. These wavelengths are not the same as the wavelengths of the same lines measured from a source in the laboratory.

Which of the following explains the difference in the wavelengths?

- A The star emits light with a shorter wavelength.
- **B** The star emits light with a longer wavelength.
- C The light arrives at the Earth with a shorter wavelength.
- **D** The light arrives at the Earth with a longer wavelength.

(Total for Question 5 = 1 mark)

6 The Moon is gradually moving away from the Earth.

The gravitational potential at the Earth due to the Moon, and the gravitational force between the Earth and the Moon, are both changing.

Which row of the table is correct?

		Gravitational potential	Gravitational force
X	A	decreases	decreases
X	В	increases	decreases
X	C	decreases	increases
X	D	increases	increases

(Total for Question 6 = 1 mark)



A small satellite of mass 122 kg is orbiting the Earth. The satellite is orbiting at a height above the surface of the Earth equal to the radius of the Earth.

Which of the following is the approximate weight of the satellite in its orbit?

- A zero
- **■ B** 300 N
- **■ D** 1200 N

X

X


(Total for Question 7 = 1 mark)

8 Which row of the table gives the relative ionising power and penetration of gamma radiation?

	Ionising power	Penetration
A	low	low
В	low	high
C	high	low
D	high	high

(Total for Question 8 = 1 mark)

9 The Hertzsprung-Russell diagram shows four areas P, Q, R and S, where different types of stars are located.



Which of the following is a possible evolutionary path for stars initially in area **P**?

- \square A $P \rightarrow Q \rightarrow S$
- \square B P \rightarrow S \rightarrow Q
- \square C $P \rightarrow R \rightarrow S$
- \square D P \rightarrow S \rightarrow R

(Total for Question 9 = 1 mark)

10 A simple pendulum is displaced through a small angle then released. The frequency of oscillation is determined.

The length of the pendulum is then doubled.

By what factor will the frequency of oscillation be multiplied?

- \triangle A $\frac{1}{\sqrt{2}}$
- \boxtimes B $\frac{1}{2}$
- \square C $\sqrt{2}$
- **■ D** 2

(Total for Question 10 = 1 mark)

TOTAL FOR SECTION A = 10 MARKS

SECTION B

Answer ALL questions in the spaces provided.

11 Air is used to inflate a balloon.

The inflated balloon has a volume of $1.77 \times 10^{-3}\,\mathrm{m}^3$ and contains 5.15×10^{22} air molecules. The air inside the balloon exerts a pressure of $1.15 \times 10^5\,\mathrm{Pa}$.


Calculate the average kinetic energy of the air molecules.

Average kinetic energy of air molecules =

(Total for Question 11 = 3 marks)

12 A student investigated the relationship between the pressure and volume of a fixed mass of gas at constant temperature.

He used his data to plot the following graph.



The student stated that his graph demonstrated that the pressure exerted by the gas was inversely proportional to the volume occupied by the gas.

Assess the validity of the student's statement.

(Total for Question 12 = 4 marks)

13 In the early part of the 20th century, Rutherford made the first observation of an element being changed into a different element.

Alpha particles were fired at nitrogen atoms.

The nuclear equation for this reaction is:

$${}_{2}^{4}\text{He} + {}_{7}^{14}\text{N} \rightarrow {}_{8}^{17}\text{O} + {}_{1}^{1}\text{H}$$

(a) Calculate the minimum energy, in MeV, required for the reaction to take place.

(4)

Nuclide	Mass / 10 ⁻²⁷ kg
¹⁷ O	28.2185
¹⁴ N	23.2451
⁴ He	6.64432
¹ H	1.67299

(b) Explain why the alpha particle must have an energy greater than the minimum energy for the reaction to take place.

/	100	. 1
1	1	

(Total for Question 13 = 6 marks)

Minimum energy =

- 14 At the beginning of the 20th century, astronomers identified that a number of galaxies were moving away from the Earth.
 - (a) State how astronomers knew that galaxies were moving away from the Earth.


(1)

(b) Hubble suggested that the recessional velocity v of a galaxy was related to the distance d of the galaxy from the Earth by the expression

$$v = H_0 d$$

where H_0 is a constant.

The graph shows some of Hubble's data.



Assess the validity of Hubble's expression based on the data shown.

(3)

|
 |
|------|------|------|------|------|------|------|------|------|
| | | | | | | | | |
|
 |
| | | | | | | | | |
|
 |
| | | | | | | | | |
|
 |
| | | | | | | | | |
| | | | | | | | | |
|
 |
| | | | | | | | | |
|
 |
| | | | | | | | | |

(Total for Question 14 = 4 marks)

'15	A student is planning an experiment to investigate how the resistance of a thermistor changes with temperature. The student wants to use the temperature range 0°C to 100°C.
	Describe a method the student could use to determine values of resistance and temperature over this range.
	You should include one precaution the student should take to ensure accurate measurements.
	You do not need to draw a circuit diagram.

	(Total for Question 15 = 6 marks)
_	(10mi 101 Question 15 0 marks)

is 22°C.					
The kettle is switched on. The water takes 85 s to reach a temperature of 100°C.					
(a) Show that the power of the kettle is about 1600 W.					
specific heat capacity of water = $4190 \mathrm{Jkg^{-1}K^{-1}}$ density of water = $998 \mathrm{kgm^{-3}}$	(4)				
(b) The water boils at 100°C.					
Calculate the time now taken for 75% of the water to boil away.					
specific latent heat of vaporisation of water = $2.26 \times 10^6 \mathrm{Jkg^{-1}}$					
	(3)				

Time taken =

(Total for Question 16 = 7 marks)



- 17 Makemake is a dwarf planet in the solar system. Makemake has a mass of $3.1 \times 10^{21} \, kg$ and a radius of $715 \, km$.
 - (a) Calculate the gravitational field strength at the surface of Makemake.

(2)

Gravitational field strength at surface =

(b) The average distance of Makemake from the Sun is similar to the average distance of the dwarf planet Pluto from the Sun.

A website states:

"The time taken by Makemake to complete one orbit of the Sun is 20% greater than the time taken by Pluto to complete one orbit of the Sun."

Assess the accuracy of the website statement.

mass of Sun =
$$1.99 \times 10^{30}$$
 kg
orbit radius of Makemake = 6.80×10^{12} m
orbit time of Pluto = 248 years
 $1 \text{ year} = 3.15 \times 10^{7}$ s

(Total for Question 17 = 8 marks)



(6)

18	In January 2023,	the asteroid	2023 BU	came o	close to	the Ea	arth. T	The closest	distance of
	the asteroid from	the surface	of the Ear	rth was	3590 km	n.			

(a)	Calculate the force between the	he asteroid	and th	e Earth	at the	e closest	distance.	You
	may assume the asteroid is a	sphere.						

asteroid diameter = $5.65 \, m$ asteroid density = $1950 \, kg \, m^{-3}$ mass of Earth = $5.98 \times 10^{24} \, kg$ radius of Earth = $6380 \, km$

	(4)
Force between asteroid and Earth =	
(b) Calculate the change in gravitational potential energy of the asteroid if it had falles to the surface of the Earth from a height of 3590 km.	
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled	
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n
(b) Calculate the change in gravitational potential energy of the asteroid if it had falled to the surface of the Earth from a height of 3590 km.	n

Change in gravitational potential energy =



Explain why the mass of the asteroid would not remain constant as the asteroid fell to the surface of the Earth.	
to the surface of the Earth.	(2)
(Total for Question 18 = 9 mar	rks)

19 In 2023, a small capsule of caesium-137 was lost when being transported from a mine in Western Australia.

The activity of the caesium in the capsule was 19 GBq.

(a) (i) Show that the decay constant for caesium-137 is about $7.3 \times 10^{-10} \, \mathrm{s}^{-1}$.

half-life of caesium-137 = 30.1 years
1 year =
$$3.15 \times 10^7$$
 s

(2)

(ii) Calculate the mass of caesium in the capsule.

/	1	ь.	
1	٠.,	ď.	


3. /	C			

Mass of caesium =

(iii) Calculate the activity of the caesium in the capsule after 2 years.	(2)
Activity after 2 years =	
(b) The capsule was found after 14 days.	
Calculate the total energy, in J, released from the capsule in 14 days.	
activity of caesium in capsule = 19 GBq energy released in each decay of caesium-137 = 1.17 MeV 1 day = 86 400 s	
1 day 00 4003	(3)
Total energy released in 14 days =	J
(Total for Question 19 =	
((((((((((((((((((((,



20 A 150 g mass was hung from a spring as shown. The spring extended by 7.5 cm.



The mass was displaced vertically 3.0 cm from its equilibrium position and then released. The mass oscillated vertically.

(a) Explain why the motion of the mass was simple harmonic motion.

(2)

(b) Calculate the maximum velocity of the oscillating mass.

(6)

Maximum velocity of mass =



(c) Explain why the maximum velocity of the oscillating mass decreased over time.	(2)
(Total for Question 20 = 10 ma	ırks)

- 21 During their life cycle, stars go through a number of stages.
 - (a) Betelgeuse is a red giant star with a luminosity of 4.49×10^{31} W.

The peak intensity of radiation from Betelgeuse occurs at a wavelength of 850 nm.

(i) A website claims that the radius of Betelgeuse is 1000 times the radius of the Sun.

Assess the accuracy of this claim.


surface temperature of Sun = $5800 \, \text{K}$ luminosity of Sun = $3.83 \times 10^{26} \, \text{W}$

(4)

(2)

(ii) The Hertzsprung-Russell diagram shows the positions of some main sequence stars.

Add the approximate positions of the Sun and Betelgeuse to the diagram. Mark the positions with labelled crosses.





(iii) State what is meant by a main sequence star.	(1)
b) The final stage of evolution for a massive star may be a rotating neutron star.	
One neutron star rotates with a period of 33.5 ms.	
As this star rotates, one side of the star moves towards the Earth while the other of the star moves away from the Earth. This causes a range of wavelengths to be received by an observer.	
Ultraviolet radiation of wavelength 91.2 nm is emitted by the star.	
Calculate the maximum and minimum wavelengths observed.	
diameter of star = $20.5 \mathrm{km}$	
	(6)
Maximum wavelength observed =	
Minimum wavelength observed =	
	marks)

TOTAL FOR SECTION B = 80 MARKS TOTAL FOR PAPER = 90 MARKS

