

# Preliminary migrate analysis of *M. californianus*

## MIGRATION RATE AND POPULATION SIZE ESTIMATION

using the coalescent and maximum likelihood or Bayesian inference

Migrate-n version 3.7.2 [April-12-18]

Program started at Tue Jun 1 10:52:05 2021

Program finished at Tue Jun 1 18:19:17 2021



## Options

Datatype:

DNA sequence data

Inheritance scalers in use for Thetas:

All loci use an inheritance scaler of 1.0

[The locus with a scaler of 1.0 used as reference]

Random number seed:

(with internal timer)

1756435143

Start parameters:

Theta values were generated

from guessed values

Theta = 0.01000

M values were generated

from guessed values

M-matrix:

100000.00 [all are the same]

Connection type matrix:

where m = average (average over a group of Thetas or M,

s = symmetric M, S = symmetric 4Nm, 0 = zero, and not estimated,

\* = free to vary, Thetas are on diagonal

Population	1	2	3	4	5	6	7	8	9	10	11	12
1 ElfinCo	*	s	0	0	0	0	0	0	0	0	0	0
2 Bamfiel	s	*	s	0	0	0	0	0	0	0	0	0
3 PortRen	0	s	*	s	0	0	0	0	0	0	0	0
4 WalkOnB	0	0	s	*	s	0	0	0	0	0	0	0
5 BodegaH	0	0	0	s	*	s	0	0	0	0	0	0
6 Davenpo	0	0	0	0	s	*	s	0	0	0	0	0
7 VistaDe	0	0	0	0	0	s	*	s	0	0	0	0
8 HazardR	0	0	0	0	0	0	s	*	s	0	0	0
9 Refugio	0	0	0	0	0	0	0	s	*	s	0	0
10 Carpint	0	0	0	0	0	0	0	0	s	*	s	0

11 WhitePo	0	0	0	0	0	0	0	0	0	s	*	s
12 LaJolla	0	0	0	0	0	0	0	0	0	0	s	*

Order of parameters:

1	$\Theta_1$		<displayed>
2	$\Theta_2$		<displayed>
3	$\Theta_3$		<displayed>
4	$\Theta_4$		<displayed>
5	$\Theta_5$		<displayed>
6	$\Theta_6$		<displayed>
7	$\Theta_7$		<displayed>
8	$\Theta_8$		<displayed>
9	$\Theta_9$		<displayed>
10	$\Theta_{10}$		<displayed>
11	$\Theta_{11}$		<displayed>
12	$\Theta_{12}$		<displayed>
13	$M_{2 \rightarrow 1}$	= $M_{2 \rightarrow 1}$ [s]	<displayed>
24	$M_{1 \rightarrow 2}$	= $M_{2 \rightarrow 1}$ [s]	
25	$M_{3 \rightarrow 2}$	= $M_{3 \rightarrow 2}$ [s]	<displayed>
36	$M_{2 \rightarrow 3}$	= $M_{3 \rightarrow 2}$ [s]	
37	$M_{4 \rightarrow 3}$	= $M_{4 \rightarrow 3}$ [s]	<displayed>
48	$M_{3 \rightarrow 4}$	= $M_{4 \rightarrow 3}$ [s]	
49	$M_{5 \rightarrow 4}$	= $M_{5 \rightarrow 4}$ [s]	<displayed>
60	$M_{4 \rightarrow 5}$	= $M_{5 \rightarrow 4}$ [s]	
61	$M_{6 \rightarrow 5}$	= $M_{6 \rightarrow 5}$ [s]	<displayed>
72	$M_{5 \rightarrow 6}$	= $M_{6 \rightarrow 5}$ [s]	
73	$M_{7 \rightarrow 6}$	= $M_{7 \rightarrow 6}$ [s]	<displayed>
84	$M_{6 \rightarrow 7}$	= $M_{7 \rightarrow 6}$ [s]	
85	$M_{8 \rightarrow 7}$	= $M_{8 \rightarrow 7}$ [s]	<displayed>
96	$M_{7 \rightarrow 8}$	= $M_{8 \rightarrow 7}$ [s]	
97	$M_{9 \rightarrow 8}$	= $M_{9 \rightarrow 8}$ [s]	<displayed>
108	$M_{8 \rightarrow 9}$	= $M_{9 \rightarrow 8}$ [s]	
109	$M_{10 \rightarrow 9}$	= $M_{10 \rightarrow 9}$ [s]	<displayed>
120	$M_{9 \rightarrow 10}$	= $M_{10 \rightarrow 9}$ [s]	
121	$M_{11 \rightarrow 10}$	= $M_{11 \rightarrow 10}$ [s]	<displayed>
132	$M_{10 \rightarrow 11}$	= $M_{11 \rightarrow 10}$ [s]	
133	$M_{12 \rightarrow 11}$	= $M_{12 \rightarrow 11}$ [s]	<displayed>
144	$M_{11 \rightarrow 12}$	= $M_{12 \rightarrow 11}$ [s]	

Mutation rate among loci:

Mutation rate is constant

Analysis strategy:

Bayesian inference

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Slice sampling

Prior distribution for parameter

Parameter	Prior	Minimum	Mean*	Maximum	Delta	Bins
Theta	Exp window	0.000010	0.010000	10.000000	1.000000	500
M	Exp window	0.000100	100000.000000	1000000.000000	100000.000000	500

Markov chain settings:

	Long chain
Number of chains	1
Recorded steps [a]	1000
Increment (record every x step [b])	100
Number of concurrent chains (replicates) [c]	3
Visited (sampled) parameter values [a*b*c]	300000
Number of discard trees per chain (burn-in)	1000

Multiple Markov chains:

Static heating scheme	4 chains with temperatures
	100000.00 3.00 1.50 1.00
	Swapping interval is 1

Print options:

Data file:	../mcalifornianus_210528.mig
Output file:	outfile.txt
Posterior distribution raw histogram file:	bayesfile
Print data:	No
Print genealogies [only some for some data type]:	None

*Data summary*

Datatype: Sequence data  
 Number of loci: 1

Population	Locus	Gene copies
1 ElfinCo	1	19
2 Bamfiel	1	23
3 PortRen	1	15
4 WalkOnB	1	16
5 BodegaH	1	7
6 Davenpo	1	17
7 VistaDe	1	19
8 HazardR	1	23
9 Refugio	1	16
10 Carpint	1	19
11 WhitePo	1	11
12 LaJolla	1	8
Total of all populations	1	193







## *Bayesian Analysis: Posterior distribution table*


Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	$\Theta_1$	0.00001	0.00001	0.01001	0.06001	0.18001	0.07001	0.02756
1	$\Theta_2$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01820
1	$\Theta_3$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01692
1	$\Theta_4$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01744
1	$\Theta_5$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.02139
1	$\Theta_6$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01661
1	$\Theta_7$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01795
1	$\Theta_8$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01128
1	$\Theta_9$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01098
1	$\Theta_{10}$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.00924
1	$\Theta_{11}$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01249
1	$\Theta_{12}$	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.00805
1	$M_{2 \rightarrow 1}$	0.0	2000.0	11000.0	18000.0	28000.0	85000.0	67904.6
1	$M_{1 \rightarrow 2}$	0.0	2000.0	11000.0	18000.0	28000.0	85000.0	67904.6
1	$M_{3 \rightarrow 2}$	30000.0	46000.0	57000.0	66000.0	80000.0	59000.0	57027.0
1	$M_{2 \rightarrow 3}$	30000.0	46000.0	57000.0	66000.0	80000.0	59000.0	57027.0
1	$M_{4 \rightarrow 3}$	28000.0	66000.0	83000.0	96000.0	110000.0	73000.0	70907.6
1	$M_{3 \rightarrow 4}$	28000.0	66000.0	83000.0	96000.0	110000.0	73000.0	70907.6
1	$M_{5 \rightarrow 4}$	22000.0	30000.0	43000.0	56000.0	62000.0	53000.0	66096.4
1	$M_{4 \rightarrow 5}$	22000.0	30000.0	43000.0	56000.0	62000.0	53000.0	66096.4
1	$M_{6 \rightarrow 5}$	50000.0	62000.0	81000.0	98000.0	112000.0	95000.0	101408.9
1	$M_{5 \rightarrow 6}$	50000.0	62000.0	81000.0	98000.0	112000.0	95000.0	101408.9
1	$M_{7 \rightarrow 6}$	10000.0	16000.0	29000.0	38000.0	46000.0	83000.0	77778.5
1	$M_{6 \rightarrow 7}$	10000.0	16000.0	29000.0	38000.0	46000.0	83000.0	77778.5
1	$M_{8 \rightarrow 7}$	62000.0	74000.0	91000.0	106000.0	120000.0	83000.0	73635.4
1	$M_{7 \rightarrow 8}$	62000.0	74000.0	91000.0	106000.0	120000.0	83000.0	73635.4
1	$M_{9 \rightarrow 8}$	0.0	4000.0	17000.0	30000.0	38000.0	27000.0	37028.6
1	$M_{8 \rightarrow 9}$	0.0	4000.0	17000.0	30000.0	38000.0	27000.0	37028.6
1	$M_{10 \rightarrow 9}$	0.0	6000.0	17000.0	24000.0	34000.0	81000.0	72370.2
1	$M_{9 \rightarrow 10}$	0.0	6000.0	17000.0	24000.0	34000.0	81000.0	72370.2
1	$M_{11 \rightarrow 10}$	32000.0	38000.0	53000.0	64000.0	70000.0	47000.0	39384.0
1	$M_{10 \rightarrow 11}$	32000.0	38000.0	53000.0	64000.0	70000.0	47000.0	39384.0
1	$M_{12 \rightarrow 11}$	78000.0	94000.0	109000.0	124000.0	140000.0	101000.0	92158.8
1	$M_{11 \rightarrow 12}$	78000.0	94000.0	109000.0	124000.0	140000.0	101000.0	92158.8

Citation suggestions:


- Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341-345
- Beerli P., 2007. Estimation of the population scaled mutation rate from microsatellite data, *Genetics*, 177:1967-1968.
- Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use? In *Population Genetics for Animal Conservation*, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli, and C. Vernesi, eds., vol. 17 of *Conservation Biology*, Cambridge University Press, Cambridge UK, pp. 42-79.


*Bayesian Analysis: Posterior distribution over all loci*











## *Log-Probability of the data given the model (marginal likelihood)*

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as  $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Method	$\ln(\text{Prob}(D \mid \text{Model}))$	Notes
Thermodynamic integration	-2270.854750	(1a)
	-2178.794854	(1b)
Harmonic mean	-1884.079694	(2)

(1a, 1b and 2) are approximations to the marginal likelihood, make sure that the program run long enough!

(1a, 1b) and (2) should give similar results, in principle.

But (2) is overestimating the likelihood, it is presented for historical reasons and should not be used

(1a, 1b) needs heating with chains that span a temperature range of 1.0 to at least 100,000.

(1b) is using a Bezier-curve to get better approximations for runs with low number of heated chains

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

*Acceptance ratios for all parameters and the genealogies*

Parameter	Accepted changes	Ratio
$\Theta_1$	1033/4517	0.22869
$\Theta_2$	1402/4354	0.32200
$\Theta_3$	683/4371	0.15626
$\Theta_4$	1398/4324	0.32331
$\Theta_5$	1350/4400	0.30682
$\Theta_6$	1615/4489	0.35977
$\Theta_7$	1496/4420	0.33846
$\Theta_8$	837/4338	0.19295
$\Theta_9$	971/4448	0.21830
$\Theta_{10}$	1395/4355	0.32032
$\Theta_{11}$	1754/4316	0.40639
$\Theta_{12}$	769/4427	0.17371
M <sub>2→1</sub>	4432/4432	1.00000
M <sub>1→2</sub>	4432/4432	1.00000
M <sub>3→2</sub>	4430/4430	1.00000
M <sub>2→3</sub>	4430/4430	1.00000
M <sub>4→3</sub>	4384/4384	1.00000
M <sub>3→4</sub>	4384/4384	1.00000
M <sub>5→4</sub>	4527/4527	1.00000
M <sub>4→5</sub>	4527/4527	1.00000
M <sub>6→5</sub>	4469/4469	1.00000
M <sub>5→6</sub>	4469/4469	1.00000
M <sub>7→6</sub>	4402/4402	1.00000
M <sub>6→7</sub>	4402/4402	1.00000
M <sub>8→7</sub>	4449/4449	1.00000
M <sub>7→8</sub>	4449/4449	1.00000
M <sub>9→8</sub>	4410/4410	1.00000
M <sub>8→9</sub>	4410/4410	1.00000
M <sub>10→9</sub>	4466/4466	1.00000
M <sub>9→10</sub>	4466/4466	1.00000
M <sub>11→10</sub>	4341/4341	1.00000
M <sub>10→11</sub>	4341/4341	1.00000
M <sub>12→11</sub>	4270/4270	1.00000
M <sub>11→12</sub>	4270/4270	1.00000
Genealogies	32788/149821	0.21885

*MCMC-Autocorrelation and Effective MCMC Sample Size*

Parameter	Autocorrelation	Effective Sample Size
$\Theta_1$	0.85826	232.29
$\Theta_2$	0.77033	388.99
$\Theta_3$	0.88784	184.91
$\Theta_4$	0.78944	353.89
$\Theta_5$	0.80355	328.03
$\Theta_6$	0.77416	383.55
$\Theta_7$	0.78145	370.50
$\Theta_8$	0.91973	127.78
$\Theta_9$	0.87983	201.55
$\Theta_{10}$	0.77731	382.42
$\Theta_{11}$	0.73890	450.43
$\Theta_{12}$	0.91410	137.99
$M_{2 \rightarrow 1}$	0.81438	309.73
$M_{1 \rightarrow 2}$	0.81438	309.73
$M_{3 \rightarrow 2}$	0.80726	321.05
$M_{2 \rightarrow 3}$	0.80726	321.05
$M_{4 \rightarrow 3}$	0.79668	339.85
$M_{3 \rightarrow 4}$	0.79668	339.85
$M_{5 \rightarrow 4}$	0.82737	287.37
$M_{4 \rightarrow 5}$	0.82737	287.37
$M_{6 \rightarrow 5}$	0.81214	310.76
$M_{5 \rightarrow 6}$	0.81214	310.76
$M_{7 \rightarrow 6}$	0.81113	320.90
$M_{6 \rightarrow 7}$	0.81113	320.90
$M_{8 \rightarrow 7}$	0.87069	208.50
$M_{7 \rightarrow 8}$	0.87069	208.50
$M_{9 \rightarrow 8}$	0.83259	278.90
$M_{8 \rightarrow 9}$	0.83259	278.90
$M_{10 \rightarrow 9}$	0.78745	357.22
$M_{9 \rightarrow 10}$	0.78745	357.22
$M_{11 \rightarrow 10}$	0.83937	262.97
$M_{10 \rightarrow 11}$	0.83937	262.97
$M_{12 \rightarrow 11}$	0.88795	178.43
$M_{11 \rightarrow 12}$	0.88795	178.43
$\text{Ln}[\text{Prob}(\text{D} \text{G})]$	0.98368	24.67

## *Potential Problems*

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is very common that some parameters for some loci will not be very informative, triggering suggestions (for example to increase the prior range) that are not sensible. This suggestion tool will improve with time, therefore do not blindly follow its suggestions. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. For example, if you run a Bayesian inference with sequence data, for macroscopic species there is rarely the need to increase the prior for Theta beyond 0.1; but if you use microsatellites it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3) it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the range will not help in such situations, reducing number of parameters may help in such situations.

No warning was recorded during the run