

Preliminary migrate analysis of *M. californianus*

MIGRATION RATE AND POPULATION SIZE ESTIMATION

using the coalescent and maximum likelihood or Bayesian inference

Migrate-n version 3.7.2 [April-12-18]

Program started at Fri May 28 15:36:32 2021

Program finished at Fri May 28 23:21:36 2021



Options

Datatype:

DNA sequence data

Inheritance scalers in use for Thetas:

All loci use an inheritance scaler of 1.0

[The locus with a scaler of 1.0 used as reference]

Random number seed:

(with internal timer)

2675007803

Start parameters:

Theta values were generated

from guessed values

Theta = 0.01000

M values were generated

from guessed values

M-matrix:

100000.00 [all are the same]

Connection type matrix:

where m = average (average over a group of Thetas or M,

s = symmetric M, S = symmetric 4Nm, 0 = zero, and not estimated,

* = free to vary, Thetas are on diagonal

Population	1	2	3	4	5	6	7	8	9	10	11	12
1 ElfinCo	*	*	0	0	0	0	0	0	0	0	0	0
2 Bamfiel	*	*	*	0	0	0	0	0	0	0	0	0
3 PortRen	0	*	*	*	0	0	0	0	0	0	0	0
4 WalkOnB	0	0	*	*	*	0	0	0	0	0	0	0
5 BodegaH	0	0	0	*	*	*	0	0	0	0	0	0
6 Davenpo	0	0	0	0	*	*	*	0	0	0	0	0
7 VistaDe	0	0	0	0	0	*	*	*	0	0	0	0
8 HazardR	0	0	0	0	0	0	*	*	*	0	0	0
9 Refugio	0	0	0	0	0	0	0	*	*	*	0	0
10 Carpint	0	0	0	0	0	0	0	0	*	*	*	0

11 WhitePo	0	0	0	0	0	0	0	0	0	*	*	*
12	0	0	0	0	0	0	0	0	0	0	*	*

Order of parameters:

1	Θ_1	<displayed>
2	Θ_2	<displayed>
3	Θ_3	<displayed>
4	Θ_4	<displayed>
5	Θ_5	<displayed>
6	Θ_6	<displayed>
7	Θ_7	<displayed>
8	Θ_8	<displayed>
9	Θ_9	<displayed>
10	Θ_{10}	<displayed>
11	Θ_{11}	<displayed>
12	Θ_{12}	<displayed>
13	$M_{2 \rightarrow 1}$	<displayed>
24	$M_{1 \rightarrow 2}$	<displayed>
25	$M_{3 \rightarrow 2}$	<displayed>
36	$M_{2 \rightarrow 3}$	<displayed>
37	$M_{4 \rightarrow 3}$	<displayed>
48	$M_{3 \rightarrow 4}$	<displayed>
49	$M_{5 \rightarrow 4}$	<displayed>
60	$M_{4 \rightarrow 5}$	<displayed>
61	$M_{6 \rightarrow 5}$	<displayed>
72	$M_{5 \rightarrow 6}$	<displayed>
73	$M_{7 \rightarrow 6}$	<displayed>
84	$M_{6 \rightarrow 7}$	<displayed>
85	$M_{8 \rightarrow 7}$	<displayed>
96	$M_{7 \rightarrow 8}$	<displayed>
97	$M_{9 \rightarrow 8}$	<displayed>
108	$M_{8 \rightarrow 9}$	<displayed>
109	$M_{10 \rightarrow 9}$	<displayed>
120	$M_{9 \rightarrow 10}$	<displayed>
121	$M_{11 \rightarrow 10}$	<displayed>
132	$M_{10 \rightarrow 11}$	<displayed>
133	$M_{12 \rightarrow 11}$	<displayed>
144	$M_{11 \rightarrow 12}$	<displayed>

Mutation rate among loci:

Mutation rate is constant

Analysis strategy:

Bayesian inference

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Slice sampling

Prior distribution for parameter

Parameter	Prior	Minimum	Mean*	Maximum	Delta	Bins
Theta	Exp window	0.000010	0.010000	10.000000	1.000000	500
M	Exp window	0.000100	100000.000000	1000000.000000	100000.000000	500

Markov chain settings:

Long chain

Number of chains	1
Recorded steps [a]	1000
Increment (record every x step [b])	100
Number of concurrent chains (replicates) [c]	3
Visited (sampled) parameter values [a*b*c]	300000
Number of discard trees per chain (burn-in)	1000

Multiple Markov chains:

Static heating scheme	4 chains with temperatures
	100000.00 3.00 1.50 1.00
	Swapping interval is 1

Print options:

Data file:	.././mcalifornianus_210528.mig
Output file:	outfile.txt
Posterior distribution raw histogram file:	bayesfile
Print data:	No
Print genealogies [only some for some data type]:	None

Data summary

Datatype: Sequence data
 Number of loci: 1

Population	Locus	Gene copies
1 ElfinCo	1	19
2 Bamfiel	1	23
3 PortRen	1	15
4 WalkOnB	1	16
5 BodegaH	1	7
6 Davenpo	1	17
7 VistaDe	1	19
8 HazardR	1	23
9 Refugio	1	16
10 Carpint	1	19
11 WhitePo	1	10
12	1	0
Total of all populations	1	184







Bayesian Analysis: Posterior distribution table


Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.00001	0.00001	0.01001	0.08001	0.18001	0.09001	0.03053
1	Θ_2	0.00001	0.00001	0.01001	0.08001	0.18001	0.09001	0.03169
1	Θ_3	0.00001	0.00001	0.01001	0.08001	0.18001	0.09001	0.03091
1	Θ_4	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01717
1	Θ_5	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01778
1	Θ_6	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01231
1	Θ_7	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01862
1	Θ_8	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01662
1	Θ_9	0.00001	0.00001	0.01001	0.06001	0.18001	0.07001	0.02699
1	Θ_{10}	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01863
1	Θ_{11}	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.00597
1	Θ_{12}	0.00001	0.00001	0.01001	0.06001	0.16001	0.07001	0.01275
1	$M_{2 \rightarrow 1}$	0.0	4000.0	25000.0	44000.0	142000.0	39000.0	49797.9
1	$M_{1 \rightarrow 2}$	0.0	26000.0	35000.0	42000.0	56000.0	29000.0	23994.6
1	$M_{3 \rightarrow 2}$	70000.0	78000.0	93000.0	106000.0	116000.0	87000.0	71611.7
1	$M_{2 \rightarrow 3}$	26000.0	42000.0	53000.0	66000.0	86000.0	57000.0	56506.1
1	$M_{4 \rightarrow 3}$	0.0	4000.0	15000.0	22000.0	40000.0	19000.0	15534.0
1	$M_{3 \rightarrow 4}$	0.0	10000.0	19000.0	28000.0	40000.0	21000.0	19357.1
1	$M_{5 \rightarrow 4}$	56000.0	70000.0	85000.0	100000.0	114000.0	79000.0	60899.8
1	$M_{4 \rightarrow 5}$	0.0	0.0	1000.0	12000.0	20000.0	83000.0	64387.5
1	$M_{6 \rightarrow 5}$	18000.0	28000.0	37000.0	46000.0	56000.0	91000.0	81814.8
1	$M_{5 \rightarrow 6}$	62000.0	70000.0	85000.0	100000.0	110000.0	79000.0	68878.2
1	$M_{7 \rightarrow 6}$	20000.0	34000.0	45000.0	54000.0	70000.0	47000.0	44965.7
1	$M_{6 \rightarrow 7}$	34000.0	44000.0	57000.0	68000.0	78000.0	53000.0	39663.3
1	$M_{8 \rightarrow 7}$	32000.0	42000.0	59000.0	76000.0	88000.0	71000.0	76430.3
1	$M_{7 \rightarrow 8}$	0.0	10000.0	21000.0	36000.0	62000.0	31000.0	30855.1
1	$M_{9 \rightarrow 8}$	0.0	28000.0	43000.0	54000.0	60000.0	39000.0	32568.9
1	$M_{8 \rightarrow 9}$	0.0	0.0	1000.0	12000.0	52000.0	39000.0	36397.5
1	$M_{10 \rightarrow 9}$	0.0	12000.0	21000.0	30000.0	40000.0	23000.0	21437.8
1	$M_{9 \rightarrow 10}$	12000.0	22000.0	45000.0	64000.0	74000.0	57000.0	86091.2
1	$M_{11 \rightarrow 10}$	0.0	26000.0	41000.0	52000.0	60000.0	35000.0	29774.1
1	$M_{10 \rightarrow 11}$	0.0	0.0	9000.0	22000.0	48000.0	35000.0	45430.1
1	$M_{12 \rightarrow 11}$	2000.0	12000.0	21000.0	30000.0	40000.0	63000.0	61995.4
1	$M_{11 \rightarrow 12}$	24000.0	36000.0	51000.0	66000.0	80000.0	139000.0	170308.0


Citation suggestions:


- Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341-345
- Beerli P., 2007. Estimation of the population scaled mutation rate from microsatellite data, *Genetics*, 177:1967-1968.
- Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use? In *Population Genetics for Animal Conservation*, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli, and C. Vernesi, eds., vol. 17 of *Conservation Biology*, Cambridge University Press, Cambridge UK, pp. 42-79.





Bayesian Analysis: Posterior distribution over all loci











Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Method	$\ln(\text{Prob}(D \mid \text{Model}))$	Notes
Thermodynamic integration	-2151.281687	(1a)
	-2068.775343	(1b)
Harmonic mean	-1818.290896	(2)

(1a, 1b and 2) are approximations to the marginal likelihood, make sure that the program run long enough!

(1a, 1b) and (2) should give similar results, in principle.

But (2) is overestimating the likelihood, it is presented for historical reasons and should not be used

(1a, 1b) needs heating with chains that span a temperature range of 1.0 to at least 100,000.

(1b) is using a Bezier-curve to get better approximations for runs with low number of heated chains

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	1311/4431	0.29587
Θ_2	813/4502	0.18059
Θ_3	753/4370	0.17231
Θ_4	1698/4422	0.38399
Θ_5	1769/4375	0.40434
Θ_6	966/4472	0.21601
Θ_7	1537/4343	0.35390
Θ_8	1016/4441	0.22878
Θ_9	681/4414	0.15428
Θ_{10}	1840/4527	0.40645
Θ_{11}	1214/4487	0.27056
Θ_{12}	2583/4361	0.59230
$M_{2 \rightarrow 1}$	4488/4488	1.00000
$M_{1 \rightarrow 2}$	4403/4403	1.00000
$M_{3 \rightarrow 2}$	4453/4453	1.00000
$M_{2 \rightarrow 3}$	4308/4308	1.00000
$M_{4 \rightarrow 3}$	4379/4379	1.00000
$M_{3 \rightarrow 4}$	4480/4480	1.00000
$M_{5 \rightarrow 4}$	4448/4448	1.00000
$M_{4 \rightarrow 5}$	4431/4431	1.00000
$M_{6 \rightarrow 5}$	4412/4412	1.00000
$M_{5 \rightarrow 6}$	4447/4447	1.00000
$M_{7 \rightarrow 6}$	4538/4538	1.00000
$M_{6 \rightarrow 7}$	4325/4325	1.00000
$M_{8 \rightarrow 7}$	4432/4432	1.00000
$M_{7 \rightarrow 8}$	4384/4384	1.00000
$M_{9 \rightarrow 8}$	4413/4413	1.00000
$M_{8 \rightarrow 9}$	4486/4486	1.00000
$M_{10 \rightarrow 9}$	4378/4378	1.00000
$M_{9 \rightarrow 10}$	4417/4417	1.00000
$M_{11 \rightarrow 10}$	4426/4426	1.00000
$M_{10 \rightarrow 11}$	4372/4372	1.00000
$M_{12 \rightarrow 11}$	4321/4321	1.00000
$M_{11 \rightarrow 12}$	4347/4347	1.00000
Genealogies	37095/149767	0.24768

MCMC-Autocorrelation and Effective MCMC Sample Size

Parameter	Autocorrelation	Effective Sampe Size
Θ_1	0.75075	514.59
Θ_2	0.85169	259.70
Θ_3	0.87770	197.49
Θ_4	0.74849	440.06
Θ_5	0.73800	461.74
Θ_6	0.81241	355.68
Θ_7	0.78983	360.09
Θ_8	0.86991	215.19
Θ_9	0.91215	141.97
Θ_{10}	0.74547	446.11
Θ_{11}	0.78630	398.84
Θ_{12}	0.60015	881.02
M _{2→1}	0.74630	452.25
M _{1→2}	0.69138	558.17
M _{3→2}	0.77894	377.64
M _{2→3}	0.77556	388.73
M _{4→3}	0.73292	469.35
M _{3→4}	0.63956	663.27
M _{5→4}	0.78262	367.89
M _{4→5}	0.75644	434.86
M _{6→5}	0.81263	311.75
M _{5→6}	0.65016	644.85
M _{7→6}	0.90414	151.68
M _{6→7}	0.83558	273.28
M _{8→7}	0.75807	417.46
M _{7→8}	0.67853	579.16
M _{9→8}	0.76616	397.36
M _{8→9}	0.80050	339.25
M _{10→9}	0.81612	313.82
M _{9→10}	0.78507	364.49
M _{11→10}	0.88978	174.85
M _{10→11}	0.79766	340.85
M _{12→11}	0.71600	501.64
M _{11→12}	0.83097	276.80
Ln[Prob(D G)]	0.97120	43.81

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is very common that some parameters for some loci will not be very informative, triggering suggestions (for example to increase the prior range) that are not sensible. This suggestion tool will improve with time, therefore do not blindly follow its suggestions. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. For example, if you run a Bayesian inference with sequence data, for macroscopic species there is rarely the need to increase the prior for Theta beyond 0.1; but if you use microsatellites it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3) it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the range will not help in such situations, reducing number of parameters may help in such situations.

No warning was recorded during the run