

# Unit 6: Introduction to linear regression

## 2. Prediction, outliers, and inference for regression

Sta 101 - Fall 2015

Duke University, Department of Statistical Science

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

- ▶ PA 6 opens today, due Nov 22, Sun
- ▶ PS 6 due Nov 20, Fri
- ▶ RA 7 (last RA!) on Wednesday
- ▶ Project 2 questions?
  - If you want to see sample posters from previous years, stop by office hours
  - Most important advice: Sketch out a meeting / working plan with your team **TODAY**, keeping in mind Thanksgiving break

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled


## 3. Summary

### Clicker question

Suppose you want to predict annual murder count (per million) for a series of districts that were not included in the dataset. For which of the following districts would you be most comfortable with your prediction?

A district where % in poverty =

- (a) 5%
- (b) 15%
- (c) 20%
- (d) 26%
- (e) 40%




### Clicker question


Suppose you want to predict annual murder count (per million) for a series of districts that were not included in the dataset. For which of the following districts would you be most comfortable with your prediction?

A district where % in poverty =

- (a) 5%
- (b) 15%
- (c) 20%
- (d) 26%
- (e) 40%



Sometimes the intercept might be an extrapolation: useful for adjusting the height of the line, but meaningless in the context of the data.





*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*In R:*

```
# load data
murder <- read.csv("https://stat.duke.edu/~mc301/data/murder.csv")
# fit model
m_mur_pov <- lm(annual_murders_per_mil ~ perc_pov, data = murder)
# create new data
newdata <- data.frame(perc_pov = 20)
# predict
predict(m_mur_pov, newdata)
```

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*In R:*

```
# load data
murder <- read.csv("https://stat.duke.edu/~mc301/data/murder.csv")
# fit model
m_mur_pov <- lm(annual_murders_per_mil ~ perc_pov, data = murder)
# create new data
newdata <- data.frame(perc_pov = 20)
# predict
predict(m_mur_pov, newdata)
```

```
1
21.28663
```

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*In R:*

```
# load data
murder <- read.csv("https://stat.duke.edu/~mc301/data/murder.csv")
# fit model
m_mur_pov <- lm(annual_murders_per_mil ~ perc_pov, data = murder)
# create new data
newdata <- data.frame(perc_pov = 20)
# predict
predict(m_mur_pov, newdata)
```

*By hand:*  $\widehat{murder} = -29.91 + 2.56 \text{ poverty}$

The predicted number of murders per million per year for a county with 20% poverty rate is:

$$\widehat{murder} = -29.91 + 2.56 \times 20 = 21.29$$

*In R:*

```
# load data
murder <- read.csv("https://stat.duke.edu/~mc301/data/murder.csv")
# fit model
m_mur_pov <- lm(annual_murders_per_mil ~ perc_pov, data = murder)
# create new data
newdata <- data.frame(perc_pov = 20)
# predict
predict(m_mur_pov, newdata)
```

```
1
21.28663
```



## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.
- ▶ If the model is good, the predictions should be close to the true value of the response variable for this observation, however it may not be exact, i.e.  $\hat{y}$  might be different than  $y$ .

- ▶ Regression models are useful for making predictions for new observations not include in the original dataset.
- ▶ If the model is good, the predictions should be close to the true value of the response variable for this observation, however it may not be exact, i.e.  $\hat{y}$  might be different than  $y$ .
- ▶ With any prediction we can (and should) also report a measure of uncertainty of the prediction.

A *prediction interval* for  $y$  for a given  $x^*$  is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where  $s$  is the standard deviation of the residuals, and  $x^*$  is a new observation.

A *prediction interval* for  $y$  for a given  $x^*$  is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where  $s$  is the standard deviation of the residuals, and  $x^*$  is a new observation.

- Interpretation: We are XX% confident that  $\hat{y}$  for given  $x^*$  is within this interval.

A *prediction interval* for  $y$  for a given  $x^*$  is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where  $s$  is the standard deviation of the residuals, and  $x^*$  is a new observation.

- ▶ Interpretation: We are XX% confident that  $\hat{y}$  for given  $x^*$  is within this interval.
- ▶ The width of the confidence interval for  $\hat{y}$  increases as
  - $x^*$  moves away from the center
  - $s$  (the variability of residuals), i.e. the scatter, increases

A *prediction interval* for  $y$  for a given  $x^*$  is

$$\hat{y} \pm t_{n-2}^* s \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2}}$$

where  $s$  is the standard deviation of the residuals, and  $x^*$  is a new observation.

- ▶ Interpretation: We are XX% confident that  $\hat{y}$  for given  $x^*$  is within this interval.
- ▶ The width of the confidence interval for  $\hat{y}$  increases as
  - $x^*$  moves away from the center
  - $s$  (the variability of residuals), i.e. the scatter, increases
- ▶ Prediction level: If we repeat the study of obtaining a regression data set many times, each time forming a XX% prediction interval at  $x^*$ , and wait to see what the future value of  $y$  is at  $x^*$ , then roughly XX% of the prediction intervals will contain the corresponding actual value of  $y$ .



*By hand:*

Don't worry about it...

*By hand:*

Don't worry about it...

*In R:*

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

*By hand:*

Don't worry about it...

*In R:*

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

	fit	lwr	upr
1	21.28663	9.418327	33.15493

*By hand:*

Don't worry about it...

*In R:*

```
# predict  
predict(m_mur_pov, newdata, interval = "prediction", level = 0.95)
```

	fit	lwr	upr
1	21.28663	9.418327	33.15493

We are 95% confident that the annual murders per million for a county with 20% poverty rate is between 9.52 and 33.15.

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.

## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression:  $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual\_murders\_per\_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		



## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression:  $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual\_murders\_per\_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}}$$

## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression:  $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual\_murders\_per\_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}}$$

## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression:  $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table

Response: annual\_murders\_per\_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}} = \frac{1308.34}{1308.34 + 546.86}$$

## (1) $R^2$ assesses model fit -- higher the better

- ▶  $R^2$ : percentage of variability in  $y$  explained by the model.
- ▶ For single predictor regression:  $R^2$  is the square of the correlation coefficient,  $R$ .

```
murder %>%  
  summarise(r_sq = cor(annual_murders_per_mil, perc_pov)^2)
```

```
      r_sq  
1 0.7052275
```

- ▶ For all regression:  $R^2 = \frac{SS_{reg}}{SS_{tot}}$

```
anova(m_mur_pov)
```

Analysis of Variance Table


Response: annual\_murders\_per\_mil

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
perc_pov	1	1308.34	1308.34	43.064	3.638e-06 ***
Residuals	18	546.86	30.38		

$$R^2 = \frac{\text{explained variability}}{\text{total variability}} = \frac{SS_{reg}}{SS_{tot}} = \frac{1308.34}{1308.34 + 546.86} = \frac{1308.34}{1855.2} \approx 0.71$$

### Clicker question


$R^2$  for the regression model for predicting annual murders per million based on percentage living in poverty is roughly 71%. Which of the following is the correct interpretation of this value?



- (a) 71% of the variability in percentage living in poverty is explained by the model.
- (b) 84% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (c) 71% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (d) 71% of the time percentage living in poverty predicts murder rates accurately.

### Clicker question

$R^2$  for the regression model for predicting annual murders per million based on percentage living in poverty is roughly 71%. Which of the following is the correct interpretation of this value?



- (a) 71% of the variability in percentage living in poverty is explained by the model.
- (b) 84% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.
- (c) *71% of the variability in the murder rates is explained by the model, i.e. percentage living in poverty.*
- (d) 71% of the time percentage living in poverty predicts murder rates accurately.

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
- 4. Inference for regression uses the  $t$ -distribution**
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

- ▶ Use a T distribution for inference on the slope, with degrees of freedom  $n - 2$ 
  - Degrees of freedom for the slope(s) in regression is  $df = n - p - 1$  where  $p$  is the number of predictors (explanatory variables) in the model.



- ▶ Use a  $T$  distribution for inference on the slope, with degrees of freedom  $n - 2$ 
  - Degrees of freedom for the slope(s) in regression is  $df = n - p - 1$  where  $p$  is the number of predictors (explanatory variables) in the model.
- ▶ Hypothesis testing for a slope:  $H_0 : \beta_1 = 0$ ;  $H_A : \beta_1 \neq 0$ 
  - $T_{n-2} = \frac{b_1 - 0}{SE_{b_1}}$
  - $p\text{-value} = P(\text{observing a slope at least as different from 0 as the one observed if in fact there is no relationship between } x \text{ and } y)$

- ▶ Use a  $T$  distribution for inference on the slope, with degrees of freedom  $n - 2$ 
  - Degrees of freedom for the slope(s) in regression is  $df = n - p - 1$  where  $p$  is the number of predictors (explanatory variables) in the model.
- ▶ Hypothesis testing for a slope:  $H_0 : \beta_1 = 0$ ;  $H_A : \beta_1 \neq 0$ 
  - $T_{n-2} = \frac{b_1 - 0}{SE_{b_1}}$
  - $p\text{-value} = P(\text{observing a slope at least as different from 0 as the one observed if in fact there is no relationship between } x \text{ and } y)$
- ▶ Confidence intervals for a slope:
  - $b_1 \pm T_{n-2}^* SE_{b_1}$

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
- 5. Conditions for regression**
6. Type of outlier determines how it should be handled

## 3. Summary

### *Important regardless of doing inference*

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

### *Important regardless of doing inference*

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

### *Important for inference*

- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals

### *Important regardless of doing inference*

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference

### *Important for inference*

- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals
- ▶ Constant variability of residuals (*homoscedasticity*) → no fan shape in the residuals plot

### *Important regardless of doing inference*

- ▶ Linearity → randomly scattered residuals around 0 in the residuals plot – important regardless of doing inference


### *Important for inference*

- ▶ Nearly normally distributed residuals → histogram or normal probability plot of residuals
- ▶ Constant variability of residuals (*homoscedasticity*) → no fan shape in the residuals plot
- ▶ Independence of residuals (and hence observations) → depends on data collection method, often violated for time-series data

## Clicker question

What condition is this linear model obviously and definitely violating?

- (a) Linear relationship
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations






## Clicker question

What condition is this linear model obviously and definitely violating?


- (a) Linear relationship
- (b) Non-normal residuals
- (c) *Constant variability*
- (d) Independence of observations



## Clicker question

What condition is this linear model obviously and definitely violating?


- (a) Linear relationship
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations



## Clicker question

What condition is this linear model obviously and definitely violating?

- (a) *Linear relationship*
- (b) Non-normal residuals
- (c) Constant variability
- (d) Independence of observations




## 1. Housekeeping

## 2. Main ideas


1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary

- ▶ *Leverage* point is away from the cloud of points horizontally, does not necessarily change the slope
- ▶ *Influential* point changes the slope (most likely also has high leverage) – run the regression with and without that point to determine



- ▶ *Leverage* point is away from the cloud of points horizontally, does not necessarily change the slope
- ▶ *Influential* point changes the slope (most likely also has high leverage) – run the regression with and without that point to determine
- ▶ *Outlier* is an unusual point without these special characteristics (this one likely affects the intercept only)
- ▶ If clusters (groups of points) are apparent in the data, it might be worthwhile to model the groups separately.



## Application exercise: 6.2 Linear regression

See course website for details

## 1. Housekeeping

## 2. Main ideas

1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit -- higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled

## 3. Summary



1. Predict, but don't extrapolate
2. Predicted values also have uncertainty around them
3.  $R^2$  assesses model fit – higher the better
4. Inference for regression uses the  $t$ -distribution
5. Conditions for regression
6. Type of outlier determines how it should be handled