

# Simplified Belief Space Planning with an Alternative Observation Space and Formal Performance Guarantees Supplementary Material

Da Kong<sup>1</sup> and Vadim Indelman<sup>2</sup>

<sup>1</sup> Technion Autonomous Systems Program

<sup>2</sup> Department of Aerospace Engineering

Technion - Israel Institute of Technology, Haifa 32000, Israel

da-kong@campus.technion.ac.il, vadim.indelman@technion.ac.il

This document provides supplementary material to [1]. Therefore, it should not be considered a self-contained document, but instead regarded as an appendix of [1]. Throughout this report, all notations and definitions are with compliance to the ones presented in [1].

## 1 Proof of Lemma 1

$$\left| \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^\tau}^\tau (r(b_i^\tau)) - \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^{\tau Z}}^{\tau Z} (r(b_i^{\tau Z})) \right| \quad (1)$$

Assume state-dependent reward

$$= \left| \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^\tau}^\tau \mathbb{E}_{x_i|b_i^\tau} r(x_i) - \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_i|b_i^{\tau Z}} (r(x_i)) \right| \quad (2)$$

$$= \left| \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^\tau}^\tau \mathbb{E}_{x_i|h_i^{\tau-} \bar{z}_i|x_i, h_i^\tau}^\tau r(x_i) - \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_i|h_i^{\tau Z-} \bar{z}_i|x_i}^{\tau Z} (r(x_i)) \right| \quad (3)$$

(Chain rule and cancel  $z_i$ )

$$= \left| \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^\tau}^\tau \mathbb{E}_{x_i|h_i^{\tau-}} r(x_i) - \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_i|h_i^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, \pi_{i-1}^{TC}}^{\tau Z} (r(x_i)) \right| \quad (4)$$

(Bayes Rule)

$$= \left| \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^\tau}^\tau \mathbb{E}_{x_i|h_i^{\tau-}} r(x_i) - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}}^{\tau Z} \mathbb{E}_{x_i|h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}}^{\tau Z} r(x_i) \right| \quad (5)$$

$$= \left| \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^\tau}^\tau \mathbb{E}_{x_i|h_i^{\tau-}} r(x_i) \right. \\ \left. - \mathbb{E}_{x_0|b_0} \mathbb{E}_{x_1|x_0, \pi_0^{TC}} \mathbb{E}_{\bar{z}_1|x_1, h_1^{p-}}^{\tau Z} \dots \mathbb{E}_{x_{i-1}|x_{i-2}, \pi_{i-2}^{TC}} \mathbb{E}_{\bar{z}_{i-1}|x_{i-1}, h_{i-1}^{p-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}}^{\tau Z} \mathbb{E}_{x_i|h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}}^{\tau Z} r(x_i) \right| \quad (6)$$

Here,  $h_1^p, \dots, h_{i-1}^p$  are propagated history from the previous expectation.

$$\leq \max_{\bar{\pi}^\tau} \left| \mathbb{E}_{\bar{z}_{1:i-1}|b_0, \pi^\tau} \mathbb{E}_{x_i|h_i^{\tau-}} r(x_i) - \mathbb{E}_{x_0|b_0} \mathbb{E}_{x_1|x_0, \bar{\pi}_0^\tau} \mathbb{E}_{\bar{z}_1|x_1, h_1^p} \mathbb{E}^{\tau} \dots \mathbb{E}_{x_{i-1}|x_{i-2}, \bar{\pi}_{i-2}^\tau} \mathbb{E}_{\bar{z}_{i-1}|x_{i-1}, h_{i-1}^p} \mathbb{E}^{\tau} \mathbb{E}_{x_i|x_{i-1}, \bar{\pi}_{i-1}^\tau} r(x_i) \right| \quad (7)$$

## 2 Proof of Lemma 2

*Proof.* We can look into the difference in the Q function between two different topologies, which is defined as below:

$$\Delta Q(b_0, a_0, \pi^{\tau Z}, \pi^\tau, \tau_Z, \tau) \triangleq |Q_{\tau_Z}^{\pi^{\tau Z}}(b_0, a_0) - Q_{\tau}^{\pi^\tau}(b_0, a_0)| \quad (8)$$

$$= \left| \sum_{i=1}^L \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^{\tau Z}}^{\tau Z} (r(b_i)) - \sum_{i=1}^L \mathbb{E}_{\bar{z}_{1:i}|b_0, \pi^\tau}^{\tau} (r(b_i)) \right| \quad (9)$$

(Assume state-dependent reward and chain rule, same as Equation (5))

$$= \left| \sum_{i=1}^L \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}} \mathbb{E}^{\tau Z} r(x_i) - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^\tau}^{\tau} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-} x_i|x_{i-1}, \pi_{i-1}^{\tau}} \mathbb{E}^{\tau} r(x_i) \right] \right| \quad (10)$$

(Take out the expected reward  $r(x_L)$  from the summation)

$$= \left| \sum_{i=1}^{L-1} \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}} \mathbb{E}^{\tau Z} r(x_i) - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^\tau}^{\tau} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-} x_i|x_{i-1}, \pi_{i-1}^{\tau}} \mathbb{E}^{\tau} r(x_i) \right] + \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau Z-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau Z-} x_L|x_{L-1}, \pi_{L-1}^{\tau Z}} \mathbb{E}^{\tau Z} r(x_L) - \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^\tau}^{\tau} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-} x_L|x_{L-1}, \pi_{L-1}^{\tau}} \mathbb{E}^{\tau} r(x_L) \right| \quad (11)$$

(Bound the expected reward  $r(x_L)$  using Lemma ??)

$$\leq \max_{\bar{\pi}_{L-1}^\tau \in \bar{\Pi}_{L-1}(\mathcal{H}_{L-1}^{\tau Z-}, \bar{\mathcal{Z}}_{L-1}^\tau)} \left| \sum_{i=1}^{L-1} \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}} \mathbb{E}^{\tau Z} r(x_i) - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^\tau}^{\tau} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-} x_i|x_{i-1}, \pi_{i-1}^{\tau}} \mathbb{E}^{\tau} r(x_i) \right] + \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau Z-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau Z-} x_L|x_{L-1}, \bar{\pi}_{L-1}^\tau} \mathbb{E}^{\tau Z} r(x_L) - \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^\tau}^{\tau} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-} x_L|x_{L-1}, \bar{\pi}_{L-1}^\tau} \mathbb{E}^{\tau} r(x_L) \right| \quad (12)$$

(Take out the expected reward  $r(x_{L-1})$  from the summation, and denote the policy maximizing the distance as  $\bar{\pi}_{L-1}^+$ )

$$= \left| \sum_{i=1}^{L-2} \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}} r(x_i) \right. \right. \\ \left. \left. - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-} x_i|x_{i-1}, \pi_{i-1}^{\tau}} r(x_i) \right] \right| \quad (13)$$

$$+ \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau Z-} \bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau Z-} x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau Z}} r(x_{L-1}) \\ - \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau-} \bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau-} x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau}} r(x_{L-1}) \quad (14)$$

$$+ \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau Z-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau Z-} x_L|x_{L-1}, \bar{\pi}_{L-1}^+} r(x_L) \\ - \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-} x_L|x_{L-1}, \pi_{L-1}^{\tau}} r(x_L) \quad (15)$$

$$= \left| \sum_{i=1}^{L-2} \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-} x_i|x_{i-1}, \pi_{i-1}^{\tau Z}} r(x_i) \right. \right. \\ \left. \left. - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-} \bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-} x_i|x_{i-1}, \pi_{i-1}^{\tau}} r(x_i) \right] \right| \quad (16)$$

$$+ \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau Z}}^{\tau Z} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau Z-} \bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau Z-} x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau Z}} \left( r(x_{L-1}) \right. \\ \left. + \mathbb{E}_{\bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau Z-} x_L|x_{L-1}, \bar{\pi}_{L-1}^+}^{\tau} (h_{L-1}^{\tau Z-}, \bar{z}_{L-1}^{\tau}) r(x_L) \right) \quad (17)$$

$$- \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau-} \bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau-} x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau}} r(x_{L-1}) \\ - \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau}}^{\tau} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau-} \bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-} x_L|x_{L-1}, \pi_{L-1}^{\tau}} r(x_L) \quad (18)$$

(Bound the term  $\mathbb{E}_{x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau Z}}(r(x_{L-1})) + \mathbb{E}_{\bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau Z-}} \mathbb{E}_{x_L|x_{L-1}, \bar{\pi}_{L-1}^+} r(x_L)$ ) using a similar method as Remark ??)

$$\leq \max_{\bar{\pi}_{L-2, L-1}} \left| \sum_{i=1}^{L-2} \left[ \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau Z}} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau Z-}} \mathbb{E}_{\bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau Z-}} \mathbb{E}_{x_i|x_{i-1}, \pi_{i-1}^{\tau Z_1}} r(x_i) \right. \right. \\ \left. \left. - \mathbb{E}_{\bar{z}_{1:i-2}|b_0, \pi^{\tau}} \mathbb{E}_{x_{i-1}|h_{i-1}^{\tau-}} \mathbb{E}_{\bar{z}_{i-1}|x_{i-1}, h_{i-1}^{\tau-}} \mathbb{E}_{x_i|x_{i-1}, \pi_{i-1}^{\tau}} r(x_i) \right] \right. \quad (19)$$

$$+ \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau Z}, \tau Z} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau Z-}} \mathbb{E}_{\bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau Z-}} \mathbb{E}_{x_{L-1}|x_{L-2}, \bar{\pi}_{L-2}} r(x_{L-1}) \\ + \mathbb{E}_{\bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-}} \mathbb{E}_{x_L|x_{L-1}, \bar{\pi}_{L-1}} r(x_L) \Big) \quad (20)$$

$$- \mathbb{E}_{\bar{z}_{1:L-3}|b_0, \pi^{\tau}} \mathbb{E}_{x_{L-2}|h_{L-2}^{\tau-}} \mathbb{E}_{\bar{z}_{L-2}|x_{L-2}, h_{L-2}^{\tau-}} \mathbb{E}_{x_{L-1}|x_{L-2}, \pi_{L-2}^{\tau}} r(x_{L-1}) \\ - \mathbb{E}_{\bar{z}_{1:L-2}|b_0, \pi^{\tau}} \mathbb{E}_{x_{L-1}|h_{L-1}^{\tau-}} \mathbb{E}_{\bar{z}_{L-1}|x_{L-1}, h_{L-1}^{\tau-}} \mathbb{E}_{x_L|x_{L-1}, \pi_{L-1}^{\tau}} r(x_L) \Big| \quad (21)$$

(We can do it in a recursive way)

$$\leq \max_{\bar{\pi}^{\tau} \in \Pi^{\tau}} |Q_{\tau Z}^{\bar{\pi}^{\tau}}(b_0, a_0) - Q_{\tau}^{\bar{\pi}^{\tau}}(b_0, a_0)| \quad (22)$$

Which completes the proof of Lemma 2.

### 3 Proof of Theorem 1

The proof can be simple.

*Proof:*

$$Q_{\tau}^{\pi^{\tau Z}}(b_k, a_k) \leq \min_{\pi^{\tau} \in \Pi^{\tau}} [Q_{\tau}^{\pi^{\tau}}(b_k, a_k) + \delta_Q(b_k, a_k, \pi^{\tau}, \tau)] \quad (23)$$

$$= \min_{\pi^{\tau} \in \Pi^{\tau}} [Q_{\tau}^{\pi^{\tau}}(b_k, a_k) + \max_{\bar{\pi}^{\tau} \in \Pi^{\tau}} |Q_{\tau}^{\bar{\pi}^{\tau}}(b_k, a_k) - Q_{\tau}^{\pi^{\tau}}(b_k, a_k)|] \quad (24)$$

$$= \max_{\pi^{\tau} \in \Pi^{\tau}} [Q_{\tau}^{\pi^{\tau}}(b_k, a_k)] \quad (25)$$

The second direction in (20) can be proved similarly.  $\square$

### 4 Proof of Theorem 2

*Proof.* Let's consider the end of the planning horizon  $L$ .

If  $\beta^{\tau}(h(b_{L-1}^-)) = 1$ , we have:

$$\int_{z_{L-1}} p(z_{L-1}|x_{L-1}) \max_{\pi_{L-1}^{\tau} \in \Pi^{\tau}} \int_{x_L} p(x_L|x_{L-1}, \pi_{L-1}^{\tau}) r(x_L) \quad (26)$$

$$= \int_{z_{L-1}} p(z_{L-1}|x_{L-1}) \int_{x_L} p(x_L|x_{L-1}, \pi_{L-1}^{\tau Z*}) r(x_L) \quad (27)$$

If  $\beta^\tau(h(b_{L-1}^-)) = 0$ , we have:

$$\begin{aligned} & \min_{\pi_{L-1}^\tau \in \Pi^\tau} \int_{x_L} p(x_L | x_{L-1}, \pi_{L-1}^\tau) r(x_L) \\ & \leq \int_{z_{L-1}} p(z_{L-1} | x_{L-1}) \int_{x_L} p(x_L | x_{L-1}, \pi_{L-1}^{\tau_{Z^*}}) r(x_L) \end{aligned} \quad (28)$$

We denote the left-hand side in Equation (27) and (28) as  $LHS1(x_{L-1}, \tau)$  and the right-hand side as  $RHS1(x_{L-1}, \tau_Z)$ . In general, we can say  $LHS1(x_{L-1}, \tau) \leq RHS1(x_{L-1}, \tau_Z)$ .

Then, we take one more step earlier. Also if  $\beta^\tau(h(b_{L-2}^-)) = 1$ , we have:

$$\begin{aligned} & \int_{z_{L-1}} p(z_{L-1} | x_{L-1}) \max_{\pi_{L-2}^\tau \in \Pi^\tau} \int_{x_{L-1}} p(x_{L-1} | x_{L-2}, \pi_{L-2}^\tau) [r(x_{L-1}) + LHS1(x_{L-1}, \tau)] \\ & \leq \int_{z_{L-2}} p(z_{L-2} | h_{L-2}) \int_{x_{L-2}} p(x_{L-2} | x_{L-2}, \pi_{L-2}^{\tau_{Z^*}}) [r(x_{L-1}) + RHS1(x_{L-1}, \tau)] \end{aligned} \quad (29)$$

And if  $\beta^\tau(h(b_{L-2}^-)) = 0$ , we have:

$$\begin{aligned} & \min_{\pi_{L-2}^\tau \in \Pi^\tau} \int_{x_{L-1}} p(x_{L-1} | x_{L-2}, \pi_{L-2}^\tau) [r(x_{L-1}) + LHS1(x_{L-1}, \tau)] \\ & \leq \int_{z_{L-2}} p(z_{L-2} | h_{L-2}) \int_{x_{L-2}} p(x_{L-2} | x_{L-2}, \pi_{L-2}^{\tau_{Z^*}}) [r(x_{L-1}) + RHS1(x_{L-1}, \tau)] \end{aligned} \quad (30)$$

We also denote the left-hand side of the inequality in Equation (29) and (30) as  $LHS2(x_{L-2}, \tau)$  and the right-hand side as  $RHS2(x_{L-2}, \tau_Z)$ . In general, we can say  $LHS2(x_{L-2}, \tau) \leq RHS2(x_{L-2}, \tau_Z)$ .

The left-hand side is the iterative way to calculate  $lb(b_t, \pi_t, \tau)$ . If we keep iterate the above process from the end to the beginning of the planning horizon, we will get the inequality at the top of the belief tree:  $lb(b_0, a_0, \tau) \leq Q_{\tau_Z}^{\pi_{\tau_Z^*}}(b_0, a_0)$ .

## 5 Sparse Sampling

For a random variable  $X \sim \mathcal{P}$ , we use another sampling-based random variable,  $Y \sim \mathcal{Q}$  to approximate the expectation of the original theoretic value. If we obtain some samples  $\{y_i\}_{i=1}^N$  from a known distribution  $\mathcal{Q}$ , the calculation will be:

$$\mathbb{E}_{X \sim \mathcal{P}}[f(X)] = \int f(x) \mathcal{P}(x) dx = \int f(x) \frac{\mathcal{P}(x)}{\mathcal{Q}(x)} \mathcal{Q}(x) dx \simeq \frac{1}{N} \sum_{i=1}^N \frac{\mathcal{P}(y_i)}{\mathcal{Q}(y_i)} f(y_i) \quad (31)$$

Usually, the proposed distribution  $\mathcal{Q}$  is the known motion model. The weighted sampling can be more efficient, where resampling will create more particles from the samples with higher weights. The belief density is approximated by  $b(x) \simeq \frac{\sum_{i=1}^N w^i \delta(x - x^i)}{\sum_{i=1}^N w^i}$ .

### 5.1 Proof of Theorem 3

**Theorem 1 (Theorem 3). Bounded Estimation Error.** For all the depth  $d = 0, \dots, L - 1$  and  $a$ , the following concentration bound holds with probability at least  $1 - 2|A|(|A|C)^L \exp(\frac{-C\lambda^2}{2V_{\max}^2})$ :

$$\Delta \hat{ub}(b_0, a_0, \tau) \leq \frac{L(L-1)}{2} \lambda, \quad \Delta \hat{lb}(b_0, a_0, \tau) \leq \frac{L(L-1)}{2} \lambda. \quad (32)$$

For the upper bound:

*Proof.* We will follow a similar way to prove it as [16, 22].

Firstly, we consider a general situation at a depth of  $d$  and try to bound the error of estimating the optimal Q function given a belief  $b_d$  and an action  $a_d$ :

$$\delta_{ub}^\tau(d) \triangleq |Q_\tau^{\pi^*}(b_d, a_d) - \hat{Q}_\tau^{\pi^*}(b_d, a_d)| \quad (33)$$

$$= \left| r(b_d, a_d) + \mathbb{E}_{b_{d+1}|b_d, a_d} [V^{\tau^*}(b_{d+1})] - r(b_d, a_d) - \frac{1}{C} \sum_{i=1}^C \hat{V}^{\tau^*}(b_{d+1}^{[I_i]}) \right| \quad (34)$$

$$= \left| \mathbb{E}_{z_{d+1}|b_d, a_d, \tau} [V^{\tau^*}(b_{d+1})] - \frac{1}{C} \sum_{i=1}^C \hat{V}^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) \right| \quad (35)$$

$$\leq \left| \mathbb{E}_{z_{d+1}|b_d, a_d, \tau} [V^{\tau^*}(b_{d+1})] - \frac{1}{C} \sum_{i=1}^C V^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) \right| + \left| \frac{1}{C} \sum_{i=1}^C V^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) - \frac{1}{C} \sum_{i=1}^C \hat{V}^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) \right| \quad (36)$$

Here, the propagated next step belief state samples for a given belief tree topology  $\tau$  is denoted as  $\bar{b}_{d+1}^{\tau, [I_i]}$ . It is updated by the motion and observation models for the given topology  $\tau$ :

$$\bar{b}_{d+1}^{\tau, [I_i]} \sim \psi^\tau(\bar{b}_{d+1}^\tau | b_d, a_d) = \mathbb{1}_{\beta(h_{d+1}^-)=1} \alpha \cdot \mathbb{P}(z_{d+1} | x_{d+1}) \mathbb{P}(x_{d+1} | b_d, a_d) + \mathbb{1}_{\beta(h_{d+1}^-)=0} \mathbb{P}(x_{d+1} | b_d, a_d) \quad (37)$$

Here,  $\alpha$  is a Bayes normalization factor. For the first term in Equation (36), we can directly use the Hoeffding's inequality:

$$\mathbb{P}\left( \left| \mathbb{E}_{z_{d+1}|b_d, a_d, \tau} [V^{\tau^*}(b_{d+1})] - \frac{1}{C} \sum_{i=1}^C V^{\tau^*}(b_{d+1}^{[I_i]}) \right| \leq \lambda \right) \geq 1 - 2 \exp\left(\frac{-C\lambda^2}{2V_{\max}^2}\right) \quad (38)$$

For the second term in Equation (36), we analyze it in an iterative way:

$$\left| \frac{1}{C} \sum_{i=1}^C V^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) - \frac{1}{C} \sum_{i=1}^C \hat{V}^{\tau^*}(\bar{b}_{d+1}^{\tau, [I_i]}) \right| = \frac{1}{C} \left| \sum_{i=1}^C Q_\tau^{\pi^*}(\bar{b}_{d+1}^{\tau, [I_i]}, a_{d+1}^*) - \sum_{i=1}^C \hat{Q}_\tau^{\pi^*}(\bar{b}_{d+1}^{\tau, [I_i]}, a_{d+1}^*) \right| \quad (39)$$

$$\leq \delta_{ub}^\tau(d+1) \quad (40)$$

So, the below iterative bound is satisfied with a probability of at least  $1 - 2|A|(|A|C)^{L-d} \exp(\frac{-C\lambda^2}{2V_{\max}^2})$ :

$$\delta_{ub}^\tau(d) \leq \lambda + \delta_{ub}^\tau(d+1) \quad (41)$$

Here, the probability is based on the worst case for iteration, where it requires all the child belief nodes generated are well estimated with the number of  $|A|C$  child nodes for each time step.

The estimation error at the end of the planning horizon  $L - 1$  is:

$$\delta_{ub}^\tau(L-1) = \lambda \quad (42)$$

Then, we can find out the estimation bound at the top of the given belief tree by calculating it from the bottom to the top by Equation (41).

For the estimation of lower bound:

*Proof.* Firstly, we consider a general situation at a depth of  $d$  and try to bound the error of estimating the lower bound function given a belief  $b_d$  and an action  $a_d$ :

$$\delta_{lb}^\tau(t) \triangleq |lb(b_t, a_t, \tau) - \hat{lb}(b_t, a_t, \tau)| \quad (43)$$

$$\begin{aligned} &= \left| \mathbb{1}_{\beta(h(b_{t+1}^-))=1} \left[ \mathbb{E}_{z_{t+1}|h(b_t), a_t, \tau} \max_{\pi_{t+1}} lb(b_{t+1}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \max_{\pi_{t+1}} \hat{lb}(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right] \right. \\ &\quad \left. + \mathbb{1}_{\beta(h(b_{t+1}^-))=0} \left[ \mathbb{E}_{z_{t+1}|h(b_t), a_t, \tau} \min_{\pi_{t+1}} lb(b_{t+1}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \min_{\pi_{t+1}} \hat{lb}(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right] \right| \quad (44) \end{aligned}$$

$$\begin{aligned} &\leq \left| \mathbb{1}_{\beta(h(b_{t+1}^-))=1} \left\{ \left| \mathbb{E}_{z_{t+1}|h(b_t), a_t, \tau} \max_{\pi_{t+1}} lb(b_{t+1}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \max_{\pi_{t+1}} lb(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right| \right. \right. \\ &\quad \left. \left. + \left| \frac{1}{C} \sum_{i=1}^C \max_{\pi_{t+1}} lb(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \max_{\pi_{t+1}} \hat{lb}(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right| \right\} \right. \\ &\quad \left. + \mathbb{1}_{\beta(h(b_{t+1}^-))=0} \left\{ \left| \mathbb{E}_{z_{t+1}|h(b_t), a_t, \tau} \min_{\pi_{t+1}} lb(b_{t+1}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \min_{\pi_{t+1}} lb(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right| \right. \right. \\ &\quad \left. \left. + \left| \frac{1}{C} \sum_{i=1}^C \min_{\pi_{t+1}} lb(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \min_{\pi_{t+1}} \hat{lb}(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right| \right\} \right| \quad (45) \end{aligned}$$

Here, we can use a technique similar to the previous proof. We can directly use the Hoeffding inequality to bound the first term in each indicator as:

$$\mathbb{P}\left( \left| \mathbb{E}_{z_{t+1}|h(b_t), a_t, \tau} \max_{\pi_{t+1}} lb(b_{t+1}, \pi_{t+1}, \tau) - \frac{1}{C} \sum_{i=1}^C \max_{\pi_{t+1}} lb(\bar{b}_{t+1}^{I_i}, \pi_{t+1}, \tau) \right| \leq \lambda \right) \geq 1 - 2 \exp\left(\frac{-C\lambda^2}{2V_{\max}^2}\right) \quad (46)$$

$$\mathbb{P}\left(\left|\mathbb{E}_{z_{t+1}|h(b_t),a_t,\tau}\min_{\pi_{t+1}}lb(b_{t+1},\pi_{t+1},\tau)-\frac{1}{C}\sum_{i=1}^C\min_{\pi_{t+1}}lb(\bar{b}_{t+1}^{I_i},\pi_{t+1},\tau)\right|\leq\lambda\right)\geq 1-2\exp\left(\frac{-C\lambda^2}{2V_{\max}^2}\right) \quad (47)$$

The second term is also iteratively bounded by  $\delta_{lb}^\tau(t+1)$ . So, combining the two probabilistic bounds together, we can get the iterative bound for the estimator satisfied with a probability of at least  $1 - 2|A|(|A|C)^{L-ts}\exp(\frac{-C\lambda^2}{2V_{\max}^2})$ :

$$\delta_{lb}^\tau(t) \leq (45) \leq \mathbb{1}_{\beta(h(b_{t+1}^-))=1}[\lambda + \delta_{lb}^\tau(t+1)] + \mathbb{1}_{\beta(h(b_{t+1}^-))=0}[\lambda + \delta_{lb}^\tau(t+1)] \quad (48)$$

$$= \lambda + \delta_{lb}^\tau(t+1) \quad (49)$$

The estimate error at the end of the planning horizon is:

$$\delta_{lb}^\tau(L-1) = \lambda \quad (50)$$

Now, we can find out the estimation bound at the top of the given belief tree by calculating it from the bottom to the top, from  $L-1$  to 0.

## 6 Experiment Settings

### 6.1 Experiment 1

In our current implementation, the belief tree  $\mathbb{T}^\tau$  that corresponds to an initial simplified topology  $\tau$  is constructed by expanding the original observation space only at randomly chosen 15% of the propagated belief nodes and switching the rest to an alternative observation space  $\mathcal{O} = \mathcal{X}$  (see (16)), i.e. considering the state space  $\mathcal{X}$  instead of the observation space  $\mathcal{Z}$ . Then, at each iteration, if the condition (12) is not satisfied, we switch to a different (less simplified) topology  $\tau'$  by turning 5 randomly-chosen propagated belief nodes in  $\tau$ , that had an alternative observation space, back to the original observation space. Once (12) is satisfied, we are guaranteed to find the optimal action  $a_k^*$ , and we terminate the process.

### 6.2 Experiment 2

The sparse sampling planner in a particle-belief POMDP setting has three parameters: planning horizon  $d$ , number of observation samples  $K$ , and number of the weighted state samples  $N$  that represent the particle belief. In our experiments, we evaluate our approach and the original sparse sampling solver using two sets of these parameters:  $d = 2, K = 50, N = 50$  and  $d = 2, K = 90, N = 90$ .

### 6.3 Experiment 3

The beacon navigation problem serves as a common benchmark for evaluating POMDP solvers [9, 10].



Displayed in Figure ??, the planning environment comprises a robot located at the initial belief, a beacon for generating observations, an obstacle, and two goals. The locations of the robot, beacon, obstacle, and goals are denoted as  $s$ ,  $x_b$ ,  $x_o$ , and  $x_{g1,g2}$ , respectively. The observation is modeled as the relative position with respect to the beacon, employing a Gaussian observation model defined as:

$$z \sim \begin{cases} \mathcal{N}(s - x_b, I/(100\|s - x_b\|)), & \text{if } \|s - x_b\| > 1, \\ \mathcal{N}(s - x_b, I/100), & \text{otherwise.} \end{cases} \quad (51)$$

The rewards are calculated based on the distance from the two goals:

$$r_g = \frac{10}{\|s - x_{g1}\| + 0.001} + \frac{20}{\|s - x_{g2}\| + 0.001} \quad (52)$$

Upon entering the nearby area of the obstacle, the robot incurs a significant penalty:

$$r_o = -100, \text{ if } \|s - x_o\| \leq 2. \quad (53)$$

The motion model follows a Gaussian distribution:  $P_T = \mathcal{N}(\mathbf{s} + \mathbf{a}, \Sigma_T)$ .

#### 6.4 Limitation

Our approach demonstrates effectiveness in scenarios characterized by a large observation space, which is consistent with the theoretical analysis of simplifying policy space. Despite requiring multiple iterations, our simplification strategy is still able to reduce computation time.

During the experiment, we also encountered failure cases in which nearly all nodes need to be un-simplified and use the original observation space so that the optimal action at the root could be distinguished. This implies that our method will eventually converge to a topology resembling the original one after numerous iterations, resulting in increased time costs. We address this dilemma as the trade-off between simplification and direct calculation, which all the simplification problems will face.

In addressing this dilemma, it is crucial to have a method that can intelligently and adaptively determine whether a situation is amenable to simplification. By evaluating the potential level of simplification before incurring high costs of explicit calculating, we can prune inadequate simplification candidates and further simplify the iteration process. We will focus on this problem in future work.

### 7 Proof of Lemma ??

*Proof.* **First, we consider the situation that the belief node uses the alternative observation space and model:**  $\beta^\tau(h_i^-) = 0$ .

For each  $x_i \in \mathcal{X}$ , we can find an action  $a_i$  that maximize and minimize the expected reward:

$$\int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \leq \max_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \quad (54)$$

$$\int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \geq \min_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \quad (55)$$

From (54), we have:

$$\begin{aligned} & \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \\ & \leq \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \max_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \end{aligned} \quad (56)$$

$$= \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau} \max_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \quad (57)$$

Similarly, from (55), we will get:

$$\begin{aligned} & \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \\ & \geq \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \min_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \end{aligned} \quad (58)$$

$$= \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau} \min_{a_i(x_i) \in \mathcal{A}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i) r(x_{i+1}) dx_{i+1} \quad (59)$$

Since we consider  $\bar{z}_i^T = o_i$  at the belief node for the topology and the alternative observation model is fully observable, the actions in Equation (57) and (59) can be viewed as a policy:  $a_i(x_i) = \pi_i, \pi_i \in \{\pi_i : \mathcal{H}_i^- \times \bar{\mathcal{Z}}_i^T \mapsto \mathcal{A}\} := \bar{\Pi}_i(\mathcal{H}_i^-, \bar{\mathcal{Z}}_i^T)$ .

$$\begin{aligned} & \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \\ & \leq \max_{\bar{\pi}_i \in \bar{\Pi}_i(\mathcal{H}_i^-, \bar{\mathcal{Z}}_i^T)} \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \bar{\pi}_i(h_i^-, \bar{z}_i^T)) r(x_{i+1}) dx_{i+1} \end{aligned} \quad (60)$$

$$\begin{aligned} & \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau_Z} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \pi_i^{\tau_Z}) r(x_{i+1}) dx_{i+1} \\ & \geq \min_{\bar{\pi}_i \in \bar{\Pi}_i(\mathcal{H}_i^-, \bar{\mathcal{Z}}_i^T)} \mathbb{E}_{x_i|h_i^-, \bar{z}_i|x_i, h_i^-} \mathbb{E}^{\tau} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \bar{\pi}_i(h_i^-, \bar{z}_i^T)) r(x_{i+1}) dx_{i+1} \end{aligned} \quad (61)$$

Since different policy  $\bar{\pi}_i$  upper and lower bound the expected closed-loop reward, we will also bound the distance between the expected closed-loop reward and a

known value by exploring the policy space:

$$\begin{aligned}
& \left| \mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau Z} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \pi_i^{\tau Z}) r(x_{i+1}) dx_{i+1} - \mathbb{E}_{x_i | h_i^{\tau-} \bar{z}_i | x_i, h_i^{\tau-} x_{i+1} | x_i, \pi_i^{\tau}} \mathbb{E}^{\tau} r(x_{i+1}) \right| \\
& \leq \max_{\bar{\pi}_i \in \bar{\Pi}_i(\mathcal{H}_i^-, \bar{\mathcal{Z}}_i^{\tau})} \left| \mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \bar{\pi}_i(h_i^-, \bar{z}_i^{\tau})) r(x_{i+1}) dx_{i+1} \right. \\
& \quad \left. - \mathbb{E}_{x_i | h_i^{\tau-} \bar{z}_i | x_i, h_i^{\tau-} x_{i+1} | x_i, \pi_i^{\tau}} \mathbb{E}^{\tau} r(x_{i+1}) \right| \tag{62}
\end{aligned}$$

**Then, we consider the belief node uses the original observation space and model:**  $\beta(h_i^-) = 1$ .

We have:

$$\mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau Z} \mathbb{E}_{x_{i+1} | x_i, \pi_i^{\tau Z}} r(x_{i+1}) = \mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau} \mathbb{E}_{x_{i+1} | x_i, \pi_i^{\tau}} r(x_{i+1}) \tag{63}$$

Here,  $\pi_i^{\tau Z}$  and  $\pi_i^{\tau}$  share the same mapping,  $\mathcal{H}_i^- \times \mathcal{Z}_i \mapsto \mathcal{A}$ . We can say that the optimal and worst  $\pi_i^{\tau}$  can upper and lower bound any  $\pi_i^{\tau Z}$ . Thus, there exists a policy  $\bar{\pi}_i^{\tau}$  that can bound the distance in Equation (62).

Here finishes the proof of Lemma ??.

## 8 Proof of Remark 2

$$\begin{aligned}
& \max_{\bar{\pi}_i \in \bar{\Pi}_i(\mathcal{H}_i^-, \bar{\mathcal{Z}}_i^{\tau})} \left| \mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \bar{\pi}_i^{\tau}) r(x_{i+1}) \right. \\
& \quad \left. - \mathbb{E}_{x_i | h_i^{\tau-} \bar{z}_i | x_i, h_i^{\tau-} x_{i+1} | x_i, \pi_i^{\tau}} \mathbb{E}^{\tau} r(x_{i+1}) \right| \tag{64}
\end{aligned}$$

$$\begin{aligned}
& \triangleq \left| \mathbb{E}_{x_i | h_i^-, \bar{z}_i | x_i, h_i^-} \mathbb{E}^{\tau} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \bar{\pi}_i^+(h_i^-, \bar{z}_i^{\tau})) r(x_i) \right. \\
& \quad \left. - \mathbb{E}_{x_i | h_i^{\tau-} \bar{z}_i | x_i, h_i^{\tau-} x_{i+1} | x_i, \pi_i^{\tau}} \mathbb{E}^{\tau} r(x_{i+1}) \right| \tag{65}
\end{aligned}$$

$$\begin{aligned}
& = \left| \mathbb{E}_{x_{i-1} | h_{i-1}^-, \bar{z}_{i-1} | x_{i-1}, h_{i-1}^-} \mathbb{E}^{\tau Z} \mathbb{E}_{x_i | x_{i-1}, \pi_{i-1}^{\tau C} \bar{z}_i | x_i, h_i^-} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \bar{\pi}_i^+(h_i^-, \bar{z}_i^{\tau})) r(x_i) \right. \\
& \quad \left. - \mathbb{E}_{x_i | h_i^{\tau-} \bar{z}_i | x_i, h_i^{\tau-} x_{i+1} | x_i, \pi_i^{\tau}} \mathbb{E}^{\tau} r(x_i) \right| \tag{66}
\end{aligned}$$

**i) If both  $\beta(h_{i-1}^-) = 0$  and  $\beta(h_i^-) = 0$ ,** for every possible  $x_i \in \mathcal{X}$  and  $x_{i-1} \in \mathcal{X}$ , we can find an action  $a_i^+(x_i)$  and  $a_{i-1}^+(x_{i-1})$  such that:

$$\begin{aligned}
& \mathbb{E}_{x_i | x_{i-1}, \pi_{i-1}^{\tau C}} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, \bar{\pi}_i^+) r(x_{i+1}) \\
& \leq \max_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i | x_{i-1}, a_{i-1}(x_{i-1})} \max_{a_i(x_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1} | x_i, a_i(x_i)) r(x_{i+1}) \tag{67}
\end{aligned}$$

Then we have:

$$\begin{aligned}
& \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^{\tau Z} \mathbb{E}_{x_{i-1},\pi_{i-1}^{\tau C}\bar{z}_i|x_i,h_i^-} \mathbb{E}^\tau \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,\bar{\pi}_i^+(h_i^-, \bar{z}_i^\tau)) r(x_{i+1}) \\
\leq & \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^{\tau Z} \max_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1},a_{i-1}(x_{i-1})\bar{z}_i|x_i,\{h_{i-1}^-,a_{i-1}\}} \mathbb{E}^\tau \\
& \max_{a_i(x_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,a_i(x_i)) r(x_{i+1}) \tag{68}
\end{aligned}$$

$$\begin{aligned}
= & \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^\tau \max_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1},a_{i-1}(x_{i-1})\bar{z}_i|x_i,\{h_{i-1}^-,z_{i-1}^\tau,a_{i-1}\}} \mathbb{E}^\tau \\
& \max_{a_i(x_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,a_i(x_i)) r(x_{i+1}) \tag{69}
\end{aligned}$$

Similarly, we can find the lower bound:

$$\mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^{\tau Z} \mathbb{E}_{x_{i-1},\pi_{i-1}^{\tau C}\bar{z}_i|x_i,h_i^-} \mathbb{E}^\tau \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,\bar{\pi}_i^+(h_i^-, \bar{z}_i^\tau)) r(x_{i+1}) \tag{70}$$

$$\begin{aligned}
\geq & \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^{\tau Z} \min_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1},a_{i-1}(x_{i-1})\bar{z}_i|x_i,\{h_i^-,a_{i-1}\}} \mathbb{E}^\tau \\
& \min_{a_i(x_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,a_i(x_i)) r(x_{i+1}) \tag{71}
\end{aligned}$$

$$\begin{aligned}
= & \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^\tau \min_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1},a_{i-1}(x_{i-1})\bar{z}_i|x_i,\{h_{i-1}^-,z_{i-1}^\tau,a_{i-1}\}} \mathbb{E}^\tau \\
& \min_{a_i(x_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,a_i(x_i)) r(x_{i+1}) \tag{72}
\end{aligned}$$

The  $a_{i-1}(x_{i-1})$  and  $a_i(x_i)$  in the above equations can be viewed as policy respectively,  $\pi_{i-1} \in \Pi_{i-1}(\mathcal{H}_{i-1}^-, \bar{\mathcal{Z}}_{i-1}^\tau)$  and  $\pi_i \in \Pi_i(\mathcal{H}_{i-1}^-, \bar{\mathcal{Z}}_{i-1}^\tau, \pi_{i-1}, \bar{\mathcal{Z}}_i^\tau)$ . Similar to Lemma ??, we can say there exists a policy to bound the below distance:

$$\begin{aligned}
& \left| \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^{\tau Z} \mathbb{E}_{x_{i-1},\pi_{i-1}^{\tau C}\bar{z}_i|x_i,h_i^-} \mathbb{E}^\tau \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,\bar{\pi}_i^+(h_i^-, \bar{z}_i^\tau)) r(x_{i+1}) \right. \\
& \left. - \mathbb{E}_{x_i|h_i^{\tau-}\bar{z}_i|x_i,h_i^{\tau-}} \mathbb{E}^\tau \mathbb{E}_{x_{i+1}|x_i,\pi_i^\tau} r(x_{i+1}) \right| \tag{73}
\end{aligned}$$

$$\begin{aligned}
\leq & \max_{\bar{\pi}_{i-1} \in \bar{\Pi}_{i-1}(\mathcal{H}_{i-1}^-, \bar{\mathcal{Z}}_{i-1}^\tau), \bar{\pi}_i(h_{i-1}^-, z_{i-1}^\tau, \bar{\pi}_{i-1}(h_{i-1}^-, z_{i-1}^\tau), z_i^\tau)} \\
& \left| \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^\tau \mathbb{E}_{x_{i-1},\bar{\pi}_{i-1}\bar{z}_i|x_i,h_i^{p-}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,\bar{\pi}_i) r(x_{i+1}) \right. \\
& \left. - \mathbb{E}_{x_i|h_i^{\tau-}\bar{z}_i|x_i,h_i^{\tau-}} \mathbb{E}^\tau \mathbb{E}_{x_{i+1}|x_i,\pi_i^\tau} r(x_{i+1}) \right| \tag{74}
\end{aligned}$$

$$\begin{aligned}
& \left| \mathbb{E}_{x_{i-1}|h_{i-1}^-\bar{z}_{i-1}|x_{i-1},h_{i-1}^-} \mathbb{E}^\tau \mathbb{E}_{x_{i-1},\bar{\pi}_{i-1}\bar{z}_i|x_i,h_i^{p-}} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i,\bar{\pi}_i) r(x_{i+1}) \right. \\
& \left. - \mathbb{E}_{x_i|h_i^{\tau-}\bar{z}_i|x_i,h_i^{\tau-}} \mathbb{E}^\tau \mathbb{E}_{x_{i+1}|x_i,\pi_i^\tau} r(x_{i+1}) \right| \tag{75}
\end{aligned}$$

ii) If  $\beta(h_{i-1}^-) = 0$  and  $\beta(h_i^-) = 1$ , for each possible  $x_{i-1} \in \mathcal{X}$  and each  $z_i \in \mathcal{Z}_i$ :

$$\begin{aligned} & \mathbb{E}_{x_i|x_{i-1}, \pi_{i-1}^{\tau C}} \mathbb{E}_{\bar{z}_i|x_i} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \bar{\pi}_i^+(h_{i-1}^-, z_{i-1}^{\tau Z}, \pi_{i-1}^{\tau C}, z_i^{\tau})) r(x_{i+1}) \\ & \leq \max_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1}, a_{i-1}(x_{i-1})} \mathbb{E}_{\bar{z}_i|x_i} \max_{a_i(z_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i(z_i)) r(x_{i+1}) \end{aligned} \quad (76)$$

$$\begin{aligned} & \mathbb{E}_{x_i|x_{i-1}, \pi_{i-1}^{\tau C}} \mathbb{E}_{\bar{z}_i|x_i} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, \bar{\pi}_i^+(h_{i-1}^-, z_{i-1}^{\tau Z}, \pi_{i-1}^{\tau C}, z_i^{\tau})) r(x_{i+1}) \\ & \geq \min_{a_{i-1}(x_{i-1})} \mathbb{E}_{x_i|x_{i-1}, a_{i-1}(x_{i-1})} \mathbb{E}_{\bar{z}_i|x_i} \min_{a_i(z_i)} \int_{x_{i+1}} \mathbb{P}(x_{i+1}|x_i, a_i(z_i)) r(x_{i+1}) \end{aligned} \quad (77)$$

The  $a_{i-1}(x_{i-1})$  and  $a_i(z_i)$  in the above equations can be viewed as policy respectively,  $\pi_{i-1}(h_{i-1}^-, z_{i-1}^{\tau})$  and  $\pi_i(h_{i-1}^-, z_{i-1}^{\tau}, \pi_{i-1}(h_{i-1}^-, z_{i-1}^{\tau}), z_i^{\tau})$ . Then we can get the same result as Equation (75).

iii) If both  $\beta(h_{i-1}^-) = 1$  and  $\beta(h_i^-) = 0$ , the result is similar.

iv) If  $\beta(h_{i-1}^-) = 1$  and  $\beta(h_i^-) = 1$ , the policy  $\bar{\pi}_{i-1,i}$  in Equation (75) shares the same mapping as the closed-loop policy  $\pi_{i-1,i}^{\tau C}$ . So there exists the optimal and worst policy that upper and lower bound the expected reward, so we can also get the same result as Equation (75).

## References

1. D. Kong and V. Indelman. Simplified belief space planning with an alternative observation space and formal performance guarantees. In *Proc. of the Intl. Symp. of Robotics Research (ISRR)*, 2024. Submitted.