

Locally Robust Policy Learning: Inequality, Inequality of Opportunity and Intergenerational Mobility*

Joël Terschuur

Universidad Carlos III de Madrid

October 12, 2023

[Click here for the latest version of the paper.](#)

Abstract

Policy makers need to decide whether to treat or not to treat different individuals. The optimal choice depends on the welfare function that the policy maker has in mind. I study a general setting for policy learning with general semiparametric Social Welfare Functions (SWFs), possibly defined by semiparametric U-statistics, which accomodate a wide range of distributional preferences. I use locally robust/orthogonal scores to provide strong statistical guarantees for the estimated policy rules even in observational settings where the propensity score is unknown. This work expands previous results in [Athey and Wager \(2021\)](#). Three main applications of the general theory motivate the paper: (i) Inequality aware SWFs, (ii) Inequality of Opportunity aware SWFs and (iii) Intergenerational Mobility SWFs. I use the Panel Study of Income Dynamics (PSID) to asses the effect of attending preschool on adult earnings and estimate optimal policy rules based on parental years of education and parental income.

JEL Classification: C13; C14; C21; D31; D63; I24

Keywords: local robustness, U-statistics, Inequality, Intergenerational mobility, empirical welfare maximization.

R package (forthcoming): <https://joelters.github.io/home/code/>

*Research funded by Ministerio de Ciencia e Innovación, grant PRE2020-092794. I am grateful to Juan Carlos Escanciano and Toru Kitagawa for their guidance and support, and to Miguel Ángel Delgado, Juan José Dolado, Ignacio Ortuño, Nazarii Salish, Jan Stuhler and Carlos Velasco. *Email address:* jrobert@eco.uc3m.es

1 Introduction

Whenever a policy or treatment has heterogeneous effects it is important to decide carefully who should be treated. In the simplest case in which we care only about the average outcome, there are no budgetary limits and the treatment effect is positive for everyone; it follows that the best policy is to treat everyone. However, in most cases we do not have such a luxury. We might have a limited budget, distributional concerns or negative treatment effect for some individuals. In these cases, it is important to decide whether *to treat or not to treat* different individuals. This is the problem of policy learning.

Such a problem is omnipresent not only in economics but in business, law, education, medicine and many others. While in economics we might want to know whether to provide training or not to unemployed, decide different rules to assign conditional cash transfers or even whether a transfer should be given unconditionally, in business we might want to know whether to provide a discount to a customer or not, or whether we should send price recommendations to some stores and not to others. Judges might have to decide whether to release someone on parole or not based on the reincidence probability of the individual. In education we might want to know whether to provide a scholarship to a student or not or whether we should provide additional extra-curricular lessons. Certain medicines might be beneficial for some but prejudicial for others or there might not be enough vaccines to cover the whole population as seen in the COVID-19 pandemic.

The inherent ethical and distributional considerations in all these examples are quite different. Hence, it is important to have a general framework which can accomodate different welfare functions. While the framework has to be as general as possible, it also needs to allow for estimation of optimal policy rules with certain statistical guarantees. In this paper I provide a framework which allows to compute such optimal rules for a rich class of semiparametric welfare functions, possibly defined as U-statistics. This includes, among many others, the standard utilitarian welfare function but also: (i) Inequality aware SWFs, (ii) Inequality of Opportunity (IOp) aware SWFs and (iii) Intergenerational Mobility SWFs. These three SWFs motivate and serve as guiding examples in this paper.

To my knowledge, there is no prior work on IOp and Intergenerational mobility aware social welfare functions in the policy learning literature. IOp is the part of inequality which is explained by circumstances X outside the control of the individual, e.g. sex, race, parental education or parental income. Based on the seminal contributions in [Van De Gaer \(1993\)](#), [Fleurbaey \(1995\)](#) and [Roemer \(1998\)](#) the IOp literature has grown and focused on how to measure IOp. A popular measure of IOp is the Gini of the best predictions (in mean squared error sense) of the outcome Y given the circumstances X , i.e. $G(\gamma(X))$ where $\gamma(X) = \mathbb{E}[Y|X]$. In order to accomodate a possibly high-dimensional set of circumstances, IOp literature has started using machine learners to estimate the predictions (e.g [Brunori et al. \(2019a\)](#), [Brunori et al. \(2019b\)](#), [Brunori et al.](#)

(2021), Brunori and Neidhöfer (2021), Rodríguez et al. (2021), Carranza (2022) or Hufe et al. (2022) among others). As usual in such two-step procedures, the bias-variance trade-off in the prediction might allow for some bias which can creep into the second stage. Escanciano and Terschuur (2022) provide locally robust IOp estimators which are robust to such biases. I use these results to construct IOp aware SWFs.

Inequality aware SWFs have been studied before in Kitagawa and Tetenov (2021). A popular welfare function for a random outcome Y is $W = \mathbb{E}[Y](1 - G(Y))$, where $G(Y)$ is the Gini of the distribution of Y . This welfare function values the average outcome but penalizes high inequality (the Gini is between 0 and 1 where 0 is complete equality and 1 complete inequality). My framework allows for a simpler way to deal with the case in which the population eligible for treatment and the population whose welfare we care about differ. For instance, we might want to study how to allocate some transfer among the poor but care about the inequality in the whole population. I do so by using the fact that the Gini coefficient can be written as a U-statistic which prevents me from having to use the c.d.f. of the outcome as in Kitagawa and Tetenov (2021).

While inequality aware SWFs look at the distribution of Y , IOp aware SWFs focus on the distribution of the predictions $\gamma(X) = \mathbb{E}[Y|X]$. So an example of a natural IOp aware SWF would be $W = \mathbb{E}[\gamma(X)](1 - G(\gamma(X))) = \mathbb{E}[Y](1 - G(\gamma(X)))$, which only penalizes inequalities explained by circumstances. This example adds an extra unknown nuisance parameter, $\gamma(X)$, on top of the conditional expectations/propensity scores needed to identify treatment effects. To my knowledge there is no previous work on policy learning with general semiparametric welfare functions which depend on additional unknown functions besides from those needed to identify treatment effects.

Intergenerational mobility is the study of the relationship between the outcomes of parents and the outcomes of their children. A popular measure of intergenerational mobility is the rank correlation between the income of parents and the income of their children. This measure is known as the Kendall- τ and is a popular measure of intergenerational mobility in the literature (see Chetty et al. (2014) or Kitagawa et al. (2018)). A natural intergenerational mobility aware SWF would be $W = -|\tau - t|$ for some target rank correlation $t \in [-1, 1]$. This example is of interest in, for instance, deciding the allocation of higher education scholarships to students based on individual characteristics whenever the treatment effect of education on long term income can be identified and there is a policy interest in reducing the association between parental income and child's income.

The technical goal in the policy learning literature, sometimes called empirical welfare maximization or offline policy learning in the computer science literature, is to find an optimal allocation rule π which maps individual characteristics to a binary decision $\{0, 1\}$ of treatment or no treatment. This optimal rule is searched in a class Π of plausible treatment rules to

maximize some welfare function. Following the seminal work in [Manski \(2004\)](#), I search for an optimal policy in the plausible class so as to minimize regret, i.e. the expected difference between the best possible welfare and the welfare evaluated at the estimated policy. Other relevant work on treatment rules in econometrics includes [Dehejia \(2005\)](#), [Hirano and Porter \(2009\)](#), [Stoye \(2009, 2012\)](#), [Chamberlain \(2011\)](#), [Bhattacharya and Dupas \(2012\)](#), [Tetenov \(2012\)](#), [Kasy \(2016\)](#), [Kitagawa and Tetenov \(2018, 2021\)](#), [Athey and Wager \(2021\)](#), or [Zhou et al. \(2023\)](#).

Non-parametric estimation of the unknown functions in semiparametric welfare functions poses a challenge to the statistical guarantees of estimated policy rules. This is due to the slow convergence rate of non-parametric estimators such as kernels or machine learners. The semiparametric literature has developed methods to overcome this problem by using locally robust/orthogonal scores. These are alternative moment conditions which identify the quantity of interest and allow for its estimation at parametric (\sqrt{n}) rate. I expand previous work by considering any semiparametric welfare function, possibly defined as a U-statistic, which can be estimated by locally robust/orthogonal scores. The main theoretical result is to provide an asymptotic upper bound to the regret of the estimated policy rule.

This paper is close to [Kitagawa and Tetenov \(2021\)](#) in taking into account other distributional aspects besides from the mean and it is also closely related to [Athey and Wager \(2021\)](#) and [Zhou et al. \(2023\)](#) in making use of the latest semiparametric literature on locally robust/orthogonal scores to obtain parametric rates of convergence even with slow nonparametric first steps. I also build upon [Escanciano and Terschuur \(2022\)](#) to expand policy learning results to welfare functions defined by U-statistics. The key result in [Athey and Wager \(2021\)](#) is to find rates of the regret which optimally depend on the complexity of Π and the sample size in observational settings where propensity scores are unknown. They do so for utilitarian average-treatment-like welfare functions. I substantially generalize this setting by allowing arbitrary semiparametric welfare functions, possibly defined as U-statistics, as long as they can be identified with locally robust/orthogonal scores.

Empirically, treatment allocation with inequality, IOp and IGM welfare functions poses many challenges. First, we need outcomes for which it makes sense to look at inequality. For instance, standardized test scores are not suitable for inequality measures while income is. Second, we need rich information on circumstances and parental income which are absent in many modern datasets. Third, we need a credible identification strategy to identify treatment effects. In this paper I tackle these challenges by looking at the effect of attending preschool on adult earnings using the Panel Study of Income Dynamics (PSID) dataset. This application has many advantages. First, any variable which induces preschool attendance can be considered a circumstance under the (very reasonable) assumptions that we cannot hold the kid responsible for these variables. Second, PSID has been following families for nearly 50 years meaning we have rich information on family background. Third, PSID allows us to look at long term outcomes

such as adult earnings. The main disadvantage is that the treatment is not randomly assigned. Hence, the identification of treatment effects relies on the assumption of selection on observable circumstances.

The effect of preschool on short/medium/long-term outcomes has been extensively studied and there is a public interest in expanding public preschool programs in the US. The share of 4 year olds in public preschool has grown from 14% in 2002 to 34% in 2019 and many states and large cities in the US now operate large-scale public preschool programs (Gray-Lobe et al. (2023)). The first popular small-scale randomized preschool experiments in the US were the High/Scope Perry Preschool project and Carolina Abecedarian project whose participants have been followed for decades leading to many studies showing positive effects (Campbell and Ramey (1994), Campbell et al. (2012), Heckman et al. (2013), García et al. (2020)). Gray-Lobe et al. (2023) use admission lotteries to study large-scale public preschool in Boston on a range of outcomes. They find positive effects and varying treatment effects based on gender. Heckman and Raut (2016) study the impact of preschool using a structural model and find that a tax-financed public preschool program targeted at children with poor socioeconomic status increases average earnings and increases intergenerational mobility.

In this paper I find that the effect of preschool attendance is heterogenous. While on average preschool has a positive effect on adult earnings, children with highly educated mother’s and high parental income are negatively affected by preschool. This is in line with results in the psychology and economics literature which document that in early educational institutions there is less interaction with adults and hence there can be a negative effect of attending such institutions if the interactions with adults in the household are of ”high-quality” (see Fort et al. (2020)).

This heterogenous effects have different implications when estimating optimal treatment rules for different welfare functions. I compute optimal treatment rules based on parental income and mother’s education in a class of decision trees of depth two. Utilitarian estimated optimal rules try to treat anyone who has a positive treatment effect. IOp and inequality aware SWFs exclude individuals from treatment even if they have a positive treatment effect if the decrease in average earnings is compensated by a decrease in inequality. An undesirable feature of these welfare functions is that they can assign children who are negatively impacted by preschool to treatment. This happens if the decrease in the average outcome is compensated by a decrease in inequality. This possibility can be ruled out by restricting to trees who do not treat groups with negative estimated treatment effects.

I start by introducing the main welfare objects which are going to serve as guiding examples of the general theory in Section 2. Section 3 elaborates on the general theory for general welfare functions identified by locally robust/orthogonal scores which are linear on the distribution of the data (i.e. not defined as U-statistics) and Section 4 expands the results to general welfare functions, possibly defined as U-statistics. Section 5 provides upper bounds on the regret of

estimated policies and Section 6 deals with the empirical application. All proofs are in the Appendix.

2 Welfare economics for inequality, IOp and rank correlations

The policy learning literature is at the intersection of welfare economics and econometrics. Before we delve into the econometric problem of computing optimal rules and evaluating their statistical performance I present in this section the main welfare objects we are going to be interested in. The most basic welfare function is that of the average outcome. Suppose we have some continuous random outcome $Y_i \in \mathbb{R}^+$. A utilitarian planner cares about

$$W = \mathbb{E}[Y_i].$$

The above welfare does not care about other distributional aspects besides from the average outcome. A first approach to include distributional concerns in our analysis is to follow Dalton (1920) and Atkinson et al. (1970) and consider increasing and concave transformations $u(\cdot)$ of the outcome¹.

$$W = \mathbb{E}[u(Y_i)].$$

This welfare function will already rank two outcome distributions in the same way for all increasing and concave $u(\cdot)$ if the Lorenz curve of one of the distributions is everywhere above the Lorenz curve of the other distribution and has equal or higher mean; equivalently, if one distribution second order stochastically dominates the other. However, if we want to obtain a complete ordering we need to specify $u(\cdot)$ further. One popular choice is

$$u(y) = \begin{cases} \frac{y^{1-\theta}}{1-\theta} & \text{if } \theta \in (0, 1) \\ \log(y) & \text{if } \theta = 1. \end{cases}$$

θ captures the concavity of $u(\cdot)$ and can therefore be interpreted as an inequality aversion parameter. This paper also focuses on welfare which is aware of Inequality of Opportunity (IOp). IOp is the part of total inequality which can be explained by circumstances, i.e. by variables that are outside the control of the individual such as parental education or parental income. Let $X_i \in \mathbb{R}^k$ be such a random vector of circumstances. Let also $\gamma(X_i) = \mathbb{E}[Y_i|X_i]$, i.e. the best predictor (in mean squared error sense) of the outcome Y_i given the circumstances X_i . By looking at the distribution of $\gamma(X_i)$ instead of that of the outcome Y_i we get IOp averse welfare functions. For instance,

$$W = \mathbb{E}[u(\gamma(X_i))].$$

¹With abuse of notation we call W to all welfare functions as they appear.

If there is no IOp, circumstances are unable to predict the outcome and we have that the best predictor is the unconditional mean: $\gamma(X_i) = \mathbb{E}[Y_i]$. In this case we have that $W = u(\mathbb{E}[Y_i])$ so we only care about the average income (with a different scale due to $u(\cdot)$). If we have maximum IOp, the outcome is a deterministic function of the circumstances and $\gamma(X_i) = Y_i$. Then, $W = \mathbb{E}[u(Y_i)]$. Since all inequality is IOp, we are back at the inequality averse welfare function.

Another option to take into account distributional concerns is to weight differently different parts of the distribution. Let F_Y be the distribution of the outcome and F_Y^{-1} be the quantiles. Then, for some weights $w(\cdot)$ a planner might have the following welfare in mind

$$W = \int_0^1 F_Y^{-1}(\tau) w(\tau) d\tau.$$

This welfare has been used in [Mehran \(1976\)](#), [Donaldson and Weymark \(1980\)](#), [Weymark \(1981\)](#), [Donaldson and Weymark \(1983\)](#) or [Aaberge et al. \(2021\)](#). If we let $w_k(\tau) = (k-1)(1-\tau)^{k-2}$ we get what is known as the extended Gini family of social welfare functions. In this paper we focus on $k = 3$ which is known as the standard Gini social welfare function and can be shown to be

$$\begin{aligned} W &= \mathbb{E}[Y_i](1 - G(Y_i)) \\ &= (1/2)\mathbb{E}[Y_i + Y_j - |Y_i - Y_j|], \end{aligned}$$

where $G(Y_i)$ is the Gini coefficient of the distribution of Y_i , the second equality follows from the fact that we can write the Gini of Y_i as $G(Y_i) = \mathbb{E}[|Y_i - Y_j|]/\mathbb{E}[Y_i + Y_j]$ where Y_j is a copy of Y_i (i.e. the Gini can be interpreted as a normalized absolute distance between the outcomes of two individuals taken at random). The welfare above is utilitarian as long as there is no inequality ($G(Y_i) = 0$) and penalizes positive values of the Gini coefficient. Again, if we do not care about inequality but only about IOp we can look at the distribution of $\gamma(X_i)$ instead of the distribution of Y_i . In that case we have

$$\begin{aligned} W &= \mathbb{E}[\gamma(X_i)](1 - G(\gamma(X_i))) \\ &= (1/2)\mathbb{E}[\gamma(X_i) + \gamma(X_j) - |\gamma(X_i) - \gamma(X_j)|]. \end{aligned}$$

If there is no IOp, then $G(\gamma(X_i)) = 0$ and we are back in the utilitarian case. If there is full IOp, then $G(\gamma(X_i)) = G(Y_i)$ and we are back to the standard Gini social welfare function of outcome Y_i . Finally, I also consider the problem of intergenerational mobility. Let $X_{1i} \in \mathbb{R}$ be the first component of X_i . A measure of rank correlation between Y_i and X_{1i} is the Kendall- τ

$$\tau = \mathbb{E}[\text{sgn}(Y_i - Y_j)\text{sgn}(X_{1i} - X_{1j})],$$

where $\text{sgn}(a) = \mathbb{1}(a > 0) - \mathbb{1}(a < 0)$. This parameter is popular in the intergenerational mobility literature (see [Chetty et al. \(2014\)](#) or [Kitagawa et al. \(2018\)](#)) where X_{1i} is parental income and

Y_i is the child's income. It takes values between 1 and -1 . $\tau = 1$ means perfect rank correlation, whenever an individual has higher income than another, she also has higher parental income and viceversa. $\tau = -1$ is the opposite, whenever someone has higher income, she has lower parental income. $\tau = 0$ means the ranks are not correlated. For some target rank correlation $t \in [-1, 1]$ an intergenerational mobility aware welfare function is

$$W = -\left| \mathbb{E} \left[\text{sgn}(Y_i - Y_j) \text{sgn}(X_{1i} - X_{1j}) \right] - t \right|.$$

3 Policy learning with general orthogonal scores

Consider random variables $(Y_i(1), Y_i(0), D_i, X_i) \sim F_0$ where $(Y_i(1), Y_i(0)) \in \mathcal{Y} \times \mathcal{Y}$ are real-valued potential outcomes, i.e. $Y_i(1)$ is the outcome of individual i under treatment and $Y_i(0)$ is the outcome of individual i in the absence of treatment. D_i is a binary treatment and $X_i \in \mathcal{X}$ is a vector of pre-treatment covariates. Let $\gamma^{(j)}(X_i) = \mathbb{E}[Y_i(j)|X_i] \in \Gamma$ for $j = 0, 1$ be potential predictions, i.e. the predictions of the potential outcomes given X_i . We observe an i.i.d. sample (Z_1, \dots, Z_n) with $Z_i = (Y_i, D_i, X_i) \in \mathcal{Z}$ and $Y_i = Y_i(1)D_i + Y_i(0)(1 - D_i) \in \mathcal{Y}$. Let $\pi : \mathcal{X} \mapsto \{0, 1\}$ be a treatment rule which indicates who receives treatment and Π be a collection of such treatment rules. We are interested in choosing a policy $\pi \in \Pi$ so as to maximize the following welfare function

$$W(\pi) = \mathbb{E}[g(Y_i(1), X_i, \gamma^{(1)})\pi(X_i) + g(Y_i(0), X_i, \gamma^{(0)})(1 - \pi(X_i))]. \quad (3.1)$$

Importantly, (3.1) depends on possibly infinite dimensional unknown nuisance parameters γ . While throughout the paper I consider γ to be a conditional expectation of the outcome given X , this framework allows for much more general first steps such as high-dimensional quantile regressions, see (2.10) in [Chernozhukov et al. \(2021\)](#).

Example 1 (IOp Atkinson) *If we are interested in an inequality aware SWF we can use Atkinson SWF, $W(\pi) = \mathbb{E}[u(Y_i(1))\pi(X_i) + u(Y_i(0))(1 - \pi(X_i))]$ with $u(\cdot)$ a concave function. In this case the optimal policy can be estimated using the methods in [Kitagawa and Tetenov \(2018\)](#) and [Athey and Wager \(2021\)](#). If we want an IOp aware SWF we can simply look at the distribution of $\gamma(X_i)$ instead of at the distribution of Y_i :*

$$W(\pi) = \mathbb{E}[u(\gamma^{(1)}(X_i))\pi(X_i) + u(\gamma^{(0)}(X_i))(1 - \pi(X_i))].$$

■

(3.1) is not observable since for a given individual we do not observe both potential outcomes. In order to identify (3.1) we first need our sample to come from an experimental or observational experiment where the policy has already been implemented and where the following holds.

Assumption 1 *i)* $(Y_i(1), Y_i(0)) \perp D_i | X_i$,

ii) *There exists $\kappa \in (0, 1/2]$ such that $e(x) \in [\kappa, 1 - \kappa]$.*

The next proposition states the first identification result. There are two ways of identifying the welfare, either using what is usually called the direct method (DM) based on conditional expectations or using Inverse Propensity Score Weighting (IPW). Let $\gamma(D_i, X_i) = \mathbb{E}[Y_i | D_i, X_i]$, $\gamma_j(X_i) = \gamma(j, X_i)$ for $j = 0, 1$ and $\varphi(D_i, X_i, \gamma) = \mathbb{E}[g(Y_i, X_i, \gamma) | D_i, X_i]$.

Proposition 3.1 *Under Assumption 1, $W(\pi)$ is identified as*

$$W(\pi) = \mathbb{E}[m_1(Z_i, \gamma, \nu)\pi(X_i) + m_0(Z_i, \gamma, \nu)(1 - \pi(X_i))],$$

with $\nu \in \{\varphi, e\}$ and where m_1 and m_0 can be any of the following

$$\begin{aligned} (DM) \quad m_1(Z_i, \gamma, \varphi) &= \varphi(1, X_i, \gamma_1), \quad m_0(Z_i, \gamma, \varphi) = \varphi(0, X_i, \gamma_0) \\ (IPW) \quad m_1(Z_i, \gamma, e) &= \frac{g(Y_i, X_i, \gamma_1)D_i}{e(X_i)}, \quad m_0(Z_i, \gamma, e) = \frac{g(Y_i, X_i, \gamma_0)(1 - D_i)}{1 - e(X_i)}. \end{aligned}$$

Note that if g only depends on potential nuisance parameters but not on actual potential outcomes directly, i.e. $g(u, X_i, \gamma^{(j)}) = g(t, X_i, \gamma^{(j)}) \equiv g(X_i, \gamma^{(j)})$ for all $u, t \in \mathcal{Y}$ then φ is known since $\varphi(D_i, X_i, \gamma) = g(X_i, \gamma)$. This is the case in all IOp examples such as Example 1. Hence, depending on which case we are, we will have either γ or (γ, φ) or (γ, e) as nuisance parameters. To enjoy local robustness to high dimensional and ML first steps I provide orthogonal scores in the next result. First I need the following assumption to take care of the nuisance parameter γ .

Assumption 2 *There exist (α_1, α_0) such that for any $\tilde{\gamma} \in L_2$ and $j = 0, 1$ and $\tau \geq 0$*

$$\left. \frac{d}{d\tau} \mathbb{E}[m_j(Z_i, \bar{\gamma}_\tau, \nu)] \right|_{\tau=0} = \left. \frac{d}{d\tau} \mathbb{E}[\alpha_j(D_i, X_i) \bar{\gamma}_\tau(D_i, X_i)] \right|_{\tau=0},$$

where $\bar{\gamma}_\tau = \gamma + \tau \tilde{\gamma}$.

This is a common assumption in the semiparametric and orthogonal moments literature (e.g. (4.1) in Newey (1994)). Since γ enters m_j only through g a sufficient condition is to assume a similar result for the function g instead of m_j and then it will be straightforward to find α . Orthogonal scores usually take the form of the original identifying score plus mean zero correction terms based on residuals which make the score locally robust to first steps.

Proposition 3.2 *The orthogonal score is given by*

$$\Gamma_i(\pi) = \Gamma_{1i}\pi(X_i) + \Gamma_{0i}(1 - \pi(X_i)),$$

where

$$\begin{aligned}\Gamma_{1i} &= \varphi(1, X_i, \gamma) + \frac{D_i}{e(X_i)}(g(Y_i, X_i, \gamma_1) - \varphi(1, X_i, \gamma)) + \alpha_1(D_i, X_i)(Y_i - \gamma(D_i, X_i)), \\ \Gamma_{0i} &= \varphi(0, X_i, \gamma) + \frac{1 - D_i}{1 - e(X_i)}(g(Y_i, X_i, \gamma_0) - \varphi(0, X_i, \gamma)) + \alpha_0(D_i, X_i)(Y_i - \gamma(D_i, X_i)).\end{aligned}$$

As expected, orthogonal scores are formed by identifying scores (φ already identifies the welfare) and correction terms for nuisance parameters φ and γ . Note that whenever g does not depend on the potential outcomes directly we have that $g(Y_i, X_i, \gamma_j) - \varphi(j, X_i, \gamma) = 0$ for $j = 0, 1$ so we have $\Gamma_{1i} = \varphi(1, X_i, \gamma) + \alpha_1(D_i, X_i, g)(Y_i - \gamma(D_i, X_i))$ and $\Gamma_{0i} = \varphi(0, X_i, \gamma) + \alpha_0(D_i, X_i, g)(Y_i - \gamma(D_i, X_i))$. To estimate the welfare for a given $\pi \in \Pi$ we employ cross-fitting as in [Chernozhukov et al. \(2021\)](#). Let the data be split in L groups I_1, \dots, I_L , then

$$\hat{W}_n(\pi) = \frac{1}{n} \sum_{l=1}^L \sum_{i \in I_l} \hat{\Gamma}_{1i,l} \pi(X_i) + \hat{\Gamma}_{0i,l} (1 - \pi(X_i)),$$

where

$$\begin{aligned}\hat{\Gamma}_{1i,l} &= \hat{\varphi}_l(1, X_i, \hat{\gamma}_l) + \frac{D_i}{\hat{e}_l(X_i)}(Y_i - \hat{\varphi}_l(1, X_i, \hat{\gamma}_l)) + \hat{\alpha}_{1,l}(D_i, X_i, \nu)(Y_i - \hat{\gamma}_l(D_i, X_i)), \\ \hat{\Gamma}_{0i,l} &= \hat{\varphi}_l(0, X_i, \hat{\gamma}_l) + \frac{1 - D_i}{1 - \hat{e}_l(X_i)}(Y_i - \hat{\varphi}_l(0, X_i, \hat{\gamma}_l)) + \hat{\alpha}_{0,l}(D_i, X_i, \nu)(Y_i - \hat{\gamma}_l(D_i, X_i)),\end{aligned}$$

and $(\hat{\varphi}_l, \hat{e}_l, \hat{\gamma}_l, \hat{\alpha}_{j,l})$, $j = 0, 1$, are estimators of the nuisance functions which do not use observations in I_l . Again, whenever g does not depend on the potential outcomes the middle term in both expressions is zero. This is the case in the example of Atkinson welfare IOp.

Example 1 (IOp Atkinson (cont.)) For $\theta \in (0, 1]$ let

$$U(\gamma(x)) = \begin{cases} \frac{\gamma(x)^{1-\theta}}{1-\theta} & \text{if } \theta \in (0, 1) \\ \log(\gamma(x)) & \text{if } \theta = 1. \end{cases}$$

θ controls the concavity of U and therefore is a parameter capturing inequality aversion which can be picked by the policy maker. In this case $g = U$ which only depends on the nuisance parameters. The orthogonal score for $\theta \in (0, 1]$ is

$$\begin{aligned}\Gamma_i(\pi) &= U(\gamma(1, X_i)) + \frac{\gamma(D_i, X_i)^{-\theta} D_i}{e(X_i)}(Y_i - \gamma(D_i, X_i))\pi(X_i) \\ &\quad + U(\gamma(0, X_i)) + \frac{\gamma(D_i, X_i)^{-\theta} (1 - D_i)}{1 - e(X_i)}(Y_i - \gamma(D_i, X_i))(1 - \pi(X_i)),\end{aligned}$$

i.e. $\alpha_1(D_i, X_i, g) = e(X_i)^{-1} \gamma(D_i, X_i)^{-\theta} D_i$ and $\alpha_0(D_i, X_i, g) = (1 - e(X_i))^{-1} \gamma(D_i, X_i)^{-\theta} (1 - D_i)$.

■

The estimator of the optimal treatment rule among a class of rules Π is

$$\hat{\pi} = \arg \max_{\pi \in \Pi} \hat{W}_n(\pi).$$

Before analysing the statistical performance of such rule let us first extend the results in this section to welfare functions based on U-statistics. This will allow us to consider inequality and IOp aware SWFs based on the Gini coefficient and also a rank correlation aware SWF which is relevant for intergenerational mobility.

4 Policy learning with U-statistics

Let now $\pi_{ab}(X_i, X_j) = \mathbb{1}(\pi(X_i) = a) \times \mathbb{1}(\pi(X_j) = b)$ with $a, b \in \{0, 1\}$. Now we consider the following SWFs

$$W(\pi) = \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} g(Y_i(a), X_i, Y_j(b), X_j, \gamma^{(a)}, \gamma^{(b)}) \pi_{ab}(X_i, X_j) \right]. \quad (4.1)$$

Example 2 (Inequality) *We can accommodate the standard Gini welfare function with*

$$g(Y_i(a), Y_j(b)) = (1/2)(Y_i(a) + Y_j(b) - |Y_i(a) - Y_j(b)|).$$

■

Example 3 (Inequality of Opportunity IOp) *We can apply the standard Gini welfare function to the distribution of the predictions to get $\mathbb{E}[\gamma(X_i)](1 - G(\gamma(X_i)))$. This fits our setting by letting*

$$g(X_i, X_j, \gamma^{(a)}, \gamma^{(b)}) = (1/2)(\gamma^{(a)}(X_i) + \gamma^{(b)}(X_j) - |\gamma^{(a)}(X_i) - \gamma^{(b)}(X_j)|).$$

■

Example 4 (Rank correlation) *If we want to allocate a treatment targeting a specific Kendall- τ , say $t \in \mathbb{R}$, we have to extend our setting to transformations of the RHS of 4.1. We can define*

$$g(Y_i(a), X_{1i}, Y_j(b), X_{1j}) = \text{sgn}(Y_i(a) - Y_j(b)) \text{sgn}(X_{1i} - X_{1j}),$$

and let

$$W(\pi) = - \left| \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} g(Y_i(a), X_{1i}, Y_j(b), X_{1j}) \pi_{ab}(X_i, X_j) \right] - t \right|.$$

■

For $a, b \in \{0, 1\}$ let now $\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) = \mathbb{E}[g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) | D_i = a, X_i, D_j = b, X_j]$ and $e_{ab}(X_i, X_j) = e_a(X_i)e_b(X_j)$ where for $c \in \{0, 1\}$, $e_c(X_i) = \mathbb{P}(D_i = c | X_i)$. Let also $D_{ij}^{ab} = \mathbb{1}(D_i = a)\mathbb{1}(D_j = b)$.

Proposition 4.1 *Under Assumption 1, $W(\pi)$ in (4.1) is identified in the following ways*

$$W(\pi) = \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} m_{ab}(Z_i, Z_j, \gamma, \nu) \pi_{ab}(X_i, X_j) \right],$$

where $\nu \in \{\varphi, e\}$ and m_{ab} can be any of the following

$$\begin{aligned} (DM) \quad m_{ab}(Z_i, Z_j, \gamma, \varphi) &= \varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) \\ (IPW) \quad m_{ab}(Z_i, Z_j, \gamma, e) &= g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab} / e_{ab}(X_i, X_j). \end{aligned}$$

Again, note that whenever g does not depend directly on the potential outcomes we have that $\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) = g(X_i, X_j, \gamma_a, \gamma_b)$ as we can see in Example 3 below. Whenever g does depend on the potential outcomes $\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b)$ must be estimated using dyadic regressions. Since not much is known about machine learning dyadic regressions and their properties I avoid using the direct method whenever φ is unknown. Now we apply Proposition 4.1 to identify the welfare in each of our three main examples.

Example 2 (Inequality (cont.)) *In this example the welfare is identified by*

$$W(\pi) = \mathbb{E} \left[\frac{1}{2} (Y_i + Y_j - |Y_i - Y_j|) \sum_{(a,b) \in \{0,1\}^2} \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \pi_{ab}(X_i, X_j) \right].$$

■

Example 3 (IOp (cont.)) *In this example the welfare is identified by*

$$W(\pi) = \frac{1}{2} \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} \left(\gamma_a(X_i) + \gamma_b(X_j) - |\gamma_a(X_i) - \gamma_b(X_j)| \right) \pi_{ab}(X_i, X_j) \right].$$

■

Example 4 (Intergenerational mobility (cont.)) *In this example the welfare is identified by*

$$W(\pi) = - \left| \mathbb{E} \left[\text{sgn}(X_{1i} - X_{1j}) \text{sgn}(Y_i - Y_j) \sum_{(a,b) \in \{0,1\}^2} \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \pi_{ab}(X_i, X_j) \right] - t \right|.$$

■

In Examples 2 and 4 we have used IPW since we will see that it makes the estimation simpler. Example 3 does not depend on the potential outcomes which makes the use of DM easier. In order to compute the orthogonal scores we need to assume a similar linearization property as the one in Assumption 2 and to the linearization assumed in Escanciano and Terschuur (2022).

Assumption 3 *There exist α_{ab} , $P < \infty$, and (c_{1p}, c_{2p}) for $p = 1, \dots, P$, such that for all $(a, b) \in \{0, 1\}^2$ the following linearization holds*

$$\frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, \nu)] = \mathbb{E} \left[\sum_{p=1}^P \alpha_{ab,p}^\gamma(D_i, X_i, D_j, X_j) (c_{1p} \bar{\gamma}_\tau(D_i, X_i) + c_{2p} \bar{\gamma}_\tau(D_j, X_j)) \right],$$

where $\bar{\gamma}_\tau$ is defined as in Assumption 2.

Again, γ enters m_{ab} only through g so a sufficient condition that would allow to compute α_{ab} is to assume a linearization like the above for g instead of for m_{ab} . Possible dependence of c_{1p} on (D_i, X_i) and c_{2p} on (D_j, X_j) is neglected in the notation for simplicity. Now we are ready to present the result of the orthogonal scores for welfare functions defined with U-statistics.

Proposition 4.2 *The orthogonal scores are given by*

$$\Gamma_{ij}(\pi) = \sum_{(a,b) \in \{0,1\}^2} \Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j),$$

where depending on whether we identify with DM or IPW we have

$$\begin{aligned} (DM) \quad \Gamma_{ij}^{ab} &= \varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) + \phi_{ab}^\varphi(D_i, X_i, D_j, X_j, \varphi, \alpha^e) + \phi_{ab}^\gamma(D_i, X_i, D_j, X_j, \gamma, \alpha^\gamma) \\ (IPW) \quad \Gamma_{ij}^{ab} &= \frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_{ab}(X_i, X_j)} + \phi_{ab}^e(D_i, X_i, D_j, X_j, e, \alpha^e) + \phi_{ab}^\gamma(D_i, X_i, D_j, X_j, \gamma, \alpha^\gamma), \end{aligned}$$

where

$$\begin{aligned} \phi_{ab}^\gamma(D_i, X_i, D_j, X_j, e, \alpha^\gamma) &= \sum_{p=1}^P \alpha_{ab,p}^\gamma(D_i, X_i, D_j, X_j, e) (c_{1p} Y_i + c_{2p} Y_j - c_{1p} \gamma(D_i, X_i) - c_{2p} \gamma(D_j, X_j)), \\ \phi_{ab}^\varphi(D_i, X_i, D_j, X_j, \varphi, \alpha^m) &= \alpha_{ab}^\varphi(D_i, X_i, D_j, X_j) (g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) - \varphi(D_i, X_i, D_j, X_j, \gamma_a, \gamma_b)), \\ \phi_{ab}^e(D_i, X_i, D_j, X_j, e, \alpha^e) &= \alpha_{ab,1}^e(X_i) (\mathbf{1}(D_i = a) - e_a(X_i)) + \alpha_{ab,2}^e(X_j) (\mathbf{1}(D_j = b) - e_b(X_j)), \end{aligned}$$

and

$$\begin{aligned} \alpha_{ab}^\varphi(D_i, X_i, D_j, X_j) &= \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)}, \\ \alpha_{ab,1}^e(X_i) &= -\mathbb{E} \left[\frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i)^2 e_b(X_j)} \middle| X_i \right], \\ \alpha_{ab,2}^e(X_j) &= -\mathbb{E} \left[\frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i) e_b(X_j)^2} \middle| X_j \right]. \end{aligned}$$

Once again, note that whenever g does not directly depend on the potential outcomes then $g = \varphi$ and we have that $\phi_{ab}^m = 0$. Now we can see how Proposition 4.2 applies to our examples.

Example 2 (Inequality (cont.)) *In this example we have that*

$$\Gamma_{ij}^{ab} = \frac{1}{2}(Y_i + Y_j - |Y_i - Y_j|) \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} + \alpha_{ab,1}^e(X_i)(\mathbb{1}(D_i = a) - e_a(X_i)) + \alpha_{ab,2}^e(X_j)(\mathbb{1}(D_j = b) - e_b(X_j)).$$

■

Example 3 (IOp (cont.)) *I introduce the orthogonal score of the IOp example as a Proposition with its proof in the Appendix.*

Proposition 4.3 *Assume for all $(a, b) \in \{0, 1\}^2$ that either (i) $\mathbb{P}(\gamma_a(X_i) - \gamma_b(X_j) = 0) = 0$ or that (ii) $x_i \neq x_j \implies \gamma_a(X_i) - \gamma_b(X_j) \neq 0$ and let $\delta_{ij}^{ab} = \text{sgn}(\gamma_a(X_i) - \gamma_b(X_j))$, then*

$$\Gamma_{ij}^{ab} = \frac{1}{2} \left(\gamma_a(X_i) + \gamma_b(X_j) - |\gamma_a(X_i) - \gamma_b(X_j)| + \frac{\mathbb{1}(D_i = a)}{e_a(X_i)} (1 - \delta_{ij}^{ab})(Y_i - \gamma(D_i, X_i)) + \frac{\mathbb{1}(D_j = b)}{e_b(X_j)} (1 + \delta_{ij}^{ab})(Y_j - \gamma(D_j, X_j)) \right).$$

These assumptions deal with the point of non-differentiability of the absolute value. For a thorough discussion see [Escanciano and Terschuur \(2022\)](#). ■

To estimate the welfare in these examples for a given $\pi \in \Pi$ I use an adaptation to U-statistics of the cross-fitting used before (see [Escanciano and Terschuur \(2022\)](#)). I split the pairs $\{(i, j) \in \{1, \dots, n\}^2 : i < j\}$ in L groups I_1, \dots, I_L , then

$$\hat{W}_n(\pi) = \binom{n}{2}^{-1} \sum_{l=1}^L \sum_{(i,j) \in I_l} \hat{\Gamma}_{ij,l}(\pi), \quad (4.2)$$

where $\hat{\Gamma}_{ij,l}$ is the same as Γ_{ij} but with all nuisance parameters replaced by estimators which do not use observations in the pairs in I_l . As before, the estimator of the optimal treatment rule among a class of rules Π is

$$\hat{\pi} = \arg \max_{\pi \in \Pi} \hat{W}_n(\pi).$$

For the Intergenerational mobility example the estimation is slightly different.

Example 4 (Intergenerational mobility (cont.)) *The orthogonal score is given by*

$$\Gamma_{ij}^{ab} = \frac{1}{2} \text{sgn}(X_{1i} - X_{1j}) \text{sgn}(Y_i - Y_j) \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} + \alpha_{ab,1}^e(X_i)(\mathbb{1}(D_i = a) - e_a(X_i)) + \alpha_{ab,2}^e(X_j)(\mathbb{1}(D_j = b) - e_b(X_j)).$$

The estimator of the welfare for a given $\pi \in \Pi$ and target t is

$$\hat{W}_n(\pi) = - \left| \binom{n}{2}^{-1} \sum_{l=1}^L \sum_{(i,j) \in I_l} \sum_{(a,b) \in \{0,1\}^2} \hat{\Gamma}_{ij,l}^{ab} \pi_{ab}(X_i, X_j) - t \right|. \quad (4.3)$$

■

5 Asymptotic statistical guarantees

Now it is useful to make clear the dependence of the scores Γ_{ij}^{ab} on the data and the nuisance parameters. Hence, I let now $\Gamma_{ij}^{ab} = \psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha)$, where ψ_{ab} is the sum of a function whose expectation identifies a term in the welfare plus other functions which are correction terms needed to achieve orthogonality with respect to the nuisance parameter γ and with respect to nuisance parameters which appear in the identification result $\nu \in \{\varphi, e\}$

$$\psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha) = m_{ab}(Z_i, Z_j, \gamma, \nu) + \phi_{ab}^\gamma(Z_i, Z_j, \gamma, \alpha^\gamma) + \phi_{ab}^\nu(Z_i, Z_j, \nu, \alpha^\nu).$$

In general we have that for a given treatment rule π , orthogonal scores are given by

$$\Gamma_{ij}(\pi) = \sum_{(a,b) \in \{0,1\}^2} \psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha) \pi_{ab}(X_i, X_j).$$

This framework accomodates also the welfare functions which are not defined as U-statistics if $\psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha)$ does not depend on Z_j and only depends on a so that we could rewrite it as $\psi_a(Z_i, \gamma, \nu, \alpha)$ for $a \in \{0,1\}$. For this reason I stick to this notation and do not state all conditions and results for welfare functions which are not U-statistics and those which are. The intergenerational mobility example does not fit in this general setting, however, the results extend easily to this example by Corollary 1 at the end of this section. In the next subsections I give conditions on the convergence of the nuisance parameters and on the complexity of the policy class Π which will allow me to prove asymptotical statistical guarantees for the estimated treatment rules.

5.1 Conditions on the nuisance parameter estimators

I give high level conditions for the estimators of all nuisance parameters which have to be used to estimate the welfare. These conditions have been showed to hold for a variety of non-parametric estimators such as kernels or sieve estimators. The assumptions below are analogous to those in [Escanciano and Terschuur \(2022\)](#).

Assumption 4 $\mathbb{E}[|\psi(Z_i, Z_j, \gamma, \nu, \alpha)|^2] < \infty$, $\omega \in \{\gamma, \nu\}$ and for $(a, b) \in \{0, 1\}^2$ and for some $a(n) \rightarrow 0$

- (i) $n^{\lambda_\gamma} \sqrt{\mathbb{E}(|m_{ab}(z_i, z_j, \hat{\gamma}_l, \nu) - m_{ab}(z_i, z_j, \gamma, \nu)|^2)} \leq a(n)$;
- (ii) $n^{\lambda_\nu} \sqrt{\mathbb{E}(|m_{ab}(z_i, z_j, \gamma, \hat{\nu}_l) - m_{ab}(z_i, z_j, \gamma, \nu)|^2)} \leq a(n)$;
- (iii) $n^{\lambda_\gamma} \sqrt{\mathbb{E}(|\phi_{ab}^\gamma(z_i, z_j, \hat{\gamma}_l, \alpha^\gamma) - \phi_{ab}^\gamma(z_i, z_j, \gamma, \alpha^\gamma)|^2)} \leq a(n)$;
- (iv) $n^{\lambda_\nu} \sqrt{\mathbb{E}(|\phi_{ab}^\nu(z_i, z_j, \hat{\nu}_l, \alpha^\nu) - \phi_{ab}^\nu(z_i, z_j, \nu, \alpha^\nu)|^2)} \leq a(n)$;

$$(v) \ n^{\lambda_\alpha} \sqrt{\mathbb{E}(|\phi_{ab}^\omega(z_i, z_j, \omega, \hat{\alpha}_l^\omega) - \phi_{ab}^\omega(z_i, z_j, \omega, \alpha^\omega)|^2)} \leq a(n),$$

where $1/4 < \lambda_\gamma, \lambda_\nu, \lambda_\alpha < 1/2$.

These are mild mean-square consistency conditions for $\hat{\gamma}_l$, $\hat{\nu}_l$ and $\hat{\alpha}_l$ separately. Assumption 4 often follows from the L2 convergence rates of the nuisance estimators. There is a large literature checking L_2 -convergence rates for different machine learners under low level sparsity or smoothness conditions on the nuisance parameters. The traditional non-parametric literature gives rates for kernel regression and sieves/series (e.g. [Chen \(2007\)](#)). For L_1 -penalty estimators such as Lasso see, e.g., [Belloni and Chernozhukov \(2011\)](#) and [Belloni and Chernozhukov \(2013\)](#). Also for low level conditions for shrinkage and kernel estimators see Appendix B in [Sasaki and Ura \(2021\)](#). Rates for L_2 -boosting in low dimensions are found in [Zhang and Yu \(2005\)](#), and more recently [Kueck et al. \(2023\)](#) find rates for L_2 -boosting with high dimensional data. For results on versions of random forests see [Wager and Walther \(2015\)](#) and [Athey et al. \(2019\)](#). Finally, for single-layer, sigmoid-based neural networks see [Chen and White \(1999\)](#) and for a modern setting of deep neural networks with rectified linear (ReLU) activation function see [Farrell et al. \(2021\)](#). Define now the following interaction terms for $\omega \in \{\gamma, \nu\}$

$$\begin{aligned} \hat{\xi}_{ij,ab,l} &= m_{ab}(Z_i, Z_j, \hat{\gamma}_l, \hat{\nu}_l) - m_{ab}(Z_i, Z_j, \hat{\gamma}_l, \nu) - m_{ab}(Z_i, Z_j, \gamma, \hat{\nu}_l) + m_{ab}(Z_i, Z_j, \gamma, \nu), \\ \hat{\xi}_{ij,ab,l}^\omega &= \phi_{ab}(z_i, z_j, \hat{\omega}_l, \hat{\alpha}_l^\omega) - \phi_{ab}(z_i, z_j, \omega, \hat{\alpha}_l^\omega) - \phi_{ab}(z_i, z_j, \hat{\omega}_l, \alpha^\omega) + \phi_{ab}(z_i, z_j, \omega, \alpha^\omega). \end{aligned}$$

Assumption 5 For each $l = 1, \dots, L$

- (i) $\int \int \phi_{ab}^\gamma(z_i, z_j, \gamma, \hat{\alpha}_l^\gamma) F(dz_i) F(dz_j) = 0$ and $\int \int \phi_{ab}^\nu(Z_i, Z_j, \nu, \hat{\alpha}_l^\nu) F(dz_i) F(dz_j) = 0$.
- (ii) $\mathbb{E}(\|\hat{\gamma}_l - \gamma\|^2) = o(n^{-2\lambda_\gamma})$, $\mathbb{E}(\|\hat{\nu}_l - \nu\|^2) = o(n^{-2\lambda_\nu})$ and

$$\begin{aligned} |\mathbb{E}[(m_{ab}(Z_i, Z_j, \tilde{\gamma}, \nu) + \phi_{ab}^\gamma(Z_i, Z_j, \tilde{\gamma}, \alpha^\gamma)) \pi_{ab}(X_i, X_j)]| &\leq C \|\tilde{\gamma} - \gamma\|^2 \\ |\mathbb{E}[(m_{ab}(Z_i, Z_j, \gamma, \tilde{\nu}) + \phi_{ab}^\nu(Z_i, Z_j, \tilde{\nu}, \alpha^\nu)) \pi_{ab}(X_i, X_j)]| &\leq C \|\tilde{\nu} - \nu\|^2. \end{aligned}$$

Assumption 5 (i) is usually easy to verify from visual inspection and (ii) requires L2 convergence rates and some smoothness.

Assumption 6 For each $l = 1, \dots, L$

$$\sqrt{n} \mathbb{E}(\hat{\xi}_{ij,ab,l}^\omega \pi_{ab}(X_i, X_j)) \leq a(n).$$

These are rate conditions on the remainder terms $\hat{\xi}_l^\omega(w_i, w_j)$. Often, Assumption 6 follows if $\sqrt{n} \|\hat{\alpha}_l^\omega - \alpha\| \|\hat{\omega}_l - \omega\| \leq a(n)$, where $\|\cdot\|$ denotes the L2 norm. For example in [Athey and Wager \(2021\)](#), who in the proof of their Lemma 4 use Cauchy-Schwarz inequality to get a bound on their interaction term which does not depend on π and only on the product of L2 norms. This is the precise assumption that allows for parametric rates even with slow nonparametric estimators. In essence, it is enough for the product of the nonparametric estimators to go at parametric rate.

5.2 Conditions on the complexity of the policy class

The complexity of the policy class must also be restricted. If all sorts of subsets of \mathcal{X} are allowed to decide who should be treated then we get overfitted policy rules. As in [Athey and Wager \(2021\)](#) I measure the policy class complexity with its VC dimension (see for instance [Wainwright \(2019\)](#)) which I allow to grow with the sample size. Hence, for now on I subscript the policy class by n , Π_n .

Assumption 7 *There are constants $0 < \beta < 1/2$ and $n^* \geq 1$ such that for all $n \geq n^*$, $VC(\Pi_n) < n^\beta$.*

Examples of finite VC-dimension classes are linear eligibility scores or generalized eligibility scores (see [Kitagawa and Tetenov \(2018\)](#)). Policy classes whose VC-dimension increases with the sample size are for example decision trees (see [Athey and Wager \(2021\)](#)).

5.3 Upper bounds

Let now

$$\begin{aligned} W(\pi) &= \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} \psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha) \pi_{ab}(X_i, X_j) \right], \\ \widetilde{W}_n(\pi) &= \binom{n}{2}^{-1} \sum_{i < j} \left[\sum_{(a,b) \in \{0,1\}^2} \psi_{ab}(Z_i, Z_j, \gamma, \nu, \alpha) \pi_{ab}(X_i, X_j) \right], \\ \hat{W}_n(\pi) &= \binom{n}{2}^{-1} \sum_{l=1}^L \sum_{(i,j) \in I_l} \left[\sum_{(a,b) \in \pi} \psi_{ab}(Z_i, Z_j, \hat{\gamma}_l, \hat{\nu}_l, \hat{\alpha}_l) \pi_{ab}(X_i, X_j) \right], \end{aligned}$$

$W(\pi)$ and $\widetilde{W}_n(\pi)$ are the welfare and the unfeasible estimator of the welfare at policy rule π when all nuisance parameters are known. $\hat{W}_n(\pi)$ is the feasible estimator which we already introduced in (4.2). Let $W_{\Pi_n}^* = \sup_{\pi \in \Pi_n} W(\pi)$ be the best possible welfare. I want to give upper bounds to the regret: $\mathbb{E}[W_{\Pi_n}^* - W(\hat{\pi})]$, i.e. the expected difference between the best possible welfare and the welfare evaluated at the estimated policy. As usual I start bounding the regret as follows

$$\begin{aligned} \mathbb{E}[W_{\Pi_n}^* - W(\hat{\pi})] &\leq 2\mathbb{E} \left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - W(\pi)| \right] \\ &\leq 2\mathbb{E} \left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - \widetilde{W}_n(\pi)| \right] + 2\mathbb{E} \left[\sup_{\pi \in \Pi_n} |\widetilde{W}_n(\pi) - W(\pi)| \right], \end{aligned} \quad (5.1)$$

where in the second inequality I have added and subtracted $\widetilde{W}_n(\pi)$ and used the triangle inequality. The second term above is just a standard centered U-process indexed by $\pi \in \Pi_n$. I

start as in [Athey and Wager \(2021\)](#) by showing the rate of convergence of this second term. I work for some fixed $(a, b) \in \{0, 1\}^2$ and define the following set

$$\Pi_{ab,n} = \{\pi_{ab} : \pi \in \Pi_n\}.$$

The first step is to bound it by the Rademacher complexity which we define as

$$\mathcal{R}_n(\Pi_{ab,n}) = \mathbb{E}_\varepsilon \left(\sup_{\pi \in \Pi_n} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \right| \right).$$

Lemma 1

$$\mathbb{E} \left[\sup_{\pi \in \Pi_n} |\widetilde{W}_n(\pi) - W(\pi)| \right] \leq \mathbb{E}[2\mathcal{R}_n(\Pi_{ab,n})].$$

Now we want an asymptotic upper bound for $\mathbb{E}[\mathcal{R}_n(\Pi_{ab})]$. Importantly, we want the bound to depend on the following variance

$$S_{ab} = \mathbb{E}[\Gamma_{i,j}^{2ab}].$$

While [Kitagawa and Tetenov \(2018\)](#) and others provide bounds in terms of the max of the scores, [Athey and Wager \(2021\)](#) provide bounds based on the variance and the efficient variance. The next result provides a bound on the Rademacher complexity based on S_{ab} .

Lemma 2 *Assume that Γ_{ij}^{ab} has bounded support for $(a, b) \in \{0, 1\}^2$. Then, under Assumptions 4 and 6*

$$\mathbb{E}[\mathcal{R}_n(\Pi_{ab,n})] = \mathcal{O} \left(\sqrt{\frac{S_{ab} \cdot VC(\Pi_{ab,n})}{\lfloor n/2 \rfloor}} \right).$$

The boundedness assumption can be generalized to sub-gaussianity. However, this generalization comes at the cost of making the (already involved) proofs substantially less tractable. Now we want to provide asymptotic upper bounds for the first term in (5.1). [Escanciano and Terschuur \(2022\)](#) show that for given $\pi \in \Pi_n$

$$\sqrt{n}(\hat{W}_n(\pi) - \widetilde{W}_n(\pi)) \rightarrow_p 0.$$

The next result makes the above uniform in $\pi \in \Pi_n$.

Lemma 3 (Uniform coupling) *Under Assumptions 4 and 6*

$$\sqrt{n} \mathbb{E} \left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - \widetilde{W}_n(\pi)| \right] = \mathcal{O} \left(1 + \frac{VC(\Pi_{ab,n})}{\lfloor n/2 \rfloor^{\min(\lambda_\gamma, \lambda_\nu, \lambda_\alpha)}} \right).$$

Finally, using Lemmas 2 and 3 the following holds.

Theorem 1 *Suppose Assumptions 4 and 6 hold, that Assumption 7 holds with $\beta < \min(\lambda_\gamma, \lambda_\nu, \lambda_\alpha)$. Then*

$$\mathbb{E}[W_{\Pi_n}^* - W(\hat{\pi})] = \mathcal{O} \left(\sqrt{\frac{S_{ab} \cdot (2VC(\Pi_n) - 1)}{\lfloor n/2 \rfloor}} \right).$$

Corollary 1 *The bound in Theorem 1 applies to the Intergenerational mobility example.*

6 Empirical application

In our empirical application we study the optimal allocation of children to preschool for our leading welfare functions. We make use of the Panel Study of Income Dynamics (PSID) database which has been following families for nearly 50 years. The nature of this survey allows us to observe a rich set of circumstances and long term outcomes. In 1995, PSID asked about the participation in preschool to adults between 18-30 years hold. Hence, we are able to track long term outcomes of these individuals. We take as outcome the average earnings from 25 to 35 years old. We assume selection on observables holds. In particular we condition on sex, birthyear, average parental income in the 5 years before birth, mother’s education, father’s education, father’s occupation and whether the individual is black. In Table 1 we see the results of estimating the Average Causal Effect (ATE), Gini, IOp and Intergenerational mobility as captured by the rank correlation of parents and child income.

	Estimate	se	pval	Gini	IOp	Kendall	n
Earnings 25-35	4147	1258	0.001	0.392	0.163	0.168	2971

Table 1: ATE, Gini, IOp and Kendal- τ

To estimate the ATE I use doubly robust scores and random forests to estimate the regression functions and logit Lasso to estimate propensity scores. To implement random forests I use the **ranger** package in R while to implement logit lasso I use the **glmnet** package in R. Under our assumption of no selection on observables we observe a sizeable and significant positive effect of attending preschool of 4,147\$ of added annual earnings. Dolars have been adjusted by the CPI to 2010 dolars. We see that the Gini coefficient is 0.39 and that IOp is 0.16, meaning that 41% of total inequality can be explained by the circumstances we observe. The Kendall- τ is 0.17 which indicates a positive association between parental and child incomes. I compute optimal treatment rules based on parental income and mother’s years of education. I set the target in the Kendall- τ welfare to zero, meaning that the aim is to completely erase intergenerational persistence. As the policy class I use 2-depth decision trees. Unfortunately, the U-statistic nature of the welfare function prevents me from using the computational shortcuts in [Athey and Wager \(2021\)](#) since the sub-trees are not independent optimization problems. To ease the computational problem I use the deciles of parental income as cutting points instead of all the observed values of parental income. I do exhaustive search meaning that I consider all possible 2-depth decision trees.

I start by showing the results for the utilitarian welfare function in Figure 1. At the terminal nodes I report the number of observations, the conditional average treatment effect (CATE) in the node and the proportion of observations in the terminal node which are treated in the

data. For the Utilitarian welfare we see that the first cutting point is whether parental income is below or above 109,785\$. If an observation is below this cut-off the tree splits according to the education of the mother. If parental income is below 109,785\$ and mother's education is below 16 years the tree allocates the observation to treatment. Most of the sample belongs in this terminal node. We see that the CATE in this node is positive so, as we would expect, a utilitarian policy maker would decide to treat these observations. If parental income is below 109,785\$ but the mother is highly educated we see that the CATE is actually negative and hence the utilitarian policy maker does not allocate the individual to treatment. This is in line with results in the psychology and economics literature which documents that in early educational institutions there is less interaction with adults and hence there can be a negative effect of attending such institutions if the interactions with adults in the household are of "high-quality" (see [Fort et al. \(2020\)](#)). However, we see that in the right-most terminal node there are only 25 observations and an extremely negative CATE, this is driven by extreme results from a few observations only.




Figure 1: Utilitarian optimal policy rule: the left branches mean that the condition in the node holds while it does not hold in the right branch.

In Figure 2 I show the result for the IOP aware welfare function. The first income cutoff is lower. Then, in both subnodes the classification is based on mother's education. In this case we only treat individuals in the lower part of the parental income distribution if their mothers do not exceed 13 years of education. However, for richer individuals we treat them if the mother has less than 16 years of education. This makes sense if children with less parental income go to worse preschool institutions. In this case, the amount of mother's education required to beat the education given in preschool is lower. Compared to Figure 1 we see that the IOP rule places more importance on mother's education. For instance, it allows to treat very rich individuals as long as their mothers have less than 16 years of education. This indicates that mother's education is a bigger contributor to inequality.

Finally, in Figure 3 we see the results for an inequality aware welfare function. The most impressive result here is that a group of individuals who is (on average) negatively affected by preschool is allocated to treatment. This is perfectly possible if by decreasing their outcomes a more equal distribution and a higher welfare is achieved at the cost of a lower average outcome. However, it poses an ethical problem. A way to circumvent this is to restrict our welfare maximization to treat only groups with a positive CATE. The tree treats individuals with mother's education below 16 years and income below 172,853 \$ and individuals with highly educated mothers and parental income above 61,996\$.




Figure 2: IOp optimal policy rule: the left branches mean that the condition in the node holds while it does not hold in the right branch.




Figure 3: Inequality optimal policy rule: the left branches mean that the condition in the node holds while it does not hold in the right branch.

7 Appendix

A Auxiliary lemmas

In this section we prove some lemmas which will be needed to prove the main results. Let us first define some important objects. For a fixed sample $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$ we have

$$\tilde{\Pi}_{ab} = \{\pi_{ab}(X_1, X_{\lfloor n/2 \rfloor + 1}), \dots, \pi_{ab}(X_{\lfloor n/2 \rfloor}, X_n) : \pi \in \Pi\}.$$

For $\pi, \pi' \in \tilde{\Pi}_{ab}$ define the following distances

$$D_n^2(\pi, \pi') = \frac{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} (\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - \pi'_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))^2}{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}},$$

$$H(\pi, \pi') = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \neq \pi'_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})).$$

Let $N_{D_n}(\varepsilon, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor})$ be the number of balls of radius ε needed to cover $\tilde{\Pi}_{ab}$ under distance D_n . Define the same object for the Hamming distance H and let

$$N_H(\varepsilon, \tilde{\Pi}_{ab}) = \sup\{N_H(\varepsilon, \tilde{\Pi}_{ab}, \{X_i\}_{i=1}^m) : X_1, \dots, X_m \in \mathcal{X}, m \geq 1\}.$$

Note $N_H(\varepsilon, \tilde{\Pi}_{ab})$ does not depend on m . It will be useful to bound N_{D_n} with N_H which is what we do in the next lemma.

Lemma 4 *For fixed $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$ we have that*

$$N_{D_n}(\varepsilon, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}) \leq N_H(\varepsilon^2, \tilde{\Pi}_{ab}).$$

Proof: Take an auxiliary sample $\{X'_j\}_{j=1}^m$ contained in $\{X_i\}_{i=1}^n$ such that

$$\left| |B_i| - \frac{m \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \right| \leq 1,$$

where $B_i = \{j \in \{1, \dots, m\} : X'_j = X_i\}$. Then, for $\pi, \pi' \in \tilde{\Pi}_{ab}$

$$D_n^2(\pi, \pi') = \frac{1}{m} \sum_{i=1}^{\lfloor n/2 \rfloor} \underbrace{\frac{m \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{\sum_{k=1}^{\lfloor n/2 \rfloor} \Gamma_{k, \lfloor n/2 \rfloor + k}^{2ab}}}_{\geq |B_i| - 1} \mathbb{1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \neq \pi'_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})).$$

So

$$\begin{aligned}
D_n^2(\pi, \pi') &\geq \sum_{i=1}^{\lfloor n/2 \rfloor} \frac{|B_i|}{m} \mathbb{1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \neq \pi'_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - O(1/m) \\
&= \sum_{i=1}^{\lfloor n/2 \rfloor} \frac{|B_i|}{m} \frac{1}{|B_i|} \sum_{j \in B_i} \mathbb{1}(\pi_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j}) \neq \pi'_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j})) - O(1/m) \\
&= \frac{1}{m} \sum_{i=1}^{\lfloor n/2 \rfloor} \sum_{j \in B_i} \mathbb{1}(\pi_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j}) \neq \pi'_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j})) - O(1/m).
\end{aligned}$$

In the first equality above we have used the fact that all sumands in the inner sum are the same since for all $j \in B_i$ we know that $(X_i, X_{\lfloor n/2 \rfloor + i}) = (X'_j, X'_{\lfloor n/2 \rfloor + j})$. Now we notice that the sum $\sum_{i=1}^{\lfloor n/2 \rfloor} \sum_{j \in B_i}$ might sum some pairs more than once (e.g. if $(X_1, X_{\lfloor n/2 \rfloor + 1}) = (X_2, X_{\lfloor n/2 \rfloor + 2})$ then $B_1 = B_2$). Using this fact and that $\{X'_j\}_{j=1}^m$ is contained in $\{X'_i\}_{i=1}^m$ we have that

$$\begin{aligned}
D_n^2(\pi, \pi') &\geq \frac{1}{m} \sum_{j=1}^m \mathbb{1}(\pi_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j}) \neq \pi'_{ab}(X'_j, X'_{\lfloor n/2 \rfloor + j})) - O(1/m) \\
&= H(\pi, \pi') - O(1/m).
\end{aligned}$$

Hence, $H(\pi, \pi') \leq D_n^2(\pi, \pi') + O(1/m)$. Since N_H does not depend on m , we can make m arbitrarily large and conclude that

$$N_{D_n}(\varepsilon, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}) \leq N_H(\varepsilon^2, \tilde{\Pi}_{ab}).$$

■

Now we prove that the sequence of covers we use in the proof of Lemma 2 exists.

Lemma 5 *There exists a sequence of covers $\{B_k\}_{k=0}^K$ with $K < \infty$ of $\tilde{\Pi}_{ab}$ with $B_k \subset \tilde{\Pi}_{ab}$ such that for $k = 0, \dots, K$*

- For all $\pi \in \tilde{\Pi}_{ab}$, there exists $b \in B_k$ such that $D_n(\pi, b) \leq 2^{-k}$,
- $|B_k| = N_{D_n}(2^{-k}, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}) \leq |\tilde{\Pi}_{ab}|$.

Proof: First note that $|\tilde{\Pi}_{ab}| < 2^{\lfloor n/2 \rfloor} < \infty$ since X_i 's are fixed. Since $\tilde{\Pi}_{ab}$ is finite and $B_k \subset \tilde{\Pi}_{ab}$ for all k , there exists finite K such that we can set $B_K = \tilde{\Pi}_{ab}$. This is because for any B_k which is a strict subset of $\tilde{\Pi}_{ab}$ there exist $\pi \in \tilde{\Pi}_{ab}$ such that for all $b \in B_k$, $D_n(b, \pi) > a > 0$ and there exists $K > 0$ such that $2^{-K} < a$. K is finite since there are only finitely many subsets of $\tilde{\Pi}_{ab}$. For B_{K-1} we can look through all possible strict subsets for one which satisfies our conditions, if we do not find any we know that $B_{K-1} = \tilde{\Pi}_{ab}$ does satisfy them. In this way we can go backwards and build the sequence of covers. ■

The next Lemma relates the VC dimension of $\tilde{\Pi}_{ab}$ to that of Π .

Lemma 6 $VC(\tilde{\Pi}_{ab}) \leq 2VC(\Pi) - 1$.

Proof: Let $\pi_t(X_i) = \mathbb{1}(\pi(X_i) = t)$ for $t \in \{0, 1\}$. Define $\Pi_t = \{\mathbb{1}(\pi(X_i) = t) : \pi \in \Pi\}$. Note that $\Pi_1 = \Pi$ and that $VC(\Pi_0) = VC(\Pi_1)$ by Lemma 9.7 in Kosorok (2008). Now note that for any $(a, b) \in \{0, 1\}^2$

$$\tilde{\Pi}_{ab} = \{\pi_a \cdot \pi_b : (\pi_a, \pi_b) \in \Pi_a \times \Pi_b\},$$

so Lemma 9.9 (ii) in Kosorok (2008) yields the desired result. ■

B Proofs of main results

Proof of Proposition 3.1: I proof only the identification of the first term of the welfare since the second one follows in the same manner.

$$\begin{aligned} \mathbb{E}[g(Y_i(1), X_i, \gamma^{(1)})\pi(X_i)] &= \mathbb{E}[\mathbb{E}(g(Y_i(1), X_i, \gamma_1)|X_i)\pi(X_i)] \\ &= \mathbb{E}[\mathbb{E}(g(Y_i(1), X_i, \gamma_1)|D_i = 1, X_i)\pi(X_i)] \\ &= \mathbb{E}[\mathbb{E}(g(Y_i, X_i, \gamma_1)|D_i = 1, X_i)\pi(X_i)] \\ &= \mathbb{E}\left[\mathbb{E}\left(\frac{g(Y_i, X_i, \gamma_1)D_i}{e(X_i)}|X_i\right)\pi(X_i)\right] \\ &= \mathbb{E}\left[\frac{g(Y_i, X_i, \gamma_1)D_i}{e(X_i)}\pi(X_i)\right], \end{aligned}$$

the first equality follows by LIE and the fact that by selection on observables and definition of Y_i we have that $\mathbb{E}[Y_i(1)|X_i] = \mathbb{E}[Y_i(1)|D_i = 1, X_i] = \mathbb{E}[Y_i|D_i = 1, X_i]$. Second equality follows from selection on observables, third equality from the definition of Y_i and already establishes the identification by the direct method. ■

Proof of Proposition 3.2: Let $d/d\tau$ be the derivative with respect to τ evaluated at $\tau = 0$, let $\varphi_\tau = \varphi + \tau\tilde{\varphi}$ for some $\tilde{\varphi}$ in the space where φ lives and \mathbb{E}_τ be the expectation with respect to $F + \tau(H - F)$ for some alternative distribution H . Then

$$\frac{d}{d\tau}\mathbb{E}[\varphi_\tau(1, X_i, \bar{\gamma}_\tau(1, X_i))\pi(X_i)] = \frac{d}{d\tau}\mathbb{E}[\varphi_\tau(1, X_i, \gamma(1, X_i))\pi(X_i)] + \frac{d}{d\tau}\mathbb{E}[\varphi(1, X_i, \bar{\gamma}_\tau(1, X_i))\pi(X_i)].$$

For the first term note that

$$\begin{aligned} \frac{d}{d\tau}\mathbb{E}[\varphi_\tau(1, X_i, \gamma(1, X_i))\pi(X_i)] &= \frac{d}{d\tau}\mathbb{E}\left[\frac{D_i}{e(X_i)}\varphi_\tau(1, X_i, \gamma(1, X_i))\pi(X_i)\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[\frac{D_i}{e(X_i)}\varphi_\tau(1, X_i, \gamma(1, X_i))\pi(X_i)\right] \\ &\quad - \frac{d}{d\tau}\mathbb{E}_\tau\left[\frac{D_i}{e(X_i)}\varphi(1, X_i, \gamma(1, X_i))\pi(X_i)\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[\frac{D_i}{e(X_i)}(g(Y_i, X_i, \gamma(1, X_i)) - \varphi(1, X_i, \gamma(1, X_i)))\pi(X_i)\right], \end{aligned}$$

where we use LIE in the first equality, then we use the chain rule and finally that $\varphi_\tau(1, X_i, \gamma(1, X_i))$ is a projection of $g(Y_i, X_i, \gamma(1, X_i))$. For the second term we have

$$\begin{aligned} \frac{d}{d\tau} \mathbb{E}[\varphi(1, X_i, \bar{\gamma}_\tau(1, X_i))\pi(X_i)] &= \frac{d}{d\tau} \mathbb{E}[\alpha_1(D_i, X_i)\bar{\gamma}_\tau(1, X_i)\pi(X_i)] \\ &= \frac{d}{d\tau} \mathbb{E}_\tau[\alpha_1(D_i, X_i)\bar{\gamma}_\tau(1, X_i)\pi(X_i)] \\ &\quad - \frac{d}{d\tau} \mathbb{E}_\tau[\alpha_1(D_i, X_i)\gamma_\tau(1, X_i)\pi(X_i)] \\ &= \frac{d}{d\tau} \mathbb{E}_\tau[\alpha_1(D_i, X_i)(Y_i - \gamma(1, X_i))\pi(X_i)], \end{aligned}$$

where we use Assumption 2 in the first equality, then the chain rule and then the fact that γ is a projection. Then, following Chernozhukov et al. (2021) we have that

$$\Gamma_{1i} = \varphi(1, X_i, \gamma) + \frac{D_i}{e(X_i)}(g(Y_i, X_i, \gamma_1) - \varphi(1, X_i, \gamma)) + \alpha_1(D_i, X_i)(Y_i - \gamma(D_i, X_i)).$$

The arguments for Γ_{0i} are the analogous. ■

Proof of Proposition 4.1:

$$\begin{aligned} W(\pi) &= \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} g(Y_i(a), X_i, Y_j(b), X_j, \gamma^{(a)}, \gamma^{(b)}) \pi_{ab}(X_i, X_j) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left(\sum_{(a,b) \in \{0,1\}^2} g(Y_i(a), X_i, Y_j(b), X_j, \gamma^{(a)}, \gamma^{(b)}) \middle| X_i, X_j \right) \pi_{ab}(X_i, X_j) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left(\sum_{(a,b) \in \{0,1\}^2} g(Y_i(a), X_i, Y_j(b), X_j, \gamma^{(a)}, \gamma^{(b)}) \middle| X_i, D_i = a, X_j, D_j = b \right) \pi_{ab}(X_i, X_j) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left(\sum_{(a,b) \in \{0,1\}^2} g(Y_i, X_i, Y_j, X_j, \gamma^{(a)}, \gamma^{(b)}) \middle| X_i, D_i = a, X_j, D_j = b \right) \pi_{ab}(X_i, X_j) \right] \\ &= \mathbb{E} \left[\mathbb{E} \left(\sum_{(a,b) \in \{0,1\}^2} \frac{g(Y_i, X_i, Y_j, X_j, \gamma^{(a)}, \gamma^{(b)}) D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \middle| X_i, X_j \right) \pi_{ab}(X_i, X_j) \right] \\ &= \mathbb{E} \left[\sum_{(a,b) \in \{0,1\}^2} \frac{g(Y_i, X_i, Y_j, X_j, \gamma^{(a)}, \gamma^{(b)}) D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \pi_{ab}(X_i, X_j) \right], \end{aligned}$$

where in the second equality I use LIE, in the third I use selection on observables, in the fourth I use the definition of Y_i . The identification by the direct method is in the fourth equality while the IPW is the last equality. ■

Proof of Proposition 4.2: Let us start with the DM identification. As usual let $d/d\tau$ be the derivative at $\tau = 0$. Let me also make the dependence on φ explicit: $m_{ab}(Z_i, Z_j, \gamma, \varphi) = \varphi(a, X_i, b, X_j, \gamma_a, \gamma_b)$, let also $\varphi_\tau = \varphi + \tau \tilde{\varphi}$ for some $\tilde{\varphi} \in L_2$. By the chain rule

$$\frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, \varphi_\tau)] = \frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, \varphi)] + \frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \gamma, \varphi_\tau)].$$

By Assumption 3 we have that the first term is

$$\frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, \varphi)] = \mathbb{E} \left[\sum_{p=1}^P \alpha_{ab,p}^\gamma(D_i, X_i, D_j, X_j) (c_{1p} \bar{\gamma}_\tau(D_i, X_i) + c_{2p} \bar{\gamma}_\tau(D_j, X_j)) \right],$$

so by Lemma 1 and equation (2.16) in Escanciano and Terschuur (2022) we have that

$$\phi_{ab}^\gamma(D_i, X_i, D_j, X_j, e, \alpha^\gamma) = \sum_{p=1}^P \alpha_{ab,p}^\gamma(D_i, X_i, D_j, X_j, e) (c_{1p} Y_i + c_{2p} Y_j - c_{1p} \gamma(D_i, X_i) - c_{2p} \gamma(D_j, X_j)).$$

For the second term notice that

$$\begin{aligned} \mathbb{E}[\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b)] &= \mathbb{E} \left[\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \right] \\ &= \mathbb{E} \left[\varphi(a, X_i, b, X_j, \gamma_a, \gamma_b) \frac{1}{e_{ab}(X_i, X_j)} \middle| D_{ij}^{ab} = 1 \right] \mathbb{P}(D_i = a, D_j = b) \\ &= \mathbb{E} \left[\varphi(D_i, X_i, D_j, X_j, \gamma_a, \gamma_b) \frac{D_{ij}^{ab}}{e_{ab}(X_i, X_j)} \right]. \end{aligned}$$

So by the same arguments

$$\phi_{ab}^\varphi(D_i, X_i, D_j, X_j, \varphi, \alpha^m) = \alpha_{ab}^\varphi(D_i, X_i, D_j, X_j) (g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) - \varphi(D_i, X_i, D_j, X_j, \gamma_a, \gamma_b)),$$

with $\alpha_{ab}^\varphi(D_i, X_i, D_j, X_j) = D_{ij}^{ab}/e_{ab}(X_i, X_j)$. For the IPW identification let me make the dependence on the propensity score explicit: $m_{ab}(Z_i, Z_j, \gamma, \varphi, e) = g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}/e_{ab}(X_i, X_j)$.

For $c \in \{0, 1\}$ let $e_{c,\tau} = e_c + \tau \tilde{e}_c$ for some $\tilde{e}_c \in L_2$ and $e_\tau = (e_{a,\tau}, e_{b,\tau})$. Then

$$\frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, e_\tau)] = \frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \bar{\gamma}_\tau, e)] + \frac{d}{d\tau} \mathbb{E}[m_{ab}(Z_i, Z_j, \gamma, e_\tau)].$$

For the first term we have the same result as above by using Assumption 3. For the second term note

$$\begin{aligned} \frac{d}{d\tau} \mathbb{E} \left[\frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_{a,\tau}(X_i) e_{b,\tau}(X_j)} \right] &= \frac{d}{d\tau} \mathbb{E} \left[\frac{-g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i)^2 e_b(X_j)} e_{a,\tau}(X_i) \right] \\ &\quad + \frac{d}{d\tau} \mathbb{E} \left[\frac{-g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i) e_b(X_j)^2} e_{b,\tau}(X_j) \right] \\ &= \frac{d}{d\tau} \mathbb{E} \left[\mathbb{E} \left(\frac{-g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i)^2 e_b(X_j)} \middle| X_i \right) e_{a,\tau}(X_i) \right] \\ &\quad + \frac{d}{d\tau} \mathbb{E} \left[\mathbb{E} \left(\frac{-g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b) D_{ij}^{ab}}{e_a(X_i) e_b(X_j)^2} \middle| X_j \right) e_{b,\tau}(X_j) \right]. \end{aligned}$$

So by the same arguments as before

$$\phi_{ab}^e(D_i, X_i, D_j, X_j, e, \alpha^e) = \alpha_{ab,1}^e(X_i) (\mathbb{1}(D_i = a) - e_a(X_i)) + \alpha_{ab,2}^e(X_j) (\mathbb{1}(D_j = b) - e_b(X_j)),$$

where

$$\begin{aligned}\alpha_{ab,1}^e(X_i) &= -\mathbb{E}\left[\frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b)D_{ij}^{ab}}{e_a(X_i)^2 e_b(X_j)} \middle| X_i\right], \\ \alpha_{ab,2}^e(X_j) &= -\mathbb{E}\left[\frac{g(Y_i, X_i, Y_j, X_j, \gamma_a, \gamma_b)D_{ij}^{ab}}{e_a(X_i) e_b(X_j)^2} \middle| X_j\right].\end{aligned}$$

■

Proof of Proposition 3: Let $\gamma_{c,\tau} = \gamma_c + \tau\tilde{\gamma}_c$ for some $\tilde{\gamma}_c \in L_2$. We have that for $(a, b) \in \{0, 1\}^2$

$$\begin{aligned}\frac{d}{d\tau}\mathbb{E}[\gamma_{a,\tau}(X_i) + \gamma_{b,\tau}(X_j)] &= \frac{d}{d\tau}\mathbb{E}\left[\gamma_{a,\tau}(X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} + \gamma_{b,\tau}(X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right] \\ &= \frac{d}{d\tau}\mathbb{E}\left[\tilde{\gamma}_\tau(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} + \tilde{\gamma}_\tau(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[\tilde{\gamma}_\tau(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} + \tilde{\gamma}_\tau(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right] \\ &\quad - \frac{d}{d\tau}\mathbb{E}_\tau\left[\gamma(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} + \gamma(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[\frac{\mathbb{1}(D_i = a)}{e_a(X_i)}(Y_i - \gamma(D_i, X_i)) + \frac{\mathbb{1}(D_j = b)}{e_b(X_j)}(Y_j - \gamma(D_j, X_j))\right].\end{aligned}$$

Also, let $\Delta_{a,b} = \gamma_a(X_i) - \gamma_b(X_j)$, then

$$\frac{d}{d\tau}\mathbb{E}[|\gamma_{a,\tau}(X_i) - \gamma_{b,\tau}(X_j)|] = \frac{d}{d\tau}\mathbb{E}[|\Delta_{ab} + \tau(\tilde{\gamma}_a(X_i) - \tilde{\gamma}_b(X_j))|].$$

As shown in [Escanciano and Terschuur \(2022\)](#), the Gateaux derivative of the mapping $\Delta \mapsto \mathbb{E}(|\Delta|)$ is some direction ν (assuming no point mass at zero, which follows from the assumptions in the Proposition) is $\mathbb{E}[sgn(\Delta)\nu]$. Hence, by the chain rule

$$\begin{aligned}\frac{d}{d\tau}\mathbb{E}[\gamma_{a,\tau}(X_i) + \gamma_{b,\tau}(X_j)] &= \frac{d}{d\tau}\mathbb{E}[sgn(\gamma_a(X_i) - \gamma_b(X_j))(\gamma_{a,\tau}(X_i) - \gamma_{b,\tau}(X_j))] \\ &= \frac{d}{d\tau}\mathbb{E}\left[sgn(\gamma_a(X_i) - \gamma_b(X_j))\left(\gamma_{a,\tau}(X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} - \gamma_{b,\tau}(X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right)\right] \\ &= \frac{d}{d\tau}\mathbb{E}\left[sgn(\gamma_a(X_i) - \gamma_b(X_j))\left(\gamma_\tau(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} - \gamma_\tau(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right)\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[sgn(\gamma_a(X_i) - \gamma_b(X_j))\left(\gamma_\tau(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} - \gamma_\tau(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right)\right] \\ &\quad - \frac{d}{d\tau}\mathbb{E}_\tau\left[sgn(\gamma_a(X_i) - \gamma_b(X_j))\left(\gamma(D_i, X_i)\frac{\mathbb{1}(D_i = a)}{e_a(X_i)} - \gamma(D_j, X_j)\frac{\mathbb{1}(D_j = b)}{e_b(X_j)}\right)\right] \\ &= \frac{d}{d\tau}\mathbb{E}_\tau\left[sgn(\gamma_a(X_i) - \gamma_b(X_j))\left(\frac{\mathbb{1}(D_i = a)}{e_a(X_i)}(Y_i - \gamma(D_i, X_i)) - \frac{\mathbb{1}(D_j = b)}{e_b(X_j)}(Y_j - \gamma(D_j, X_j))\right)\right].\end{aligned}$$

So by the results in [Escanciano and Terschuur \(2022\)](#), the locally robust score is given by

$$\begin{aligned}
2\Gamma_{ij}^{ab} &= \gamma_a(X_i) + \gamma_b(X_j) - |\gamma_a(X_i) - \gamma_b(X_j)| \\
&+ \frac{\mathbf{1}(D_i = a)}{e_a(X_i)}(Y_i - \gamma(D_i, X_i)) + \frac{\mathbf{1}(D_j = b)}{e_b(X_j)}(Y_j - \gamma(D_j, X_j)) \\
&- \operatorname{sgn}(\gamma_a(X_i) - \gamma_b(X_j)) \left(\frac{\mathbf{1}(D_i = a)}{e_a(X_i)}(Y_i - \gamma(D_i, X_i)) - \frac{\mathbf{1}(D_j = b)}{e_b(X_j)}(Y_j - \gamma(D_j, X_j)) \right) \\
&= \gamma_a(X_i) + \gamma_b(X_j) - |\gamma_a(X_i) - \gamma_b(X_j)| \\
&+ (1 - \operatorname{sgn}(\gamma_a(X_i) - \gamma_b(X_j))) \frac{\mathbf{1}(D_i = a)}{e_a(X_i)}(Y_i - \gamma(D_i, X_i)) \\
&+ (1 + \operatorname{sgn}(\gamma_a(X_i) - \gamma_b(X_j))) \frac{\mathbf{1}(D_j = b)}{e_b(X_j)}(Y_j - \gamma(D_j, X_j)).
\end{aligned}$$

■

Before proving the rest of the main results I introduce a representation of U-statistics which will be very useful for the coming proofs. For any function $f : \mathcal{Z}^2 \rightarrow \mathbb{R}$ let $\mathbb{U}_n f(X_i, X_j) = \binom{n}{2}^{-1} \sum_{i < j} f(X_i, X_j)$. Let κ be the permutations of $\{1, \dots, n\}$, then, as in [Cléménçon et al. \(2008\)](#), we can rewrite

$$\mathbb{U}_n f(Z_i, Z_j) = \frac{1}{n!} \sum_{\kappa} \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} f(Z_{\kappa(i)}, Z_{\kappa(\lfloor n/2 \rfloor + i)}). \quad (7.1)$$

This expresses $\mathbb{U}_n f(Z_i, Z_j)$ as a (dependent) sum of averages of i.i.d. random variables (i.e. $f(Z_{\kappa(i)}, Z_{\kappa(\lfloor n/2 \rfloor + i)})$ are i.i.d. for $i = 1, \dots, \lfloor n/2 \rfloor$).

Proof of Lemma 1: Using the definition of $W(\pi)$ and $\widetilde{W}_n(\pi)$ and the triangle inequality we know that

$$\begin{aligned}
\mathbb{E} \left[\sup_{\pi \in \Pi} |\widetilde{W}_n(\pi) - W(\pi)| \right] &= \mathbb{E} \left[\sup_{\pi \in \Pi} \left| \mathbb{U}_n \sum_{(a,b) \in \{0,1\}^2} \left(\Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j) - \mathbb{E}[\Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j)] \right) \right| \right] \\
&\leq \sum_{(a,b) \in \{0,1\}^2} \mathbb{E} \left[\sup_{\pi \in \Pi} \left| \mathbb{U}_n \left(\Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j) - \mathbb{E}[\Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j)] \right) \right| \right].
\end{aligned}$$

By the representation used in (7.1) we can rewrite the above as

$$\begin{aligned}
&\sum_{(a,b) \in \{0,1\}^2} \mathbb{E} \left[\sup_{\pi \in \Pi} \left| \frac{1}{n!} \sum_{\kappa} \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \left(\Gamma_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)}^{ab} \pi_{ab}(X_{\kappa(i)}, X_{\kappa(\lfloor n/2 \rfloor + i)}) \right. \right. \right. \\
&\quad \left. \left. \left. - \mathbb{E}[\Gamma_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)}^{ab} \pi_{ab}(X_{\kappa(i)}, X_{\kappa(\lfloor n/2 \rfloor + i)})] \right) \right| \right]. \quad (7.2)
\end{aligned}$$

Introduce an independent ghost sample (Z'_1, \dots, Z'_n) which is distributed as (Z_1, \dots, Z_n) , Rademacher random variables ε_i , $i = 1, \dots, n$, such that $\mathbb{P}(\varepsilon_i = 1) = \mathbb{P}(\varepsilon_i = -1) = 1/2$ and construct ghost

scores $\Gamma'_{ij}{}^{ab}$ using the ghost sample. Let \mathbb{E}_Z is the expectation with respect to the distribution of the sample (Z_1, \dots, Z_n) and define $\mathbb{E}_{Z'}$ and \mathbb{E}_ε similarly. Define the Rademacher complexity as

$$\mathcal{R}_n(\Pi) = \mathbb{E}_\varepsilon \left(\sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \right| \right).$$

Again the key here is that the summands of the sum inside the expectation in $\mathcal{R}_n(\Pi)$ are independent. We are now ready to use a classical symmetrization argument, since Z'_i has the same distribution as Z_i we have that (7.2) is equal to

$$\begin{aligned} & \sum_{(a,b) \in \{0,1\}^2} \mathbb{E}_Z \left[\sup_{\pi \in \Pi} \left| \frac{1}{n!} \sum_{\kappa} [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \left(\Gamma_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)}^{ab} \pi_{ab}(X_{\kappa(i)}, X_{\kappa(\lfloor n/2 \rfloor + i)}) \right. \right. \right. \\ & \quad \left. \left. \left. - \Gamma'_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)} \pi_{ab}(X'_{\kappa(i)}, X'_{\kappa(\lfloor n/2 \rfloor + i)}) \right) \right| \right] \\ & \leq \frac{1}{n!} \sum_{\kappa} \sum_{(a,b) \in \{0,1\}^2} \mathbb{E}_{Z,Z'} \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \left(\Gamma_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)}^{ab} \pi_{ab}(X_{\kappa(i)}, X_{\kappa(\lfloor n/2 \rfloor + i)}) \right. \right. \right. \\ & \quad \left. \left. \left. - \Gamma'_{\kappa(i)\kappa(\lfloor n/2 \rfloor + i)} \pi_{ab}(X'_{\kappa(i)}, X'_{\kappa(\lfloor n/2 \rfloor + i)}) \right) \right| \right] \\ & = \sum_{(a,b) \in \{0,1\}^2} \mathbb{E}_{Z,Z',\varepsilon} \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \left(\Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \right. \right. \right. \\ & \quad \left. \left. \left. - \Gamma'_{i, \lfloor n/2 \rfloor + i} \pi_{ab}(X'_i, X'_{\lfloor n/2 \rfloor + i}) \right) \right| \right] \\ & \leq \sum_{(a,b) \in \{0,1\}^2} \mathbb{E}_{Z,Z',\varepsilon} \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) \right| \right. \\ & \quad \left. + \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma'_{i, \lfloor n/2 \rfloor + i} \pi_{ab}(X'_i, X'_{\lfloor n/2 \rfloor + i}) \right| \right] \\ & = \sum_{(a,b) \in \{0,1\}^2} \mathbb{E}[2\mathcal{R}_n(\Pi)]. \end{aligned}$$

The first inequality follows from Jensen's and triangle inequalities, the second equality uses the fact that the vector $(Z_{\pi(i)}, Z_{\pi(\lfloor n/2 \rfloor + i)}, Z'_{\pi(i)}, Z'_{\pi(\lfloor n/2 \rfloor + i)})$ is identically distributed across $i = 1, \dots, \lfloor n/2 \rfloor$ for all permutations in κ (so we can just take the permutation $\kappa(i) = i$) and the fact that $\varepsilon_i(\Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - \Gamma'_{i, \lfloor n/2 \rfloor + i} \pi_{ab}(X'_i, X'_{\lfloor n/2 \rfloor + i}))$ and $\Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - \Gamma'_{i, \lfloor n/2 \rfloor + i} \pi_{ab}(X'_i, X'_{\lfloor n/2 \rfloor + i})$ have the same distribution, the third inequality uses the triangle inequality and the last equality uses that $Z_i \sim Z'_i$ and the definition of the Rademacher complexity. ■

Proof of Lemma 2: Note that Lemma 5 gives us a sequence of covers B_k for $k = 0, \dots, K$ of $\tilde{\Pi}_{ab}$ of radius less than 2^{-k} for some K . For any $j = 1, \dots, J$ with $J = \lceil \log_2(\lfloor n/2 \rfloor)(1 - \beta) \rceil$

and $\pi \in \tilde{\Pi}_{ab}$ let $b_j : \tilde{\Pi}_{ab} \mapsto \tilde{\Pi}_{ab}$ be an operator such that $b_j(\pi)$ is an approximating policy from the cover B_j such that $D_n(\pi, b_j(\pi)) \leq 2^{-j}$, such an approximation exists by Lemma 5. By the same Lemma we also know that $|\{b_j(\pi) : \pi \in \tilde{\Pi}_{ab}\}| \leq N_{D_n}(2^{-j}, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor})$. Let $\underline{J} = \lceil 1/2 \log_2(\lfloor n/2 \rfloor)(1 - \beta) \rceil$. By using a telescope sum and the approximations b_0, \dots, b_J we can decompose the Rademacher complexity as

$$\begin{aligned} \mathcal{R}_n(\Pi) = \mathbb{E}_\varepsilon \left\{ \sup_{\pi \in \Pi} \left| \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \left[b_0(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) \right. \right. \right. \\ + \sum_{j=1}^{\underline{J}} \left(b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) \right) \\ + (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{\underline{J}}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \\ \left. \left. \left. + (\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right] \right| \right\}. \end{aligned}$$

Note that since the distance D_n is bounded by 1, by the second property in Lemma 5 we have that b_0 can be any policy in $\tilde{\Pi}_{ab}$. Hence, we can set $b_0(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) = 0$ for all $i = 1, \dots, \lfloor n/2 \rfloor$. We approach each of the terms above in turn. Note that b_0, \dots, b_J is a sequence of increasingly accurate approximations. The first step is to notice that the last term above is negligible, i.e. the term involving the closest approximation vanishes at a \sqrt{n} rate. By using Cauchy-Schwarz and multiplying and dividing we get

$$\begin{aligned} & \sqrt{\lfloor n/2 \rfloor} \sup_{\pi \in \Pi} \left| \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \\ & \leq \sqrt{\lfloor n/2 \rfloor} \sup_{\pi \in \Pi} \frac{\sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \left| \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right|^2}}{\sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}} \\ & \times \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \\ & = \sqrt{\lfloor n/2 \rfloor} \sup_{\pi \in \Pi} D_n(\pi_{ab}, b_J(\pi_{ab})) \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \\ & \leq \sqrt{\lfloor n/2 \rfloor} 2^{-J} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \\ & = \frac{M}{\lfloor n/2 \rfloor^{1/2-\beta}} \rightarrow 0, \end{aligned}$$

where in the last inequality we use Lemma 5 and in the last inequality we use the fact that $J = \lceil \log_2(\lfloor n/2 \rfloor)(1 - \beta) \rceil$ and the boundedness assumption. Now we show that the second to last term

of the Rademacher decomposition is also negligible. Notice that $\{\varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{\underline{J}}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})))\}_{i=1}^{\lfloor n/2 \rfloor}$ are zero mean (conditional on $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$) i.i.d. random variables. They are also bounded below by $a_i = -|\Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{\underline{J}}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})))|$ and above by $b_i = a_i$. Hence, by Hoeffding's inequality

$$\begin{aligned} & \mathbb{P}_\varepsilon \left(\left| \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{\underline{J}}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \geq t \right) \\ & \leq 2 \exp \left(- \frac{2t^2}{\sum_{i=1}^{\lfloor n/2 \rfloor} (b_i - a_i)^2} \right) \\ & = 2 \exp \left(- \frac{t^2}{D_n^2(b_J(\pi_{ab}), b_{\underline{J}}(\pi_{ab})) \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \right). \end{aligned}$$

Hence, for any $a > 0$ we have that

$$\begin{aligned} & \mathbb{P}_\varepsilon \left(\left| \sqrt{\lfloor n/2 \rfloor} \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{\underline{J}}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right. \\ & \geq a 2^{2-\underline{J}} \sqrt{\frac{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{\lfloor n/2 \rfloor}} \Bigg) \\ & \leq 2 \exp \left(- \frac{a^2 4^{2-\underline{J}}}{D_n^2(b_J(\pi_{ab}), b_{\underline{J}}(\pi_{ab}))} \right) \\ & \leq 2 \exp \left(- \frac{a^2 4^{2-\underline{J}}}{\sum_{j=\underline{J}}^{J-1} D_n^2(b_j(\pi_{ab}), b_{j+1}(\pi_{ab}))} \right) \\ & \leq 2 \exp \left(- \frac{a^2 4^{2-\underline{J}}}{\left(\sum_{j=\underline{J}}^{J-1} 2^{-(j-1)} \right)^2} \right) \\ & \leq 2 \exp(-a^2), \end{aligned}$$

where we have used triangle inequality in the second inequality and the fact that $\sum_{j=\underline{J}}^{J-1} 2^{-(j-1)} =$

$2^{2-J} - 2^{2-J} \leq 2^{2-J}$ in the last inequality. This holds for any policy, hence

$$\begin{aligned}
& \mathbb{P}_\varepsilon \left(\sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right. \\
& \quad \left. \geq a 2^{2-J} \sqrt{\frac{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{\lfloor n/2 \rfloor}} \right) \\
& \leq 2 |\{b_J(\pi_{ab}), b_J(\pi_{ab})\}| \exp(-a^2) \\
& \leq 2 N_{D_n}(2^{-J}, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}) \exp(-a^2) \\
& \leq 2 N_H(2^{-2J}, \tilde{\Pi}_{ab}) \exp(-a^2) \\
& = 2 \exp(\log(N_H(2^{-2J}, \tilde{\Pi}_{ab}))) \exp(-a^2) \\
& \leq 2 \exp(5VC(\tilde{\Pi}_{ab}) \log(2^{2J}) - a^2) \\
& \leq 2 \exp(5VC(\tilde{\Pi}_{ab}) \log(2^{-2(1-\beta) \log_2(\lfloor n/2 \rfloor)}) - a^2),
\end{aligned}$$

where in the first inequality I use the union bound, in the second inequality I use properties of the approximations (see [Zhou et al. \(2023\)](#)), in the third I use Lemma 4 and in the fourth inequality I bound the log of the Hamming covering number by the VC dimension using a result in [Haussler \(1995\)](#). Let now

$$a = \frac{2^J}{\sqrt{\log(\lfloor n/2 \rfloor) [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}},$$

so that

$$a 2^{2-J} \sqrt{\frac{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{\lfloor n/2 \rfloor}} = \frac{4}{\sqrt{\log(\lfloor n/2 \rfloor)}}.$$

Finally,

$$\begin{aligned}
& \mathbb{P}_\varepsilon \left(\sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right. \\
& \quad \left. \geq \frac{4}{\sqrt{\log(\lfloor n/2 \rfloor)}} \right) \\
& \leq 2 \exp \left(5VC(\tilde{\Pi}_{ab}) \log(\lfloor n/2 \rfloor^{-2(1-\beta)}) - \frac{\lfloor n/2 \rfloor^{-\beta}}{\log(\lfloor n/2 \rfloor) \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \right) \\
& \leq 2 \exp \left\{ -5 \lfloor n/2 \rfloor^\beta \log \left(\lfloor n/2 \rfloor^{2(1-\beta)} \right) - \frac{1}{\lfloor n/2 \rfloor^\beta \log(\lfloor n/2 \rfloor) M^2} \right\} \rightarrow 0,
\end{aligned}$$

where I have used Assumption 7 and the boundedness assumption.

$$\mathbb{E} \left(\sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_J(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right) \rightarrow 0,$$

since for any sequence of random variables X_n and sequence of real numbers a_n if $\lim_{n \rightarrow \infty} \mathbb{P}(X_n \leq a_n) = 1$ and $\lim_{n \rightarrow \infty} a_n = 0$, then $\lim_{n \rightarrow \infty} \mathbb{E}(X_n) = 0$ (proof of this fact uses $\mathbb{E}(X_n) = \int_0^\infty \mathbb{P}(X_n > u) du$). Hence, we have proven that

$$\mathbb{E}[\mathcal{R}_n(\Pi)] = \mathbb{E} \left\{ \sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \left[\sum_{j=1}^J \left(b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) \right) \right] \right| \right\} + o\left(\frac{1}{\sqrt{n}}\right).$$

Hence I have left what [Zhou et al. \(2023\)](#) call the effective regime. Let $j \in \{1, \dots, J\}$ and a_j be some constant depending on j . As before, conditional on $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$ we can apply Hoeffding inequality and then use the definition of D_n to get

$$\begin{aligned} \mathbb{P}_\varepsilon \left(\left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right. \\ \geq a_j 2^{2-j} \sqrt{\frac{\sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}}{[n/2]}} \\ \leq 2 \exp \left(-\frac{a_j^2 4^{2-j}}{D_n^2(b_j(\pi_{ab}), b_{j-1}(\pi_{ab}))} \right) \\ \leq 2 \exp \left(\frac{-a_j^2 4^{2-j}}{4^{-(j-1)}} \right) \\ = 2 \exp \left(-4a_j^2 \right), \end{aligned}$$

where in the last inequality we have used the fact that $D_n(b_j(\pi_{ab}), b_{j-1}(\pi_{ab})) \leq 2^{-(j-1)}$ by Lemma 5. Now we let

$$a_j^2(k) = 2 \log \left(\frac{2j^2}{\delta_k} N_H(4^{-j}, \tilde{\Pi}_{ab}) \right),$$

where δ_k is some sequence of real numbers indexed by $k \in \mathbb{N}$. For notational convenience define

$$R_j = \sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} (b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right|.$$

Then we have that

$$\begin{aligned} \mathbb{P} \left(R_j \geq a_j(k) 2^{-j} \sqrt{[n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \right) &\leq 2 |\{b_j(\pi_{ab}), b_{j-1}(\pi_{ab})\}| \exp(-a_j^2(k)/2) \\ &\leq 2 N_{D_n}(2^{-j}, \tilde{\Pi}_{ab}, \{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}) \exp(-a_j^2(k)/2) \\ &\leq 2 N_H(2^{-2j}, \tilde{\Pi}_{ab}) \exp(-a_j^2(k)/2) \\ &= 2 N_H(4^{-j}, \tilde{\Pi}_{ab}) \exp(-\log(N_H(4^{-j}, \tilde{\Pi}_{ab}) 2j^2 / \delta_k)) \\ &= \frac{\delta_k}{j^2}. \end{aligned}$$

Sum across j and apply this bound with $\delta_k = 1/2^k$ to note that

$$\begin{aligned}
\sum_{j=1}^J \mathbb{P} \left(R_j \geq a_j(k) 2^{-j} \sqrt{ \lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} } \right) &\leq \sum_{j=1}^J \frac{\delta_k}{j^2} \\
&\leq \sum_{j=1}^{\infty} \frac{\delta_k}{j^2} \\
&\leq \frac{1.7}{2^k}.
\end{aligned}$$

Let F_{R_j} be the cumulative distribution function of R_j (conditional on $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$). We

can bound the following object of interest in the following way

$$\begin{aligned}
& \mathbb{E}_\varepsilon \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \sum_{j=1}^J (b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right] \\
& \leq \sum_{j=1}^J \mathbb{E}_\varepsilon [R_j] \\
& = \int_0^\infty \sum_{j=1}^J (1 - F_{R_j}(r)) dr \\
& \leq \int_0^\infty \sum_{j=1}^J \mathbb{P}(R_j \geq r) dr \\
& \leq \sum_{k=0}^\infty \sum_{j=1}^J \frac{1.7}{2^k} a_j(k) 2^{-j} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \\
& \leq \sum_{k=0}^\infty \sum_{j=1}^J \frac{1.7}{2^k} \sqrt{2} \sqrt{\log(2^{k+1} j^2 N_H(4^{-j}, \tilde{\Pi}_{ab}))} 2^{-j} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \\
& \leq 1.7\sqrt{2} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \sum_{k=0}^\infty 2^{-k} \sum_{j=1}^J 2^{-j} \sqrt{(k+1) \log 2 + 2 \log j + \log N_H(4^{-j}, \tilde{\Pi}_{ab})}} \\
& \leq 1.7\sqrt{2} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \sum_{k=0}^\infty 2^{-k} \sum_{j=1}^J 2^{-j} \left(\sqrt{k+1} + \sqrt{2 \log j} + \sqrt{5VC(\tilde{\Pi}_{ab}) \log(4^j)} \right)} \\
& \leq 1.7\sqrt{2} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \sum_{k=0}^\infty 2^{-k} \left(\sqrt{k+1} \sum_{j=1}^\infty 2^{-j} + \sqrt{2} \sum_{j=1}^\infty 2^{-j} \sqrt{\log j} \right.} \\
& \quad \left. + \sqrt{5VC(\tilde{\Pi}_{ab})} \sum_{j=1}^\infty 2^{-j} \sqrt{\log 4^j} \right)} \\
& \leq 1.7\sqrt{2} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \left(\sum_{k=0}^\infty 2^{-k} \sqrt{k+1} + \frac{\sqrt{2}}{2} \sum_{k=0}^\infty 2^{-k} + \sqrt{5VC(\tilde{\Pi}_{ab})} 1.6 \sum_{k=0}^\infty 2^{-k} \right)} \\
& \leq 1.7\sqrt{2} \sqrt{\lfloor n/2 \rfloor^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \left(5 + 3.2 \sqrt{5VC(\tilde{\Pi}_{ab})} \right)}.
\end{aligned}$$

So taking expectations over $\{X_i, \Gamma_{i, \lfloor n/2 \rfloor + i}\}_{i=1}^{\lfloor n/2 \rfloor}$, using this bound and the Jensen's inequality

we get

$$\begin{aligned}
& \mathbb{E} \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1/2} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \sum_{j=1}^J (b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right] \\
& \leq 1.7\sqrt{2} \left(5 + 8\sqrt{5VC(\tilde{\Pi}_{ab})} \right) \mathbb{E} \left[\sqrt{[n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab}} \right] \\
& \leq 1.7\sqrt{2} \left(5 + 8\sqrt{5VC(\tilde{\Pi}_{ab})} \right) \sqrt{\mathbb{E} \left[\Gamma_{i, \lfloor n/2 \rfloor + i}^{2ab} \right]} \\
& = 1.7\sqrt{2} \left(5 + 8\sqrt{5VC(\tilde{\Pi}_{ab})} \right) \sqrt{S_{ab}} \\
& \leq C\sqrt{VC(\tilde{\Pi}_{ab})S_{ab}},
\end{aligned}$$

for some constant $C > 0$. Dividing both sides by $\sqrt{\lfloor n/2 \rfloor}$ we get

$$\begin{aligned}
& \mathbb{E} \left[\sup_{\pi \in \Pi} \left| [n/2]^{-1} \sum_{i=1}^{\lfloor n/2 \rfloor} \varepsilon_i \Gamma_{i, \lfloor n/2 \rfloor + i}^{ab} \sum_{j=1}^J (b_j(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i})) - b_{j-1}(\pi_{ab}(X_i, X_{\lfloor n/2 \rfloor + i}))) \right| \right] \\
& \leq C\sqrt{\frac{VC(\tilde{\Pi}_{ab})S_{ab}}{\lfloor n/2 \rfloor}},
\end{aligned}$$

and hence

$$\begin{aligned}
\mathbb{E}[\mathcal{R}_n(\Pi)] & \leq C\sqrt{\frac{VC(\tilde{\Pi}_{ab})S_{ab}}{\lfloor n/2 \rfloor}} + o\left(\frac{1}{\sqrt{n}}\right) \\
& = \mathcal{O}\left(\sqrt{\frac{VC(\tilde{\Pi}_{ab})S_{ab}}{\lfloor n/2 \rfloor}}\right).
\end{aligned}$$

■

Proof of Lemma 3: Define the following random variables

$$\begin{aligned}
\hat{R}_{ij,ab,l}^{(1)} &= m_{ab}(Z_i, Z_j, \hat{\gamma}_l, \nu) - m_{ab}(Z_i, Z_j, \gamma, \nu) \\
\hat{R}_{ij,ab,l}^{(2)} &= m_{ab}(Z_i, Z_j, \gamma, \hat{\nu}_l) - m_{ab}(Z_i, Z_j, \gamma, \nu) \\
\hat{R}_{ij,ab,l}^{(3)} &= \phi_{ab}^\gamma(Z_i, Z_j, \hat{\gamma}_l, \alpha^\gamma) - \phi_{ab}^\gamma(Z_i, Z_j, \gamma, \alpha^\gamma) \\
\hat{R}_{ij,ab,l}^{(4)} &= \phi_{ab}^\gamma(Z_i, Z_j, \gamma, \hat{\alpha}_l^\gamma) - \phi_{ab}^\gamma(Z_i, Z_j, \gamma, \alpha^\gamma) \\
\hat{R}_{ij,ab,l}^{(5)} &= \phi_{ab}^\nu(Z_i, Z_j, \hat{\nu}_l, \alpha^\nu) - \phi_{ab}^\nu(Z_i, Z_j, \nu, \alpha^\nu) \\
\hat{R}_{ij,ab,l}^{(6)} &= \phi_{ab}^\nu(Z_i, Z_j, \nu, \hat{\alpha}_l^\nu) - \phi_{ab}^\nu(Z_i, Z_j, \nu, \alpha^\nu).
\end{aligned}$$

Then,

$$\mathbb{E} \left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - \widetilde{W}_n(\pi)| \right] = \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{l=1}^L \sum_{(i,j) \in I_l} \sum_{(a,b) \in \pi} \sum_{k=1}^6 \hat{R}_{ij,ab,l}^{(k)} + \hat{\xi}_{ij,ab,l} + \hat{\xi}_{ij,ab,l}^\gamma + \hat{\xi}_{ij,ab,l}^\nu \right| \pi_{ab}(X_i, X_j) \right).$$

By repeated use of the triangle inequality

$$\begin{aligned}
(\dagger) \quad \mathbb{E} \left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - \widetilde{W}_n(\pi)| \right] &\leq \sum_{l=1}^L \sum_{(a,b) \in \pi} \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{R}_{ij,l}^{(1)} + \hat{R}_{ij,l}^{(3)}) \pi_{ab}(X_i, X_j) \right| \right) \\
&\quad + \sum_{l=1}^L \sum_{(a,b) \in \pi} \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{R}_{ij,l}^{(2)} + \hat{R}_{ij,l}^{(5)}) \pi_{ab}(X_i, X_j) \right| \right) \\
&\quad + \sum_{l=1}^L \sum_{(a,b) \in \pi} \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{R}_{ij,l}^{(4)} + \hat{R}_{ij,l}^{(6)}) \pi_{ab}(X_i, X_j) \right| \right) \\
&\quad + \sum_{l=1}^L \sum_{(a,b) \in \pi} \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{\xi}_{ij,l} + \hat{\xi}_{ij,l}^\gamma + \hat{\xi}_{ij,l}^\nu) \pi_{ab}(X_i, X_j) \right| \right).
\end{aligned}$$

I will bound each of the terms separately. The same arguments apply for all $l = 1, \dots, L$ and $(a, b) \in \pi$, hence we focus on some fixed (a, b) and l . Let N_l^c be the observations not in I_l . By adding and subtracting $\mathbb{E}[(\hat{R}_{ij,ab,l}^{(1)} + \hat{R}_{ij,ab,l}^{(3)}) \pi_{ab}(X_i, X_j) | N_l^c]$ and applying the triangle inequality we get that the summands of the first term are bounded by

$$\begin{aligned}
&\mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{R}_{ij,l}^{(1)} + \hat{R}_{ij,l}^{(3)}) \pi_{ab}(X_i, X_j) - \mathbb{E}[(\hat{R}_{ij,ab,l}^{(1)} + \hat{R}_{ij,ab,l}^{(3)}) \pi_{ab}(X_i, X_j) | N_l^c] \right| \right) \quad (\star) \\
&\quad + \mathbb{E} \left(\sup_{\pi \in \Pi_n} \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} |\mathbb{E}[(\hat{R}_{ij,ab,l}^{(1)} + \hat{R}_{ij,ab,l}^{(3)}) \pi_{ab}(X_i, X_j) | \hat{\gamma}_l]| \right). \quad (\star\star)
\end{aligned}$$

By Assumption 5 we know that

$$\begin{aligned}
|\mathbb{E}[\hat{R}_{ij,ab,l}^{(1)} + \hat{R}_{ij,ab,l}^{(3)} | N_l^c]| &= |\mathbb{E}[m_{ab}(Z_i, Z_j, \hat{\gamma}_l, \nu) + \phi_{ab}^\gamma(Z_i, Z_j, \hat{\gamma}_l, \alpha^\gamma) | \hat{\gamma}_l]| \\
&\leq C \|\hat{\gamma}_l - \gamma\|^2.
\end{aligned}$$

Applying the conditional Jensen's inequality (on the absolute value) in $(\star\star)$ and noting that the resulting expression is maximized by treating everybody we get that

$$(\star\star) \leq C \underbrace{\mathbb{E}[\|\hat{\gamma}_l - \gamma\|^2]}_{\leq 1} \binom{n}{2}^{-1} |I_l| = o(n^{-2\lambda_\gamma}) = o(1/\sqrt{n}),$$

where the last equality follows since $2\lambda_\gamma \geq 1/2$. For (\star) , note that

$$\begin{aligned}
(\star) &\leq \binom{n}{2}^{-1} |I_l| \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| |I_l|^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \right) \\
&+ \binom{n}{2}^{-1} |I_l| \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| |I_l|^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(3)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(3)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \right) \\
&= \binom{n}{2}^{-1} |I_l| \mathbb{E} \left[\mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| |I_l|^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \middle| N_l^c \right) \right] \\
&+ \binom{n}{2}^{-1} |I_l| \mathbb{E} \left[\mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| |I_l|^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(3)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(3)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \middle| N_l^c \right) \right].
\end{aligned}$$

The inner expectations are the expected supremum of centered U-processes. Using Lemma 1 We can bound these inner expectations with Rademacher complexities. However, in the same way we used the construction in Appendix ?? in Lemma 1 to be able to bound the U-process with a Rademacher complexity which involves a sum of independent terms, we need to use such a construction for each fold I_l . From Appendix ?? we know that I_l can be either a triangle ($I_l \in T$) or a square ($I_l \in S$) and that in each case we can bound the U-process with the following Rademacher complexities

$$\mathcal{R}_{n,l}(\Pi_{ab}) = \begin{cases} \mathbb{E}_\varepsilon \left(\sup_{\pi \in \Pi} \left| |C_k|^{-1} \sum_{i=1}^{|C_k|} \varepsilon_i \hat{R}_{\rho(i,k),|C_k|+i}^{(q)} \pi_{ab}(X_{\rho(i,k)}, X_{|C_k|+i}) \right| \right) & \text{if } I_l \in S \\ \mathbb{E}_\varepsilon \left(\sup_{\pi \in \Pi} \left| \lfloor |C_k|/2 \rfloor^{-1} \sum_{i=1}^{\lfloor |C_k|/2 \rfloor} \varepsilon_i \hat{R}_{i, \lfloor |C_k|/2 \rfloor + i, l}^{(q)} \pi_{ab}(X_i, X_{\lfloor |C_k|/2 \rfloor + i}) \right| \right) & \text{if } I_l \in T, \end{cases}$$

for $q = 1, 3$. Hence, by Lemmas 1 and 2 we have that

$$\binom{n}{2}^{-1} |I_l| \mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| |I_l|^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \middle| N_l^c \right) = \mathcal{O} \left(\sqrt{\frac{S_{ab,l}^{(1)} VC(\Pi_{ab,n})}{\lfloor |C_k|/2 \rfloor}} \right),$$

where $S_{ab,l}^{(1)} = \mathbb{E}[\hat{R}_{ij,l}^{(1)2} | N_l^c]$. Noting that $\mathbb{E}[S_{ab,l}^{(1)}] = \mathbb{E}[(m_{ab}(Z_i, Z_j, \hat{\gamma}_l, \nu) - m_{ab}(Z_i, Z_j, \gamma, \nu))^2]$ and using Assumption 4, Jensen's inequality, the fact that $|I_l| = |C_k| \times |C_m|$ if $I_l = I(C_k, C_m)$ and $|I_l| = |C_k| \times |C_k - 1|/2$ if $I_l = I(C_k, C_k)$ and that for evenly sized folds $|C_k|/(n-1) \leq 1$ for all $k = 1, \dots, K$ we have that

$$\begin{aligned}
&\mathbb{E} \left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) - \mathbb{E}[\hat{R}_{ij,l}^{(1)} \pi_{ab}(X_i, X_j) | N_l^c] \right| \right) \\
&= \mathcal{O} \left(\sqrt{VC(\Pi_{ab,n}) \frac{a((1-K^{-1})n)^2}{n^{1+2\lambda_\gamma}}} \right).
\end{aligned}$$

The same bound applies by using the same arguments when we replace $\hat{R}_{ij,l}^{(1)}$ by $\hat{R}_{ij,l}^{(3)}$. Also, this bound applies to all folds I_l , hence, summing across all folds gives us the same asymptotic

bound. As a result, we have bounded the first term in the RHS in (\dagger) . For the second term we can follow exactly the same steps as with the first term to get the same bounds with λ_γ replaced by λ_ν . For the third term in (\dagger) note that by Assumption 5 (i) (global robustness of α), we have that $\mathbb{E}[\hat{R}_{ij,ab,l}^{(4)}|N_l^c] = \mathbb{E}[\hat{R}_{ij,ab,l}^{(6)}|N_l^c] = 0$. Hence, we do not need to add and subtract anything and we can apply Lemmas 1 and 2 directly to get that for $q = 4, 6$

$$\mathbb{E}\left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} \hat{R}_{ij,l}^{(q)} \pi_{ab}(X_i, X_j) \right| \right) = \mathcal{O}\left(\sqrt{VC(\Pi_{ab,n}) \frac{a((1-K^{-1})n)^2}{n^{1+2\lambda_\alpha}}}\right).$$

Finally, the bound for the last term in (\dagger) follows directly from Assumption 6

$$\mathbb{E}\left(\sup_{\pi \in \Pi_n} \left| \binom{n}{2}^{-1} \sum_{(i,j) \in I_l} (\hat{\xi}_{ij,l} + \hat{\xi}_{ij,l}^\gamma + \hat{\xi}_{ij,l}^\nu) \pi_{ab}(X_i, X_j) \right| \right) = \mathcal{O}\left(\frac{a(1-K^{-1})}{\sqrt{n}}\right).$$

Putting everything together we know that

$$\begin{aligned} \sqrt{n} \mathbb{E}\left[\sup_{\pi \in \Pi_n} |\hat{W}_n(\pi) - \widetilde{W}_n(\pi)|\right] &= \mathcal{O}\left(\sqrt{VC(\Pi_{ab,n}) \frac{a((1-K^{-1})n)^2}{n^{2\lambda_\gamma}}}\right) \\ &\quad + \mathcal{O}\left(\sqrt{VC(\Pi_{ab,n}) \frac{a((1-K^{-1})n)^2}{n^{2\lambda_\nu}}}\right) \\ &\quad + \mathcal{O}\left(\sqrt{VC(\Pi_{ab,n}) \frac{a((1-K^{-1})n)^2}{n^{2\lambda_\alpha}}}\right) \\ &\quad + \mathcal{O}\left(a(1-K^{-1})\right) + o(1) \\ &= \mathcal{O}\left(a((1-K^{-1})n) \left(1 + \sqrt{\frac{VC(\Pi_{ab,n})}{n^{2\min(\lambda_\gamma, \lambda_\nu, \lambda_\alpha)}}}\right)\right). \end{aligned}$$

■

Proof of Theorem 1: Follows from Lemmas 2, 3 and 6. ■

Proof of Corollary 1: Let Γ_{ij}^{ab} and $\hat{\Gamma}_{ij,l}^{ab}$ be defined as in the Intergenerational mobility example and let

$$\begin{aligned} K(\pi) &= \mathbb{E}\left[\sum_{(a,b) \in \{0,1\}^2} \Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j)\right], \\ \widetilde{K}_n(\pi) &= \binom{n}{2}^{-1} \sum_{i < j} \left[\sum_{(a,b) \in \{0,1\}^2} \Gamma_{ij}^{ab} \pi_{ab}(X_i, X_j) \right], \\ \hat{K}_n(\pi) &= \binom{n}{2}^{-1} \sum_{l=1}^L \sum_{(i,j) \in I_l} \left[\sum_{(a,b) \in \pi} \hat{\Gamma}_{ij,l}^{ab} (Z_i, Z_j, \hat{\gamma}_l, \hat{\nu}_l, \hat{\alpha}_l) \pi_{ab}(X_i, X_j) \right]. \end{aligned}$$

Note also that $W(\pi) = -|K(\pi) - t|$. Hence, we can write the regret as

$$\mathbb{E}\left[\sup_{\pi \in \Pi_n} -|K(\pi) - t| + |K(\hat{\pi}) - t|\right] \leq \mathbb{E}\left[\sup_{\pi \in \Pi_n} |K(\pi) - K(\hat{\pi})|\right].$$

The result follows from applying Theorem 1 with W replaced by K . ■

References

- AABERGE, R., T. HAVNES, AND M. MOGSTAD (2021): “Ranking intersecting distribution functions,” *Journal of Applied Econometrics*, 36, 639–662.
- ATHEY, S., J. TIBSHIRANI, AND S. WAGER (2019): “Generalized random forests,” *The Annals of Statistics*, 47, 1148–1178.
- ATHEY, S. AND S. WAGER (2021): “Policy learning with observational data,” *Econometrica*, 89, 133–161.
- ATKINSON, A. B. ET AL. (1970): “On the measurement of inequality,” *Journal of economic theory*, 2, 244–263.
- BELLONI, A. AND V. CHERNOZHUKOV (2011): “l1-penalized quantile regression in high-dimensional sparse models,” .
- (2013): “Least squares after model selection in high-dimensional sparse models,” .
- BHATTACHARYA, D. AND P. DUPAS (2012): “Inferring welfare maximizing treatment assignment under budget constraints,” *Journal of Econometrics*, 167, 168–196.
- BRUNORI, P., P. HUFÉ, AND D. MAHLER (2021): “The roots of inequality: Estimating inequality of opportunity from regression trees and forests,” *IZA Discussion Paper No. 14689*.
- BRUNORI, P. AND G. NEIDHÖFER (2021): “The evolution of inequality of opportunity in Germany: A machine learning approach,” *Review of Income and Wealth*, 67, 900–927.
- BRUNORI, P., F. PALMISANO, AND V. PERAGINE (2019a): “Inequality of opportunity in sub-Saharan Africa,” *Applied Economics*, 51, 6428–6458.
- BRUNORI, P., V. PERAGINE, AND L. SERLENGA (2019b): “Upward and downward bias when measuring inequality of opportunity,” *Social Choice and Welfare*, 52, 635–661.
- CAMPBELL, F. A., E. P. PUNGELLO, M. BURCHINAL, K. KAINZ, Y. PAN, B. H. WASIK, O. A. BARBARIN, J. J. SPARLING, AND C. T. RAMEY (2012): “Adult outcomes as a function of an early childhood educational program: an Abecedarian Project follow-up,” *Developmental psychology*, 48, 1033.
- CAMPBELL, F. A. AND C. T. RAMEY (1994): “Effects of early intervention on intellectual and academic achievement: a follow-up study of children from low-income families,” *Child development*, 65, 684–698.

- CARRANZA, R. (2022): “Upper and Lower Bound Estimates of Inequality of Opportunity: A Cross-National Comparison for Europe,” *Review of Income and Wealth*.
- CHAMBERLAIN, G. (2011): “1011 Bayesian Aspects of Treatment Choice,” in *The Oxford Handbook of Bayesian Econometrics*, Oxford University Press.
- CHEN, X. (2007): “Large sample sieve estimation of semi-nonparametric models,” *Handbook of econometrics*, 6, 5549–5632.
- CHEN, X. AND H. WHITE (1999): “Improved rates and asymptotic normality for nonparametric neural network estimators,” *IEEE Transactions on Information Theory*, 45, 682–691.
- CHERNOZHUKOV, V., J. C. ESCANCIANO, H. ICHIMURA, W. K. NEWEY, AND J. M. ROBINS (2021): “Locally robust semiparametric estimation,” *Forthcoming Econometrica*.
- CHETTY, R., N. HENDREN, P. KLINE, AND E. SAEZ (2014): “Where is the land of opportunity? The geography of intergenerational mobility in the United States,” *The Quarterly Journal of Economics*, 129, 1553–1623.
- CLÉMENÇON, S., G. LUGOSI, AND N. VAYATIS (2008): “Ranking and empirical minimization of U-statistics,” *The Annals of Statistics*, 36, 844–874.
- DALTON, H. (1920): “The measurement of the inequality of incomes,” *The Economic Journal*, 30, 348–361.
- DEHEJIA, R. H. (2005): “Program evaluation as a decision problem,” *Journal of Econometrics*, 125, 141–173.
- DONALDSON, D. AND J. A. WEYMARK (1980): “A single-parameter generalization of the Gini indices of inequality,” *Journal of economic Theory*, 22, 67–86.
- (1983): “Ethically flexible Gini indices for income distributions in the continuum,” *Journal of Economic Theory*, 29, 353–358.
- ESCANCIANO, J. C. AND J. R. TERSCHUUR (2022): “Debiased Semiparametric U-Statistics: Machine Learning Inference on Inequality of Opportunity,” *arXiv preprint arXiv:2206.05235*.
- FARRELL, M. H., T. LIANG, AND S. MISRA (2021): “Deep neural networks for estimation and inference,” *Econometrica*, 89, 181–213.
- FLEURBAEY, M. (1995): “Equal opportunity or equal social outcome?” *Economics & Philosophy*, 11, 25–55.

- FORT, M., A. ICHINO, AND G. ZANELLA (2020): “Cognitive and noncognitive costs of day care at age 0–2 for children in advantaged families,” *Journal of Political Economy*, 128, 158–205.
- GARCÍA, J. L., J. J. HECKMAN, D. E. LEAF, AND M. J. PRADOS (2020): “Quantifying the life-cycle benefits of an influential early-childhood program,” *Journal of Political Economy*, 128, 2502–2541.
- GRAY-LOBE, G., P. A. PATHAK, AND C. R. WALTERS (2023): “The long-term effects of universal preschool in Boston,” *The Quarterly Journal of Economics*, 138, 363–411.
- HAUSSLER, D. (1995): “Sphere packing numbers for subsets of the Boolean n-cube with bounded Vapnik-Chervonenkis dimension,” *Journal of Combinatorial Theory, Series A*, 69, 217–232.
- HECKMAN, J., R. PINTO, AND P. SAVELYEV (2013): “Understanding the mechanisms through which an influential early childhood program boosted adult outcomes,” *American Economic Review*, 103, 2052–2086.
- HECKMAN, J. J. AND L. K. RAUT (2016): “Intergenerational long-term effects of preschool-structural estimates from a discrete dynamic programming model,” *Journal of econometrics*, 191, 164–175.
- HIRANO, K. AND J. R. PORTER (2009): “Asymptotics for statistical treatment rules,” *Econometrica*, 77, 1683–1701.
- HUFE, P., A. PEICHL, P. SCHÜLE, AND J. TODOROVIĆ (2022): “Fairness in Europe: A Multidimensional Comparison,” in *CESifo Forum*, München: ifo Institut-Leibniz-Institut für Wirtschaftsforschung an der ..., vol. 23, 45–51.
- KASY, M. (2016): “Partial identification, distributional preferences, and the welfare ranking of policies,” *Review of Economics and Statistics*, 98, 111–131.
- KITAGAWA, T., M. NYBOM, AND J. STUHLER (2018): “Measurement error and rank correlations,” Tech. rep., cemmap working paper.
- KITAGAWA, T. AND A. TETENOV (2018): “Who should be treated? empirical welfare maximization methods for treatment choice,” *Econometrica*, 86, 591–616.
- (2021): “Equality-minded treatment choice,” *Journal of Business & Economic Statistics*, 39, 561–574.
- KOSOROK, M. R. (2008): *Introduction to empirical processes and semiparametric inference.*, Springer.

- KUECK, J., Y. LUO, M. SPINDLER, AND Z. WANG (2023): “Estimation and inference of treatment effects with L2-boosting in high-dimensional settings,” *Journal of Econometrics*, 234, 714–731.
- MANSKI, C. F. (2004): “Statistical treatment rules for heterogeneous populations,” *Econometrica*, 72, 1221–1246.
- MEHRAN, F. (1976): “Linear measures of income inequality,” *Econometrica: Journal of the Econometric Society*, 805–809.
- NEWHEY, W. K. (1994): “The asymptotic variance of semiparametric estimators,” *Econometrica: Journal of the Econometric Society*, 1349–1382.
- RODRÍGUEZ, L. J. C., G. A. MARRERO, J. G. R. HERNÁNDEZ, AND P. S. ROJO (2021): “Inequality of Opportunity in Spain,” *Hacienda Pública Española/Review of Public Economics*, 237, 153–185.
- ROEMER, J. E. (1998): *Equality of opportunity*, Harvard University Press.
- SASAKI, Y. AND T. URA (2021): “Estimation and inference for policy relevant treatment effects,” *Journal of Econometrics*.
- STOYE, J. (2009): “Minimax regret treatment choice with finite samples,” *Journal of Econometrics*, 151, 70–81.
- (2012): “Minimax regret treatment choice with covariates or with limited validity of experiments,” *Journal of Econometrics*, 166, 138–156.
- TETENOV, A. (2012): “Statistical treatment choice based on asymmetric minimax regret criteria,” *Journal of Econometrics*, 166, 157–165.
- VAN DE GAER (1993): “Equality of opportunity and investment in human capital,” Phd thesis, Katholieke Universiteit Leuven.
- WAGER, S. AND G. WALTHER (2015): “Adaptive concentration of regression trees, with application to random forests,” *arXiv preprint arXiv:1503.06388*.
- WAINWRIGHT, M. J. (2019): *High-dimensional statistics: A non-asymptotic viewpoint*, vol. 48, Cambridge university press.
- WEYMARK, J. A. (1981): “Generalized Gini inequality indices,” *Mathematical Social Sciences*, 1, 409–430.

ZHANG, T. AND B. YU (2005): “Boosting with early stopping: Convergence and consistency,”

.

ZHOU, Z., S. ATHEY, AND S. WAGER (2023): “Offline multi-action policy learning: Generalization and optimization,” *Operations Research*, 71, 148–183.