

Maximum Likelihood Estimation

So far we have assumed knowledge of the true population parameters:

$$\vec{\theta} = (c, \phi_1, \phi_2, \dots, \phi_p, \theta_1, \dots, \theta_q \sigma^2)'$$

- In reality, the true parameters are unknown.
- We will estimate them with maximum likelihood

Maximum likelihood has two steps:

1. Express the likelihood

- This is the joint probability density function

$$f_{\vec{y}_T, \dots, \vec{y}_1}(y_T, \dots, y_1 | \vec{\theta}) \text{ where } (y_1, \dots, y_T)' = \vec{y}_T \text{ is a realization of } \vec{y}_T = (\vec{y}_1, \dots, \vec{y}_T)$$

- $f_{\vec{y}_T}$ can be viewed (loosely) as the joint probability of observing \vec{y}_T given $\vec{\theta}$.
- Alternatively,

$$L(\vec{\theta} | y_1, \dots, y_T) = f_{\vec{y}_T, \dots, \vec{y}_1}(y_T, \dots, y_1 | \vec{\theta}) \text{ is viewed as the "likelihood" of } \vec{\theta} \text{ given } \vec{y}_T.$$

2. Finding the value of $\vec{\theta}$ that maximizes $L(\vec{\theta} | \vec{y}_T)$.

Expressing the joint density/likelihood requires making a distributional assumption about ε_t .

- Typically, we'll assume $\varepsilon_t \stackrel{iid}{\sim} N(0, \sigma^2)$.

Exact Likelihood of AR(1):

Recall a Gaussian AR(1) process is:

$$Y_t = c + \phi Y_{t-1} + \varepsilon_t, \quad \varepsilon_t \stackrel{iid}{\sim} N(0, \sigma^2).$$

- In this case $\hat{\theta} = (c, \phi, \sigma^2)$
- Suppose $|\phi| < 1$ (stationary and causal)

We know that the first and second unconditional moments are:

$$E[Y_t] = \frac{c}{1-\phi}, \quad Var(Y_t) = \frac{\sigma^2}{1-\phi^2}, \quad \forall t.$$

Thus, $Y_1 \sim N\left(\frac{c}{1-\phi}, \frac{\sigma^2}{1-\phi^2}\right)$, or

$$f_{Y_1}(y_1 | \vec{\theta}) = \frac{1}{\sqrt{2\pi} \frac{\sigma^2}{1-\phi^2}} \cdot \exp\left\{-\frac{1}{2} \frac{y_1 - \frac{c}{1-\phi}}{\frac{\sigma^2}{1-\phi^2}}\right\}$$

Now, conditional of $Y_1 = y_1$,

$$Y_2 = c + \phi y_1 + \varepsilon_2$$

So, $Y_2 | Y_1 = y_1 \sim N(c + \phi y_1, \sigma^2)$ or

$$f_{Y_2 | Y_1 = y_1}(y_2 | y_1, \vec{\theta}) = \frac{1}{\sqrt{2\pi} \sigma^2} \cdot \exp\left\{-\frac{1}{2} \frac{(y_2 - c - \phi y_1)}{\sigma^2}\right\}$$

The same is true for all $t > 1$

$$f_{Y_t|Y_{t-1}=y_t}(y_t|y_{t-1}, \vec{\theta}) = \frac{1}{\sqrt{2\pi\sigma^2}} \cdot \exp \left\{ -\frac{1}{2} \frac{(y_t - c - \phi y_{t-1})^2}{\sigma^2} \right\}$$

The joint density is the product of marginal + conditionals:

$$f_{\vec{Y}}(\vec{y}|\vec{\theta}) = f_{Y_1|Y_0}(y_1|y_0, \vec{\theta}) f_{Y_2|Y_1}(y_2|y_1, \vec{\theta}) \cdots f_{Y_T|Y_{T-1}}(y_T|y_{T-1}, \vec{\theta})$$

This is analogous to:

$$P(A \cap B \cap C) = P(A|BC)P(B|C)P(C)$$

Bayes Rule

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(A \cap B) = P(A|B)P(B)$$

$$P(\underbrace{A \cap B}_{\text{Bob}} \cap C) = P(\text{Bob} \cap C)$$

$$= P(\text{Bob}|C)P(C)$$

$$= P(A \cap B|C)P(C)$$

$$= P(A|B, C)P(B|C)P(C)$$

$$f_{\vec{Y}}(\vec{y}|\vec{\theta}) = f_{Y_1}(y_1|\vec{\theta}) \prod_{t=2}^T f_{Y_t|Y_{t-1}}(y_t|y_{t-1}, \vec{\theta}) = L(\vec{\theta}|\vec{y})$$

The log likelihood is:

$$\begin{aligned} l(\vec{\theta}|\vec{y}) &= \log(L(\vec{\theta}|\vec{y})) = \log(f_{Y_1}(y_1|\vec{\theta})) + \sum_{t=2}^T \log(f_{Y_t|Y_{t-1}}(y_t|y_{t-1}, \vec{\theta})) \\ &= -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log \left(\frac{\sigma^2}{1 - \phi^2} \right) - \frac{1}{2} \frac{\left(y_1 - \frac{c}{1 - \phi} \right)^2}{\frac{\sigma^2}{1 - \phi^2}} + \sum_{t=2}^T \left(-\frac{1}{2} \log(2\pi) - \frac{1}{2} \log(\sigma^2) - \frac{1}{2} \frac{(y_t - c - \phi y_{t-1})^2}{\sigma^2} \right) \\ &= -\frac{T}{2} \log(2\pi) - \frac{T}{2} \log(\sigma^2) + \frac{1}{2} (1 - \phi^2) - \frac{1}{2} \frac{\left(y_1 - \frac{c}{1 - \phi} \right)^2}{\frac{\sigma^2}{1 - \phi^2}} - \frac{1}{2} \sum_{t=2}^T \frac{(y_t - c - \phi y_{t-1})^2}{\sigma^2} \end{aligned}$$

Expressing in vector form - we can view $\vec{y}' = (y_1, \dots, y_T)'$ as a single observation of a vector RV

$\vec{Y}' = (Y_1, \dots, Y_T)'$ with Gaussian errors $Y \sim MVN(\vec{\mu}, \Omega)$

$$\vec{\mu} = \begin{bmatrix} \mu \\ \vdots \\ \mu \end{bmatrix}_{Tx1} \quad \Omega = \begin{bmatrix} \gamma_0 & \cdots & \gamma_{T-1} \\ \vdots & \ddots & \vdots \\ \gamma_{T-1} & \cdots & \gamma_0 \end{bmatrix} = \frac{\sigma^2}{1 - \phi^2} \begin{bmatrix} 1 & \phi & \cdots & \phi^{T-1} \\ \vdots & \ddots & \ddots & \vdots \\ \phi^{T-1} & \cdots & \cdots & 1 \end{bmatrix}$$

Thus, the joint density of \vec{Y} is:

$$f_{\vec{Y}}(\vec{y}|\vec{\theta}) = (2\pi)^{-T/2} |\Omega|^{1/2} \exp \left\{ -\frac{1}{2} (\vec{y} - \vec{\mu})' \Omega^{-1} (\vec{y} - \vec{\mu}) \right\}$$

$$\implies \log \text{ likelihood is } l(\vec{\theta}|\vec{y}) = -\frac{T}{2} \log(2\pi) + \frac{1}{2} \log(|\Omega^{-1}|) - \frac{1}{2} (\vec{y} - \vec{\mu})' \Omega^{-1} (\vec{y} - \vec{\mu})$$

If we factor $\Omega^{-1} = LL^{-1}$, we can manipulate the vector equation above to show it is equivalent to the scalar equation.

If we differentiate the exact log likelihood w.r.t. μ or θ & set = 0, we will have a non-linear system of equations with $\vec{\theta}$ and \vec{y} as variables. This would require a numerical solution.

Conditional Maximum Likelihood for Gaussian AR(1)

Suppose we neglect the marginal density of Y_1 and only consider the conditional joint density:

$$f_{Y_t, \dots, Y_2 | Y_1}(y_t, \dots, y_2 | y_1, \vec{\theta}) = \prod_{t=2}^T f_{Y_t | Y_{t-1}}(y_t | y_{t-1}, \vec{\theta})$$

We can approximate the likelihood as:

$$L(\vec{\theta} | \vec{y}) \approx f_{Y_t, \dots, Y_2 | Y_1}(y_t, \dots, y_2 | y_1, \vec{\theta})$$

When $\varepsilon_t \stackrel{i.i.d.}{\sim} N(0, \sigma^2)$

$$l(\vec{\theta} | \vec{y}) \approx \log(f_{Y_t, \dots, Y_2 | Y_1}) = -\frac{T-1}{2} \log(2\pi) - \frac{T-1}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=2}^T (y_t - c - \phi y_{t-1})^2$$

Maximizing with respect to (c, ϕ) involves only maximizing:

$$-\sum_{t=2}^T (y_t - c - \phi y_{t-1})^2$$

which is equivalent to minimizing:

$$\sum_{t=2}^T (y_t - c - \phi y_{t-1})^2 \text{ (OLS)}$$

The values of c and ϕ that minimize the equation above are identical to the least squares estimates of the following:

$$y_t = c + \phi y_{t-1} + \varepsilon_t$$

These are:

$$\hat{\beta} = (X^T X)^{-1} X^T y$$

where

$$\beta = \begin{bmatrix} c \\ \phi \end{bmatrix}, \quad X = \begin{bmatrix} 1 & y_{T-1} \\ \vdots & \vdots \\ 1 & y_1 \end{bmatrix} \quad \text{and} \quad \vec{y} = \begin{bmatrix} y_T \\ \vdots \\ y_2 \end{bmatrix}$$

Differentiating the log likelihood with respect to σ^2 and setting equal to zero:

$$\frac{\partial l}{\partial \sigma^2} \Big|_{\hat{\sigma}^2} = \frac{T-1}{2\hat{\sigma}^2} + \frac{1}{2\hat{\sigma}^4} \sum_{t=2}^T (y_t - c - \phi y_{t-1})^2 = 0$$

$$\implies \frac{1}{2\hat{\sigma}^4} \sum_{t=2}^T (y_t - c - \phi y_{t-1})^2 = \frac{T-1}{2\hat{\sigma}^2}$$

$$\implies \sigma^2 = \frac{1}{T-1} \sum_{t=2}^T (y_t - c - \phi y_{t-1})^2 \approx \frac{1}{T-1} \sum_{t=2}^T (y_t - \hat{c} - \hat{\phi} y_{t-1})^2$$

This is the average of the squared regression residuals starting at $t = 2$

Exact Likelihood of Gaussian AR(p)

Now $\vec{\theta} = (c, \phi_1, \dots, \phi_p, \sigma^2)'$. Let:

$$\vec{Y}_p = (Y_1, \dots, Y_p)' \quad \text{and} \quad \vec{Y} = (Y_1, \dots, Y_T)'$$

Where $T > p$. Then $Y_p \sim N(\vec{\mu}, \Omega_p)$

where

$$\vec{\mu}_p = \begin{bmatrix} \mu \\ \vdots \\ \mu \end{bmatrix}_{p \times 1} \quad \text{and} \quad \Omega_p = \begin{bmatrix} \gamma_0 & \cdots & \gamma_{p-1} \\ \vdots & \ddots & \vdots \\ \gamma_{p-1} & \cdots & \gamma_0 \end{bmatrix}$$

Thus, the joint marginal density of \vec{Y}_p is

$$f_{\vec{y}_p}(\vec{y}_p|\vec{\theta}) = (2\pi)^{-\frac{p}{2}} |\Omega^{-1}|^{\frac{1}{2}} \exp \left\{ -\frac{1}{2} (y_p - \mu_p)' \Omega^{-1} (y_p - \mu_p) \right\}$$

Now, given $Y_{t-1} = y_{t-1}, \dots, Y_{t-p} = y_{t-p}$ for $t > p$

$$Y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_p y_{t-p} + \varepsilon_t$$

This means $Y_t|Y_{t-1} = y_{t-1}, \dots, Y_{t-p} = y_{t-p} \sim N(c + \phi_1 y_{t-1} + \dots + \phi_p y_{t-p}, \sigma^2)$.

Thus,

$$f_{Y_t|Y_{t-1}, \dots, Y_{t-p}} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{1}{2\sigma^2} (y_t - c - \phi_1 y_{t-1} + \dots + \phi_p y_{t-p})^2 \right\}$$

The complete joint density is

$$\begin{aligned} f_{\vec{y}_p}(\vec{y}_p|\vec{\theta}) &= \prod_{t=p+1}^T f_{Y_t|Y_{t-1}, \dots, Y_{t-p}}(y_t|y_{t-1}, \dots, y_{t-p}, \vec{\theta}) \\ \implies l(\vec{\theta}|\vec{y}) &= -\frac{T}{2} \log(2\pi) - \frac{T-P}{2} \log(\sigma^2) + \frac{1}{2} \log(|\Omega^{-1}|) - \frac{1}{2} (\vec{y}_p - \vec{\mu}_p)' \Omega^{-1} (\vec{y}_p - \vec{\mu}_p) - \frac{1}{2\sigma^2} \sum_{t=p+1}^T (y_t - c - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p})^2 \end{aligned}$$

This involves inverting a $P \times P$ matrix, Ω .

As with AR(1), maximizing the exact lg likelihood results in a set of nonlinear equations in $\vec{\theta}$ and \vec{y} , which must be solved numerically.

Conditional Likelihood of Gaussian AR(p)

As with the AR(1), we neglect the initial marginal likelihood of \vec{y}_p

$$l(\vec{\theta}|\vec{y}) \approx -\frac{T-P}{2} \log(2\pi) - \frac{T-P}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=p+1}^T (y_t - c - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p})^2$$

Maximizing the conditional log likelihood with respect to c, ϕ_1, \dots, ϕ_p is the same as minimizing

$$\sum_{t=p+1}^T (y_t - c - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p})^2$$

These are identical to the LS estimates of

$$\vec{y} = X\vec{\beta} + \vec{\varepsilon}$$

where

$$\vec{\beta} = \begin{bmatrix} c \\ \phi_1 \\ \vdots \\ \phi_p \end{bmatrix}, \quad X = \begin{bmatrix} 1 & y_{T-1} & y_{T-2} & \dots & y_{T-p} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & y_p & y_{p-1} & \dots & y_1 \end{bmatrix} \quad \text{and} \quad \vec{\varepsilon} = \begin{bmatrix} \varepsilon_T \\ \vdots \\ \varepsilon_{p+1} \end{bmatrix} \quad \vec{y} = \begin{bmatrix} y_T \\ \vdots \\ y_{p+1} \end{bmatrix}$$

$$\hat{\vec{\beta}} = (X^T X)^{-1} X^T \vec{y}$$

Differentiating the log likelihood with respect to σ^2 , we get

$$\frac{\partial l}{\partial \sigma^2} \Big|_{\hat{\sigma}^2} = -\frac{T-P}{2\hat{\sigma}^2} + \frac{1}{2\hat{\sigma}^4} \sum_{t=p+1}^T (y_t - c - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p})^2 = 0$$

$$\implies \sigma^2 = \frac{1}{T-p} \sum_{t=p+1}^T (y_t - c - \phi_1 y_{t-1} - \dots - \phi_p y_{t-p})^2 \approx \frac{1}{T-1} \sum_{t=2}^T (y_t - \hat{c} - \hat{\phi}_1 y_{t-1})^2$$

So assuming a Gaussian model doesn't impact the consistency of our estimates

That is $\hat{\beta}$ is consistent for the linear projection of Y_t on Y_{t-1}, \dots, Y_{t-p} even if ε isn't Gaussian.

If $\hat{\varepsilon}$ is not Gaussian, then $\hat{\beta}$ are the Quasi Maximum Likelihood Estimates because the model is misspecified.

Conditional Gaussian MA(1) Likelihood

Recall the form of a Gaussian MA(1) process

$$Y_t = \mu + \varepsilon_t + \psi\varepsilon_{t-1}, \quad \varepsilon_t \stackrel{i.i.d}{\sim} N(0, \sigma^2)$$

In this case $\vec{\theta} = (\mu, \psi, \sigma^2)'$.

Suppose we know ε_{t-1} , then

$$Y_t | \varepsilon_{t-1} \sim N(\mu + \psi\varepsilon_{t-1}, \sigma^2)$$

$$\Rightarrow f_{Y_t | \varepsilon_{t-1}(y_t | \varepsilon_{t-1}, \vec{\theta})} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{1}{2\sigma^2} (\overbrace{y_t - \mu - \psi\varepsilon_t}^{\varepsilon_t})^2 \right\}$$

Suppose $\varepsilon_0 = 0$. Then $Y_1 | \varepsilon_0 = 0 \sim N(\mu, \sigma^2)$.

With an observation y_1 , we can compute,

$$\varepsilon_1 = y_1 - \mu$$

Thus,

$$f_{Y_2 | Y_1, \varepsilon_0} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{1}{2\sigma^2} (\overbrace{y_2 - \mu - \psi\varepsilon_1}^{\varepsilon_2})^2 \right\}$$

We can continue to iterate

$$\varepsilon_t = y_t - \mu - \psi\varepsilon_{t-1}$$

$$\Rightarrow f_{Y_t | Y_{t-1}, \dots, Y_1, \varepsilon_0} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{1}{2\sigma^2} (\overbrace{y_t - \mu - \psi\varepsilon_{t-1}}^{\varepsilon_t})^2 \right\}$$

The joint density is

$$f_{Y_1 | \varepsilon_0} \prod_{t=2}^T f_{Y_t | Y_{t-1}, \dots, Y_1, \varepsilon_0} = \left(\frac{1}{2\pi\sigma^2} \right)^{\frac{T}{2}} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{t=1}^T \varepsilon_t^2 \right\}$$

The log likelihood is

$$l(\vec{\theta} | \vec{y}) = -\frac{T}{2} \log(2\pi) - \frac{T}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=1}^T \varepsilon_t^2$$

The conditional log likelihood of an MA(1) is:

$$l(\vec{\theta} | \vec{y}) = -\frac{T}{2} \log(2\pi) - \frac{T}{2} \log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=1}^T \varepsilon_t^2$$

where: $\vec{\theta} = (\mu, \psi, \sigma^2)'$.

In this case the conditional log likelihood cannot be solved analytically for μ, ψ, σ^2 .

The rough numerical algorithm would be:

1. Guess values for $\vec{\theta} = (\mu, \psi, \sigma^2)'$
2. Assume $\varepsilon_0 = 0$
3. Iteratively compute ε_t
4. Evaluate the likelihood for $\{\varepsilon_t\}_{t=1}^T$
5. Update $\vec{\theta}$ and return to step 2 until convergence

Using backward recursion, we can express

$$\varepsilon_t = \mu - y_t - \psi\varepsilon_{t-1}$$

$$\begin{aligned}
&= y_t - \mu - \psi(y_{t-1} - \mu - \psi\varepsilon_{t-2}) \\
&= (y_t - \mu) - \psi(y_{t-1} - \mu) + \psi^2\varepsilon_{t-2} \\
&= (y_t - \mu) - \psi(y_{t-1} - \mu) + \psi^2(y_{t-2} - \mu) - \psi^3\varepsilon_{t-3} \\
&= (y_t - \mu) - \psi(y_{t-1} - \mu) + \psi^2(y_{t-2} - \mu) - \dots + (-1)^{t-1}\psi^{t-1}(y_1 - \mu) + \underbrace{(-1)^t\psi^t\varepsilon_0}_{=0}
\end{aligned}$$

If $|\psi| < 1$, the effect of assuming $\varepsilon_0 = 0$ will die out with a large sample size.

- Conditional likelihood will be a good approximation of exact likelihood
- The approximation will be better for smaller $|\psi|$.
- If $|\psi| > 1$, the assumption of ε_0 is explosive as the sample size increases
- We can re-express as and invertible MA(1) and compute the new log likelihood

Gaussian MA(q) Conditional Likelihood

Recall the form of a Gaussian MA(q) is:

$$y_t = \mu + \varepsilon_t + \psi_1\varepsilon_{t-1} + \psi_2\varepsilon_{t-2} + \dots + \psi_q\varepsilon_{t-q}, \quad \varepsilon_t \stackrel{i.i.d.}{\sim} N(0, \sigma^2)$$

Now, $\vec{\theta} = (\mu, \psi_1, \dots, \psi_q, \sigma^2)'$.

We will now assume $\varepsilon_0 = \varepsilon_{-1} = \varepsilon_{-2} = \dots = \varepsilon_{-q+1} = 0$ and iteratively compute $\{\varepsilon_t\}_{t=1}^T$ using \vec{y} .

- The conditional likelihood will have the exact same form as for the MA(1), but with these new $\{\varepsilon_t\}_{t=1}^T$
- It must be computed numerically, in a fashion that is almost identical to the MA(1).
- The conditional likelihood can only be used with the invertible form of the MA(q).

Gaussian ARMA(p,q) Conditional Likelihood

Recall the form of a Gaussian ARMA(p,q):

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_p y_{t-p} + \varepsilon_t + \psi_1 \varepsilon_{t-1} + \dots + \psi_q \varepsilon_{t-q}$$

To form the conditional likelihood, we combine the methods of AR(p) and MA(q)

- Condition on $y_0 = y_{-1} = \dots = y_{-p+1} = \frac{c}{1 - \phi_1 - \phi_2 - \dots - \phi_p}$
- Condition on $\varepsilon_0 = \varepsilon_{-1} = \dots = \varepsilon_{-q+1} = 0$
- Recursively compute $\{\varepsilon_t\}_{t=1}^T$ using $\{y_t\}_{t=1}^T, \{\varepsilon_t\}_{t=-q+1}^0, \{y_t\}_{t=-p+1}^0$
- Form the log likelihood as:

$$l(\vec{\theta}|\vec{y}) = -\frac{T}{2}\log(2\pi) - \frac{T}{2}\log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=1}^T \varepsilon_t^2$$

Alternatively, we could start the recursions at $t = p + 1$ without an initial condition on $\{y_t\}_{t=-p+1}^0$.

- Condition on $\varepsilon_p = \varepsilon_{p-1} = \dots = \varepsilon_{p-q+1} = 0$
- Recursively compute $\{\varepsilon_t\}_{t=p+1}^T$ using past values of $y_t + \varepsilon_t$

- Form the log likelihood as:

$$l(\vec{\theta}|\vec{y}) = -\frac{T-p}{2}\log(2\pi) - \frac{T-p}{2}\log(\sigma^2) - \frac{1}{2\sigma^2} \sum_{t=p+1}^T \varepsilon_t^2$$

- The MA polynomial must be invertible for us to use the conditional log likelihood in estimation
- The Kalman filter can be used to compute the exact likelihood