

NAME OF THE STUDENT:

REGISTRATION NO OF STUDENT:

SLOT:

DATE:

LAB EXPERIMENT 3

RANDOM VARIABLE AND PROBABILITY DISTRIBUTION

AIM:

Conduction random experiments with probability concepts;

QUESTION:

1. Select 20 numbers at random from the set 1 to 100.

Code:

```
2 #q.1
3 print(sample(1:100,20))
4 print(sample(1:100,20,replace=TRUE))
```

Output:

```
> #q.1
> print(sample(1:100,20))
[1] 24 38 9 4 26 3 2 12 1 58 73 84 87 6 13 53 80 32 60 63
> print(sample(1:100,20,replace=TRUE))
[1] 25 21 66 53 47 17 69 35 12 57 88 15 15 24 97 26 57 73 69 26
```

2. Sampling with replacement is suitable for throwing a die.

Code:

```
5 #q.2
6 print(sample(1:6,6,replace=TRUE))
```

Output:

```
> #q.2
> print(sample(1:6,6,replace=TRUE))
[1] 6 5 5 5 3 4
```

3. Roll 2 dice and find the product of the faces values when rolling two dice.

Code:

```
7 #q.3
8 print(as.vector(outer(1:6,1:6))))
```

Output:

```
> #q.3
> print(as.vector(outer(1:6,1:6)))
[1] 1 2 3 4 5 6 2 4 6 8 10 12 3 6 9 12 15 18 4 8 12 16 20 24 5 10 15
[28] 20 25 30 6 12 18 24 30 36
```

4. Combination for nCr (assign values for n and r).

Code:

```
9 #q.4
10 n=12;r=3;
11 print(choose(n,r))|
```

Output:

```
> #q.4
> n=12;r=3;
> print(choose(n,r))
[1] 220
```

5. Permutation (assign values for n and r).

Code:

```
12 #q.5
13 n=10;r=4;
14 print(factorial(n)/factorial(n-r))
```

Output:

```
> #q.5
> n=10;r=4;
> print(factorial(n)/factorial(n-r))
[1] 5040
```

6. To find the binomial coefficient of 8th and 10th terms.

Code:

```
15 #q.6
16 print(choose(10,8))
17 print(choose(10,10))|
```

Output:

```
> #q.6
> print(choose(10,8))
[1] 45
> print(choose(10,10))
[1] 1
```

7. Pascal's triangle.

Code:

```
18 #q.7
19 for(n in 0:7) print(choose(n,0:n))
```

Output:

```
> #q.7
> for(n in 0:7) print(choose(n,0:n))
[1] 1
[1] 1 1
[1] 1 2 1
[1] 1 3 3 1
[1] 1 4 6 4 1
[1] 1 5 10 10 5 1
[1] 1 6 15 20 15 6 1
[1] 1 7 21 35 35 21 7 1
```

8. Tossing '3' coins with library 'prob'.

Code:

```
20 #q.8
21 library(prob)
22 tosscoin(3)
```

Output:

```
> tosscoin(3)
  toss1 toss2 toss3
1      H      H      H
2      T      H      H
3      H      T      H
4      T      T      H
5      H      H      T
6      T      H      T
7      H      T      T
8      T      T      T
```

9. Roll '4' dice with library 'prob'.

Code:

```
23 #q.9
24 rolldie(4)
```

Output:

```
> #q.9
> rolldie(4)
```

	x1	x2	x3	x4
1	1	1	1	1
2	2	1	1	1
3	3	1	1	1
4	4	1	1	1
5	5	1	1	1
6	6	1	1	1
7	1	2	1	1
8	2	2	1	1
9	3	2	1	1
10	4	2	1	1
11	5	2	1	1
12	6	2	1	1
13	1	3	1	1
14	2	3	1	1
15	3	3	1	1
16	4	3	1	1
17	5	3	1	1
18	6	3	1	1
19	1	4	1	1
20	2	4	1	1
21	3	4	1	1
22	4	4	1	1
23	5	4	1	1
24	6	4	1	1
25	1	5	1	1

26	2	5	1	1
27	3	5	1	1
28	4	5	1	1
29	5	5	1	1
30	6	5	1	1
31	1	6	1	1
32	2	6	1	1
33	3	6	1	1
34	4	6	1	1
35	5	6	1	1
36	6	6	1	1
37	1	1	2	1
38	2	1	2	1
39	3	1	2	1
40	4	1	2	1
41	5	1	2	1
42	6	1	2	1
43	1	2	2	1
44	2	2	2	1
45	3	2	2	1
46	4	2	2	1
47	5	2	2	1
48	6	2	2	1
49	1	3	2	1
50	2	3	2	1
51	3	3	2	1
52	4	3	2	1
53	5	3	2	1

54	6	3	2	1
55	1	4	2	1
56	2	4	2	1
57	3	4	2	1
58	4	4	2	1
59	5	4	2	1
60	6	4	2	1
61	1	5	2	1
62	2	5	2	1
63	3	5	2	1
64	4	5	2	1
65	5	5	2	1
66	6	5	2	1
67	1	6	2	1
68	2	6	2	1
69	3	6	2	1
70	4	6	2	1
71	5	6	2	1
72	6	6	2	1
73	1	1	3	1
74	2	1	3	1
75	3	1	3	1
76	4	1	3	1
77	5	1	3	1
78	6	1	3	1
79	1	2	3	1
80	2	2	3	1
81	3	2	3	1

82	4	2	3	1
83	5	2	3	1
84	6	2	3	1
85	1	3	3	1
86	2	3	3	1
87	3	3	3	1
88	4	3	3	1
89	5	3	3	1
90	6	3	3	1
91	1	4	3	1
92	2	4	3	1
93	3	4	3	1
94	4	4	3	1
95	5	4	3	1
96	6	4	3	1
97	1	5	3	1
98	2	5	3	1
99	3	5	3	1
100	4	5	3	1
101	5	5	3	1
102	6	5	3	1
103	1	6	3	1
104	2	6	3	1
105	3	6	3	1
106	4	6	3	1
107	5	6	3	1
108	6	6	3	1
109	1	1	4	1

110	2	1	4	1
111	3	1	4	1
112	4	1	4	1
113	5	1	4	1
114	6	1	4	1
115	1	2	4	1
116	2	2	4	1
117	3	2	4	1
118	4	2	4	1
119	5	2	4	1
120	6	2	4	1
121	1	3	4	1
122	2	3	4	1
123	3	3	4	1
124	4	3	4	1
125	5	3	4	1
126	6	3	4	1
127	1	4	4	1
128	2	4	4	1
129	3	4	4	1
130	4	4	4	1
131	5	4	4	1
132	6	4	4	1
133	1	5	4	1
134	2	5	4	1
135	3	5	4	1
136	4	5	4	1
137	5	5	4	1

138	6	5	4	1
139	1	6	4	1
140	2	6	4	1
141	3	6	4	1
142	4	6	4	1
143	5	6	4	1
144	6	6	4	1
145	1	1	5	1
146	2	1	5	1
147	3	1	5	1
148	4	1	5	1
149	5	1	5	1
150	6	1	5	1
151	1	2	5	1
152	2	2	5	1
153	3	2	5	1
154	4	2	5	1
155	5	2	5	1
156	6	2	5	1
157	1	3	5	1
158	2	3	5	1
159	3	3	5	1
160	4	3	5	1
161	5	3	5	1
162	6	3	5	1
163	1	4	5	1
164	2	4	5	1
165	3	4	5	1

166	4	4	5	1	194	2	3	6	1
167	5	4	5	1	195	3	3	6	1
168	6	4	5	1	196	4	3	6	1
169	1	5	5	1	197	5	3	6	1
170	2	5	5	1	198	6	3	6	1
171	3	5	5	1	199	1	4	6	1
172	4	5	5	1	200	2	4	6	1
173	5	5	5	1	201	3	4	6	1
174	6	5	5	1	202	4	4	6	1
175	1	6	5	1	203	5	4	6	1
176	2	6	5	1	204	6	4	6	1
177	3	6	5	1	205	1	5	6	1
178	4	6	5	1	206	2	5	6	1
179	5	6	5	1	207	3	5	6	1
180	6	6	5	1	208	4	5	6	1
181	1	1	6	1	209	5	5	6	1
182	2	1	6	1	210	6	5	6	1
183	3	1	6	1	211	1	6	6	1
184	4	1	6	1	212	2	6	6	1
185	5	1	6	1	213	3	6	6	1
186	6	1	6	1	214	4	6	6	1
187	1	2	6	1	215	5	6	6	1
188	2	2	6	1	216	6	6	6	1
189	3	2	6	1	217	1	1	1	2
190	4	2	6	1	218	2	1	1	2
191	5	2	6	1	219	3	1	1	2
192	6	2	6	1	220	4	1	1	2
193	1	3	6	1	221	5	1	1	2

222	6	1	1	2
223	1	2	1	2
224	2	2	1	2
225	3	2	1	2
226	4	2	1	2
227	5	2	1	2
228	6	2	1	2
229	1	3	1	2
230	2	3	1	2
231	3	3	1	2
232	4	3	1	2
233	5	3	1	2
234	6	3	1	2
235	1	4	1	2
236	2	4	1	2
237	3	4	1	2
238	4	4	1	2
239	5	4	1	2
240	6	4	1	2
241	1	5	1	2
242	2	5	1	2
243	3	5	1	2
244	4	5	1	2
245	5	5	1	2
246	6	5	1	2
247	1	6	1	2
248	2	6	1	2
249	3	6	1	2
250	4	6	1	2

[reached 'max' / getopt("max.print") -- omitted 1046 rows]

10. To find expectation and variance for Discrete Random Variable.

A discrete random variable X has the following probability distribution:

x	13	18	20	24	27
$P(x)$	0.22	0.25	0.20	0.17	0.16

Compute expectation and variance.

Code:

```
25 #q.10
26 x=c(13,18,20,24,27)
27 px=c(0.22,0.25,0.20,0.17,0.16)
28 ex2=sum(x^2*px)
29 print(ex2)
30 mean=sum(x*px)
31 print(mean)
32 var=ex2-(mean)^2
33 print(var)
```

Output:

```
> #q.10
> x=c(13,18,20,24,27)
> px=c(0.22,0.25,0.20,0.17,0.16)
> ex2=sum(x^2*px)
> print(ex2)
[1] 412.74
> mean=sum(x*px)
> print(mean)
[1] 19.76
> var=ex2-(mean)^2
> print(var)
[1] 22.2824
```