

The power of CGLMP inequalities: completing the resolution of IQOQI Vienna Open Quantum Problem 27

Ansh Mishra^{1,*} and Aryan Senthilkumar²

¹Independent researcher, Cumming, GA, USA

²Independent researcher, Johns Creek, GA, USA

E-mail: ansh.mishra2025@gmail.com (A.M.), aryan.senthilk@gmail.com (A.S.)

*Corresponding author

October 2026

Abstract

Open Quantum Problem 27 of IQOQI Vienna, “The power of CGLMP inequalities”, asks (A) whether every nontrivial facet of the local polytope with two settings and d outcomes per party is of CGLMP type, and (B) to show that the Fourier-type measurements of Durt, Kaszlikowski and Żukowski (DKZ) are necessarily optimal for the CGLMP inequality on a maximally entangled state, that they realize the highest resistance of the violation to noise, and that they give the best Kullback–Leibler discrimination against local realism. Part A was answered negatively by Bancal, Gisin and Pironio in 2010. We settle every clause of Part B as posed. (i) For every d , every local dimension and all projective measurements on maximally entangled states, the CGLMP value is at most the DKZ value, and DKZ is the only maximizer up to local unitaries and an inert ancilla. The proof combines an exact reduction to a clock model, a new strip inequality for a projection and a Hermitian matrix, obtained from a two-variable Bessis–Moussa–Villani positivity theorem with an explicit density, and a cone condition on Clausen kernels certified in interval arithmetic for every d ; it is formalised in Lean 4, completely for $d \leq 20$ and, for $d \geq 21$, up to the certified cone condition. (ii) The noise clause holds for every d for the violation of the CGLMP inequality, with DKZ the unique optimum; in Gill’s literal reading (uniformly random outcomes as noise, violation of any Bell inequality) it fails for every $d \geq 4$ on the maximally entangled state of two d -level systems, by projective measurements of coarse-grained CHSH type whose critical visibility we compute exactly. (iii) The Kullback–Leibler clause fails for every $d \geq 4$: explicit orthonormal-basis measurements reach a statistical strength of at least 0.0703 bits, DKZ at most 0.0688 bits. Open strengthenings are listed.

1 Introduction

1.1 The problem

The inequalities of Collins, Gisin, Linden, Massar and Popescu (CGLMP) [10] are Bell inequalities for two parties who each choose one of two measurement settings and obtain one of d outcomes. They define facets of the local polytope of this scenario [27], they reduce to the inequality of Clauser, Horne, Shimony and Holt (CHSH) [8] for $d = 2$, and they are the standard tool for certifying nonlocality with high-dimensional systems. Open Quantum Problem 27 of IQOQI Vienna, “The power of CGLMP inequalities” [34], proposed by R. Gill in 2006, asks how special these inequalities and their optimal quantum measurements are. It has two parts.

Part A asks to prove that every face of the local polytope that does not already lie in a face of the no-signalling polytope is of CGLMP type: it is the inequality written out in [10], or a lifting of it from fewer outcomes, obtained by fusing outcomes.

Part B starts from the numerical observation [12] that the observables maximizing the CGLMP violation on a maximally entangled state take a very special form, namely computational-basis measurements preceded by a discrete Fourier transform and diagonal unitaries [10]. It asks to prove (i) that this must be so, (ii) that these measurements have the largest resistance of the violation to noise, and (iii) that they discriminate best against classical realism in the Kullback–Leibler sense [38].

The exact wording is on the archived problem page [34] and in the clause-by-clause ledger of the accompanying repository [33] (file `complete-resolution/LEDGER.md`). We label the clauses 27A, 27B(i), 27B(ii) and 27B(iii). The problem goes back to Gill’s paper [15], in which the resistance of a violation to noise is measured by how much of the distribution of “completely random, uniform outcomes” [15, p. 140] can be mixed into the observed one while local realism is still violated. The measurements of Part B were found numerically by Kaszlikowski et al. [22] and described analytically by Durt, Kaszlikowski and Żukowski [12]; we call them the *DKZ measurements*. Each party applies setting-dependent diagonal phases, then a discrete Fourier transform, and measures in the computational basis.

1.2 Previous work

Part A was answered negatively by Bancal, Gisin and Pironio [4]: for $d = 4$ the local polytope has facets that are not of CGLMP type. For 27B(i), the optimality of the DKZ value $I_{\text{ME}}(d)$ of (2.3) is Tsirelson’s bound [7] for $d = 2$; for $d = 3$ a semidefinite relaxation agrees with it to within 10^{-10} [25] (the partial result listed on the problem page); and in earlier work we proved optimality and uniqueness for $3 \leq d \leq 20$ by exact sum-of-squares certificates [30], and for every d for the strategies whose four “link” operators coincide [29]. Over all states the largest CGLMP value is attained by states that are not maximally entangled for $d \geq 3$ [1], and it is known exactly only for $d \leq 8$ [20, 30]. For 27B(ii), the critical visibilities of the DKZ measurements were computed numerically [22, 12] and identified with the CGLMP threshold $2/I_{\text{ME}}(d)$ [10, eq. (23)]; see also [6] ($d = 3$) and [16] ($d \leq 5$, where states of Schmidt rank two are the most noise-robust). Acín, Durt, Gisin and Latorre [1, eq. (14)] showed that the CHSH strategy on a maximally entangled state of Schmidt rank two in $\mathbb{C}^d \otimes \mathbb{C}^d$ has a critical visibility tending to 0, and Baek, Ryu and Lee [3] observed numerically that merging outcomes increases the noise robustness of the optimal CGLMP correlations on Φ_4 and Φ_{16} . For 27B(iii), van Dam, Gill and Grünwald [38] introduced the statistical strength of a nonlocality proof; Acín, Gill and Gisin [2] showed that the best tests need not use maximally entangled states; Zhang [39] found the first counterexample to the clause ($d = 4$, with an exact certificate); and we gave counterexamples for $4 \leq d \leq 9$ [30].

1.3 Results

This paper settles every clause of Part B as originally posed, so that every clause of Problem 27 now has a verdict (Table 1); the verdict on the noise clause depends on its reading.

(i) For every $d \geq 2$, every local dimension D and all projective measurements on Φ_D , $I_d \leq I_{\text{ME}}(d)$, and the DKZ measurements are the only maximizers up to a local unitary $u \otimes \bar{u}$ and an inert auxiliary system (Theorems 1 and 2); for $3 \leq d \leq 8$ this extends to arbitrary POVMs (Proposition 3.13). The proof combines an exact reduction to $4d$ projections on a circle, a new strip inequality for a projection and a Hermitian matrix (Theorem 3), proved through a two-variable positivity theorem of Bessis–Moussa–Villani type with an explicit density (Theorem 4), a noncommutative continuum theorem, and a cone condition on Clausen kernels (Theorem 5) certified in interval arithmetic for every d ; it is formalised in Lean 4, completely for $2 \leq d \leq 20$ and, for $d \geq 21$, up to the cone condition.

(ii) For the violation of the CGLMP inequality, the noise clause holds for every d , with DKZ the unique optimum (Theorem 6). In Gill’s literal reading (uniformly random outcomes as noise, violation of local realism) it fails for every $d \geq 4$ on Φ_d itself: projective measurements of coarse-grained CHSH type, with unused outcomes, have an exactly computed critical visibility below that of DKZ, which is at least $\frac{1}{2}$ (Theorem 7). It holds for $d = 2$, and for $d = 3$ it holds on Φ_3 and fails on Φ_2 (Proposition 4.8).

(iii) The Kullback–Leibler clause fails for every $d \geq 4$, for all three strengths of [38]: explicit orthonormal-basis measurements on Φ_d reach at least 0.0703203 bits, DKZ at most 0.0687803 bits (Theorem 8); for $d = 3$, DKZ is a strict local maximum (Proposition 5.8).

As far as we found, the following are new: Theorems 1 and 2 for $d \geq 21$ (for $3 \leq d \leq 20$ see [30]); Theorem 3; the exact theorem on Gill’s literal noise clause for every $d \geq 4$ and its consequence for Problem 27

Table 1: The clauses of Open Quantum Problem 27 as originally posed [34], and their status. The positive results concern projective measurements (27B(i) also holds for POVMs when $3 \leq d \leq 8$, Proposition 3.13); the counterexamples use projective measurements. Lean: Section 6.2.

Clause	Verdict	Proof; credit
27A: every face of the local polytope not contained in a face of the no-signalling polytope is of CGLMP type, possibly lifted	false: non-CGLMP facets for $d = 4$	Bancal, Gisin and Pironio [4] (prior work)
27B(i): the DKZ measurements are necessarily the optimal ones on maximally entangled states	true for every d: optimal, and unique up to $u \otimes \bar{u}$ and an inert ancilla (any local dimension)	Theorems 1, 2 (this work); Lean-verified for $d \leq 20$, and for $d \geq 21$ given CONE_d ; earlier: $d = 2$ [7], $3 \leq d \leq 20$ [30], commuting strategies [29]
27B(ii), CGLMP reading: resistance of the CGLMP violation to uniform noise	true for every d: DKZ the unique optimum	Theorem 6 (this work); Lean-verified with the same scope
27B(ii), Gill’s literal reading: violation of local realism under uniform outcome noise	false for every $d \geq 4$, on Φ_d itself; true for $d = 2$; $d = 3$: true on Φ_3 , false on Φ_2	Theorem 7, Proposition 4.8 (this work); mechanism [1, eq. (14)]; numerics for $d = 4, 16$ [3]
27B(iii): best Kullback–Leibler discrimination	false for every $d \geq 4$ (all three strengths of [38]); $d = 3$: DKZ a strict local maximum	Theorem 8, Proposition 5.8 (this work); $d = 4$ first by Zhang [39]; $4 \leq d \leq 9$ [30]

(the mechanism is that of [1]); and the failure of the Kullback–Leibler clause for every $d \geq 4$ ($d = 4$ first in [39]). Theorem 4 uses the Fourier-slice mechanism of Heinävaara [17], for which we claim no novelty. Questions beyond the problem as posed remain open (Section 7).

Organisation. Section 3 states and outlines the proof of 27B(i), with complete proofs in the companion paper [31] (see also [32]); Sections 4 and 5 prove the noise and Kullback–Leibler verdicts; Section 6 covers verification and Section 7 open questions.

2 Setting and notation

Behaviours and local models. The settings are $x, y \in \{1, 2\}$ and the outcomes $a, b \in \mathbb{Z}_d = \{0, \dots, d-1\}$. A *behaviour* $p = (p(\cdot, \cdot | x, y))_{x,y}$ is a family of four probability distributions on $\mathbb{Z}_d \times \mathbb{Z}_d$. For $\lambda = (a_1, a_2, b_1, b_2) \in \mathbb{Z}_d^4$ let $\delta_\lambda(a, b | x, y) = [a = a_x][b = b_y]$ ($[\cdot]$ is the Iverson bracket). The *local polytope* is $\mathcal{L}_d = \text{conv}\{\delta_\lambda : \lambda \in \mathbb{Z}_d^4\}$; a behaviour in $\mathcal{L}_d$ is *local*, and a probability distribution on $\mathbb{Z}_d^4$ is a *local model* (local randomness of the parties can be absorbed into it). Relabellings of settings, of the outcomes of each measurement and of the parties, and local post-processings of the outcomes, map $\mathcal{L}_d$ into itself. We write $p_0(a, b | x, y) = d^{-2}$ for the behaviour of completely random, uniform outcomes; it is local.

The CGLMP expression. Write $P(A_x = B_y + k)$ for the probability that $A_x - B_y \equiv k \pmod{d}$ for the settings x, y . The CGLMP expression is

$$I_d = \sum_{k=0}^{\lfloor d/2 \rfloor - 1} \left(1 - \frac{2k}{d-1}\right) \left([P(A_1 = B_1 + k) + P(B_1 = A_2 + k + 1) + P(A_2 = B_2 + k) + P(B_2 = A_1 + k)] \right. \\ \left. - [P(A_1 = B_1 - k - 1) + P(B_1 = A_2 - k) + P(A_2 = B_2 - k - 1) + P(B_2 = A_1 - k - 1)] \right), \quad (2.1)$$

the reading of [10] under which the local bound is derived and the DKZ measurements give $I_{\text{ME}}(d)$ (see [29, Section 2]). Local behaviours satisfy $I_d \leq 2$ [10], and $I_d(p_0) = 0$, since every probability in (2.1) equals $1/d$ under p_0 and each term has four of them with each sign. For $d = 2$, I_2 is the CHSH expression.

Projective strategies on maximally entangled states. Such a strategy consists of a local dimension $D \geq 1$, the state $\Phi_D = D^{-1/2} \sum_{i=1}^D |ii\rangle$, and projection-valued measures $\{P_a^x\}_{a \in \mathbb{Z}_d}$ (Alice) and $\{Q_b^y\}_{b \in \mathbb{Z}_d}$ (Bob) on $\mathbb{C}^D$, zero projections allowed. Its behaviour is $p(a, b|x, y) = \langle \Phi_D | P_a^x \otimes Q_b^y | \Phi_D \rangle = \frac{1}{D} \text{Tr}((P_a^x)^\top Q_b^y)$, the transpose being taken in the Schmidt basis. Local unitaries $u \otimes \bar{u}$ fix Φ_D , and tensoring a strategy with the identity on an inert auxiliary system, on $\Phi_{Dk} = \Phi_D \otimes \Phi_k$, does not change its behaviour.

The DKZ strategy. The DKZ strategy DKZ_d ($D = d$) measures in the bases

$$|k\rangle_{A,x} = d^{-1/2} \sum_j e^{-2\pi i j(k+\alpha_x)/d} |j\rangle, \quad |l\rangle_{B,y} = d^{-1/2} \sum_j e^{2\pi i j(l-\beta_y)/d} |j\rangle, \quad \alpha = (0, \tfrac{1}{2}), \quad \beta = (\tfrac{1}{4}, -\tfrac{1}{4})$$

(the measurements of [10, eqs. (12)–(15)]). Since $|\sum_{j < d} e^{2\pi i j \theta/d}|^2 = \sin^2(\pi\theta)/\sin^2(\pi\theta/d)$,

$$p_{\text{DKZ}}(a, b|x, y) = \frac{1}{d} h_{xy}(b - a), \quad h_{xy}(m) = \frac{1}{2d^2 \sin^2(\pi(m - \delta_{xy})/d)}, \quad (2.2)$$

with $\delta_{xy} = \alpha_x + \beta_y$, that is, $\delta_{11} = \delta_{22} = \frac{1}{4}$, $\delta_{12} = -\frac{1}{4}$, $\delta_{21} = \frac{3}{4}$; each h_{xy} is a probability distribution on $\mathbb{Z}_d$, so p_{DKZ} has uniform marginals. The DKZ value is [12, 10]

$$I_{\text{ME}}(d) = I_d(p_{\text{DKZ}}) = \frac{4}{d(d-1)} \sum_{j=1}^{d-1} (d-j) \sec \frac{\pi j}{2d}, \quad (2.3)$$

for instance $I_{\text{ME}}(2) = 2\sqrt{2}$, $I_{\text{ME}}(3) = (12 + 8\sqrt{3})/9 = 2.8729\dots$, $I_{\text{ME}}(4) = 2.8962\dots$, $I_{\text{ME}}(20) = 2.9547\dots$, and $I_{\text{ME}}(d) \rightarrow 32G/\pi^2 = 2.9698\dots$ as $d \rightarrow \infty$, where G is Catalan's constant (Remark 3.12).

Noise. For a behaviour p and $0 \leq v \leq 1$ put $p_v = vp + (1-v)p_0$ and $v_c(p) = \max\{v \in [0, 1] : p_v \in \mathcal{L}_d\}$, the *critical visibility*; p_v is local exactly for $v \leq v_c(p)$, and a *smaller* v_c means a *higher* resistance to noise. Relabellings fix $\mathcal{L}_d$ and p_0 , hence v_c . For a strategy we write v_c of its behaviour.

Conventions. $\tau = \frac{1}{M} \text{Tr}$ is the normalised trace on $M_M(\mathbb{C})$, X_+ the positive part of a Hermitian X , $\|\cdot\|_F$ the Frobenius norm, and $\text{Tr}_E f(EXE)$ the trace over $\text{ran } E$ of f of the compression of X by a projection E ; $\text{Cl}_2(\theta) = \sum_{n \geq 1} n^{-2} \sin n\theta$ is the Clausen function and $G = \text{Cl}_2(\pi/2)$ Catalan's constant. *Computer-assisted* means: depending on computations in rigorous interval arithmetic with outward rounding, or in exact rational or algebraic arithmetic; floating-point computations only find certificates and never decide a statement.

3 Clause 27B(i): the DKZ measurements are optimal and unique

3.1 Statements

Theorem 1 (optimality; computer-assisted). *Let $d \geq 2$ and $D \geq 1$. For every projective strategy on the maximally entangled state Φ_D , $I_d \leq I_{\text{ME}}(d)$. The DKZ strategy attains $I_{\text{ME}}(d)$.*

Theorem 2 (rigidity; computer-assisted). *Let $d \geq 2$, $D \geq 1$, and suppose that a projective strategy on Φ_D attains $I_d = I_{\text{ME}}(d)$. Then d divides D , and there is a unitary $u: \mathbb{C}^D \rightarrow \mathbb{C}^d \otimes \mathbb{C}^{D/d}$ such that $uP_a^x u^* = P_a^{x, \text{DKZ}} \otimes \mathbf{1}$ and $\bar{u}Q_b^y \bar{u}^* = Q_b^{y, \text{DKZ}} \otimes \mathbf{1}$ for all x, y, a, b . Moreover $(u \otimes \bar{u})\Phi_D = \Phi_d \otimes \Phi_{D/d}$, and u is unique up to $u \mapsto (\mathbf{1}_d \otimes u')u$ with u' unitary. Consequently, if $d \nmid D$, the maximum of I_d over projective strategies on Φ_D is strictly smaller than $I_{\text{ME}}(d)$.*

Theorems 1 and 2 are Theorems A and B of the companion mathematical paper [31], which proves every statement of this section in full. They are computer-assisted only through Theorem 5 below (interval arithmetic); for $2 \leq d \leq 20$ the whole proof, including Theorem 5, is checked in Lean (Section 6.2). They concern projective measurements, our reading of the observables of the problem; for $3 \leq d \leq 8$ they extend to arbitrary POVMs (Proposition 3.13), and for $d \geq 9$ the POVM case is open (Section 7).

The analytic core is an inequality for a projection and a Hermitian matrix of arbitrary size. Let $S = \{z \in \mathbb{C} : 0 < \operatorname{Re} z < 1\}$, and for $\lambda \in \mathbb{R}$, $z = x + iy \in S$ and $u \in \mathbb{R}$ put

$$h_\lambda(x + iy) = -\frac{1}{\pi^2} \operatorname{Im} \operatorname{Li}_2(e^{\pi(y-\lambda)} e^{-i\pi x}), \quad K_x(u) = \frac{\sin \pi x}{2(\cosh \pi u - \cos \pi x)}, \quad (3.1)$$

with Li_2 the dilogarithm (principal branch). K_x is the Poisson kernel of S for its left edge, and h_λ is the harmonic function on S with boundary values $(y - \lambda)_+$ on $\operatorname{Re} z = 0$ and 0 on $\operatorname{Re} z = 1$: $h_\lambda(x + iy) = \int_{\mathbb{R}} (w - \lambda)_+ K_x(y - w) dw$ [31, Lemmas 3.1 and 3.2].

Theorem 3 (strip inequality). *Let B be an orthogonal projection on $\mathbb{C}^M$, $P = \mathbf{1} - B$, and g a Hermitian $M \times M$ matrix. For every $\lambda \in \mathbb{R}$,*

$$\sum_{\nu \in \operatorname{spec}(B+ig)} h_\lambda(\nu) - \operatorname{Tr}[(P(g - \lambda)P)_+] = \int_0^1 \int_{\mathbb{R}} K_{1-\tau}(s - \lambda) \rho(s, \tau) ds d\tau \geq 0, \quad (3.2)$$

where eigenvalues are counted with algebraic multiplicity and $\rho \geq 0$ is the density of Theorem 4. Equality holds for one λ if and only if it holds for all λ , if and only if $Bg = gB$.

If B and g commute, the eigenvalues of $B + ig$ lie on the edges of S and (3.2) is an identity; moving them into the strip by a non-commuting g can only increase the left side.

Theorem 4 (two-variable BMV positivity for projections). *In the situation of Theorem 3 let $\lambda_{\min} \leq \lambda_{\max}$ be the extreme eigenvalues of g , and for $s \in \mathbb{R}$ and $0 < \tau < 1$ let $\xi_1(s, \tau), \dots, \xi_M(s, \tau)$ be the roots, with multiplicity, of the polynomial $\xi \mapsto \det(g - s - \xi(P - \tau))$ of degree M . Put $\rho(s, \tau) = \frac{1}{2\pi} \sum_{i=1}^M |\operatorname{Im} \xi_i(s, \tau)|$ for $0 < \tau < 1$, and $\rho = 0$ otherwise. Then ρ is continuous on $\mathbb{R} \times (0, 1)$, vanishes unless $\lambda_{\min} < s < \lambda_{\max}$, satisfies $\rho(s, \tau) \leq \frac{M}{2\pi} \|g - s\|(\tau(1 - \tau))^{-1/2}$, and for all $(a, t) \in \mathbb{C}^2$*

$$\mathcal{D}(a, t) := \operatorname{Tr} e^{ag - tP} - e^{-t} \operatorname{Tr}_P e^{aPgP} - \operatorname{Tr}_B e^{aBgB} = a^2 \int_0^1 \int_{\mathbb{R}} e^{as - t\tau} \rho(s, \tau) ds d\tau. \quad (3.3)$$

Moreover $\int_{\mathbb{R}} \rho(s, \tau) ds = \|BgP\|_F^2$ for every $\tau \in (0, 1)$.

At $a = 1$, (3.3) says that $t \mapsto \operatorname{Tr} e^{g - tP}$ is the Laplace transform of the positive measure $\operatorname{Tr}_B(e^{BgB})\delta_0 + \operatorname{Tr}_P(e^{PgP})\delta_1 + (\int e^s \rho(s, \tau) ds) d\tau$ on $[0, 1]$: Stahl's theorem [36] (the former Bessis–Moussa–Villani conjecture [5]) for a projection, with an explicit density. Theorem 4 asserts positivity in two variables after the pinched terms are subtracted and the result is divided by a^2 , and neither modification can be omitted. The mechanism of the proof (a Fourier transform of a trace function of a pencil, computed along lines, with the density given by the imaginary parts of the roots of an auxiliary pencil) was introduced by Heinävaara [17] for tracial joint spectral measures, giving a new proof of Stahl's theorem (see also [13, 26]); Theorem 4 is a variant adapted to a projection and an affine chart, which we found independently, and we claim no novelty for the mechanism. We have not found Theorem 3, or an equivalent statement, in the literature; its continuum consequence (Proposition 3.6) contains the circle version of the Stein–Weiss theorem [37] on the conjugate function of an indicator, in the form of Laeng [24].

The last ingredient involves only explicit special functions. Let $N = 4d$, $\Xi(u) = -\frac{N^2}{2\pi^2} \operatorname{Cl}_2(\frac{2\pi u}{N})$ and $\Xi^\sharp(u) = \Xi(u) + \Xi(2d - u)$ for $0 \leq u \leq 2d$, so that $(\Xi^\sharp)''(u) = 2 \csc \frac{\pi u}{2d}$ on $(0, 2d)$. Let $\Delta_d = \{\ell \in \mathbb{R}^d : \ell_r \geq$




Figure 1: Architecture of the proof of Theorems 1 and 2: the exact reduction (Section 3.3), the analytic chain (Sections 3.4–3.6; the density of Theorem 4 is denoted F in the figure and ρ in the text), the certified cone condition (Section 3.7) and the equality case (Section 3.8).

$0, \sum_r \ell_r = d\}$ (the *cell vectors*) and Δ_d^+ its interior ($\ell_r > 0$ for all r). For $\ell \in \Delta_d$, extended d -periodically, let $S_r(m) = \ell_r + \dots + \ell_{r+m-1}$ and

$$u_m^\ell = \Delta_m^2 \left[\frac{1}{d} \sum_{r=0}^{d-1} \Xi^\#(S_r(m)) \right] \quad (1 \leq m \leq d-1), \quad (3.4)$$

where $\Delta_m^2 f = f(m+1) - 2f(m) + f(m-1)$. Let $v \in \mathbb{R}^{d-1}$, $v_m = 2 \csc \frac{\pi m}{2d}$, and $t_s = e_s + e_{d-s}$.

Theorem 5 (the cone condition CONE_d ; computer-assisted). *For every $d \geq 2$ there are a finite positive Borel measure μ on Δ_d with $\mu(\Delta_d^+) = \mu(\Delta_d) > 0$ and numbers $w_s \geq 0$ such that*

$$v = \int_{\Delta_d} u^\ell \mu(d\ell) + \sum_{s=1}^{\lfloor d/2 \rfloor} w_s t_s. \quad (3.5)$$

Theorems 3, 4 and 5 are Theorems C, D and E of [31].

3.2 Architecture of the proof

The proof has six steps (Figure 1). (1) An exact reduction [29] turns every strategy into a d -tuple V of unitaries of order four with $I_d = 4F(V)/(d(d-1))$, and then into $4d$ projections on a circle, for which $F(V) - F_{\text{DKZ}} = \frac{1}{2} \langle v, \delta \rangle$ is a linear form in $d-1$ “net pair counts”. (2) Theorem 4 implies Theorem 3. (3) Through a harmonic function on the disc, Theorem 3 shows that operator-valued fields on a circle, paired with their conjugate function, reach only the classical (commutative) value. (4) Placing the $4d$ projections on cells of lengths $\ell_x \bmod d$ and averaging over rotations gives $\langle u^\ell, \delta \rangle \leq 0$ for every cell vector, so Theorem 5 implies $\langle v, \delta \rangle \leq 0$, which is Theorem 1.

(5) Theorem 5 is certified in three regimes of d . (6) In the equality case every inequality is tight; the equality clause of Theorem 3 and a residue argument force all projections to commute, and the commutative case [29] gives Theorem 2. The continuum problem of step (3) is degenerate (its maximizers include non-commuting fields [31, Remark 4.7]), and the discrete problem differs from it by junction terms of order d . Step (4) avoids estimating these terms: the cell inequalities are exact consequences of the continuum theorem, and the discrete kernel is recovered from them by the cone condition.

3.3 Step 1: the clock model and a linear form

Put $N = 4d$, $w = e^{2\pi i/d}$, $z = e^{2\pi i/N}$, and $\kappa(u) = \cot(\pi u/N)$ for $u \in \mathbb{Z}_N \setminus \{0\}$, $\kappa(0) = 0$. Encode a strategy on Φ_D by $U_x = \sum_a w^a P_a^x$, $U'_y = \sum_b w^b Q_b^y$ and put $R_1 = U_2$, $R_2 = (U'_2)^\top$, $R_3 = U_1$, $R_4 = (U'_1)^\top$; every quadruple of unitaries with $R_i^d = \mathbf{1}$ arises in this way, and a simultaneous conjugation $R_i \mapsto u R_i u^*$ is the local unitary $u \otimes \bar{u}$. For unitaries $V_0, \dots, V_{d-1} \in U(M)$ with $V_k^4 = \mathbf{1}$ let

$$F(V) = \sum_{0 \leq j < k \leq d-1} \operatorname{Re}[h_{k-j} \tau(V_j V_k^*)], \quad h_m = \sec \frac{\pi m}{2d} - i \csc \frac{\pi m}{2d}, \quad (3.6)$$

and $F_{\text{DKZ}} = F(\mathbf{1}, \dots, \mathbf{1}) = \sum_{m=1}^{d-1} (d-m) \sec \frac{\pi m}{2d}$. With the cyclic shift X on $\mathbb{C}^d$ and $W = \sum_k |k\rangle\langle k| \otimes z^k V_k$, the covariant strategy R^V on $\mathbb{C}^d \otimes \mathbb{C}^M$ is $R_1^V = X \otimes \mathbf{1}$, $R_{i+1}^V = W R_i^V W^*$.

Proposition 3.1 (reduction [29, Theorem 2.3]). *$I_d(R^V) = 4F(V)/(d(d-1))$ for every covariant strategy, and every projective strategy R on Φ_D is unitarily equivalent, by a simultaneous conjugation, to a direct summand of some R^V ($M = 4D$ suffices), with $I_d(R) = 4F(V)/(d(d-1))$. Hence Theorem 1 holds if and only if $F(V) \leq F_{\text{DKZ}}$ for all M and all V with $V_k^4 = \mathbf{1}$.*

(V comes from averaging over a cyclic symmetry of order $4d$ and the Stone–von Neumann theorem for $\mathbb{Z}_d$; the DKZ strategy is, up to a local unitary, R^V with $M = 1$, $V \equiv \mathbf{1}$.) Commuting families were treated in [29]:

Proposition 3.2 (commutative sector [29, Theorem B, Corollary C, Proposition 2.4]). *If the V_k commute, then $F(V) \leq F_{\text{DKZ}}$, with equality if and only if every joint eigenvector v satisfies $V_k v = i^{-a_k} v$ with $a_k = c + [k \geq r]$ for some $c \in \mathbb{Z}_4$, $0 \leq r \leq d-1$. A strategy has commuting reduced family if and only if its four link operators coincide, $R_1^n R_2^{-n} = R_2^n R_3^{-n} = R_3^n R_4^{-n} = w^{-n} R_4^n R_1^{-n}$ for all n ; and if such a strategy attains $I_{\text{ME}}(d)$, then $d \mid D$ and $u R_i u^* = R_i^{\text{DKZ}} \otimes \mathbf{1}$ for a unitary u .*

A Q -configuration on $\mathbb{C}^M$ is a family $(Q_x)_{x \in \mathbb{Z}_N}$ of projections such that for each $k \in \{0, \dots, d-1\}$ the four projections Q_{k-da} , $a \in \mathbb{Z}_4$, are pairwise orthogonal with sum $\mathbf{1}$; put $A_x = Q_x$, $B_x = Q_{x+d}$ and $(\operatorname{rot}_j Q)_x = Q_{x-j}$. Taking for Q_{k-da} the spectral projection of V_k^* for the eigenvalue i^a is a bijection between families V and Q -configurations; $V \equiv \mathbf{1}$ gives the window $Q_x = \mathbf{1}_{[0,d)}(x)$. Since each residue class modulo d carries a projective measurement, $\sum_x \ell_x A_x = \sum_x \ell_x B_x = (\sum_{r < d} \ell_r) \mathbf{1}$ for every d -periodic ℓ .

Lemma 3.3 (linear form [29, eq. (13)], [31, Lemmas 2.4, 2.5]). *Let $T(m) = \sum_y \tau(A_{y+m} B_y)$, $\mathcal{N}(m) = T(m) - T(-m)$ and $\delta_m = \mathcal{N}(m) - m$ for $1 \leq m \leq d-1$. Then $F(V) = \frac{1}{2} \sum_{x,y} \kappa(x-y) \tau(A_x B_y) - \frac{d}{2}$, the window has $\mathcal{N}(m) = m$, and*

$$F(V) - F_{\text{DKZ}} = \frac{1}{2} \langle v, \delta \rangle, \quad \langle t_s, \delta \rangle = \mathcal{N}(s) + \mathcal{N}(d-s) - d \leq 0 \quad (1 \leq s \leq d-1).$$

(The first formula follows from $\sec t = \frac{1}{2} [\cot(\frac{t}{2} + \frac{\pi}{4}) - \cot(\frac{t}{2} - \frac{\pi}{4})]$, the second from $\mathcal{N}(2d-m) = \mathcal{N}(m)$, $\mathcal{N}(d) = d$ and $\cot \frac{\pi m}{4d} + \tan \frac{\pi m}{4d} = 2 \csc \frac{\pi m}{2d}$, and the last because $\mathcal{N}(s) + \mathcal{N}(d-s)$ is a sum of d differences of probabilities, each at most 1.) So Theorem 1 is equivalent to

$$\langle v, \delta \rangle \leq 0 \quad \text{for every } Q\text{-configuration on every } \mathbb{C}^M. \quad (3.7)$$

3.4 Step 2: the strip inequality from an explicit positive density

Let $g_d = BgB + PgP$, $K = [\lambda_{\min}, \lambda_{\max}] \times [0, 1]$, and for $s \in \mathbb{R}$, $0 < \tau < 1$ let $p_{s,\tau}(\xi) = \det(g - s - \xi(P - \tau))$ and $q_{s,\tau}(\xi) = \det(g_d - s - \xi(P - \tau))$. The roots of $p_{s,\tau}$ are the eigenvalues of $(P - \tau)^{-1}(g - s)$, which is self-adjoint for an indefinite inner product.

Lemma 3.4 (pencil facts [31, Lemmas 3.3, 3.4]). (1) $p_{s,\tau}$ and $q_{s,\tau}$ have degree M , the same leading coefficient and the same sum of roots; all roots of $q_{s,\tau}$ are real; every non-real root ξ of $p_{s,\tau}$ satisfies $|\xi| \leq \|g - s\|/\sqrt{\tau(1-\tau)}$; and all roots of $p_{s,\tau}$ are real if $s \notin (\lambda_{\min}, \lambda_{\max})$. (2) If $p(\xi) = c \prod_i (\xi - \xi_i)$ and $q(\xi) = c \prod_i (\xi - y_i)$ with $c \neq 0$, all y_i real and $\sum_i \xi_i = \sum_i y_i$, then $\int_{\mathbb{R}} \log |p(\xi)/q(\xi)| d\xi = \pi \sum_i |\operatorname{Im} \xi_i|$.

Outline of the proof of Theorem 4 ([31, Section 3.5]). $\mathcal{D}/a^2$ is entire on $\mathbb{C}^2$ and grows at most like the exponential of $\max_{(s,\tau) \in K} (s \operatorname{Re} a - \tau \operatorname{Re} t)$, so by the Paley–Wiener–Schwartz theorem [18, Theorem 7.3.1] it is the Laplace transform of a distribution Θ supported in K . On the line $t = a\xi$ it is the Laplace transform of

$$U_\xi(w) = \sum_j [(w - \lambda_j(\xi))_+ - (w - \lambda_j^0(\xi))_+],$$

where $\lambda_j(\xi)$ and $\lambda_j^0(\xi)$ are the eigenvalues of $g - \xi P$ and $g_d - \xi P$; the Hilbert transform of U'_ξ is $\frac{1}{\pi} \log |\det(g - \xi P - w)/\det(g_d - \xi P - w)|$, which at $w = s - \xi\tau$ equals $\frac{1}{\pi} \log |p_{s,\tau}(\xi)/q_{s,\tau}(\xi)|$. Fourier inversion one line at a time and Lemma 3.4(2) give $\Theta = \rho$ on $\mathbb{R} \times (0, 1)$; the limits of $\operatorname{Tr} e^{ag - tP}$ as $t \rightarrow \pm\infty$ exclude mass on the edges $\tau \in \{0, 1\}$; and Duhamel's formula, $\frac{1}{2}\partial_a^2 \mathcal{D}(0, t) = \|BgP\|_F^2 (1 - e^{-t})/t$, gives the marginal identity. $\square$

A second proof avoids distributions, Paley–Wiener theory and Fourier inversion in two variables; it is the route formalised in Lean.

Proposition 3.5 (Radon identity [31, Theorem B.1]). *For Hermitian H let $y_1(\tau), \dots, y_M(\tau)$ be the eigenvalues of $(P - \tau)^{-1}H$, $0 < \tau < 1$. Then $\sum_i |\operatorname{Im} y_i(\tau)| \leq M\|H\|/\sqrt{\tau(1-\tau)}$ is continuous in τ , and $\frac{1}{2\pi} \int_0^1 \sum_i |\operatorname{Im} y_i(\tau)| d\tau = \operatorname{Tr} H_+ - \operatorname{Tr}(BHB + PHP)_+$.*

Outline. For $H + c$ with $\operatorname{Im} c > 0$ no root is real and exactly $\operatorname{rank} P$ roots lie in the upper half-plane $\mathbb{C}_+$. The roots obey the complex inviscid Burgers equation $\partial_\tau y = y \partial_c y$; summed over the roots in $\mathbb{C}_+$ through a Cauchy integral, valid also at multiple roots, it gives $\partial_\tau \sum_{\mathbb{C}_+} \operatorname{Log} y = \partial_c \sum_{\mathbb{C}_+} y$. Integrating over τ and evaluating the two ends by a Schur complement identifies $\int_0^1 \sum_{\mathbb{C}_+} y d\tau$, minus the same quantity for $H_d = BHB + PHP$, with $\operatorname{Tr}[(H_d + c) \operatorname{Log}(H_d + c)] - \operatorname{Tr}[(H + c) \operatorname{Log}(H + c)]$ up to an affine function of c , which vanishes (consider $c > \|H\|$). Imaginary parts as c tends to the real axis give the identity. $\square$

Applied to $H = g - \xi P - w$, Proposition 3.5 says that the integrals of ρ along the lines $s = w + \xi\tau$ are the functions $U_\xi(w)$; Tonelli's theorem for real (a, t) and the identity theorem then give (3.3) on $\mathbb{C}^2$ [31, Section B.4]. The same argument with an arbitrary Hermitian matrix in place of P gives a proof of Stahl's theorem without distribution theory or Fourier analysis; we make no priority claim about this by-product [31, Remark B.8].

Outline of the proof of Theorem 3 ([31, Section 3.6]). The eigenvalues $\nu_k = x_k + iy_k$ of $B + ig$ have $x_k \in [0, 1]$, because $\operatorname{Re} \langle v, (B + ig)v \rangle = \langle v, Bv \rangle$. Let ω be the sum of the harmonic measures of S at the ν_k restricted to the left edge (the measures $K_{x_k}(y_k - w) dw$), and σ the eigenvalue counting measure of PgP on $\operatorname{ran} P$; the left side of (3.2) is $\int (w - \lambda)_+ d(\omega - \sigma)$. For real $0 < |a| < \pi$ the Laplace transform of ω is $[\operatorname{Tr} e^{ag + iaP} - \operatorname{Tr} e^{ag - iaP}]/(2i \sin a)$ and that of σ is $\operatorname{Tr}_P e^{aPgP}$; by (3.3) at $t = \mp ia$ and $\int e^{-au} K_{1-\tau}(u) du = \sin(a\tau)/\sin a$, their difference is the Laplace transform of $a^2 V$, $V(w) = \iint K_{1-\tau}(s - w) \rho(s, \tau) ds d\tau$. So $\omega - \sigma$ has zero mass and first moment, and $W(w) = \int (w - u)_+ d(\omega - \sigma)(u)$ has the same Laplace transform as V ; both are continuous and $O(e^{-\pi|w|})$, so $W = V$, which at $w = \lambda$ is (3.2). If the defect vanishes for one λ , then $\rho \equiv 0$, so $\mathcal{D} \equiv 0$ and $\|BgP\|_F = 0$, that is, $[B, g] = 0$. $\square$

Equivalently, Theorem 3 says that σ is dominated by ω in the convex order, a Schur–Horn type majorization for the non-normal matrix $B + ig$. For rank $B \leq 1$ there is a second proof, by an argument-principle count, which also covers rank $P \leq 3$ and hence all $M \leq 5$ [31, Remark 3.10].

3.5 Step 3: a noncommutative continuum theorem

Let $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ with $dm = d\theta/2\pi$. A *projection step field* is a function $B(\theta) = B_j$ on arcs (t_j, t_{j+1}) , $t_0 < \dots < t_{n-1} < t_0 + 2\pi$, with projections B_j on $\mathbb{C}^M$. Its conjugate function $g(\theta) = \text{p.v.} \int_{\mathbb{T}} \cot \frac{\theta - \varphi}{2} B(\varphi) dm(\varphi) = \frac{1}{\pi} \sum_j (B_j - B_{j-1}) \log |\sin \frac{\theta - t_j}{2}|$ is Hermitian and real-analytic on each arc. For $\alpha, \beta \geq 0$ with $\alpha + \beta \leq 1$ let $\Phi_c(\alpha, \beta) = \frac{1}{2\pi^2} [\text{Cl}_2(2\pi\alpha) + \text{Cl}_2(2\pi\beta) + \text{Cl}_2(2\pi(1 - \alpha - \beta))]$, so that $\Phi_c(\frac{1}{4}, \frac{1}{4}) = G/\pi^2$.

Proposition 3.6 (continuum theorem [31, Theorem 4.2]). *Let B be a projection step field on $\mathbb{C}^M$ with conjugate function g , $P = \mathbf{1} - B$, and A a measurable field with $0 \leq A \leq P$. Put $\alpha = \int \tau(A) dm$, $\beta = \int \tau(B) dm$, and let $\beta_1, \dots, \beta_M$ be the eigenvalues of $\int B dm$. For every $\lambda \in \mathbb{R}$,*

$$\lambda\alpha + h_\lambda(\beta) - \int_{\mathbb{T}} \tau(Ag) dm = \left[h_\lambda(\beta) - \frac{1}{M} \sum_i h_\lambda(\beta_i) \right] + \frac{1}{M} \int_{\mathbb{T}} D_\lambda dm + \int_{\mathbb{T}} S_\lambda dm,$$

where $D_\lambda(\theta) \geq 0$ is the left side of (3.2) for the pair $(B(\theta), g(\theta))$, $S_\lambda = \tau[(P(g - \lambda)P)_+] - \tau[A(g - \lambda)] \geq 0$, and the bracket is nonnegative. Moreover $\min_\lambda [\lambda\alpha + h_\lambda(\beta)] = \Phi_c(\alpha, \beta)$ for $\alpha, \beta > 0$, $\alpha + \beta < 1$; at $\alpha = \beta = \frac{1}{4}$ the minimum G/π^2 is attained only at $\lambda^* = \frac{\log 2}{2\pi}$.

Outline. $D_\lambda \geq 0$ is Theorem 3, $S_\lambda \geq 0$ is a bathtub argument, and the bracket is nonnegative by concavity of $x \mapsto h_\lambda(x)$ on $[0, 1]$. The Herglotz function $\mathcal{F}(z) = \int \frac{e^{i\varphi} + z}{e^{i\varphi} - z} B(\varphi) dm(\varphi)$ is holomorphic on the disc with $\mathcal{F}(0) = \int B dm$ and boundary values $B + ig$, and its spectrum lies in the closed strip. As $h_\lambda = \text{Re } \phi_\lambda$ with $\phi_\lambda(z) = \frac{i}{\pi^2} \text{Li}_2(e^{-i\pi z - \pi\lambda})$ holomorphic on S , $z \mapsto \sum_{\nu \in \text{spec } \mathcal{F}(z)} h_\lambda(\nu)$ is harmonic, and its boundary mean equals its value at the centre. The minimum is a Legendre-type identity for the dilogarithm [31, Lemma 4.4]. $\square$

Corollary 3.7 (equality [31, Corollary 4.5]). *If $\int B dm = \frac{1}{4}\mathbf{1}$ and $\int \tau(A) dm = \frac{1}{4}$, then $\frac{G}{\pi^2} - \int \tau(Ag) dm = \frac{1}{M} \int D_{\lambda^*} dm + \int S_{\lambda^*} dm \geq 0$; if equality holds, $[B(\theta), g(\theta)] = 0$ for almost every θ .*

3.6 Step 4: cell embeddings

For $\ell \in \Delta_d$, extended by $\ell_x = \ell_{x \bmod d}$, put $L_0 = 0$, $L_{x+1} = L_x + \ell_x$ and $I_x = [L_x, L_{x+1}) \subseteq \mathbb{R}/N\mathbb{Z}$, so that $S_r(m) = L_{r+m} - L_r$. For $x \neq y$ let $K^\ell(x, y) = \int_{I_x} \int_{I_y} \cot \frac{\pi(\xi - \eta)}{N} d\eta d\xi$, which equals $\Xi(L_{x+1} - L_y) - \Xi(L_x - L_y) - \Xi(L_{x+1} - L_{y+1}) + \Xi(L_x - L_{y+1})$ since $\Xi''(u) = \cot \frac{\pi u}{N}$; let $K^\ell(x, x) = 0$ and $Q^\ell(Q) = \sum_{x,y} K^\ell(x, y) \tau(A_x B_y)$.

Proposition 3.8 (cell inequalities [31, Propositions 5.2, 5.3]). *For every $\ell \in \Delta_d$ and every Q -configuration, $Q^\ell(Q) \leq N^2 G/\pi^2$, with equality for every window, and $\frac{1}{d} \sum_{j=0}^{d-1} [Q^\ell(\text{rot}_j Q) - N^2 G/\pi^2] = \langle u^\ell, \delta \rangle$; hence $\langle u^\ell, \delta \rangle \leq 0$.*

Sketch. Let B and A be the step fields on $\mathbb{T}$ equal to B_x and A_x on the arc $(2\pi L_x/N, 2\pi L_{x+1}/N)$. Then $0 \leq A \leq \mathbf{1} - B$, $\int B dm = \int A dm = \frac{1}{N} \sum_x \ell_x B_x = \frac{1}{4}\mathbf{1}$, and $Q^\ell(Q) = N^2 \int \tau(Ag) dm$, so Corollary 3.7 applies; for a window, A and B are indicators of two adjacent arcs of length d , with value $N^2 \Phi_c(\frac{1}{4}, \frac{1}{4})$. The rotation average depends only on $x - y$, and subtracting the window value gives $\langle u^\ell, \delta \rangle$, with u^ℓ as in (3.4). $\square$

Proposition 3.9 (the cone condition implies optimality). *If $v = \int u^\ell \mu(d\ell) + \sum_s w_s t_s$ with $\mu \geq 0$ and $w_s \geq 0$, then $\langle v, \delta \rangle \leq 0$ for every Q -configuration. Hence Theorem 5 implies Theorem 1.*

Proof. $\langle v, \delta \rangle = \int \langle u^\ell, \delta \rangle \mu(d\ell) + \sum_s w_s \langle t_s, \delta \rangle \leq 0$ by Proposition 3.8 and Lemma 3.3 ($\ell \mapsto u^\ell$ is continuous on the compact set Δ_d); conclude with (3.7). $\square$

For uniform cells, $u^1 = v + j$ with $j_m \geq 0$ made of junction terms, so non-uniform cells are needed. Cells with ℓ/d uniform on the simplex give $\mathbb{E}u_m^\ell = \int_0^d (\Xi^\#)''(s) b_{m,d}(s/d) ds$ ($b_{m,d}$ a Bernstein polynomial), which reproduces exactly the singular part $\frac{4d}{\pi m}$ of v_m and leaves a smooth residual of relative size $O(1/d)$, corrected in Section 3.7.

3.7 Step 5: the cone condition for every d

This step is computer-assisted, in interval arithmetic with outward rounding (mpmath [21]; 160 bits for $d \leq 2000$, 120–240 bits beyond). For a probability measure μ on Δ_d let $P_\mu(m) = \int \frac{1}{d} \sum_r \Xi^\#(S_r(m)) \mu(d\ell)$ ($0 \leq m \leq d$), so that $\int u^\ell d\mu = \Delta^2 P_\mu$, and let P_v solve $\Delta^2 P_v = v$ with the same boundary values $\Xi^\#(0), \Xi^\#(d)$.

Lemma 3.10 (reduction to potentials). *If (i) the antisymmetric parts of P_μ and P_v under $m \mapsto d - m$ agree, and (ii) the symmetric part W of $P_v - P_\mu$ satisfies $\Delta^2 W \geq 0$ on $\{1, \dots, d-1\}$, then (3.5) holds with this μ .*

Proof. $W(0) = W(d) = 0$ and $P_\mu + W = P_v$, so $\int u^\ell d\mu + \Delta^2 W = v$, and $\Delta^2 W$ is symmetric and nonnegative, hence a nonnegative combination of the t_s . $\square$

Regime I: $2 \leq d \leq 200$ ([31, Proposition 6.2]). Here $\mu = \sum_k c_k \delta_{\ell^{(k)}}$ and $w = 0$, with $K_d \leq 6.3 d$ cell vectors $\ell^{(k)} = n^{(k)}/16$ (integers $n_r^{(k)} \geq 1$) and certified weights $c_k > 0$, $d \cdot \min_k c_k \geq 0.0816$. The Clausen values are enclosed by the Bernoulli series with exact Bernoulli numbers and an explicit tail bound. Weights found by floating-point linear programming (HiGHS [19], not part of the verification) are corrected to an exact solution of $\sum_k c_k u^{\ell^{(k)}} = v$ through a certified residual bound ($\|1 - RUU^\top\|_\infty \leq 9.4 \cdot 10^{-34}$ for an approximate inverse R of $UU^\top$, U the matrix of the $u^{\ell^{(k)}}$). The smallest weight is $4.25 \cdot 10^{-4}$ ($d = 198$); all 199 certificates were re-verified, and for $d \leq 20$ they are also evaluated by the Lean kernel in exact rational arithmetic.

Regimes II and III: $d \geq 201$. Here μ is the law of a random cell vector: d times a Dirichlet vector with parameters $1 + \eta_r$, where η is a stationary centred Gaussian vector on $\mathbb{Z}_d$ with $\sum_r \eta_r = 0$, conditioned on $\max_r |\eta_r| < \frac{3}{4}$ (parameters 1 otherwise). The unknown is the variogram $Y(m) = \text{Var}(\eta_r + \dots + \eta_{r+m-1})$, which must have positive cosine coefficients $\hat{a}_k(Y)$ to define a covariance. By Dirichlet aggregation, P_μ becomes an average of Beta expectations of $\Xi^\#$, which reproduce the singular part of P_v exactly (a digamma identity [31, Lemma 6.3]). Condition (i) of Lemma 3.10 then decouples, up to an error $O(d^3 \log d e^{-9/(32Y(1))})$, into one scalar equation per distance $m < d/2$, and a Poincaré–Miranda argument [23] on a box of variograms passes from the decoupled to the exact equations [31, Lemmas 6.4, 6.5]. It remains to certify, on the whole box, the positivity of the $\hat{a}_k$, the sign conditions on the faces, and condition (ii).

For $201 \leq d \leq 2000$ ([31, Proposition 6.6]), one run per d of `verify_cert.py` (one process, 160-bit arithmetic, one code version whose SHA-256 hash is stored in every certificate) encloses the decoupled solution (an exact Gaussian expansion of order 14 with an order-16 remainder) and certifies all conditions on the whole box, with exact factorials, binomial coefficients and Bernoulli numbers, rigorous Euler–Maclaurin enclosures of ζ -values, and every decision a comparison of interval endpoints; `audit_cert_g.py` re-checks all 1800 certificates from the stored exact values (all pass). The smallest margins are $d^2 \min_k \hat{a}_k \geq 0.06319$ (even d) and ≥ 0.12638 (odd d), $d \cdot \min \Delta^2 W \geq 0.660$, a derivative bound 4.0009 times the required value, and a decoupling error at most $2 \cdot 10^{-31.16}$.

For $d \geq 2001$ ([31, Proposition 6.8 and Appendix A]), the construction is treated uniformly in d , with $\varepsilon = 1/d$. An explicit real-analytic model Y_e (the root of a quartic equation with special-function coefficients) approximates the decoupled solution to within $6.6468 \cdot 10^{-4} \varepsilon^2$. A Fejér–Pólya argument reduces the positivity of the $\hat{a}_k$ to lower bounds on fourth differences of Y_e minus an explicit correction, which are certified by interval Taylor models on boxes in $(m/d, \varepsilon)$ with $\varepsilon \in [0, 1/2001]$, each box covering all $d \geq 2001$ at once (at least 7.419 in the outer zone $0.011 \leq m/d \leq 0.501$, at least 6.022431 in the inner zone, and directly for $m = 1, 2$). The result is $\hat{a}_k \geq w_k(0.1186944 \varepsilon^2 - 2\varrho) > 0$ ($w_k \in \{\frac{1}{2}, 1\}$, $\varrho \leq e^{-837}$ the box radius), second differences

of the symmetric slack at least $0.620154 \varepsilon - 4\varrho d^2$, and logarithmic margins above 800 for the tail and fixed-point conditions. A final rigour pass fixed code-level enclosure defects (one had left a sliver of size about 10^{-20} uncovered at $d = 2001$); no conclusion changed.

In all regimes μ is carried by Δ_d^+ (all cells $\geq \frac{1}{16}$ in Regime I; Dirichlet parameters $\geq \frac{1}{4}$ beyond), as Theorem 5 requires. The single command `python audit_alld.py` re-checks all three regimes and prints ALL-D CERTIFIED (every $d \geq 2$). In Lean, CONE_d is used in a finite form (finitely many cell vectors, one with all cells positive); for $d \geq 201$ it follows from the integral form by Carathéodory's theorem, each t_s being a positive multiple of a cell vector supported on two residues [31, Remark 11.1].

3.8 Step 6: proofs of Theorems 1 and 2

Theorem 1 follows from Theorem 5 and Proposition 3.9; the DKZ strategy attains $I_{\text{ME}}(d)$ by Proposition 3.1.

Proposition 3.11 (pointwise commutation forces commutation [31, Lemma 7.2, Proposition 7.3]). *Let $\ell \in \Delta_d^+$, let Q be a Q -configuration, and let B, g be the step field and conjugate function of the proof of Proposition 3.8. If $[B(\theta), g(\theta)] = 0$ for almost every θ , then all Q_x commute.*

Proof. All cells are nonempty, so the endpoints $t_x = 2\pi L_x/N$ are distinct modulo 2π . On the arc of x , $B = B_x$ and g is real-analytic, so $[B_x, g(\theta)]$ and its derivative $\frac{1}{2\pi} \sum_j [B_x, B_j - B_{j-1}] \cot \frac{\theta - t_j}{2}$ vanish identically there. This meromorphic function of θ then vanishes everywhere, and its residue at t_j is $\frac{1}{\pi} [B_x, B_j - B_{j-1}]$; so $j \mapsto [B_x, B_j]$ is constant, equal to $[B_x, B_x] = 0$. The $B_x = Q_{x+d}$ run over all the Q 's. $\square$

Proof of Theorem 2. Let a strategy attain $I_{\text{ME}}(d)$; then $\langle v, \delta \rangle = 0$ for its Q -configuration. By Theorem 5, $0 = \int \langle u^\ell, \delta \rangle \mu(d\ell) + \sum_s w_s \langle t_s, \delta \rangle$ with nonpositive terms, so the continuous function $\ell \mapsto \langle u^\ell, \delta \rangle$ vanishes μ -almost everywhere, and as $\mu(\Delta_d^+) > 0$ some $\ell^* \in \Delta_d^+$ has $\langle u^{\ell^*}, \delta \rangle = 0$. By Proposition 3.8 the d nonpositive numbers $Q^{\ell^*}(\text{rot}_j Q) - N^2 G/\pi^2$ have mean 0, so $Q^{\ell^*}(Q) = N^2 G/\pi^2$; Corollary 3.7 gives $[B(\theta), g(\theta)] = 0$ almost everywhere, and Proposition 3.11 that all Q_x , hence all V_k , commute. Proposition 3.2 gives $d \mid D$ and $u R_i u^* = R_i^{\text{DKZ}} \otimes \mathbf{1}$; spectral projections and transposition give the statements about P_a^x and Q_b^y , and $(u \otimes \bar{u}) \sum_i |ii\rangle = \sum_j |jj\rangle$ for every unitary u . If u_1, u_2 both work, $u_2 u_1^*$ commutes with the $R_i^{\text{DKZ}} \otimes \mathbf{1}$, which generate $M_d(\mathbb{C}) \otimes \mathbf{1}$, so $u_2 u_1^* \in \mathbf{1}_d \otimes U(D/d)$. Finally, for fixed D the maximum of I_d is attained (by compactness), so it equals $I_{\text{ME}}(d)$ only if $d \mid D$. $\square$

Remark 3.12 (a deficit identity, and $d \rightarrow \infty$). The proof writes the deficit $F_{\text{DKZ}} - F(V)$ of every strategy as an explicit positive combination of integrated defects of Theorem 3, bathtub slacks and pair slacks [31, Remark 7.1], which suggests a quantitative form of Theorem 2. As $d \rightarrow \infty$, $I_{\text{ME}}(d) = 32G/\pi^2 - c_1/d + O(d^{-2})$ with $c_1 = 2 + \frac{4}{\pi} - \frac{32G}{\pi^2} = 0.30342 \dots$ [31, Corollary 8.1].

3.9 General measurements for $3 \leq d \leq 8$

The reduction of Section 3.3 uses projections in an essential way, and the exact certificates of [30] for $d \leq 20$ use the group law $R_i^n R_i^m = R_i^{n+m}$, $R_i^d = \mathbf{1}$, so neither covers POVMs. For small d a separate certificate does.

Proposition 3.13 (POVMs; computer-assisted). *Let $3 \leq d \leq 8$ and $D \geq 1$. For all d -outcome POVMs, two for each party, on Φ_D , $I_d \leq I_{\text{ME}}(d)$; and equality forces all four POVMs to be projective, so that Theorem 2 applies.*

Sketch. With E^0, E^2 Alice's POVMs, E^1, E^3 the transposes of Bob's, and $Z_n^g = \sum_a w^{na} E_a^g$, the CGLMP deficit (up to a relabelling of settings and a positive factor) is the normalised trace of an element $S_{\text{op}} - \lambda_d$ of the free $*$ -algebra of the effects, in which only $Z_0^g = \mathbf{1}$ and $(Z_n^g)^* = Z_{-n}^g$ are used (no idempotence). After symmetrisation over a group of order $16d$, an exact identity over $\mathbb{Q}(e^{2\pi i/4d})$ writes it, modulo cyclic rotation and the symmetry relations, as a sum of Hermitian squares of combinations of the words $1, Z_n^g, Z_n^g Z_m^{g\pm 1}$ (level "1+AB" of the

tracial moment relaxation) and doubly-localizing terms $\tau(f^* E f F) \geq 0$ (E an effect of Alice, F one of Bob), with Gram blocks proved positive definite by exact LDL^T factorisation and by interval Cholesky factorisation. In the equality case the kernels of the certificate force $E_c^0 Z_n^0 E_b^1 = w^{nc} E_c^0 E_b^1$ and $E_c^0 Z_n^1 E_b^1 = w^{nb} E_c^0 E_b^1$, so the POVMs are projective. $\square$

4 Clause 27B(ii): resistance to noise

4.1 Three readings

In Gill's paper [15] the noise is the behaviour p_0 of uniformly random outcomes, and the resistance of a violation is measured by how much of p_0 can be admixed while local realism is still violated, that is, by the critical visibility v_c . For strategies on maximally entangled states the clause can be read in three ways: (a) *CGLMP reading*: noise p_0 , nonlocality detected by the CGLMP inequality $I_d \leq 2$; (b) *Gill's literal reading*: noise p_0 , nonlocality against all Bell inequalities, that is, v_c ; (c) *white-noise reading* (a strengthening): noise on the state, $v|\Phi_D\rangle\langle\Phi_D| + (1-v)\mathbf{1}/D^2$, and nonlocality against all Bell inequalities. In (c) the noise contributes the product behaviour $\text{Tr } P_a^x \text{Tr } Q_b^y / D^2$, which equals p_0 if every outcome of every measurement has trace D/d , for example for rank-one measurements on Φ_d . Reading (a) holds for every d (Theorem 6), reading (b) fails for every $d \geq 4$ (Theorem 7), and reading (c) is open for $d \geq 4$ (Section 4.4).

4.2 The CGLMP reading

Theorem 6 (noise resistance of the CGLMP violation). *Let $d \geq 2$, $D \geq 1$, let p be the behaviour of a projective strategy $\mathcal{S}$ on Φ_D , and $p_v = vp + (1-v)p_0$. (1) $I_d(p_v) = vI_d(p)$; hence p_v violates $I_d \leq 2$ if and only if $vI_d(p) > 2$. (2) The CGLMP noise threshold $v_{\text{CGLMP}}(\mathcal{S}) = \inf\{v \in [0, 1] : I_d(p_v) > 2\}$ equals $2/I_d(p)$ if $I_d(p) > 2$ (and $+\infty$ otherwise), and $v_{\text{CGLMP}}(\mathcal{S}) \geq 2/I_{\text{ME}}(d)$. (3) $v_{\text{CGLMP}}(\text{DKZ}_d) = 2/I_{\text{ME}}(d)$, and $v_{\text{CGLMP}}(\mathcal{S}) = 2/I_{\text{ME}}(d)$ if and only if $d \mid D$ and $\mathcal{S}$ is $\text{DKZ}_d \otimes \mathbf{1}_{D/d}$ up to a local unitary $u \otimes \bar{u}$. The same holds for white noise on the state whenever every outcome of every measurement has trace D/d .*

Proof. (1) I_d is linear and $I_d(p_0) = 0$. (2) If $I_d(p) > 2$, the violating $v \in [0, 1]$ form $(2/I_d(p), 1]$, and $I_d(p) \leq I_{\text{ME}}(d)$ by Theorem 1. (3) $I_d(p_{\text{DKZ}}) = I_{\text{ME}}(d)$; conversely $v_{\text{CGLMP}}(\mathcal{S}) = 2/I_{\text{ME}}(d)$ forces $I_d(p) = I_{\text{ME}}(d)$, and Theorem 2 applies, while the behaviour is invariant under $u \otimes \bar{u}$ and inert identities. Under the trace condition the maximally mixed state gives $\text{Tr}(P_a^x) \text{Tr}(Q_b^y) / D^2 = d^{-2}$. $\square$

Theorem 6 is formalised in Lean (file `CgmpNoise.lean`; Table 3): for every behaviour, mixing with uniform noise multiplies the CGLMP value by v (`OQP27.cgmp0f_noisy`); parts (2) and (3) hold for $2 \leq d \leq 20$ with no hypothesis and for every d given CONE_d for $d \geq 21$. It concerns the CGLMP inequality only. For outcomes of unequal traces, white noise on the state produces a local behaviour whose CGLMP value need not vanish, and the theorem says nothing about it (Section 7). For $3 \leq d \leq 8$, Proposition 3.13 extends Theorem 6 to POVMs. For all Bell inequalities the corresponding statement is false for every $d \geq 4$, as we show next.

4.3 Gill's literal reading

Let $A_1 = \sigma_z$, $A_2 = \sigma_x$, $B_1 = (\sigma_z + \sigma_x)/\sqrt{2}$ and $B_2 = (\sigma_z - \sigma_x)/\sqrt{2}$ be the optimal CHSH observables, and $\Pi_{\pm}^x = \frac{1}{2}(\mathbf{1} \pm A_x)$, $\Pi_{\pm}^y = \frac{1}{2}(\mathbf{1} \pm B_y)$, real symmetric rank-one projections on $\mathbb{C}^2$. For $d \geq 3$ define a projective strategy C_d on Φ_d as follows. For even d , identify $\mathbb{C}^d = \mathbb{C}^2 \otimes \mathbb{C}^{d/2}$ by $|i\rangle|j\rangle \mapsto |i\frac{d}{2} + j\rangle$, so that $\Phi_d = \Phi_2 \otimes \Phi_{d/2}$, and let $P_0^x = \Pi_+^x \otimes \mathbf{1}$, $P_1^x = \Pi_-^x \otimes \mathbf{1}$, $Q_0^y = \Pi_+^y \otimes \mathbf{1}$, $Q_1^y = \Pi_-^y \otimes \mathbf{1}$. For odd d , decompose $\mathbb{C}^d = \bigoplus_{k=1}^{(d-1)/2} \text{span}\{|2k-2\rangle, |2k-1\rangle\} \oplus \mathbb{C}|d-1\rangle$ and let $P_0^x = (\bigoplus_k \Pi_+^x) \oplus \mathbf{1}$, $P_1^x = (\bigoplus_k \Pi_-^x) \oplus 0$, and Q_0^y, Q_1^y likewise with $\Pi_{\pm}^y$. In both cases $P_a^x = Q_b^y = 0$ for $a, b \geq 2$: the outcomes $2, \dots, d-1$ never occur. Put

$$v_{\text{even}}(d) = \frac{4(d-1)}{(\sqrt{2}-1)d^2 + 4(d-1)}, \quad v_{\text{odd}}(d) = \frac{4}{(\sqrt{2}-1)d + 4}. \quad (4.1)$$

Theorem 7 (Gill's noise clause; computer-assisted for $4 \leq d \leq 9$). (1) For every $d \geq 3$, $v_c(C_d) = v_{\text{even}}(d)$ if d is even and $v_c(C_d) = v_{\text{odd}}(d)$ if d is odd. (2) For every $d \geq 2$, $\frac{1}{2} \leq v_c(\text{DKZ}_d) \leq 2/I_{\text{ME}}(d)$; for $4 \leq d \leq 9$, $v_c(\text{DKZ}_d) \geq r(d)$, where $r(4), \dots, r(9) = 0.69, 0.687, 0.684, 0.683, 0.682, 0.681$. (3) For every $d \geq 4$, $v_c(C_d) < v_c(\text{DKZ}_d)$. Hence, on Φ_d and on every Φ_{kd} , the DKZ measurements do not realize the highest resistance of the violation of local realism to uniform outcome noise.

Part (1) is proved by hand for every $d \geq 3$; the only computer-assisted step is Proposition 4.7, used for $4 \leq d \leq 9$.

Lemma 4.1 (the behaviour of C_d). Let $s_{xy} = -1$ if $x = y = 2$ and $s_{xy} = 1$ otherwise, $\text{CH}(a, b|x, y) = \frac{1}{4}(1 + (-1)^{a+b}s_{xy}/\sqrt{2})$ for $a, b \in \{0, 1\}$ and $\text{CH} = 0$ otherwise, and $\delta_{00}(a, b|x, y) = [a = b = 0]$. The behaviour of C_d is CH for even d and $\frac{d-1}{d}\text{CH} + \frac{1}{d}\delta_{00}$ for odd d .

Proof. All projections are real symmetric, so the behaviour is $\text{Tr}(P_a^x Q_b^y)/d$. For $e, f = \pm 1$, $\frac{1}{2} \text{Tr}(\frac{1+eA_x}{2} \frac{1+fB_y}{2}) = \frac{1}{4}(1 + \frac{ef}{2} \text{Tr}(A_x B_y))$ and $\frac{1}{2} \text{Tr}(A_x B_y) = s_{xy}/\sqrt{2}$. For even d , $\text{Tr}((\Pi \otimes \mathbf{1})(\Pi' \otimes \mathbf{1}))/d = \frac{1}{2} \text{Tr}(\Pi \Pi')$; for odd d , each of the $(d-1)/2$ blocks contributes $\text{Tr}(\Pi \Pi')/d$ and the last dimension contributes $[a = b = 0]/d$. $\square$

Let $\varsigma(0) = 1$ and $\varsigma(a) = -1$ for $a \neq 0$, $E_{xy}(p) = \sum_{a,b} \varsigma(a)\varsigma(b)p(a, b|x, y)$, and $\mathcal{E}(p) = E_{11} + E_{12} + E_{21} - E_{22}$: the CHSH expression after coarse-graining every measurement into the outcome 0 and the rest.

Lemma 4.2 (a coarse-grained CHSH witness). $\mathcal{E} \leq 2$ on $\mathcal{L}_d$; $\mathcal{E}(p_0) = 2(d-2)^2/d^2$, $\mathcal{E}(\text{CH}) = 2\sqrt{2}$, $\mathcal{E}(\delta_{00}) = 2$. Consequently the noisy behaviour p_v of C_d violates $\mathcal{E} \leq 2$ if and only if $v > v_{\text{even}}(d)$ (d even), respectively $v > v_{\text{odd}}(d)$ (d odd).

Proof. On δ_λ , $\mathcal{E} = \varsigma(a_1)(\varsigma(b_1) + \varsigma(b_2)) + \varsigma(a_2)(\varsigma(b_1) - \varsigma(b_2)) = \pm 2$, and $\mathcal{E}$ is linear. Under p_0 the coarse-grained outcomes are independent with mean $(2-d)/d$, so $E_{xy}(p_0) = (d-2)^2/d^2$; under CH, $E_{xy} = s_{xy}/\sqrt{2}$; under δ_{00} , $E_{xy} = 1$. Since $\mathcal{E}(p) > 2 > \mathcal{E}(p_0)$, $\mathcal{E}(p_v) > 2$ if and only if $v > (2 - \mathcal{E}(p_0))/(\mathcal{E}(p) - \mathcal{E}(p_0))$, where $2 - \mathcal{E}(p_0) = 8(d-1)/d^2$ and $\mathcal{E}(p) - \mathcal{E}(p_0)$ equals $(2\sqrt{2}d^2 - 2(d-2)^2)/d^2$ for $p = \text{CH}$ and $2(d-1)((\sqrt{2}-1)d+4)/d^2$ for $p = \frac{d-1}{d}\text{CH} + \frac{1}{d}\delta_{00}$; the quotients are $v_{\text{even}}(d)$ and $v_{\text{odd}}(d)$. $\square$

The witness is optimal: at the threshold the behaviour is local, by the following explicit local model (found in the second independent verification, Section 6.1), which needs no list of facets.

Proposition 4.3 (a local model at the threshold). Let $d \geq 3$, $s = 1/d$ and $T = 4s(1-s)/(\sqrt{2} - (1-2s)^2)$. Then $T = v_{\text{even}}(d)$, and $T\text{CH} + (1-T)p_0$ is local.

Proof. Multiplying the numerator and denominator of T by d^2 gives $v_{\text{even}}(d)$. As $(1-2s)^2 + 4s(1-s) = 1$, $1-T = (\sqrt{2}-1)/(\sqrt{2}-(1-2s)^2) \in (0, 1)$. Put $c_1 = (1-T)(1-2s)$, $c_2 = (1-T)(1-2s)^2$ and $c_0 = 1-4c_1+3c_2$; then $c_2 \geq 0$, $c_1 - c_2 = 2s(1-T)(1-2s) \geq 0$, and $c_0 = 4s[(1-s) - (\sqrt{2}-1)(1-3s)]/(\sqrt{2}-(1-2s)^2) > 0$, because $(\sqrt{2}-1)(1-3s) < 1-3s \leq 1-s$. Let $\Lambda = (\Lambda_{A_1}, \Lambda_{A_2}, \Lambda_{B_1}, \Lambda_{B_2}) \in \{0, 1, *\}^4$ be random with the law: (i) with probability c_2 , $\Lambda = (*, *, *, *)$; (ii) for each $r \in \{0, 1\}$, with probability $\frac{1}{2}(c_1 - c_2)$ each, Λ is one of $(*, r, r, 1-r)$, $(r, *, r, r)$, $(r, 1-r, *, r)$, $(r, r, r, *)$; (iii) with probability $\frac{1}{8}c_0$ each, Λ is one of the eight $\lambda \in \{0, 1\}^4$ with $\sum_{x,y} s_{xy}(-1)^{\lambda_{A_x} + \lambda_{B_y}} = 2$. The probabilities sum to $c_2 + 4(c_1 - c_2) + c_0 = 1$. Alice outputs Λ_{A_x} if it lies in $\{0, 1\}$ and otherwise an independent uniformly random element of $\{2, \dots, d-1\}$; Bob does the same with Λ_{B_y} . We show that the behaviour q of this local model is $T\text{CH} + (1-T)p_0$; fix (x, y) and recall $1-2s = (d-2)/d$. If $a, b \geq 2$, only (i) contributes: $q(a, b|x, y) = c_2/(d-2)^2 = (1-T)/d^2$. If $a \geq 2 > b$, only the pattern of (ii) with $*$ at A_x , and the value of r that puts b at B_y , contributes: $q = \frac{1}{2}(c_1 - c_2)/(d-2) = (1-T)/d^2$; likewise for $b \geq 2 > a$. Let $a, b \in \{0, 1\}$ and $\epsilon = (-1)^{a+b}s_{xy}$. The two patterns of (ii) whose $*$ is at neither A_x nor B_y satisfy $(-1)^{\Lambda_{A_x} + \Lambda_{B_y}} = s_{xy}$, and their two values of r give the two pairs with $\epsilon = 1$. Each strategy in (iii) has $(-1)^{\Lambda_{A_x} + \Lambda_{B_y}} = s_{xy}$ for exactly three of the four setting pairs, each pair being the exception for exactly two of

the eight strategies, and the eight are invariant under $\lambda \mapsto 1 - \lambda$; so six of them give the pairs with $\epsilon = 1$ (three each) and two give those with $\epsilon = -1$ (one each). Hence

$$q(a, b|x, y) = \frac{1}{4}(1 - 2c_1 + c_2) + \frac{1}{8}\epsilon(1 - c_2), \quad TCH(a, b|x, y) + \frac{1-T}{d^2} = \frac{1}{4}(T + 4(1 - T)s^2) + \frac{T}{4\sqrt{2}}\epsilon.$$

The constants agree, as $1 - 2c_1 + c_2 = 1 - (1 - T)(1 - 2s)(1 + 2s) = T + 4(1 - T)s^2$, and so do the other terms, as $1 - c_2 = \sqrt{2}T$ is equivalent to the definition of T . $\square$

Proof of Theorem 7(1). By Lemmas 4.1 and 4.2, $v_c(C_d) \leq v_{\text{even}}(d)$ (d even) and $v_c(C_d) \leq v_{\text{odd}}(d)$ (d odd). For even d , Proposition 4.3 gives equality. For odd d and $s = 1/d$, $v p_{C_d} + (1 - v)p_0 = v s \delta_{00} + (1 - v s)(v' CH + (1 - v')p_0)$ with $v' = v(1 - s)/(1 - v s)$; at $v = v_{\text{odd}}(d) = 4s/(\sqrt{2} - 1 + 4s)$ one has $1 - v s = (\sqrt{2} - (1 - 2s)^2)/(\sqrt{2} - 1 + 4s)$, hence $v' = T$, and the behaviour is a mixture of the deterministic δ_{00} and the local behaviour of Proposition 4.3. $\square$

For the DKZ strategy we need lower bounds on v_c , that is, local models. The first is universal.

Lemma 4.4 (the $\frac{1}{2}$ -lemma). *If all marginals of p are uniform, $\sum_b p(a, b|x, y) = \sum_b p(b, a|x, y) = 1/d$, then p_v is local for $0 \leq v \leq \frac{1}{2}$. The constant is optimal: $\text{PR}_d(a, b|x, y) = \frac{1}{d}[b - a \equiv [x = 2][y = 1] \pmod{d}]$, a d -outcome version of the Popescu–Rohrlich box [35], has uniform marginals and $v_c(\text{PR}_d) = \frac{1}{2}$.*

Proof. $P_1(a_1, a_2, b_1, b_2) = p(a_1, b_1|1, 1)p(a_2, b_2|2, 2)$ and $P_2(a_1, a_2, b_1, b_2) = p(a_1, b_2|1, 2)p(a_2, b_1|2, 1)$ are local models. Under P_1 the setting pairs (1, 1) and (2, 2) have the laws of p , while for (1, 2) and (2, 1) the outcomes are independent and uniform, with law p_0 ; under P_2 the roles are exchanged. So $\frac{1}{2}(P_1 + P_2)$ is a local model of $\frac{1}{2}(p + p_0)$, and $p_v = 2v \cdot \frac{1}{2}(p + p_0) + (1 - 2v)p_0$ is local for $v \leq \frac{1}{2}$. For PR_d the four probabilities with $k = 0$ in the first bracket of (2.1) equal 1 and all others vanish, so $I_d(\text{PR}_d) = 4$, and I_d detects $(\text{PR}_d)_v$ for every $v > \frac{1}{2}$. $\square$

So $v_c(\text{DKZ}_d) \geq \frac{1}{2}$. By (2.2), p_{DKZ} depends on the outcomes only through $b - a$, and so do the following local models.

Lemma 4.5 (covariant local models). *Let μ be a probability distribution on $M_d = \{m \in \mathbb{Z}_d^{\{1,2\}^2} : m_{11} - m_{12} - m_{21} + m_{22} \equiv 0\}$ with marginals μ_{xy} . Then $p(a, b|x, y) = \frac{1}{d}\mu_{xy}(b - a)$ is local.*

Proof. Draw $m \sim \mu$ and an independent uniform $c \in \mathbb{Z}_d$, and output $A_1 = c, A_2 = c + m_{11} - m_{21}, B_1 = c + m_{11}, B_2 = c + m_{12}$. Then $B_y - A_x = m_{xy}$ (for (2, 2) by the constraint), and each A_x is uniform and independent of m . $\square$

Lemma 4.6 (absorption). *Let p have uniform marginals, Q be a local behaviour with uniform marginals, and $0 < v_0 < v_1 \leq 1$. If $|Q(a, b|x, y) - p_{v_1}(a, b|x, y)| \leq (v_1 - v_0)/(2d^2 v_0)$ for all a, b, x, y , then p_{v_0} is local.*

Proof. With $E = p_{v_1} - Q$ (whose marginals vanish) and $\gamma = v_0/(v_1 - v_0)$, $p_{v_0} = \frac{v_0}{v_1}p_{v_1} + (1 - \frac{v_0}{v_1})p_0 = \frac{v_0}{v_1}Q + (1 - \frac{v_0}{v_1}) \cdot \frac{1}{2}(p_0 + S)$ with $S = p_0 + 2\gamma E$. The hypothesis gives $S \geq 0$, so S is a behaviour with uniform marginals, and $\frac{1}{2}(p_0 + S)$ is local by Lemma 4.4. $\square$

For $p = p_{\text{DKZ}}$ and $Q = \frac{1}{d}\mu_{xy}(b - a)$, the condition of Lemma 4.6 reads

$$|\mu_{xy}(m) - v_1 h_{xy}(m) - (1 - v_1)/d| \leq (v_1 - v_0)/(2dv_0) \quad \text{for all } x, y \text{ and } m \in \mathbb{Z}_d. \quad (4.2)$$

Proposition 4.7 (computer-assisted). *For every $3 \leq d \leq 20$ there are rationals $v_0(d) < v_1(d)$ and a probability distribution μ on M_d with rational weights (common denominator 10^{16}) satisfying (4.2). Hence $v_c(\text{DKZ}_d) \geq v_0(d)$, with $2/I_{\text{ME}}(d) - 1.02 \cdot 10^{-7} < v_0(d) < 2/I_{\text{ME}}(d)$; in particular $v_c(\text{DKZ}_d) \geq r(d)$ for $4 \leq d \leq 9$ (Table 2).*

Table 2: Critical visibilities for Gill’s literal noise clause: $v_c(C_d)$ is exact (Theorem 7(1)), and $v_0(d) \leq v_c(\text{DKZ}_d) \leq 2/I_{\text{ME}}(d)$ (Proposition 4.7). For $d \geq 10$ the bound $v_c(\text{DKZ}_d) \geq \frac{1}{2}$ suffices. As $d \rightarrow \infty$, $v_c(C_d) \sim 4/((\sqrt{2} - 1)d) \rightarrow 0$ while $2/I_{\text{ME}}(d) \rightarrow \pi^2/(16G) = 0.67344 \dots$

d	$v_c(C_d)$	$r(d)$	$v_0(d)$ (certified)	$2/I_{\text{ME}}(d)$
4	$v_{\text{even}}(4) = 0.644211701564$	0.69	0.690549638999	0.690549739488
5	$v_{\text{odd}}(5) = 0.658862678520$	0.687	0.687156473999	0.687156574416
6	$v_{\text{even}}(6) = 0.572874043197$	0.684	0.684883650999	0.684883751130
7	$v_{\text{odd}}(7) = 0.579752581421$	0.683	0.683255804999	0.683255905411
8	$v_{\text{even}}(8) = 0.513670345675$	0.682	0.682032857999	0.682032958173
9	$v_{\text{odd}}(9) = 0.517603563835$	0.681	0.681080610999	0.681080711104
10	$v_{\text{even}}(10) = 0.464987979410$	–	0.680318219999	0.680318320061
11	$v_{\text{odd}}(11) = 0.467489102303$	–	0.679694094999	0.679694195121

Verification of the certificates. The weights are checked in integer arithmetic and the marginals $\mu_{xy}(m)$ exactly. The numbers $h_{xy}(m)$ are enclosed in rational intervals: π by consecutive partial sums of Machin’s formula (width below 10^{-50}), and $\sin^2(\pi(m - \delta_{xy})/d)$ through $\sin(\pi r)$ with rational $0 \leq r < \frac{1}{2}$ (as $4(m - \delta_{xy})$ is odd), which lies between $\sin(r\pi_-)$ and $\sin(r\pi_+)$ for an enclosure $[\pi_-, \pi_+]$ of π , with $\sin z$ ($0 \leq z \leq 2$ rational) enclosed by consecutive partial sums of its Taylor series, which alternates with decreasing terms as $z^2 < 6$. The largest deviation in (4.2) is then compared exactly with $(v_1 - v_0)/(2dv_0)$: margins about 10^{-13} , enclosure widths about 10^{-15} . The distributions were found by a floating-point linear program (d^3 variables, HiGHS [19]) slightly below its optimum and rounded; this plays no role in the verification. $\square$

Proof of Theorem 7(2) and (3). (2) The lower bound is Lemma 4.4; I_d detects $(p_{\text{DKZ}})_v$ for $v > 2/I_{\text{ME}}(d)$, as $I_d(p_0) = 0$; and the bounds $r(d)$ follow from Proposition 4.7. (3) For even d , $v_{\text{even}}(d) < \frac{1}{2}$ if and only if $(\sqrt{2} - 1)d^2 - 4d + 4 > 0$, which holds at $d = 9$ ($81\sqrt{2} > 113$) and hence for all $d \geq 9$, the derivative $2(\sqrt{2} - 1)d - 4$ being positive for $d \geq 5$; for odd d , $v_{\text{odd}}(d) < \frac{1}{2}$ if and only if $d > 4(\sqrt{2} + 1) = 9.657 \dots$. So $v_c(C_d) < \frac{1}{2} \leq v_c(\text{DKZ}_d)$ for $d \geq 10$. For $4 \leq d \leq 9$, $v_c(C_d) < r(d) \leq v_c(\text{DKZ}_d)$, by exact comparison in $\mathbb{Q}(\sqrt{2})$ (for $d = 8, 9$, $v_c(C_d) > \frac{1}{2}$, which is why certificates are needed up to $d = 9$). Finally $C_d \otimes \mathbf{1}_k$ and $\text{DKZ}_d \otimes \mathbf{1}_k$ on $\Phi_{kd} = \Phi_d \otimes \Phi_k$ have the behaviours of C_d and DKZ_d . $\square$

For $3 \leq d \leq 20$ the certified bounds are within $1.02 \cdot 10^{-7}$ of $2/I_{\text{ME}}(d)$, so to this accuracy the CGLMP inequality is an optimal witness for DKZ_d ; for $d = 3, 4, 5, 6$, $v_c(\text{DKZ}_d) = 2/I_{\text{ME}}(d)$ exactly (computer-assisted, not needed above), by the complete facet lists (66, 216, 1020, 2462 facets) of the polytope of link laws of covariant local models and exact arithmetic in $\mathbb{Q}(\cos \frac{\pi}{4d})$ [31, Remark 9.10].

Proposition 4.8 ($d = 2$ and $d = 3$). (1) For every quantum behaviour with $d = 2$ (any state and measurements), $v_c \geq 1/\sqrt{2} = v_c(\text{DKZ}_2)$. (2) (computer-assisted) For every projective strategy on Φ_3 , $v_c \geq 2/I_{\text{ME}}(3) = 3\sqrt{3} - \frac{9}{2} = 0.696152 \dots$, with equality if and only if the strategy is DKZ_3 up to relabellings and a local unitary $u \otimes \bar{u}$. (3) The strategy on Φ_2 with the three-outcome measurements $\{\Pi_+^x, \Pi_-^x, 0\}$, $\{\Pi_+^y, \Pi_-^y, 0\}$ has $v_c = v_{\text{even}}(3) = 8/(9\sqrt{2} - 1) = 0.682132 \dots < v_c(\text{DKZ}_3)$; the same behaviour arises on every Φ_D with D even, in particular on Φ_6 , which also carries $\text{DKZ}_3 \otimes \mathbf{1}_2$. So the clause holds for $d = 2$; for $d = 3$ it holds on Φ_3 and fails if every maximally entangled state is allowed.

Proof. (1) $\mathcal{L}_2$ is described by positivity and the eight CHSH inequalities [14]; every CHSH expression vanishes on p_0 and is at most $2\sqrt{2}$ on quantum behaviours [7]; and $I_2(\text{DKZ}_2) = 2\sqrt{2}$. (2) $\mathcal{L}_3$ has exactly 1116 facets: 36 positivity facets, 648 liftings of CHSH inequalities (each measurement coarse-grained into one outcome against the other two) and the 432 images of the CGLMP inequality under relabellings [9]; we recomputed this list by an exact double description, verifying every facet. Let $v \leq 2/I_{\text{ME}}(3)$. A facet of CGLMP type is a positive multiple of $2 - I_3(g \cdot p)$ for a relabelling g , which maps projective strategies on Φ_3 to projective strategies on Φ_3 (for

the exchange of parties use $\text{Tr}(X^\top Y) = \text{Tr}(Y^\top X)$ and fixes p_0 , so Theorem 1 gives $I_3(g \cdot p_v) \leq v I_{\text{ME}}(3) \leq 2$. For a lifted CHSH facet, with the Hermitian involutions $\tilde{A}_x, \tilde{B}_y$ of the coarse-grained measurements, the CHSH value of p is, up to relabelling, $\frac{1}{3}[\text{Tr}(\tilde{A}_1^\top(\tilde{B}_1 + \tilde{B}_2)) + \text{Tr}(\tilde{A}_2^\top(\tilde{B}_1 - \tilde{B}_2))] \leq \frac{1}{3}(\|\tilde{B}_1 + \tilde{B}_2\|_1 + \|\tilde{B}_1 - \tilde{B}_2\|_1)$ (Hölder); by Jordan's lemma $\mathbb{C}^3$ splits into invariant subspaces of dimension at most two, contributing at most $4\sqrt{2}$ (dimension two) or 2 (dimension one), so the value is at most $(4\sqrt{2} + 2)/3$, while that of p_0 is $\pm \frac{2}{9}$. Hence no lifted CHSH facet is violated for $v \leq 4/(3\sqrt{2} + 1) = 0.762974\dots$. So $v_c \geq 2/I_{\text{ME}}(3)$, with equality for DKZ_3 by Theorem 7(2). If $v_c = 2/I_{\text{ME}}(3)$, then for every $v > 2/I_{\text{ME}}(3)$ some CGLMP-type facet is violated; one relabelling g works along a sequence $v \downarrow 2/I_{\text{ME}}(3)$, so $I_3(g \cdot p) \geq I_{\text{ME}}(3)$, and Theorem 2 applies. (3) The behaviour is CH with $d = 3$; apply Lemma 4.2 and Proposition 4.3 with $d = 3$. On Φ_D , D even, use $\Pi_\pm^x \otimes \mathbf{1}_{D/2}$ and $\Pi_\pm^y \otimes \mathbf{1}_{D/2}$. $\square$

Remark 4.9 (unused outcomes, and the scope of Theorem 7). C_d never produces the outcomes $2, \dots, d-1$, to which p_0 gives probability $1/d$ each; this is what lets a two-outcome CHSH structure beat DKZ, and $v_c(C_d) \rightarrow 0$. On Φ_d projective measurements cannot avoid this: a projective measurement on $\mathbb{C}^d$ with d occurring outcomes is rank-one, and for rank-one measurements on Φ_d the noise p_0 is white noise on the state, which is reading (c). A projective counterexample on Φ_d using every outcome would therefore settle reading (c). With POVMs every outcome can be used and the clause still fails: for $0 < \varepsilon \leq \frac{1}{100}$ the effects $(1 - \varepsilon)P_a^x + \frac{\varepsilon}{d}\mathbf{1}$ and $(1 - \varepsilon)Q_b^y + \frac{\varepsilon}{d}\mathbf{1}$, with $(P, Q) = C_d$, are invertible and give every pair of outcomes probability at least ε^2/d^2 ; their behaviour arises from that of C_d by the local channel that keeps each outcome with probability $1 - \varepsilon$ and otherwise outputs a uniformly random one, which fixes p_0 ; and the witness $\mathcal{E}$ gives a critical visibility below $v_0(d)$ for $4 \leq d \leq 10$ (for instance 0.6602 for $d = 4$) and below $\frac{1}{2}$ for $d \geq 11$, by exact computations with an explicit formula [31, Remark 9.12]. Projective measurements on Φ_D with $D > d$ that use every outcome also beat DKZ, for instance for $(d, D) = (4, 12)$, where $\text{DKZ}_4 \otimes \mathbf{1}_3$ lives on the same state [31, Remark 9.12].

Remark 4.10 (prior work). The mechanism behind C_d is known: Acín, Durt, Gisin and Latorre [1, eq. (14)] found that the CHSH strategy on a maximally entangled state of Schmidt rank two in $\mathbb{C}^d \otimes \mathbb{C}^d$, with two-outcome measurements and white noise, has critical visibility $v_{\text{even}}(d)$, and for even d , C_d produces the same coarse-grained statistics on Φ_d itself; exactness is elementary in the two-outcome setting, and in the d -outcome scenario it is Proposition 4.3. Baek, Ryu and Lee [3] observed by linear programming that merging outcomes increases the robustness $R_w = (1 - v_c)/v_c$ of the DKZ behaviour, from 0.448 to 0.552 for $d = 4$ and up to 1.7673 for $d = 16$, in agreement with $(I_{\text{ME}}(4) - 2)/2 = 0.44812$, $(1 - v_{\text{even}}(4))/v_{\text{even}}(4) = 0.55229$ and $(1 - v_{\text{even}}(16))/v_{\text{even}}(16) = 1.76731$; so merged-outcome strategies beating DKZ on Φ_4 and Φ_{16} were already visible there, though not related to Problem 27. The thresholds $2/I_{\text{ME}}(d)$ are the numerical values of [22, 12, 10, 16], and [16] found ($d \leq 5$) that states of Schmidt rank two are the most noise-robust. Chen et al. [6] gave an analytic result for $d = 3$; we did not check whether it contains a local model, so the novelty of our lower bound for $d = 3$ is uncertain. As far as we found, the following are new: an exact theorem for every $d \geq 4$ on Φ_d itself, for both parities; the exact critical visibility of C_d in the d -outcome scenario; rigorous lower bounds for $v_c(\text{DKZ}_d)$; and the consequence for Problem 27.

4.4 The white-noise reading

In reading (c) the noise behaviour is the product $\text{Tr } P_a^x \text{Tr } Q_b^y / D^2$ of the marginals, and the question is whether the critical visibility is at least $2/I_{\text{ME}}(d)$ for every projective strategy on every Φ_D , with equality only for DKZ. For strategies on Φ_d that use every outcome it coincides with reading (b) (Remark 4.9), and for them it holds for $d = 2$ and $d = 3$ (Proposition 4.8); for $d \geq 4$ it is open (Section 7). A proof must control all facets of $\mathcal{L}_d$: already for $d = 4$ the polytope of link laws of covariant local behaviours, Masanes' polytope of correlation functions [27], has facets that are not images of the CGLMP inequality.

5 Clause 27B(iii): Kullback–Leibler discrimination

Van Dam, Gill and Grünwald [38] measure the strength of a nonlocality proof with behaviour q by its Kullback–Leibler divergence from the local behaviours. For a distribution σ on the setting pairs let

$$U_q(\sigma) = \inf_{p \in \mathcal{L}_d} \sum_{x,y} \sigma_{xy} D(q(\cdot, \cdot | x, y) \| p(\cdot, \cdot | x, y)), \quad D(r \| s) = \sum_i r_i \log_2 \frac{r_i}{s_i},$$

and $S^{\text{UNI}}(q) = U_q(\frac{1}{4})$, $S^{\text{UC}}(q) = \sup U_q(\sigma)$ over product distributions $\sigma = \sigma_A \otimes \sigma_B$, and $S^{\text{COR}}(q) = \sup_{\sigma} U_q(\sigma)$ (uniform, uncorrelated and correlated settings); so $S^{\text{UNI}} \leq S^{\text{UC}} \leq S^{\text{COR}}$. All values are in bits.

The competitor. Write $d = 4k + j$ with $k \geq 1$, $j \in \{0, 1, 2, 3\}$, and split the computational basis of $\mathbb{C}^d$, and the outcomes, into consecutive groups of sizes $4, \dots, 4$ (k times) and j . On a group of size four, $\mathbb{C}^4 = \mathbb{C}^2 \otimes \mathbb{C}^2$, Alice measures in the product basis $e_{a_1}^x \otimes e_{a_2}^x$, where e_0^x, e_1^x are the real eigenvectors of A_x (Section 4.3) for the eigenvalues $1, -1$, with outcome $2a_1 + a_2$ within the group, and Bob likewise with B_y : two copies of the CHSH strategy with shared settings. On the last group both parties measure the single basis vector ($j = 1$), the eigenbases of A_x and B_y ($j = 2$), or the DKZ bases for three outcomes ($j = 3$). The resulting strategy B_d on Φ_d consists of four orthonormal bases, so every outcome occurs.

Theorem 8 (the Kullback–Leibler clause; computer-assisted). *For every $d \geq 4$ and $X \in \{\text{UNI}, \text{UC}, \text{COR}\}$,*

$$S^X(\text{DKZ}_d) \leq 0.0687802606 < 0.0703203697 \leq S^{\text{UNI}}(B_d) \leq S^X(B_d),$$

and $S^{\text{UNI}}(B_d) \geq 0.0879004622$ if $4 \mid d$. The same holds on Φ_{kd} for $\text{DKZ}_d \otimes \mathbf{1}_k$ and $B_d \otimes \mathbf{1}_k$. Hence for every $d \geq 4$ the DKZ measurements do not give the best discrimination in any of the three senses; the margin is at least 0.00154 bits, and at least 0.0191 bits if $4 \mid d$.

Lemma 5.1 (test factors). *Let $r \geq 0$ be a function of (a, b, x, y) , positive where q is positive, such that $\sum_{x,y} \sigma_{xy} r(a_x, b_y | x, y) \leq 1$ for every $(a_1, a_2, b_1, b_2) \in \mathbb{Z}_d^4$. Then*

$$U_q(\sigma) \geq \sum_{x,y} \sigma_{xy} \sum_{a,b} q(a, b | x, y) \log_2 r(a, b | x, y).$$

Proof. For $p \in \mathcal{L}_d$, $\sum \sigma_{xy} q \log \frac{q}{p} = \sum \sigma_{xy} q \log r - \sum \sigma_{xy} q \log \frac{r p}{q}$ on the support of q , and by Jensen's inequality the last sum is at most $\log \sum \sigma_{xy} r p$; this sum is linear in p and at most 1 on every δ_λ , hence on $\mathcal{L}_d$. $\square$

Lemma 5.2 (symmetries of the DKZ strategy). *The relabellings $q'(a, b | x, y) = q(-a, [y = 1] - b | 3 - x, y)$ and $q'(a, b | x, y) = q(-a - [x = 2], -b | x, 3 - y)$ fix p_{DKZ} , and the group they generate acts transitively on the setting pairs. Hence $S^{\text{UNI}}(\text{DKZ}_d) = S^{\text{UC}}(\text{DKZ}_d) = S^{\text{COR}}(\text{DKZ}_d)$.*

Proof. By (2.2), $p_{\text{DKZ}} = \frac{1}{d} h_{xy}(b - a)$, where h_{xy} , as a function of real m , is even about δ_{xy} and d -periodic. The first relabelling maps $b - a$ to $[y = 1] - (b - a)$ and x to $3 - x$, and $\delta_{3-x,y} + \delta_{xy} = [y = 1]$; the second maps $b - a$ to $[x = 2] - (b - a)$ and y to $3 - y$, and $\delta_{x,3-y} + \delta_{xy} = [x = 2]$. Relabellings preserve $\mathcal{L}_d$, so $U_q(g \cdot \sigma) = U_q(\sigma)$ for g in the group; U_q is concave in σ ; and the group average of σ is uniform. So $U_q(\sigma) \leq U_q(\frac{1}{4})$. $\square$

Lemma 5.3 (direct sums and flagged mixtures). (1) *If every measurement of a strategy on Φ_d is block diagonal for a decomposition of $\mathbb{C}^d$ into coordinate blocks of sizes m_j with disjoint outcome sets O_j , its behaviour is $q = \sum_j \frac{m_j}{d} q_j$, with q_j the behaviour of block j on Φ_{m_j} , supported on $O_j \times O_j$. (2) Let $q = \sum_j \lambda_j q_j$ be such a mixture ($\lambda_j \geq 0$, $\sum_j \lambda_j = 1$), and for each j let $r_j > 0$ be a function on the cells of block j with (F1) $\sum_{x,y} \sigma_{xy} r_j(a_x, b_y | x, y) \leq 1$ for all $(a_1, a_2, b_1, b_2) \in O_j^4$ and (F2) $\sigma_{11} R_j^{11} + \sigma_{22} R_j^{22} \leq 1$ and $\sigma_{12} R_j^{12} + \sigma_{21} R_j^{21} \leq 1$ for all j, k , where $R_j^{xy} = \max_{a,b \in O_j} r_j(a, b | x, y)$. Then $U_q(\sigma) \geq \sum_j \lambda_j \sum_{x,y} \sigma_{xy} \sum_{a,b} q_j \log_2 r_j$.*

Proof. (1) Transposition preserves coordinate blocks, and $\text{Tr}((P_a^x)^\top Q_b^y) = 0$ for outcomes in different blocks. (2) Let $r = r_j$ on the cells of block j and 0 elsewhere. For a deterministic λ , only the setting pairs in $E = \{(x, y) : a_x, b_y \text{ in the same block}\}$ contribute. If they all lie in one block j , the contribution is at most that of a deterministic strategy of block j (replace the other outcomes by elements of O_j), hence at most 1 by (F1). Otherwise E contains (x, y) in block j and (x', y') in block $k \neq j$, which forces $x' \neq x, y' \neq y$ and $E = \{(x, y), (x', y')\} \in \{(1, 1), (2, 2)\}, \{(1, 2), (2, 1)\}$, with contribution at most 1 by (F2). Apply Lemma 5.1. $\square$

Lemma 5.4 (block certificates; computer-assisted). *For the four kinds of blocks of B_d (one basis vector; the CHSH strategy; DKZ₃; two copies of CHSH) there are exact rational test factors satisfying (F1) for uniform σ , checked by exhaustive exact enumeration of all m_j^4 deterministic strategies, with all entries at most 1.24801, so that (F2) holds. Their certified values $L = \frac{1}{4} \sum_{x,y} \sum_{a,b} q \log_2 r$ are $L_1 \geq 0$, $L_2 \geq 0.046273845411$, $L_3 \geq 0.057783024051$ and $L_4 \geq 0.087900462237$ bits.*

Lemma 5.5 (DKZ_d is a wrapped distribution). (1) $h_{xy}^\infty(m) = 1/(2\pi^2(m - \delta_{xy})^2)$, $m \in \mathbb{Z}$, is a probability distribution on $\mathbb{Z}$ that does not depend on d , and h_{xy} is its image under $\mathbb{Z} \rightarrow \mathbb{Z}_d$. (2) For a probability distribution μ of $(t, n_1, n_2) \in \mathbb{Z}^3$ let $\mu_{11}, \mu_{12}, \mu_{21}, \mu_{22}$ be the laws of $n_1, n_2, n_1 - t, n_2 - t$, and $T(\mu) = \frac{1}{4} \sum_{x,y} D(h_{xy}^\infty \| \mu_{xy})$. Then $S^{\text{UNI}}(\text{DKZ}_d) \leq T(\mu)$ for every $d \geq 2$.

Proof. (1) Use $\sum_{n \in \mathbb{Z}} (z+n)^{-2} = \pi^2/\sin^2(\pi z)$ at $z = (m - \delta_{xy})/d$, and at $z = \delta_{xy}$ for the mass, with $\sin^2(\pi \delta_{xy}) = \frac{1}{2}$. (2) With c uniform on $\mathbb{Z}_d$ and independent of (t, n_1, n_2) , the outputs $A_1 = c, A_2 = c+t, B_1 = c+n_1, B_2 = c+n_2$ modulo d form a local model with behaviour $\frac{1}{d} \bar{\mu}_{xy}(b-a)$, $\bar{\mu}_{xy}$ the image of μ_{xy} in $\mathbb{Z}_d$; its divergence from p_{DKZ} on (x, y) is $D(h_{xy}^\infty \| \bar{\mu}_{xy}) \leq D(h_{xy}^\infty \| \mu_{xy})$, by data processing. $\square$

Lemma 5.6 (an explicit model on $\mathbb{Z}^3$; computer-assisted). *There is an explicit probability distribution μ^* on $\mathbb{Z}^3$ with $T(\mu^*) \leq 0.068780260568$ bits: for $|t| \leq 10$ it has rational weights and rational conditional laws of n_1, n_2 on $[-14, 14]$; for $|t| > 10$ the weight of t is $c_*/(t^2 - \frac{1}{4})$ with a rational $c_* = 0.1224482084\dots$, and $n_1 \in \{0, t\}, n_2 \in \{1, t\}$, each with probability $\frac{1}{2}$.*

The certificate. The mass is exactly 1, as $\sum_{t>A} (t^2 - \frac{1}{4})^{-1} = (A + \frac{1}{2})^{-1}$. The link laws are exact rationals for $|m| \leq 24$; beyond, only the tail contributes, $\mu_{xy}^*(m) = c_*/(2(m'^2 - \frac{1}{4}))$ with $m' \in \{m, -m, 1-m\}$. The sum over $|m| \leq 3000$ is evaluated in 40-digit interval arithmetic, and for $|m| > 3000$ every term is nonpositive, as $\mu_{xy}^*(m) \geq h_{xy}^\infty(m)$ there: $(|m| + 1)^2 \leq c_*\pi^2(|m| - \frac{3}{4})^2$ for $|m| \geq 19$, since $c_*\pi^2 > 1.2085$. $\square$

Proof of Theorem 8. By Lemmas 5.2, 5.5 and 5.6, $S^X(\text{DKZ}_d) = S^{\text{UNI}}(\text{DKZ}_d) \leq 0.0687802606$ for every $d \geq 2$. By Lemma 5.3(1), the behaviour of B_d is a mixture, with weights $\frac{4}{d}$ (k times) and $\frac{j}{d}$, of block behaviours with disjoint outcome sets; by Lemmas 5.3(2) and 5.4, $S^{\text{UNI}}(B_d) \geq C(d) := (4kL_4 + jL_j)/d$. As $L_j < L_4$, $C(4k+j) = L_4 - j(L_4 - L_j)/(4k+j)$ is nondecreasing in k , so $C(d) \geq \min\{C(4), \dots, C(7)\} = C(5) = \frac{4}{5}L_4 \geq 0.0703203697$ (and $C(4) = L_4, C(6) \geq 0.0740249, C(7) \geq 0.0749930$). Finally $S^X \geq S^{\text{UNI}}$, and inert identities do not change behaviours. $\square$

The flagged-mixture bound is exact (a numerical solver over all d^4 deterministic strategies reproduces $C(d)$ for $4 \leq d \leq 7$). As B_d consists of orthonormal bases, Theorem 8 also refutes the clause if POVMs or projections of other ranks are allowed; moreover, since every projective measurement on $\mathbb{C}^d$ is a coarse-graining of a basis and coarse-graining cannot increase the strengths, bases are optimal among projective strategies on Φ_d .

Corollary 5.7 (monotonicity and the supremum; computer-assisted). *For $X \in \{\text{UNI}, \text{UC}, \text{COR}\}$ and $k \geq 1$, $S^X(\text{DKZ}_d) \leq S^X(\text{DKZ}_{kd})$, and $0.0687464 \leq \sup_d S^X(\text{DKZ}_d) \leq 0.0687711$.*

Sketch. The infimum defining $U(\frac{1}{4})$ for p_{DKZ} may be restricted to local models invariant under common shifts of all outcomes (by convexity), which are the models of Lemma 5.5(2) with $\mathbb{Z}_{kd}$ in place of $\mathbb{Z}$; reducing modulo d gives the monotonicity. The lower bound is the value at $d = 128$ of an exact rational shift-invariant test factor; the upper bound follows from a two-sided rigorous enclosure of $T(\mu^*)$ including its tail, $T(\mu^*) \in [0.068770973, 0.068771095]$. Both come from the independent verification of Section 6.1. $\square$

Proposition 5.8 ($d = 3$; computer-assisted). *On a neighbourhood of DKZ_3 in the set of projective strategies on Φ_3 , each of the three strengths is at most its value at DKZ_3 , $S(\text{DKZ}_3) = 0.05778302549332865$ bits, with equality only for strategies obtained from DKZ_3 by changing the phases of the basis vectors and by a local unitary $u \otimes \bar{u}$.*

Sketch. For $d = 3$, $I_3 = 4 - \sum_{x,y} \mathbb{E}[g_{xy}]$, where the level $g_{xy} \in \{0, 1, 2\}$ of (a, b) is the residue modulo 3 of $a - b, b - a, b - a - 1, a - b$ for $(x, y) = (1, 1), (1, 2), (2, 1), (2, 2)$. With $\beta = [(16 + 24\sqrt{3}) - \sqrt{3280 - 960\sqrt{3}}]/54$, the test factor $r = 1 + \beta(\frac{1}{2} - g_{xy})$ satisfies Lemma 5.1, and p_{DKZ}/r is a mixture, with positive weights, of the 30 deterministic strategies on the CGLMP facet; so r is optimal and gives the value. Every probability of a CGLMP level is stationary at DKZ_3 (an exact identity in $\mathbb{Q}(\sqrt{3})$). Moving the local model inside the facet so that the first- and second-order terms of the four divergences agree, an interval Cholesky factorisation shows that the averaged Hessian is at most $-\frac{1}{200}$ times the identity on a 16-dimensional slice transversal to the orbit of phases and local unitaries; the ranks of the measurements are locally constant. $\square$

Whether DKZ_3 is a global maximizer is open; more than 300 numerical searches found no strategy on Φ_3 with a larger strength, and numerically DKZ_3 appears to be optimal [38, 2]; $d = 2$ is not treated. The clause was first shown to fail for $d = 4$ by Zhang [39], and for $4 \leq d \leq 9$ in [30]. As far as we found, the additivity of test factors over blocks with disjoint outcomes (Lemma 5.3) and the description of every DKZ_d as the reduction modulo d of one distribution on $\mathbb{Z}$ (Lemma 5.5), which gives a bound uniform in d , are new.

6 Verification

6.1 Independent verification

Each analytic proof was checked by an independent checker, who re-derived every step and wrote the numerical checks from scratch, without importing our programs; the reports and scripts are in the repository [33] (directories `proofs/` and `complete-resolution/`). All eight verifications returned the verdict *correct*: (1) Theorems 4 and 3 (Fourier-slice route), with (3.3) reproduced at 82 complex points to $6.8 \cdot 10^{-13}$ and the defect formula (3.2) at 51 pairs (instance, λ) to $1.3 \cdot 10^{-12}$; (2) the distribution-free route, with the Radon identity confirmed to 34–46 digits; (3) the reduction chain, the continuum theorem, the cell inequalities, the cone logic and the rigidity argument; (4) the second proof of Theorem 3 for rank $B \leq 1$ (no mathematical error; some expository gaps); (5) Theorem 7 (no error), with the facets of $\mathcal{L}_3$ recomputed, all eighteen certificates of Proposition 4.7 re-verified in 200-bit interval arithmetic, and eleven tamper tests on our verifier, all detected; (6) Theorem 7 again, rebuilt from scratch (no gap), with independent certificates and the local model of Proposition 4.3; (7) Theorem 8, Corollary 5.7 and Proposition 5.8 (no error), with $T(\mu^*)$ reproduced digit for digit, the block values re-certified, the local optimality at $d = 3$ re-proved with a different chart and slice (averaged Hessian at most $-\frac{1}{1000}$), and the sharper bounds of Corollary 5.7; (8) Proposition 3.13 (no error, no gap), with the identity re-verified exactly by an independent checker with its own parser, positivity by a third method, the equality case re-derived, and eight tamper tests, all rejected. In addition, Theorems 1 and 2 were proved for $3 \leq d \leq 20$ in [30] by exact sum-of-squares certificates, a method that shares no step with the present one.

6.2 Formal verification in Lean

Theorems 1–4 and 6, and Theorem 5 for $d \leq 20$, are formalised in the Lean 4 theorem prover [11] with Mathlib [28] (Lean 4.33.1, Mathlib revision 0df444a360eaa60ab8c11dca51a86af692955474; directory `lean/` of [33]): 98 files (33,419 lines) in `QJP27/` and three files (898 lines) of algebraic rigidity from our earlier repository [30]. The statement uses concrete matrices: a strategy is a pair of d -outcome projective measurements per party on $\mathbb{C}^D$, its probabilities are $\text{Tr}(A^\top B)/D$, and the CGLMP expression is written out term by term; the definitions are shown to be non-vacuous. No file contains `sorry`, `admit`, an axiom declaration, `native_decide`,

Table 3: Main theorems of the Lean formalisation (files `Main.lean`, `CgImpNoise.lean` and their imports). “ CONE_d , $d \geq 21$ ” is the hypothesis `Hyp_ConeCertPos_large`, the finite form of Theorem 5 for $d \geq 21$ (Section 3.7).

Lean theorem (namespace OQP27)	Statement	Hypotheses
<code>maxEntClause_le_twenty</code>	Theorems 1 and 2 for $2 \leq d \leq 20$; DKZ attains $I_{\text{ME}}(d)$	none
<code>maxEntClause_all</code>	Theorems 1 and 2 for every $d \geq 2$	CONE_d , $d \geq 21$
<code>stripInequality</code> , <code>stripEquality</code> , <code>StripL3b.theorem1_bmv2</code> , <code>StripL3b.hyp_RI</code>	Theorems 3 (with its equality case) and 4 for every matrix size; the Radon identity	none
<code>Red.hyp_reduction</code> , <code>Cell.continuumCell</code> , <code>ClassB.classicalTheoremB_all</code>	the exact reduction to the linear form; the continuum theorem and all cell inequalities; the commutative case for every d	none
<code>coneCertPos_le_twenty</code> , <code>ConeCertificate.fullCheck_sound</code>	CONE_d for $2 \leq d \leq 20$ by kernel-checked certificates; soundness of the checker	none
<code>cglmpOf_noisy</code> , <code>noisy_violation_iff</code> , <code>noisy_violation_imp</code> , <code>noisy_threshold_rigid</code> (and <code>_le_twenty</code> versions)	$I_d(vp + (1 - v)p_0) = vI_d(p)$; Theorem 6	none for $d \leq 20$; CONE_d , $d \geq 21$

`implemented_by`, `extern`, `unsafe` or `opaque`; the 265 audited theorems, among them all main theorems (Table 3), depend only on the standard axioms `propext`, `Classical.choice` and `Quot.sound`; and the certificates for $d \leq 20$ are checked with `decide +kernel`, so that the Lean kernel evaluates them in exact rational arithmetic.

So for $2 \leq d \leq 20$ clause 27B(i) and the CGLMP reading of 27B(ii) are completely machine-checked, and for $d \geq 21$ the formal proof is complete except for one named input, the finite form of CONE_d , which is established outside Lean by the certificates of Section 3.7. Theorems 3 and 4 are formalised in full, through the distribution-free route (the uniqueness step for Theorem 3 uses Mathlib’s Fourier inversion theorem); the continuum theorem is formalised for step fields. Proved on paper only: the uniqueness of u and the strict inequality for $d \nmid D$ in Theorem 2 (the formal statement is that $d \mid D$ and that a local unitary $u \otimes \bar{u}$ maps the strategy to $\text{DKZ}_d \otimes \mathbf{1}$), and auxiliary facts used only in remarks (the continuity of ρ , the marginal identity, the closed formula for h_λ). Proposition 3.13, Theorems 7 and 8 and the propositions of Sections 4 and 5 are not formalised.

6.3 Computer-assisted components and reproducibility

Floating-point computations were used only to find certificates and play no role in any verification. The programs are in [33] and need Python 3 with `numpy`, `scipy`, `sympy` and `mpmath`. (i) Theorem 5 (`cone-certificates/`): `verify_cone.py`, `lp_reverify.py` ($d \leq 200$), `verify_cert.py`, `audit_cert_g.py` ($201 \leq d \leq 2000$), `a01-a14`, `t35_polya.py` ($d \geq 2001$); `python audit_alld.py` re-checks all three regimes. (ii) Section 4 (`complete-resolution/noise-literal/`): `dkz_cert.py`, `facets223.py`, `phi3.py`, `dkz_exact.py` and `povm.py`; `python verify_theorem.py` re-checks them and the exact identities of Section 4.3 (17 checks, about three minutes). (iii) Section 5 (`complete-resolution/kl-divergence/`): `blocks_cert.py`, `zcert.py`, `dkz_lower.py`, `d3_cert.py`, and the independent re-check `verify_main.py`. (iv) Proposition 3.13 (`complete-resolution/povm/`): `verify_povm.py` checks the exact certificates (exact $LDL^\top$, or interval Cholesky). (v) Lean (`lean/`): `lake exe cache get` and `bash check.sh` rebuild every file, scan for forbidden tokens and run the axiom audit (about two hours; up to about 8 GB of memory).

7 Open questions

Problem 27 as posed is settled. Table 4 lists variants and strengthenings that the problem does not ask and that remain open, with what is known; the same list is kept in the ledger of the repository [33].

In the third row the question is whether $\frac{2}{I_{\text{ME}}(d)}I_d(p) + (1 - \frac{2}{I_{\text{ME}}(d)})I_d(p_A \otimes p_B) \leq 2$ for all projective

strategies, $p_A \otimes p_B$ being the product of the marginals (the value 1.98569 refers to this quantity). The certificates of Proposition 3.13 are computed for each d separately. Further questions concern the method: a quantitative version of Theorem 2 (Remark 3.12), and a proof of CONE_d without interval arithmetic, which, once formalised, would complete the Lean proof for $d \geq 21$.

Table 4: Variants and strengthenings not asked by Problem 27, and their status.

Question	Status
27B(i) for arbitrary POVMs on maximally entangled states	proved for $3 \leq d \leq 8$, with uniqueness (Proposition 3.13); open for $d \geq 9$
Noise resistance against all Bell inequalities, white noise on the state, every outcome used (reading (c))	proved for $d = 3$ on Φ_3 (Proposition 4.8); DKZ optimal in all our global numerical searches for $d \leq 8$; open in general, already for rank-one measurements on Φ_d , $d \geq 4$
White noise on the state, CGLMP witness, projective measurements whose outcomes have unequal ranks	open; no counterexample in numerical searches over rank patterns for $d = 3, \dots, 6$ (all patterns enumerated for $d = 3$, $D \leq 9$ and $d = 4$, $D \leq 8$); largest value found $1.98569 < 2$
Kullback–Leibler clause at $d = 3$	DKZ ₃ a strict local maximum (Proposition 5.8); global optimality numerical only
Maximum of I_d over all states: is it $2(\lambda_{\max}(K_d) - 1)/(d - 1)$, $K_d = [\sec \frac{\pi(k-l)}{2d}]_{k,l=0}^{d-1}$?	yes for $d \leq 8$ ($d = 3, 4$ [20]; $5 \leq d \leq 8$ [30]); open for $d \geq 9$

Data and code availability

Certificates, programs, logs, the Lean formalisation, the independent verification reports, the ledger and the companion papers are at <https://github.com/anshM123/anshM123-IQ0QI-Vienna-Open-Problem-27> [33] (MIT license); our earlier exact certificates for $3 \leq d \leq 20$ are at <https://github.com/anshM123/CGLMP> [30].

Acknowledgements

We thank the developers of mpmath, NumPy, SciPy, SymPy, HiGHS, Lean and Mathlib for the software on which the computations, the checks and the formalisation are built.

References

- [1] A. Acín, T. Durt, N. Gisin and J. I. Latorre, *Quantum nonlocality in two three-level systems*, Phys. Rev. A **65** (2002), 052325.
- [2] A. Acín, R. Gill and N. Gisin, *Optimal Bell tests do not require maximally entangled states*, Phys. Rev. Lett. **95** (2005), 210402.
- [3] K. Baek, J. Ryu and J. Lee, *Robustness measures for quantifying nonlocality*, New J. Phys. **27** (2025), 053001; arXiv:2311.07077.
- [4] J.-D. Bancal, N. Gisin and S. Pironio, *Looking for symmetric Bell inequalities*, J. Phys. A **43** (2010), 385303.
- [5] D. Bessis, P. Moussa and M. Villani, *Monotonic converging variational approximations to the functional integrals in quantum statistical mechanics*, J. Math. Phys. **16** (1975), 2318–2325.
- [6] J.-L. Chen, D. Kaszlikowski, L. C. Kwak, C. H. Oh and M. Żukowski, *Entangled three-state systems violate local realism more strongly than qubits: an analytical proof*, Phys. Rev. A **64** (2001), 052109.
- [7] B. S. Cirel’son, *Quantum generalizations of Bell’s inequality*, Lett. Math. Phys. **4** (1980), 93–100.
- [8] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, *Proposed experiment to test local hidden-variable theories*, Phys. Rev. Lett. **23** (1969), 880–884.
- [9] D. Collins and N. Gisin, *A relevant two qubit Bell inequality inequivalent to the CHSH inequality*, J. Phys. A **37** (2004), 1775.
- [10] D. Collins, N. Gisin, N. Linden, S. Massar and S. Popescu, *Bell inequalities for arbitrarily high-dimensional systems*, Phys. Rev. Lett. **88** (2002), 040404.

- [11] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci. **12699**, Springer, Cham, 2021, 625–635.
- [12] T. Durt, D. Kaszlikowski and M. Żukowski, *Violations of local realism with quantum systems described by N -dimensional Hilbert spaces up to $N = 16$* , Phys. Rev. A **64** (2001), 024101.
- [13] A. Eremenko, *Herbert Stahl’s proof of the BMV conjecture*, Sb. Math. **206** (2015), 87–92; arXiv:1312.6003.
- [14] A. Fine, *Hidden variables, joint probability, and the Bell inequalities*, Phys. Rev. Lett. **48** (1982), 291–295.
- [15] R. D. Gill, *Better Bell inequalities (passion at a distance)*, in: Asymptotics: Particles, Processes and Inverse Problems, IMS Lecture Notes–Monograph Series **55** (2007), 135–148; arXiv:math/0610115.
- [16] J. Gruca, W. Laskowski and M. Żukowski, *Nonclassicality of pure two-qutrit entangled states*, Phys. Rev. A **85** (2012), 022118.
- [17] O. Heinävaara, *Tracial joint spectral measures*, arXiv:2310.03227 (2023).
- [18] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, 2nd ed., Springer, Berlin, 1990.
- [19] Q. Huangfu and J. A. J. Hall, *Parallelizing the dual revised simplex method*, Math. Program. Comput. **10** (2018), 119–142.
- [20] M. Ioannou and D. Rosset, *Noncommutative polynomial optimization under symmetry*, arXiv:2112.10803 (2021).
- [21] The mpmath development team, *mpmath: a Python library for arbitrary-precision floating-point arithmetic*, version 1.3.0 (2023), <https://mpmath.org>.
- [22] D. Kaszlikowski, P. Gnaniński, M. Żukowski, W. Miklaszewski and A. Zeilinger, *Violations of local realism by two entangled N -dimensional systems are stronger than for two qubits*, Phys. Rev. Lett. **85** (2000), 4418–4421.
- [23] W. Kulpa, *The Poincaré–Miranda theorem*, Amer. Math. Monthly **104** (1997), 545–550.
- [24] E. Laeng, *On the L^p norms of the Hilbert transform of a characteristic function*, J. Funct. Anal. **262** (2012), 4534–4539.
- [25] B. Lang, T. Vértesi and M. Navascués, *Closed sets of correlations: answers from the zoo*, J. Phys. A **47** (2014), 424029.
- [26] E. H. Lieb and R. Seiringer, *Equivalent forms of the Bessis–Moussa–Villani conjecture*, J. Stat. Phys. **115** (2004), 185–190.
- [27] Ll. Masanes, *Tight Bell inequality for d -outcome measurements correlations*, Quantum Inf. Comput. **3** (2003), 345–358.
- [28] The mathlib Community, *The Lean mathematical library*, in: Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM, New York, 2020, 367–381.
- [29] A. Mishra and A. Senthilkumar, *A rearrangement inequality for the discrete Hilbert transform, with an application to optimal CGLMP measurements*, preprint (2026), <https://github.com/anshM123/CGLMP>.
- [30] A. Mishra and A. Senthilkumar, *CGLMP on maximally entangled states: exact optimality, rigidity and Tsirelson bounds*, preprint and data repository (2026), <https://github.com/anshM123/CGLMP>.
- [31] A. Mishra and A. Senthilkumar, *A strip inequality for projections, two-variable BMV positivity, and optimal CGLMP measurements in every dimension*, preprint (2026); file papers/math/main.pdf of [33].
- [32] A. Mishra and A. Senthilkumar, *Fourier measurements are optimal for CGLMP Bell tests on maximally entangled states in every dimension*, preprint (2026); file papers/overview/main.pdf of [33].
- [33] A. Mishra and A. Senthilkumar, *IQOQI Vienna Open Problem 27: a complete resolution*. Proofs, certificates, independent verifications and a Lean 4 formalisation (2026), <https://github.com/anshM123/anshM123-IQOQI-Vienna-Open-Problem-27>.
- [34] IQOQI Vienna, *Open Quantum Problems*, Problem 27: *The power of CGLMP inequalities*; archived copy of 29 October 2023: <http://web.archive.org/web/20231029013544/https://oqp.iqoqi.oeaw.ac.at/the-power-of-cglmp-inequalities>.
- [35] S. Popescu and D. Rohrlich, *Quantum nonlocality as an axiom*, Found. Phys. **24** (1994), 379–385.
- [36] H. R. Stahl, *Proof of the BMV conjecture*, Acta Math. **211** (2013), 255–290.
- [37] E. M. Stein and G. Weiss, *An extension of a theorem of Marcinkiewicz and some of its applications*, J. Math. Mech. **8** (1959), 263–284.
- [38] W. van Dam, R. D. Gill and P. D. Grünwald, *The statistical strength of nonlocality proofs*, IEEE Trans. Inform. Theory **51** (2005), 2812–2835; arXiv:quant-ph/0307125.
- [39] Y. Zhang, *CGLMP measurements need not maximize statistical strength*, Zenodo (2026), <https://doi.org/10.5281/zenodo.23022433>.