


The Derivative of the Sigmoid Function

Contents

1. Overview	2
2. Two rules we need	2
3. The derivation	2
3.1 Differentiate	3
3.2 Split into two fractions	3
3.3 The result	3
4. The shape of the derivative	3
5. Summary	6
6. Sources	6
7. Key terms	6

Where this fits: Pipeline map step 8 (Model). See the [Course map](#).



- **Leads to:** Logistic regression ([Note 75](#)); Gradient boosting ([Note 120](#)); Multi-layer perceptron (MLP) ([Note 1003](#)); Backpropagation ([Note 1015](#)); Vanishing gradient ([Note 1018](#)).

1. Overview

Key point: The derivative of the sigmoid can be written using the sigmoid itself: $\sigma'(z) = \sigma(z)(1 - \sigma(z))$.

Gradient descent on the log loss (the next Note) needs the derivative of the sigmoid function. The same derivative appears again in neural networks, where the sigmoid is used inside every neuron. This short Note derives it once.

As a reminder, the sigmoid is

$$\sigma(z) = \frac{1}{1 + e^{-z}}$$

It squeezes any number into the range 0 to 1 (the sigmoid Note).

2. Two rules we need

Key point: The chain rule, and the derivative of e^{-z} , which is $-e^{-z}$.

Chain rule: to differentiate a function of a function, differentiate the outer one, keep the inside, and multiply by the derivative of the inside. For a reciprocal:

$$\frac{d}{dz} \left(\frac{1}{u} \right) = -\frac{1}{u^2} \cdot \frac{du}{dz}$$

Exponential: the derivative of e^z is e^z itself. By the chain rule, the derivative of e^{-z} is e^{-z} times the derivative of $-z$, which is -1 :

$$\frac{d}{dz} e^{-z} = -e^{-z}$$

3. The derivation

Key point: Differentiate, then split the fraction into $\sigma(z)$ times $(1 - \sigma(z))$.

3.1 Differentiate

Key point: Treat $\sigma(z)$ as $1/u$ with $u = 1 + e^{-z}$.

Write $u = 1 + e^{-z}$, so $\sigma(z) = 1/u$. The derivative of u is $0 + (-e^{-z}) = -e^{-z}$. Using the reciprocal rule:

$$\sigma'(z) = -\frac{1}{(1 + e^{-z})^2} \cdot (-e^{-z}) = \frac{e^{-z}}{(1 + e^{-z})^2}$$

The two minus signs cancel.

3.2 Split into two fractions

Key point: One fraction is $\sigma(z)$ itself; the other turns out to be $1 - \sigma(z)$.

Split the result into a product of two fractions:

$$\sigma'(z) = \frac{1}{1 + e^{-z}} \cdot \frac{e^{-z}}{1 + e^{-z}}$$

The first fraction is $\sigma(z)$. For the second, add and subtract 1 in the numerator:

$$\frac{e^{-z}}{1 + e^{-z}} = \frac{(1 + e^{-z}) - 1}{1 + e^{-z}} = 1 - \frac{1}{1 + e^{-z}} = 1 - \sigma(z)$$

3.3 The result

Key point: $\sigma'(z) = \sigma(z)(1 - \sigma(z))$.

Putting the two pieces together:

$$\sigma'(z) = \sigma(z)(1 - \sigma(z))$$

With numbers: at $z = 2$, $\sigma(2) = 0.88$, so $\sigma'(2) = 0.88 \times 0.12 = 0.105$.

This form is very convenient. When a model has already computed $\sigma(z)$ for its prediction, the derivative costs just one subtraction and one multiplication.

4. The shape of the derivative

Key point: The derivative is largest, 0.25, at $z = 0$ and falls towards 0 on both sides.




Figure 1: The sigmoid and its derivative

z	-4	-2	0	2	4
$\sigma(z)$	0.018	0.119	0.5	0.881	0.982
$\sigma'(z)$	0.018	0.105	0.25	0.105	0.018

- At $z = 0$ the sigmoid is steepest: $0.5 \times 0.5 = 0.25$. This is the largest value the derivative can take.
- Far from 0, the sigmoid is almost flat, near 0 or 1, so its slope is almost 0.
- The curve is symmetric: $\sigma'(-z) = \sigma'(z)$.

Python: The derivative, checked numerically.

```
def sigmoid(z):
    return 1 / (1 + np.exp(-z))

def sigmoid_derivative(z):
    return sigmoid(z) * (1 - sigmoid(z))
```

```
h = 1e-5
(sigmoid(2 + h) - sigmoid(2 - h)) / (2 * h)    # 0.104994
sigmoid_derivative(2)                           # 0.104994
```

The numerical slope (rise over a tiny run) matches the formula to six decimals.

Extra: Because the derivative is at most 0.25, and almost 0 for large or small z , stacking many sigmoid layers in a deep neural network multiplies many small numbers together. The gradients become tiny and learning stalls: the **vanishing gradient problem**. Saturating sigmoids make gradient-based learning very difficult, so the sigmoid is now discouraged inside the hidden layers of a network, and ReLU is the default choice

(Goodfellow et al. §6.1, §6.3.2).

5. Summary

- $\sigma(z) = 1/(1 + e^{-z})$.
- $\sigma'(z) = \sigma(z)(1 - \sigma(z))$, found with the chain rule and one algebra step.
- The derivative peaks at 0.25 when $z = 0$ and approaches 0 far from it.
- The next Note uses it to derive the gradient of the log loss.

6. Sources

- **Goodfellow et al.:** Goodfellow, I., Bengio, Y. and Courville, A. *Deep Learning*. MIT Press, 2016. Sections 6.1 and 6.3.2.

7. Key terms

Term	Meaning
Derivative	The rate of change, or slope, of a function at a point
Chain rule	To differentiate a function of a function, multiply the outer derivative by the inner derivative
Vanishing gradient	Gradients shrinking towards 0 as they pass through many layers, which slows learning