

What Is Statistics: Population, Sample and Types of Data

Contents

1. Overview	2
2. What statistics is	2
3. Descriptive and inferential statistics	3
4. Population and sample	3
4.1 What makes a good sample	4
4.2 Parameters and statistics	4
5. The tools of inferential statistics	5
6. Types of data	5
6.1 Nominal and ordinal data	6
6.2 Discrete and continuous data	6
6.3 Why the type matters	6
7. Summary	7
8. Key terms	7

1. Overview

Key point: Statistics summarises the data we have (descriptive) and uses a sample to draw conclusions about a whole population (inferential); the type of each column decides which tools we can use.




Figure 1: A sample is drawn from the population, and conclusions about the population are inferred from it

Figure 1 shows the central idea of statistics: we rarely see the whole group we care about, so we study a part of it and reason back to the whole. This Note defines statistics and its two branches, explains population and sample, and sorts data into four types.

2. What statistics is

Key point: Statistics is the branch of mathematics for collecting, organising, summarising and drawing conclusions from data.

Statistics is the branch of mathematics that deals with collecting, analysing, interpreting and presenting data. It gives us tools to make sense of large amounts of data, to draw conclusions and to make decisions based on data.

Few branches of mathematics are applied as widely. Some examples:

- **Business and finance:** companies study customer data to find buying patterns, then set prices, plan stock and forecast demand.
- **Medicine:** a new medicine goes through clinical trials, and a statistical test decides whether it works better than the old one.
- **Government and politics:** surveys and opinion polls estimate what a whole country thinks from a few thousand answers.
- **Environmental science:** climate research uses past data to predict future change.

- **Machine learning:** linear regression, logistic regression and Naive Bayes all grow out of statistics.

3. Descriptive and inferential statistics

Key point: Descriptive statistics describes the data in hand; inferential statistics predicts things about a larger group from it.

Statistics has two branches:

- Descriptive statistics (see the [understanding your data Note](#)) summarises and describes the data we already have, without drawing conclusions beyond it. The mean, median, standard deviation, five-number summary and every graph of our data belong here. Descriptive statistics studies the past: what the data says.
- **Inferential statistics** uses the data we have to make predictions, or **inferences**, about a larger group we cannot fully observe.

Every exploratory data analysis so far, such as the summaries in the [understanding your data Note](#), was descriptive statistics. Inferential statistics needs one more idea first: population and sample.

4. Population and sample

Key point: The population is every individual we want to study; a sample is the part of it we actually measure.

Suppose the government asks us for the average salary in India. India has about 140 crore people. To get the exact answer, we would have to ask every person their salary, add them up and divide by 140 crore. That is impossible in practice.

So we pick, say, 50,000 people from different states, regions, genders and age groups. We ask them, compute their average salary, and use it to **infer** the average salary of the whole country (Figure 1).

- The **population** is the entire group of individuals or objects we are interested in studying: here, every person in India.
- A **sample** is a subset of the population, the part we actually measure: here, the 50,000 people.

Inferential statistics is exactly this step: from a sample, say something about the population. Two more examples:

Population	Sample
Every Indian cricket fan who watches a match	The fans watching in the stadium
Every student of a college: on campus, online, distance and past students	The students who attend lectures in person

4.1 What makes a good sample

Key point: A sample must be large enough, random and representative, or its conclusions will be wrong.

A badly made sample gives wrong conclusions about the population, however carefully we analyse it. A good sample is:

- **Large enough:** a tiny sample makes the result depend on luck. This is the sampling noise of the [challenges in ML Note](#).
- **Random:** every member of the population has a fair chance of being picked. If our salary survey somehow favours rich people, its average is too high. This is sampling bias, from the same Note.
- **Representative:** every kind of member is present. For India's average salary we need people from every state, men and women, every age group, business owners and employees.

Different ways of building samples are called **sampling techniques**; a later maths Note covers them.

4.2 Parameters and statistics

Key point: A number that describes the population is a parameter (Greek letters); the same number computed from a sample is a statistic (Latin letters).

A number computed from the whole population, such as India's true average salary, is a **parameter**. The same number computed from a sample is a **statistic**, and we use it as an estimate of the parameter.

The two are generally different. The average salary of 50,000 random people is close to the national average, but there is no guarantee it is equal. So we always keep track of which one we have, and write them differently:

Quantity	Population (parameter)	Sample (statistic)
Number of items	N	n
Mean	μ ("mu")	$\bar{x}$ ("x bar")
Variance	σ^2 ("sigma squared")	s^2
Standard deviation	σ	s

The mean and variance are covered in the [measures of central tendency Note](#) and the [measures of dispersion Note](#). For some of them the formula itself changes between population and sample.

5. The tools of inferential statistics

Key point: Hypothesis tests, confidence intervals, ANOVA, regression, chi-square tests, sampling and Bayesian statistics are the main tools for reasoning from a sample.

Inferential statistics has a set of standard tools. Each gets its own Note later:

- **Hypothesis testing:** we make a claim about a population parameter and use a sample to decide whether it holds. For example: is the mean height of a population different from 170 cm? The procedures are called **statistical tests**.
- **Confidence intervals:** we estimate a population parameter from a sample and give a range that most likely contains it.
- **ANOVA (analysis of variance):** compares the means of several groups at once.
- **Regression:** models how one variable depends on others. The [linear regression Notes](#) use it as an ML algorithm.
- **Chi-square test:** a test designed for categorical variables.
- **Sampling techniques:** ways of drawing a good sample from a population.
- **Bayesian statistics:** reasoning that updates beliefs with data, built on [Bayes' theorem](#).

6. Types of data

Key point: Every column is categorical (nominal or ordinal) or numerical (discrete or continuous); the type decides which statistics and graphs make sense.

Descriptive statistics starts by asking what type of data each column holds (Figure 2). We mostly work with tables, so “data” here means one column of a table.




Figure 2: The four types of data

The first split, into categorical and numerical data, is the one of the [types of ML Note](#): categorical data is labels, numerical data is numbers that measure an amount. Each splits

once more.

6.1 Nominal and ordinal data

Key point: Nominal categories have no order; ordinal categories do.

Categorical data is nominal or ordinal, as the [ordinal and label encoding Note](#) explains:

- **Nominal:** the categories have no order. Gender, state, religion and favourite IPL team are nominal: no state is “more” than another.
- **Ordinal:** the categories have a natural order. Course feedback of bad, average and good is ordinal: $\text{bad} < \text{average} < \text{good}$.

6.2 Discrete and continuous data

Key point: Discrete data takes separate values that we count; continuous data can take any value in a range, decimals included.

Numerical data is discrete or continuous:

- **Discrete data** can take only separate values, usually whole numbers that come from counting. A rank is 1, 2 or 3, never 1.5. The number of children in a family, or of siblings on board the Titanic, is discrete.
- **Continuous data** can take any value in a range, including every decimal in between. Weight (35.3 kg), height and a ticket fare are continuous.

Extra: Age is a borderline case. We usually record it in whole years (26, 27, 35), which makes the column discrete. Age itself is continuous, though: the Titanic data stores a baby of five months as 0.42 years. What matters is how the column is recorded and used.

6.3 Why the type matters

Key point: The type of a column decides which measures and graphs we can apply to it.

Before applying any measure or graph, we ask two questions of a column: categorical or numerical? Then nominal or ordinal, discrete or continuous? The answers decide what we can do:

- The mean of a nominal column such as state makes no sense; its most frequent category (the mode) does.
- A median needs an order, so it works for ordinal and numerical data, but not nominal data.
- Categorical columns get bar charts and pie charts; numerical columns get histograms and box plots, as the [frequency tables and graphs Note](#) shows.

7. Summary

Idea	Meaning	Example
Descriptive statistics	Summarising the data we have	Mean fare of the Titanic passengers
Inferential statistics	Concluding about a population from a sample	India's average salary from 50,000 people
Population	Everyone we want to study	Every person in India
Sample	The part we measure	50,000 people from every state
Parameter / statistic	A number of the population / of a sample	μ / $\bar{x}$
Nominal / ordinal	Categories without / with an order	state / feedback
Discrete / continuous	Separate values / any value in a range	rank / weight

- A good sample is large enough, random and representative.
- Population numbers use Greek letters (μ , σ); sample numbers use Latin letters ($\bar{x}$, s).
- Check each column's type before choosing a measure or a graph.

8. Key terms

Term	Meaning
Statistics	The branch of mathematics for collecting, analysing, interpreting and presenting data
Inferential statistics	Statistics that draws conclusions about a population from a sample
Inference	A conclusion about a population drawn from a sample
Population	The entire group of individuals or objects we want to study
Parameter	A number that describes the population, such as μ
Statistic	A number computed from a sample, such as $\bar{x}$; an estimate of a parameter
Sampling techniques	Ways of drawing a good sample from a population
Hypothesis testing	Checking a claim about a population parameter with a sample
Statistical test	A procedure for hypothesis testing
ANOVA	Analysis of variance: a test comparing the means of several groups

Term	Meaning
Chi-square test	A statistical test for categorical variables
Discrete data	Numerical data that takes only separate values, usually counts
Continuous data	Numerical data that can take any value in a range