

Null and Alternative Hypotheses

Contents

1. Overview	2
2. The problem hypothesis testing solves	3
2.1 A YouTube channel tries a new style	3
2.2 The same question in business	3
2.3 Definition	3
3. The null hypothesis	4
4. The alternative hypothesis	4
5. Three rules for H_0 and H_1	5
5.1 How to tell which statement is H_0	5
5.2 The goal: evidence against H_0	5
5.3 Failing to reject H_0 does not prove it	6
6. The eight steps of a hypothesis test	6
7. Summary	9
8. Key terms	9

Where this fits: Pipeline map step 0 (Foundations). See the [Course map](#).



- **Builds on:** Inferential statistics ([Note 220](#)); Sampling distribution and standard error ([Note 271](#)).
- **Leads to:** Z-test and rejection regions ([Note 291](#)); Beta and A/B testing ([Note 292](#)); T-tests: one-sample, two-sample, paired ([Note 301](#)).
- **Compare with:** Confidence intervals ([Note 282](#)).

1. Overview

Key point: A hypothesis test turns an idea into two opposite statements, the null and the alternative hypothesis, and asks whether a sample gives strong enough evidence against the null.




Figure 1: The idea behind every hypothesis test

Hypothesis testing checks a claim about a population parameter with a sample (see the [what is statistics Note](#)). Figure 1 shows the whole idea on one example: we write the claim as two statements, look at the data, and either reject the first statement or fail to reject it.

This Note covers:

- the problem hypothesis testing solves;

- the null hypothesis and the alternative hypothesis, and how to write them;
- three rules for using them, with the courtroom analogy;
- the eight steps of a hypothesis test.

The next Notes run these steps on real numbers: the [rejection region Note](#) with a z-test, the [p-values Note](#) with the method used in practice.

2. The problem hypothesis testing solves

Key point: One result, or a few, can be good or bad by chance; hypothesis testing tells us whether a sample is strong enough evidence for a claim about the whole population.

2.1 A YouTube channel tries a new style

Suppose a YouTube channel records its videos on a digital writing pad. Its analytics show that, over all its videos, people watch for **6 minutes** on average. Successful channels seem to film in front of a whiteboard instead, so the owner has an idea: the whiteboard style would raise the average view duration.

The obvious test is to try it. The first whiteboard video is watched for 30 minutes on average. Can we conclude that the new style works?

No. One video proves very little:

- its topic may have been popular (say, a video on ChatGPT that the whole world wants to watch);
- the presenter may have looked especially good that day;
- any other random reason may have lifted that one video.

So the channel makes five more videos in the new style. Their average view durations are 7, 9, 5, 11 and 13 minutes. Their mean, 9 minutes, is above 6. Is that enough? The question has not changed: are five videos enough evidence, or could the old style give such numbers by chance?

2.2 The same question in business

A chips company sells a green packet and launches a purple one. Before replacing green with purple everywhere, it wants to know whether the purple packet really sells better. It can ask customers for feedback, but then: how many customers, and how good must the feedback be before we believe it?

A small channel can afford to try and see. A company such as Amazon or Flipkart, where one wrong decision costs a lot of money, needs a method. That method is hypothesis testing.

2.3 Definition

A **statistical hypothesis test** is a method of statistical inference used to decide whether the data at hand sufficiently supports a particular hypothesis. It lets us make probabilistic

statements about population parameters.

In the channel example the population parameter is μ , the mean view duration of all videos the channel could make in the new style. The new videos are a sample. Hypothesis testing tells us what the sample can say about μ .

This is inferential statistics: reasoning from a sample to a population (see the [what is statistics Note](#)). It is used constantly in business, finance and economics.

3. The null hypothesis

Key point: The null hypothesis H_0 says that nothing has changed: no effect, no difference, no relationship.

Every hypothesis test starts with two statements. The first is the **null hypothesis**, written H_0 :

a statement that assumes there is no significant effect or relationship between the variables being studied. It is the starting point of the test and stands for the status quo, “no effect until proven otherwise”.

Two examples:

- **The channel.** We want to show that the new style raises the view duration. The null hypothesis says the opposite: the old and the new style make no difference, and the mean view duration is still 6 minutes.
- **A chips packet.** A packet says “100 g” on the back, and we suspect the true mean weight is different, either more or less. The null hypothesis says the mean weight of the packets is exactly 100 g.

In both cases H_0 is the boring statement: whatever was true before is still true. The test then tries to prove it wrong.

4. The alternative hypothesis

Key point: The alternative hypothesis H_1 contradicts H_0 and claims there is an effect; exactly one of the two is true.

The second statement is the **alternative hypothesis**, written H_1 or H_a :

a statement that contradicts the null hypothesis and claims there is a significant effect or relationship between the variables being studied.

The two statements are mutually exclusive: they cannot both be true, and the test ends by siding with one of them. For our examples:

Example	H_0 (no effect)	H_1 (an effect)
Channel: mean view duration in minutes	$\mu = 6$	$\mu > 6$

Example	H_0 (no effect)	H_1 (an effect)
Chips: mean packet weight in grams	$\mu = 100$	$\mu \neq 100$

The channel's H_1 has a direction: the new style should **increase** the duration. The chips' H_1 has none: the weight is wrong, whether too high or too low. This difference decides between a one-tailed and a two-tailed test (see the [errors, power and tails Note](#)).

Two other names are common in books and exam questions:

- H_0 is called the **status quo**: the current state of things;
- H_1 is called the **research hypothesis**: the idea that comes out of research. The channel owner studied other channels and concluded that the whiteboard style might help.

Extra: The hypotheses are always about the **population** parameter (μ), never about the sample mean ($\bar{x}$). We already know $\bar{x} = 9$ for the five new videos; there is nothing to test about it. The open question is μ , the mean of all future videos. Writing $H_0 : \bar{x} = 6$ is a common slip.

5. Three rules for H_0 and H_1

Key point: H_0 says nothing new is happening; the test collects evidence against H_0 ; and failing to reject H_0 does not prove it true.

5.1 How to tell which statement is H_0

Faced with a new problem, we may wonder which statement should be H_0 . The rule of thumb: **H_0 is the statement that nothing new is happening.**

- We changed the filming style, and the mean view duration is still 6 minutes: H_0 .
- We opened packets and weighed them, and the mean is still 100 g: H_0 .

The statement with the change, the difference or the effect is H_1 .

Extra: The equals sign always goes in H_0 . The test needs one exact value to compute with ($\mu = 6$), and only H_0 provides it. Some books write a one-tailed null as $H_0 : \mu \leq 6$; the calculation is the same, because $\mu = 6$ is the case in H_0 closest to H_1 .

5.2 The goal: evidence against H_0

A hypothesis test does not try to prove H_1 directly. It assumes H_0 and then uses the data to collect **evidence against H_0** . If the evidence is strong enough, we reject H_0 , and H_1 is what remains.

5.3 Failing to reject H_0 does not prove it

The most important rule: **failing to reject the null hypothesis does not mean the null hypothesis is true.** It only means the data did not bring enough evidence against it.

A courtroom makes this clear:

Court	Hypothesis test
The accused is presumed innocent	H_0 is assumed true: no crime
The prosecutor claims the accused is guilty	H_1 : there was a crime
The prosecutor brings evidence	We collect sample data
Strong evidence: verdict “guilty”	Reject H_0
Weak evidence: verdict “not guilty”	Fail to reject H_0

A “not guilty” verdict does not prove that no crime happened. It says the prosecutor could not prove it. In the same way, if the five videos do not let us reject H_0 , the new style may still work: our evidence was simply too weak to show it.

For the same reason, rejecting H_0 for the channel would not show that the whiteboard style is the best possible. It would only show that it beats the old average of 6 minutes.

Extra: This is why careful writers say “fail to reject H_0 ” and avoid “accept H_0 ”. With a small sample almost nothing gets rejected, so “accepting” H_0 would turn weak evidence into a false certainty. Absence of evidence is not evidence of absence.

6. The eight steps of a hypothesis test

Key point: State H_0 and H_1 , choose α , check assumptions, choose the test and its statistic, compute it, decide, and interpret.




Figure 2: The eight steps of a hypothesis test (rejection region approach)

There are two ways to carry out a test. The **rejection region approach** compares the test statistic with a fixed boundary (see the [rejection region Note](#)). The **p-value approach** computes one extra number that also measures how strong the evidence is (see the [p-values Note](#)); it is the approach used in practice. Both follow the steps in Figure 2.

1. **State H_0 and H_1 .** For the channel: $H_0 : \mu = 6$, $H_1 : \mu > 6$.
2. **Choose a significance level α .** Usually 0.05 (5%), sometimes 0.01 (1%). It is the probability of rejecting H_0 when H_0 is actually true: at 5%, about 5 tests in 100 with a true H_0 would wrongly reject it. The [rejection region Note](#) explains it fully.
3. **Check the assumptions about the data.** Is the data normally distributed? Do we know the population standard deviation σ ? Is the data numerical or categorical? Do we have one column or several, one group or several?
4. **Choose the test.** The assumptions decide it. For example:
 - normal data (or a large sample) and σ known: the one-sample **z-test**;
 - σ unknown: the **t-test** (see the [one-sample t-test Note](#));
 - categorical data: the chi-square test;
 - several group means: ANOVA.
5. **Name the test statistic.** The **test statistic** is the number the test computes from the sample. A z-test computes a z-score, called the z statistic; a t-test computes the t statistic (as in the [t-procedure Note](#)).
6. **Conduct the test.** Compute the test statistic from the sample.
7. **Decide.** Based on the statistic, reject H_0 or fail to reject it.

8. **Interpret the result.** Translate the decision back into the real question. If the channel rejected H_0 , the interpretation would be: “filming in the new style increases the mean view duration”.

Steps 1 and 2 come before looking at the data. Choosing α or the direction of H_1 after seeing the results would let us tune the test until it says what we want.

Extra: Are the five videos (7, 9, 5, 11, 13 minutes) enough evidence that $\mu > 6$? The [one-sample t-test Note](#) answers it: at the 5% level, just barely not.

7. Summary

	Null hypothesis H_0	Alternative hypothesis H_1
Says	no effect, no difference	there is an effect or difference
Other name	status quo	research hypothesis
Contains	the equals sign ($\mu = 6$)	$>$, $<$ or $\neq$
Role in the test	assumed true; we look for evidence against it	what remains if H_0 is rejected
Channel example	$\mu = 6$ minutes	$\mu > 6$ minutes
Chips example	$\mu = 100$ g	$\mu \neq 100$ g

- A single result, or a few, can be lucky; a hypothesis test asks whether a sample is strong enough evidence about a population parameter.
- H_0 says nothing new is happening; H_1 contradicts it. Exactly one is true.
- The test collects evidence against H_0 . The two outcomes are “reject H_0 ” and “fail to reject H_0 ”.
- Failing to reject H_0 does not prove H_0 : “not guilty” is not “innocent”.
- Eight steps: hypotheses, α , assumptions, test, statistic, compute, decide, interpret.

8. Key terms

Term	Meaning
Statistical hypothesis test	A method of statistical inference that decides whether the data sufficiently supports a hypothesis about a population parameter
Null hypothesis (H_0)	The statement of no effect, no difference or no relationship; assumed true until the data gives strong evidence against it
Alternative hypothesis (H_1 , H_a)	The statement that contradicts H_0 and claims an effect, difference or relationship
Status quo	Another name for the null hypothesis: the current state of things
Research hypothesis	Another name for the alternative hypothesis: the idea that came out of research
Reject H_0	The decision that the data gives strong enough evidence against H_0
Fail to reject H_0	The decision that the evidence against H_0 is not strong enough; it does not prove H_0
Test statistic	The number a test computes from the sample to make its decision, such as z or t
Rejection region approach	Carrying out a test by checking whether the test statistic falls beyond a fixed boundary

Term	Meaning
P-value approach	Carrying out a test by computing a p-value, which also measures the strength of the evidence