

Probability vs Likelihood

Contents

1. Overview	2
2. A coin	3
2.1 Probability: from the parameter to the event	3
2.2 Likelihood: from the event back to the parameter	3
3. A bag of balls	4
3.1 Probability	4
3.2 Likelihood	4
4. Heights: a continuous example	5
4.1 Probability is an area	5
4.2 Likelihood is a height	5
4.3 The likelihood function: slide the parameter	6
4.4 The same picture with mouse weights	7
5. The two definitions	8
6. A likelihood is not a probability	9
7. Summary	10
8. Sources	10
9. Key terms	10

Where this fits: Pipeline map step 0 (Foundations). See the [Course map](#).



- **Builds on:** Probability distributions ([Note 240](#)); Probability density function (PDF) ([Note 242](#)).
- **Leads to:** Maximum likelihood estimation (MLE) ([Note 631](#)); Multivariate normal distribution ([Note 640](#)).
- **Compare with:** Probability mass function (PMF) ([Note 241](#)); Probability density function (PDF) ([Note 242](#)); Student's t-distribution ([Note 282](#)).

1. Overview

Key point: Probability starts from known parameters and asks how likely an event is. Likelihood starts from an observed event and asks how plausible a parameter value is. Both use the same formula, read in opposite directions.




Figure 1: Probability goes from parameters to an event; likelihood goes from an observed event back to the parameters

The two words mean the same thing in everyday speech, which is why the difference is confusing. Picture a tunnel with a parameter at one end and an event at the other (Figure 1). Looking from the parameter end towards the event is probability. Looking from the event end back towards the parameter is likelihood.

This Note builds the idea from three examples, each read both ways:

- tossing a coin (Section 2);
- drawing balls from a bag (Section 3);
- people's heights, a continuous example (Section 4).

The Note then states both definitions (Section 5) and shows why a likelihood is not a probability (Section 6). The [maximum likelihood estimation Note](#) uses likelihood to fit parameters.

2. A coin

Key point: Fair coin known: the probability of tails is 0.5. Five heads observed: the likelihood that the coin is fair is $0.5^5 = 0.031$, lower than the likelihood $0.7^5 = 0.168$ of a coin biased towards heads.

2.1 Probability: from the parameter to the event

Key point: The coin follows a Bernoulli distribution with $p = 0.5$; plugging in the event “tails” gives its probability.

Tossing a coin once has two outcomes, heads or tails, so it follows a **Bernoulli distribution** with one parameter p , the probability of heads (see the [Bernoulli and binomial Note](#), Section 2.1). For a fair coin, $p = 0.5$.

1. **In words:** the Bernoulli PMF gives the probability of each outcome k (1 for heads, 0 for tails) once p is known.
2. **Formula:**

$$P(X = k) = p^k(1 - p)^{1-k}, \quad k \in \{0, 1\}$$

3. **Example:** for tails, $k = 0$: $P(X = 0) = 0.5^0 \times 0.5^1 = 0.5$.

Common sense gives the same answer: if heads has probability 0.5, tails has $1 - 0.5 = 0.5$. The pattern is what matters. We knew the distribution (Bernoulli) and its parameter ($p = 0.5$), and we computed the chance of an event (tails). Computing such a chance is probability.

2.2 Likelihood: from the event back to the parameter

Key point: We toss the coin five times and get five heads. The likelihood of “the coin is fair” given this data is $0.5^5 = 0.031$.

Now turn it around. We are told the coin is fair, we toss it five times, and all five come up heads. Is “fair” still believable? The tosses are independent, so the probability of five heads under p is a product of five equal factors (see the [independent events Note](#)).

1. **In words:** the likelihood of a parameter value is the probability of the observed data computed with that value; the data stays fixed and we change the parameter.
 2. **Formula:**
- $$L(p \mid 5 \text{ heads}) = p \times p \times p \times p \times p = p^5$$
3. **Example:** for a fair coin, $L(0.5) = 0.5^5 = 0.031$. For a coin with $p = 0.7$, $L(0.7) = 0.7^5 = 0.168$.

Five heads are more than five times as plausible under $p = 0.7$ as under $p = 0.5$ (Figure 2, left). The data supports a coin biased towards heads more than a fair one.




Figure 2: Likelihoods of two parameter values for each observed event: 5 heads in 5 tosses (left), 5 green balls in 5 draws (right)

3. A bag of balls

Key point: A bag holds 3 red and 2 green balls. Probability: a drawn ball is red with probability $3/5$. Likelihood: after five green draws in a row, the value $p = 2/5$ has likelihood 0.010, far below the 0.328 of a bag with 4 green balls out of 5.

3.1 Probability

Key point: One draw is again a Bernoulli trial, with p = probability of green = $2/5$.

A closed bag holds 3 red and 2 green balls. Drawing one ball has two outcomes, so it is again a Bernoulli trial; take green as the outcome counted by p , so $p = 2/5$.

1. **In words:** red is $k = 0$, so its probability is $1 - p$.

2. **Formula:**

$$P(\text{red}) = p^0(1 - p)^1 = 1 - p$$

3. **Example:** $1 - 2/5 = 3/5 = 0.6$.

Again we knew the parameter and computed the chance of an event: probability.

3.2 Likelihood

Key point: Five green draws make $p = 2/5$ implausible: $L(2/5) = (2/5)^5 = 0.010$.

We draw five times, putting each ball back before the next draw so that the draws are independent. All five are green. We were told that $p = 2/5$; how plausible is that now?

1. **In words:** multiply the probability of green, under the value of p being questioned, once per green draw.
2. **Formula:**

$$L(p \mid 5 \text{ green}) = p^5$$

3. **Example:** $L(2/5) = 0.4^5 = 0.010$. For a bag with 4 green balls out of 5, $L(4/5) = 0.8^5 = 0.328$, about 32 times higher.

So five green draws point to a bag with mostly green balls (Figure 2, right). The direction is the same as for the coin: the event is fixed, and we compare parameter values by how well each one explains it.

4. Heights: a continuous example

Key point: For a continuous variable, probability is an area under the density curve, and likelihood is the height of the curve at the observed value.

4.1 Probability is an area

Key point: With heights following $N(150, 10^2)$, the probability that a person is between 170 and 180 cm is the area under the curve there: 0.021.

Heights of people are continuous: any value is possible. Assume they follow a normal distribution with mean $\mu = 150$ cm and standard deviation $\sigma = 10$ cm (see the [normal distribution Note](#)).

1. **In words:** the probability that a randomly picked person's height falls in a range is the area under the PDF over that range (the [PDF and continuous CDF Note](#)).
2. **Formula:**

$$P(170 \leq X \leq 180) = F(180) - F(170)$$

where F is the CDF of $N(150, 10^2)$.

3. **Example:** 170 and 180 are 2 and 3 standard deviations above the mean, so the area is $0.99865 - 0.97725 = 0.021$ (values from the z-table, see the [standard normal and z-table Note](#)).

The parameters were known and the event was a range of heights: probability.

4.2 Likelihood is a height

Key point: We measure one person at 100 cm. The likelihood of $\mu = 150$, $\sigma = 10$ is the height of the $N(150, 10^2)$ curve at 100: 1.49×10^{-7} , almost nothing.

Now we pick one person at random and measure 100 cm. How plausible are $\mu = 150$ and $\sigma = 10$ for this **observation** (one recorded value)?

1. **In words:** put the observed value into the normal PDF, with the parameter values being questioned.

2. Formula:

$$L(\mu, \sigma \mid x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

3. **Example:** for $x = 100$, $\mu = 150$, $\sigma = 10$, the exponent is $-(100 - 150)^2/200 = -12.5$:

$$L = \frac{1}{10\sqrt{2\pi}} e^{-12.5} = 0.0399 \times 3.73 \times 10^{-6} = 1.49 \times 10^{-7}$$

Changing the observation changes the verdict on the same parameters:

Observed height x (cm)	Likelihood of $\mu = 150$, $\sigma = 10$
100	1.5×10^{-7}
130	0.0054
140	0.024
150	0.040
200	1.5×10^{-7}

An observation of 140 cm makes $\mu = 150$ quite plausible; 100 cm or 200 cm make it almost impossible. Only the distance between x and μ enters the formula, through $(x - \mu)^2$, so 100 and 200 cm, both 50 cm away, get the same likelihood.

4.3 The likelihood function: slide the parameter

Key point: Keep the observation fixed and slide μ : the heights trace the likelihood function, which peaks at the μ equal to the observation.

Because only $x - \mu$ matters, moving the observation away from μ is the same as moving μ away from the observation. The natural way to use likelihood is the second one: the data is what we have, and we try parameter values. Figure 3 fixes one person at 130 cm and slides the candidate mean from 100 to 200 cm.




Figure 3: One person measured at 130 cm. Top: the candidate curve $N(\mu, 10^2)$ slides; the orange bar is its height at 130. Bottom: those heights against μ , the likelihood function, highest at $\mu = 130$

The likelihood is highest, 0.040, when the curve is centred on the observation. Picking the parameter value at this peak is maximum likelihood estimation, the subject of the [maximum likelihood estimation Note](#).

4.4 The same picture with mouse weights

Key point: Probabilities are areas under a fixed distribution; likelihoods are heights at fixed data under a distribution that can move.

A second example (Starmer, StatQuest) draws the same contrast with mouse weights following $N(32, 2.5^2)$ (Figure 4):

- **probability** (left): the distribution is fixed; the chance that a mouse weighs 32 to 34 grams is the shaded area, 0.29;
- **likelihood** (right): we weighed one mouse at 34 grams; the likelihood of $N(32, 2.5^2)$ is its height at 34, 0.12. Shifting the mean to 34 raises the likelihood to 0.16.




Figure 4: Mouse weights. Left: probability as the area between 32 and 34 grams under $N(32, 2.5^2)$. Right: likelihood as the height at the observed 34 grams, for two candidate distributions

The notation keeps the two apart:

$$P(\text{data} \mid \text{distribution}) \quad \text{versus} \quad L(\text{distribution} \mid \text{data})$$

In $P(32 \leq \text{weight} \leq 34 \mid \mu = 32, \sigma = 2.5) = 0.29$ we change the left side to ask about other weights. In $L(\mu = 32, \sigma = 2.5 \mid \text{weight} = 34) = 0.12$ the right side, the data, stays fixed, and we change the left side to try other distributions.

Extra: A density height is not itself a probability; the [PDF and continuous CDF Note](#) (Section 5) shows that it is probability per unit of x . Because a density is a rate, a likelihood for continuous data can be larger than 1. Only comparisons between likelihoods are meaningful, never a single value on its own.

5. The two definitions

Key point: Probability: parameters known, chance of an event. Likelihood: data observed, plausibility of parameter values.

All three examples follow one pattern:

	Probability	Likelihood
What is known	the distribution and its parameters	the observed data
What varies	the event	the parameter values

	Probability	Likelihood
Question	how likely is this event?	how well do these parameter values explain the data?
Coin	$P(\text{tails} \mid p = 0.5) = 0.5$	$L(p = 0.5 \mid 5 \text{ heads}) = 0.031$
Bag	$P(\text{red} \mid p = 2/5) = 0.6$	$L(p = 2/5 \mid 5 \text{ green}) = 0.010$
Heights	$P(170 \leq X \leq 180 \mid 150, 10) = 0.021$	$L(150, 10 \mid x = 100) = 1.5 \times 10^{-7}$

- **Probability** measures the chance that a certain event will occur out of all possible events. A probability is a number from 0 to 1.
- **Likelihood** is a function that measures how plausible a particular parameter value is, given some observed data. The likelihood says how well an observed outcome supports a parameter value.

Said another way: a probability tells us how often to expect an outcome when we understand the process that produces the data; a likelihood tells us how good a model is, given data we have already seen.

6. A likelihood is not a probability

Key point: Probabilities of all possible events add up to 1. Likelihoods of all parameter values do not: for five heads, the area under $L(p) = p^5$ is $1/6$.

Probability and likelihood share a formula, so it is tempting to treat the likelihood as a probability distribution over the parameter. *Mathematics for Machine Learning* (MML §9.2.1, remark) warns that it is not: the likelihood is a probability distribution in the data, but not in the parameters.

The coin shows both sides. With p fixed at 0.5, the probabilities of all possible numbers of heads in five tosses, 0 to 5, add up to 1 (they form the binomial distribution of the [Bernoulli and binomial Note](#)). With the data fixed at five heads, the likelihood $L(p) = p^5$ over all values of p from 0 to 1 does not:

1. **In words:** the area under the likelihood curve, over every possible value of the parameter, is not 1.
2. **Formula:**

$$\int_0^1 p^5 dp = \left[\frac{p^6}{6} \right]_0^1 = \frac{1}{6}$$

3. **Example:** the area is 0.167, not 1. The Notebook confirms both sums: 1.000 for the probabilities, 0.167 for the likelihood area.

So a likelihood value only means something next to another likelihood value for the same data: 0.168 against 0.031 says $p = 0.7$ explains five heads better than $p = 0.5$.

7. Summary

	Probability	Likelihood
Direction	parameters $\rightarrow$ event	data $\rightarrow$ parameters
Fixed	parameters	data
Varies	event	parameters
Discrete data	PMF value or sum of PMF values	PMF value(s) of the observed data
Continuous data	area under the PDF	height of the PDF at the observed value
Adds up to 1?	yes, over all events	no, over all parameter values
Notation	$P(\text{data} \mid \theta)$	$L(\theta \mid \text{data})$

- The same formula gives a probability when the parameters are fixed and a likelihood when the data is fixed.
- For independent observations, the likelihood is a product: five heads give p^5 .
- Comparing likelihoods tells which parameter value explains the data better: 0.168 for $p = 0.7$ against 0.031 for $p = 0.5$.
- For continuous data, likelihood is the height of the density at the observation; probability is an area.
- A likelihood is not a probability distribution over the parameters.

8. Sources

Built from

- CampusX, “Probability vs Likelihood | Machine Learning Interview Question”, YouTube, <https://www.youtube.com/watch?v=QBFVcBXRzu4>
- Starmer, J. (StatQuest), “In Statistics, Probability is not Likelihood.”, YouTube, <https://www.youtube.com/watch?v=pYxNSUDSFH4>. Used in Section 4.4.

Other references

- Deisenroth, M. P., Faisal, A. A. and Ong, C. S. (2020). *Mathematics for Machine Learning*. Cambridge University Press. Free PDF at mml-book.github.io. §9.2.1 (remark: the likelihood is not a probability distribution in the parameters). Used in Section 6.

9. Key terms

Term	Meaning
Probability	The chance of an event when the distribution and its parameters are known; between 0 and 1

Term	Meaning
Likelihood	How plausible a parameter value is, given observed data: the probability (or density) of the data computed with that value
Likelihood function $L(\theta \mid \text{data})$	The likelihood as a function of the parameters, with the data held fixed
Observation	One recorded value (one row of the data table)
Plausibility	How believable a parameter value is in the light of the data; measured by its likelihood relative to other values