

The Quantum Ontology of Light: From Wave-Particle Duality to Quantum Field Theory and Beyond

Anshuman Singh

Independent Physics Research Aspirant

anshumansingh3697@gmail.com

REVIEW PAPER - COMPREHENSIVE SYNTHESIS

This is a comprehensive review paper compiling and synthesizing existing research.
All experimental data referenced are from published literature.
Original research papers cited throughout.

DATA SOURCES:

Experimental Data:	**Theoretical Framework:**
Physical Review Letters	Quantum Electrodynamics (QED)
Nature Physics	Quantum Field Theory
Science	Quantum Optics
Physical Review A/D	Mathematical Physics
Optics Express	Gauge Theory
arXiv:quant-ph	Standard Model

Compiled: January 10, 2026

Version: 1.0

Total Pages: 28 (estimated)

This review paper synthesizes existing knowledge from published research.
All figures are illustrative representations based on established physics principles.
Mathematical derivations follow standard textbooks and peer-reviewed publications.

Abstract

This comprehensive review presents a complete synthesis of light's quantum ontology, tracing the conceptual evolution from the classical wave-particle dichotomy to its ultimate resolution within quantum field theory (QFT). We demonstrate that the apparent paradox of wave-particle duality dissolves completely in the framework of quantum electrodynamics (QED), where the photon emerges as an excitation of the quantized electromagnetic field—a massless gauge boson with helicity ± 1 satisfying the $U(1)$ gauge symmetry. The review begins with a detailed historical analysis of classical optics, proceeds through the quantum revolution, establishes the complete mathematical framework of QED, and explores modern quantum optics experiments that probe the limits of our understanding. Special attention is given to the photon localization problem, the mathematical representation of photon wavefunctions, and the operational meaning of complementarity in contemporary quantum information theory. We further examine advanced topics including squeezed light states, quantum eraser experiments, photon antibunching, and the role of vacuum fluctuations. The review concludes with frontier questions at the interface of quantum optics and quantum gravity, including modifications to the photon dispersion relation in Planck-scale physics and photon propagation in curved spacetime. Throughout, we emphasize that light's behavior is completely and consistently described by QED, with wave and particle aspects representing complementary experimental manifestations of a single quantum entity. This work aims to provide both a comprehensive reference for researchers and a pedagogical synthesis of one of physics' most profound conceptual journeys.

Keywords: Wave-particle duality, Quantum Electrodynamics, Photon ontology, Quantum field theory, Quantum optics, Photon localization, Gauge invariance, Quantum information, Photon statistics, Vacuum fluctuations

Review Scope: This paper synthesizes research from approximately 150 key publications spanning 1905-2024. Primary sources include Physical Review journals, Nature/Science family journals, and foundational textbooks in quantum optics and quantum field theory.

Contents

1 Introduction: The Historical and Conceptual Landscape **5**
1.1 The Enduring Mystery of Light 5
1.2 Thesis Statement and Philosophical Orientation 6
1.3 Review Methodology and Data Sources 6

2 Historical Development: From Corpuscles to Quanta **6**
2.1 Ancient and Classical Theories 6
2.2 The Wave Theory Ascendant 7
2.2.1 Interference and Diffraction 7
2.2.2 Polarization and Transverse Waves 7
2.2.3 The Velocity of Light 7
2.3 Maxwell's Electromagnetic Synthesis 8
2.4 Three Anomalies and the Quantum Revolution 8
2.5 Early Quantum Mechanics and Wave-Particle Duality 9

3 Quantum Mechanical Foundations: Mathematical Formalism **9**
3.1 The Photon in Non-Relativistic QM 9
3.1.1 Quantization of the Electromagnetic Field 9
3.1.2 Photon States and Statistics 10
3.2 Complementarity Principle: Mathematical Formulation 11
3.3 Photon Spin and Helicity 11

4 Quantum Electrodynamics: Complete Mathematical Framework **12**
4.1 Covariant Quantization of the Electromagnetic Field 12
4.1.1 Lagrangian Formulation 12
4.1.2 Gauge Invariance and Constraints 12
4.1.3 Canonical Quantization with Constraints 12
4.2 Photon Propagator 12
4.3 Feynman Rules and Perturbation Theory 13
4.4 Renormalization and Precision Tests 14
4.4.1 Renormalization Constants 15
4.4.2 Running Coupling 15
4.4.3 Precision Tests 15

5 Modern Quantum Optics: Probing Photon Nature **15**
5.1 Non-Classical States of Light 15
5.1.1 Squeezed States 15
5.1.2 Photon-Number States 16
5.1.3 Schrödinger Cat States 16

5.2 Quantum Interference and Indistinguishability 16
5.2.1 Hong-Ou-Mandel Effect 16
5.2.2 Quantum Eraser Mathematics 17
5.3 Photon Statistics and Antibunching 17
5.4 Cavity Quantum Electrodynamics 18

6 Photon Localization and Wavefunction Representations 18
6.1 The Localization Problem 18
6.2 Photon Wavefunction Formulations 19
6.2.1 Bialynicki-Birula-Sipe Wavefunction 19
6.2.2 Landau-Peierls Wavefunction 19
6.2.3 Momentum-Space Representation 20
6.3 Operational Localization via Detection 20
6.4 Photon Position Operators 20
6.4.1 Newton-Wigner Operator 20
6.4.2 Hawton Operator 20

7 Advanced Topics and Frontiers 21
7.1 Photon in Curved Spacetime 21
7.1.1 Covariant Quantization in Curved Backgrounds 21
7.1.2 Particle Concept Ambiguity 21
7.1.3 Gravitational Redshift and Lensing 21
7.2 Quantum Gravity and Photon Dispersion 21
7.2.1 Lorentz Violation 22
7.2.2 Quantum Foam Models 22
7.2.3 Double Special Relativity 22
7.3 Photon-Photon Scattering 22
7.4 Topological Photonics and Synthetic Gauge Fields 22
7.4.1 Quantum Hall Effect for Photons 23
7.4.2 Topological Edge States 23
7.5 Quantum Information with Photons 23
7.5.1 Encoding Schemes 23
7.5.2 Quantum Key Distribution 23
7.5.3 Linear Optical Quantum Computing 23

8 Conclusion: Synthesis and Open Questions 23
8.1 The Resolution of Wave-Particle Duality 23
8.2 Open Theoretical Questions 24
8.3 Experimental Frontiers 24
8.4 Concluding Reflections 25

A Mathematical Appendices **25**
A.1 Quantization in Lorenz Gauge 25
A.2 Coherent State Properties 26
A.3 Photon Detection Theory 26
A.4 Casimir Effect Derivation 26
A.5 Photon Angular Momentum Decomposition 26

1 Introduction: The Historical and Conceptual Landscape

1.1 The Enduring Mystery of Light

The nature of light represents perhaps the most persistent and productive conceptual tension in the history of physics. From antiquity's visual rays to Newton's corpuscles, Young's waves, Maxwell's fields, Einstein's photons, and finally to quantum field theory's excitations, each paradigm shift has simultaneously resolved existing anomalies while revealing deeper mysteries. This review aims not merely to recount this history but to demonstrate how modern quantum field theory—specifically quantum electrodynamics (QED)—provides a complete, consistent, and paradox-free description of light's behavior.



Figure 1: Comprehensive historical timeline of light theories spanning four centuries of conceptual evolution. Notable transitions include: (1) Newton's corpuscular theory (1672) explaining reflection and refraction through particle mechanics; (2) Huygens' wave principle (1678) with secondary wavelet construction; (3) Young's double-slit experiment (1803) demonstrating interference; (4) Fresnel's mathematical wave theory (1818) explaining diffraction; (5) Maxwell's electromagnetic theory (1865) unifying electricity, magnetism, and optics; (6) Planck's blackbody quantization (1900); (7) Einstein's photon hypothesis (1905); (8) Dirac's quantum electrodynamics (1927); (9) Feynman-Schwinger-Tomonaga renormalized QED (1940s); (10) Modern quantum optics and photonics. Each paradigm addressed limitations of its predecessor while maintaining continuity in empirical predictions.

1.2 Thesis Statement and Philosophical Orientation

This review advances the following central thesis: *The wave-particle duality of light is not a fundamental property of nature but rather a historical artifact of attempting to describe quantum phenomena using classical concepts. In the complete framework of quantum field theory, light is described by a quantized gauge field whose excitations (photons) exhibit particle-like properties in interactions and wave-like properties in propagation, with no contradiction between these aspects.*

We adopt a *scientific realist* perspective tempered by operationalism: while the mathematical structures of QED provide our most fundamental description, the interpretation of these structures remains connected to experimental arrangements. The photon is not a "little ball of light" nor a "vibration in a medium," but rather an irreducible entity described completely by its transformation properties under the Poincaré group and gauge group $U(1)$.

1.3 Review Methodology and Data Sources

This review synthesizes data from multiple sources:

Table 1: Data sources and methodology for this review

Source Type	**Description**	**Time Period**
Experimental Papers	PRL, Nature Physics, Science, Physical Review A	1905-2024
Theoretical Papers	Annals of Physics, JHEP, Nuclear Physics B	1927-2024
Review Articles	Reviews of Modern Physics, Physics Reports	1950-2024
Textbooks	Quantum Optics, QFT textbooks	Various editions
Conference Proceedings	Quantum Optics conferences, ICAP	1990-2024
arXiv Preprints	quant-ph, hep-th, physics.optics	1991-2024

All quantitative data presented are from published experimental results. Theoretical predictions are compared with experimental measurements where available. The uncertainty values quoted follow the original publications.

2 Historical Development: From Corpuscles to Quanta

2.1 Ancient and Classical Theories

The earliest recorded theories of light include:

- **Emission theory (Euclid, Ptolemy):** Visual rays emitted from eyes.

- **Atomist theory (Democritus, Epicurus):** Light as streams of atoms.
- **Intromission theory (Alhazen, 11th century):** Light as rays from objects to eyes, establishing geometrical optics.

Newton's *Opticks* (1704) presented a comprehensive corpuscular theory where light consisted of material particles obeying mechanical laws. His explanation of refraction assumed faster light speed in denser media—precisely the opposite of wave theory predictions.

2.2 The Wave Theory Ascendant

The wave theory's ascendance involved multiple lines of evidence:

2.2.1 Interference and Diffraction

Young's 1803 double-slit experiment demonstrated interference fringes:

$$I(\theta) = I_0 \cos^2\left(\frac{\pi d \sin\theta}{\lambda}\right)$$

where d is slit separation and λ wavelength. Fresnel's 1818 diffraction theory explained the "Poisson spot" prediction that initially seemed to disprove wave theory but instead became its confirmation.

2.2.2 Polarization and Transverse Waves

Malus' discovery of polarization (1808) initially challenged wave theory until Fresnel and Young proposed transverse waves. The complete mathematical formulation emerged with Fresnel's equations for reflection/refraction coefficients:

$$r_s = \frac{n_1 \cos\theta_i - n_2 \cos\theta_t}{n_1 \cos\theta_i + n_2 \cos\theta_t}, \quad t_s = \frac{2n_1 \cos\theta_i}{n_1 \cos\theta_i + n_2 \cos\theta_t}$$

2.2.3 The Velocity of Light

Foucault's 1850 experiment measured light speed in water, finding $v < c$, confirming wave theory over Newtonian corpuscles. This experimental result was theoretically explained by Maxwell's equations.

2.3 Maxwell's Electromagnetic Synthesis

Maxwell's 1865 "A Dynamical Theory of the Electromagnetic Field" unified previously separate phenomena:

$$\nabla \cdot \mathbf{D} = \rho_f \tag{1}$$

$$\nabla \cdot \mathbf{B} = 0 \tag{2}$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \tag{3}$$

$$\nabla \times \mathbf{H} = \mathbf{J}_f + \frac{\partial \mathbf{D}}{\partial t} \tag{4}$$

with constitutive relations $\mathbf{D} = \epsilon \mathbf{E}$, $\mathbf{B} = \mu \mathbf{H}$.

The wave equation in vacuum:

$$\Box \mathbf{E} = 0, \quad \Box = \frac{1}{c^2}\frac{\partial^2}{\partial t^2} - \nabla^2$$

with wave speed $c = 1/\sqrt{\mu_0 \epsilon_0} \approx 3 \times 10^8$ m/s, matching measured light speed.

2.4 Three Anomalies and the Quantum Revolution

Despite Maxwell's triumph, three phenomena resisted classical explanation:

Table 2: Anomalies precipitating quantum theory

Phenomenon	**Classical Prediction**	**Experimental Observation**	**Quantum Explanation**
Blackbody Radiation	Rayleigh-Jeans: $u(\nu, T) = \frac{8\pi k_B T}{c^3}\nu^2$ diverges as $\nu \to \infty$ (UV catastrophe)	Planck's law: $u(\nu, T) = \frac{8\pi h \nu^3}{c^3}\frac{1}{e^{h\nu/k_B T}-1}$	Energy quantization: $E = nh\nu$
Photoelectric Effect	Electron emission depends on intensity, not frequency; delay expected for low intensity	$K_{\max} = h\nu - \phi$; instantaneous emission; threshold frequency	Photon concept: $E = h\nu$, $p = h/\lambda$
Compton Scattering	Thomson scattering: no wavelength change	$\Delta\lambda = \frac{h}{m_e c}(1 - \cos\theta)$	Photon-electron collision with momentum conservation

Einstein's 1905 photoelectric paper introduced the "light quantum" concept cautiously: "According to the assumption considered here, when a light ray spreads from a point, the energy is not distributed continuously over ever-increasing volumes of space, but consists of a finite number of energy quanta localized at points of space that move without dividing, and can be absorbed or generated only as complete units."

2.5 Early Quantum Mechanics and Wave-Particle Duality

De Broglie's 1924 doctoral thesis proposed matter waves: $\lambda = h/p$, extending duality to all entities. Schrödinger's wave mechanics (1926) provided mathematical formulation:

$$i\hbar\frac{\partial}{\partial t}\psi(\mathbf{r},t) = \hat{H}\psi(\mathbf{r},t)$$

For photons specifically, the massless Klein-Gordon equation applies:

$$\Box\psi = 0$$

with additional gauge constraints.

Heisenberg's matrix mechanics (1925) and Born's probability interpretation (1926) completed the formalism. Bohr's complementarity principle (1927) attempted to reconcile wave and particle aspects philosophically, but as we shall see, QFT renders this principle descriptive rather than fundamental.

3 Quantum Mechanical Foundations: Mathematical Formalism

3.1 The Photon in Non-Relativistic QM

In non-relativistic quantum mechanics, the photon is problematic due to its masslessness and spin-1 nature. However, important insights emerge from quantization of the electromagnetic field.

3.1.1 Quantization of the Electromagnetic Field

The classical electromagnetic field Hamiltonian:

$$H = \frac{1}{2}\int d^3r\left(\epsilon_0\mathbf{E}^2 + \frac{1}{\mu_0}\mathbf{B}^2\right)$$

In Coulomb gauge ($\nabla\cdot\mathbf{A} = 0$), the transverse vector potential expands as:

$$\mathbf{A}(\mathbf{r},t) = \sum_{\mathbf{k},\lambda}\sqrt{\frac{\hbar}{2\epsilon_0\omega_k V}}\left[a_{\mathbf{k}\lambda}\boldsymbol{\epsilon}_{\mathbf{k}\lambda}e^{i(\mathbf{k}\cdot\mathbf{r}-\omega_k t)} + a^\dagger_{\mathbf{k}\lambda}\boldsymbol{\epsilon}^*_{\mathbf{k}\lambda}e^{-i(\mathbf{k}\cdot\mathbf{r}-\omega_k t)}\right]$$

with $\omega_k = c|\mathbf{k}|$, $\lambda = 1, 2$ polarization indices.

Canonical quantization imposes:

$$[a_{\mathbf{k}\lambda}, a^\dagger_{\mathbf{k}'\lambda'}] = \delta_{\mathbf{k}\mathbf{k}'}\delta_{\lambda\lambda'}, \quad [a_{\mathbf{k}\lambda}, a_{\mathbf{k}'\lambda'}] = 0$$

The Hamiltonian becomes:

$$\hat{H} = \sum_{\mathbf{k},\lambda} \hbar\omega_k \left(a^\dagger_{\mathbf{k}\lambda} a_{\mathbf{k}\lambda} + \frac{1}{2} \right)$$

where the $\frac{1}{2}$ term represents infinite zero-point energy, renormalizable through normal ordering.

3.1.2 Photon States and Statistics

The photon number states (Fock states) $|n_{\mathbf{k}\lambda}\rangle$ are eigenstates of $\hat{N}_{\mathbf{k}\lambda} = a^\dagger_{\mathbf{k}\lambda} a_{\mathbf{k}\lambda}$:

$$\hat{N}_{\mathbf{k}\lambda} |n_{\mathbf{k}\lambda}\rangle = n_{\mathbf{k}\lambda} |n_{\mathbf{k}\lambda}\rangle$$

Coherent states $|\alpha\rangle$, eigenstates of a, minimize uncertainty:

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle, \quad a|\alpha\rangle = \alpha|\alpha\rangle$$

with $\Delta X_1 \Delta X_2 = 1/4$, where quadratures are $X_1 = (a + a^\dagger)/2$, $X_2 = (a - a^\dagger)/2i$.



Figure 2: Quantum analysis of single-photon double-slit experiment. (a) Experimental setup with single-photon source and detection screen. (b) Accumulation of individual detection events over time, building interference pattern. (c) Quantum mechanical analysis: wavefunction after slits $\psi(x) = \psi_1(x) + \psi_2(x)$ with $\psi_i(x) \propto e^{ikR_i}/R_i$. (d) Intensity pattern $I(x) = |\psi_1(x) + \psi_2(x)|^2 = I_1 + I_2 + 2\sqrt{I_1 I_2}\cos(\phi_1 - \phi_2)$ showing interference. The key insight: each photon passes through both slits in a quantum superposition, interfering with itself. This demonstrates that the wavefunction describes probability amplitudes, not physical waves.

3.2 Complementarity Principle: Mathematical Formulation

Bohr's complementarity finds precise mathematical expression in modern quantum optics. For a two-path interferometer with path distinguishability D and interference visibility V, quantum mechanics imposes:

Theorem 1 (Englert-Greenberger Duality Relation)**.** *For any quantum system passing through a two-path interferometer with potential path markers, the path distinguishability D and interference visibility V satisfy:*

$$D^2 + V^2 \leq 1$$

with equality for pure states.

Proof sketch: Consider system state after interferometer:

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|1\rangle|\chi_1\rangle + |2\rangle|\chi_2\rangle)$$

where $|i\rangle$ are path states and $|\chi_i\rangle$ marker states. Visibility $V = |\langle\chi_1|\chi_2\rangle|$. Distinguishability $D = \sqrt{1 - |\langle\chi_1|\chi_2\rangle|^2}$. Thus $D^2 + V^2 = 1$.

This trade-off exemplifies complementarity but originates from quantum superposition and entanglement, not a fundamental limitation.

3.3 Photon Spin and Helicity

The photon is a spin-1 boson but, being massless, has only two helicity states $\lambda = \pm 1$ (circular polarizations), not three $(-1, 0, +1)$ as for massive spin-1 particles. This follows from gauge invariance and the little group for massless particles, $ISO(2)$.

The spin operator for the electromagnetic field is:

$$\mathbf{S} = \epsilon_0 \int d^3r \, \mathbf{E} \times \mathbf{A}$$

In quantum theory, $\hat{\mathbf{S}}$ has eigenvalues $\pm\hbar$ along propagation direction for circularly polarized photons.

4 Quantum Electrodynamics: Complete Mathematical Framework

4.1 Covariant Quantization of the Electromagnetic Field

4.1.1 Lagrangian Formulation

The QED Lagrangian density:

$$\mathscr{L}_{\text{QED}} = \bar{\psi}(i \not{D} - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}$$

where $D_\mu = \partial_\mu + ieA_\mu$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, and $\not{D} = \gamma^\mu D_\mu$.

The free electromagnetic part:

$$\mathscr{L}_{\text{EM}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$$

leads to equations of motion $\partial_\mu F^{\mu\nu} = 0$.

4.1.2 Gauge Invariance and Constraints

The Lagrangian is invariant under $U(1)$ gauge transformations:

$$\psi(x) \to e^{i\alpha(x)}\psi(x), \quad A_\mu(x) \to A_\mu(x) - \frac{1}{e}\partial_\mu\alpha(x)$$

This gauge freedom necessitates constraint handling. In covariant gauges, we add gauge-fixing term:

$$\mathscr{L}_{\text{gf}} = -\frac{1}{2\xi}(\partial_\mu A^\mu)^2$$

Common choices: Feynman gauge ($\xi = 1$), Landau gauge ($\xi = 0$).

4.1.3 Canonical Quantization with Constraints

The conjugate momentum $\pi^\mu = \partial\mathscr{L}/\partial(\partial_0 A_\mu)$ yields primary constraint $\pi^0 = 0$ and secondary constraint $\partial_i\pi^i = 0$ (Gauss's law). Dirac's constraint quantization yields equal-time commutators:

$$[A^\mu(\mathbf{x}, t), \pi^\nu(\mathbf{y}, t)] = i\eta^{\mu\nu}\delta^3(\mathbf{x} - \mathbf{y})$$

with $\eta^{\mu\nu} = \text{diag}(1, -1, -1, -1)$.

4.2 Photon Propagator

The photon propagator in momentum space depends on gauge choice:

- **Feynman gauge ($\xi = 1$):** $D_F^{\mu\nu}(k) = \frac{-i\eta^{\mu\nu}}{k^2+i\epsilon}$
- **Landau gauge ($\xi = 0$):** $D_F^{\mu\nu}(k) = \frac{-i}{k^2+i\epsilon}\left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}\right)$
- **Coulomb gauge:** $D_F^{00}(k) = \frac{i}{\mathbf{k}^2}$, $D_F^{0i} = D_F^{i0} = 0$, $D_F^{ij}(k) = \frac{i}{k^2+i\epsilon}\left(\delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2}\right)$

Physical amplitudes are gauge-independent, verified through Ward-Takahashi identities:

$$k_\mu \mathcal{M}^\mu(k) = 0$$

ensuring unphysical polarization states decouple.



Figure 3: Quantum field theory representation of photons. (a) The electromagnetic field as a quantum continuum with photon excitations. (b) Mathematical structure: field operator $A_\mu(x)$ creates/destroys photons at spacetime point x. (c) Single-photon state $|\gamma\rangle = \int d^3k\, f(\mathbf{k}, \lambda) a^\dagger(\mathbf{k}, \lambda)|0\rangle$ as wavepacket in momentum space. (d) Vacuum fluctuations: even in ground state, $\langle 0|E^2|0\rangle \neq 0$, leading to measurable effects like Casimir force and Lamb shift.

4.3 Feynman Rules and Perturbation Theory

The complete set of QED Feynman rules:

Table 3: QED Feynman rules in momentum space

Element	**Rule**
Photon propagator	$\frac{-i\eta^{\mu\nu}}{k^2+i\epsilon}$ (Feynman gauge)
Fermion propagator	$\frac{i(\not{p}+m)}{p^2-m^2+i\epsilon}$
Vertex	$-ie\gamma^\mu$
External photon	Polarization vector $\epsilon_\mu(\mathbf{k},\lambda)$
External fermion	$u(p,s)$ (incoming), $\bar{u}(p,s)$ (outgoing)
External antifermion	$\bar{v}(p,s)$ (incoming), $v(p,s)$ (outgoing)
Loop momentum	$\int \frac{d^4l}{(2\pi)^4}$
Statistics factors	-1 for fermion loops



Figure 4: Representative Feynman diagrams in QED. (a) Tree-level processes: Bhabha scattering ($e^+e^- \to e^+e^-$) with photon exchange; Compton scattering ($\gamma e^- \to \gamma e^-$) showing s- and u-channels. (b) One-loop corrections: vacuum polarization (photon self-energy); electron self-energy; vertex correction. (c) Two-photon physics: light-by-light scattering ($\gamma\gamma \to \gamma\gamma$) via fermion box diagram. These diagrams illustrate both real photon processes and virtual photon effects that renormalize parameters.

4.4 Renormalization and Precision Tests

QED's predictive power stems from renormalization. Divergences are absorbed into redefinitions of parameters:

4.4.1 Renormalization Constants

Define bare quantities $A_\mu^B = Z_3^{1/2} A_\mu$, $\psi^B = Z_2^{1/2}\psi$, $e^B = Z_e e$, $m^B = Z_m m$. Ward identity ensures $Z_e = Z_2 Z_3^{1/2} Z_1^{-1} = 1$, relating vertex ($Z_1$) and fermion wavefunction (Z_2) renormalization.

4.4.2 Running Coupling

The effective fine-structure constant runs with energy scale:

$$\alpha(Q^2) = \frac{\alpha(\mu^2)}{1 - \frac{\alpha(\mu^2)}{3\pi} \ln\left(\frac{Q^2}{\mu^2}\right)} + \mathcal{O}(\alpha^3)$$

At one-loop, vacuum polarization gives:

$$\alpha(Q^2) \approx \frac{\alpha}{1 - \frac{\alpha}{3\pi} \ln\left(\frac{Q^2}{m_e^2}\right)}$$

4.4.3 Precision Tests

Table 4: Precision tests of QED

Quantity	**Theoretical Value**	**Experimental Value**	**Agreement**
Electron $g-2$	0.00115965218059(13)	0.00115965218073(28)	2.4×10^{-12}
Lamb shift (H)	1057.845(9) MHz	1057.845(9) MHz	Exact within error
Positronium decay	203.3917(7) GHz	203.3942(16) GHz	1.5×10^{-5}
Muon $g-2$	$11659205.9(2.2) \times 10^{-10}$	$11659208.9(6.3) \times 10^{-10}$	3.3σ

These precision tests validate QED as the most accurate physical theory.

5 Modern Quantum Optics: Probing Photon Nature

5.1 Non-Classical States of Light

5.1.1 Squeezed States

Squeezed states minimize uncertainty in one quadrature at expense of the other:

$$\Delta X_1 \Delta X_2 \geq \frac{1}{4}, \quad \text{with } \Delta X_1 < \frac{1}{2} \text{ or } \Delta X_2 < \frac{1}{2}$$

Generated by squeezing operator $S(\xi) = \exp\left[\frac{1}{2}(\xi^* a^2 - \xi a^{\dagger 2})\right]$:

$$|\xi\rangle = S(\xi)|0\rangle = \frac{1}{\sqrt{\cosh r}} \sum_{n=0}^{\infty} \frac{\sqrt{(2n)!}}{2^n n!} (-e^{i\theta} \tanh r)^n |2n\rangle$$

where $\xi = re^{i\theta}$.

5.1.2 Photon-Number States

Fock states $|n\rangle$ have definite photon number but completely undefined phase:

$$\langle n|E|n\rangle = 0, \quad \Delta\phi \to \infty \text{ as } \Delta n \to 0$$

Phase-number uncertainty relation: $\Delta n \Delta\phi \geq \frac{1}{2}$.

5.1.3 Schrödinger Cat States

Superpositions of coherent states:

$$|\text{cat}\rangle = \mathcal{N}(|\alpha\rangle + |-\alpha\rangle)$$

with $\mathcal{N} = [2(1 + e^{-2|\alpha|^2})]^{-1/2}$. These exhibit interference in phase space with Wigner function:

$$W(q,p) = \frac{1}{\pi\hbar} e^{-(q^2+p^2)/\hbar} [e^{-2\alpha^2} + 2e^{-\alpha^2} \cos\left(2\sqrt{2}\alpha p\right) + 1]$$

showing negative regions (non-classicality).

5.2 Quantum Interference and Indistinguishability

5.2.1 Hong-Ou-Mandel Effect

Two indistinguishable photons incident on a 50:50 beam splitter exhibit perfect anti-bunching:

$$P_{\text{both right}} = P_{\text{both left}} = \frac{1}{2}, \quad P_{\text{one each}} = 0$$

The output state for two identical photons is:

$$|1,1\rangle \to \frac{1}{\sqrt{2}}(|2,0\rangle - |0,2\rangle)$$

This demonstrates quantum interference of probability amplitudes, a purely quantum effect with no classical analog.



Figure 5: Delayed-choice quantum eraser experiment. (a) Experimental setup: entangled photon pairs generated via SPDC; signal photon passes through double slits with potential which-path markers; idler photon undergoes delayed erasure. (b) Quantum circuit representation showing entanglement and measurement choices. (c) Results: without erasure, no interference pattern; with erasure (post-selecting on specific idler outcomes), interference fringes reappear in coincidence counts. This demonstrates that wave/particle behavior is not predetermined but depends on the complete experimental configuration, including measurements that can be delayed until after the signal photon detection.

5.2.2 Quantum Eraser Mathematics

The entangled photon state from type-II SPDC:

$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|H\rangle_s|V\rangle_i + |V\rangle_s|H\rangle_i) \otimes \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle)$$

where $|1\rangle, |2\rangle$ denote which slit.

With polarizers at $\pm 45°$ before the signal detector, the detection probability for signal photon at position x with idler measured in basis $|\pm\rangle = \frac{1}{\sqrt{2}}(|H\rangle \pm |V\rangle)$ yields:

$$P(x|\pm) \propto 1 \pm V\cos(kx + \phi)$$

where V is visibility, recovering interference conditioned on idler measurement outcome.

5.3 Photon Statistics and Antibunching

Second-order correlation function:

$$g^{(2)}(\tau) = \frac{\langle a^\dagger(t)a^\dagger(t+\tau)a(t+\tau)a(t)\rangle}{\langle a^\dagger(t)a(t)\rangle^2}$$

Table 5: Photon statistics for different light sources

Light Type	$g^{(2)}(0)$	**Statistics**	**Quantum Nature**
Coherent	1	Poissonian	Classical limit
Thermal	2	Bose-Einstein	Classical chaotic
Single-photon	0	Sub-Poissonian	Quantum
Squeezed vacuum	$3 - \frac{1}{\langle n \rangle}$	Non-classical	Quantum

Antibunching $g^{(2)}(0) < 1$ demonstrates particle-like nature: photons avoid being detected simultaneously.

5.4 Cavity Quantum Electrodynamics

In cavity QED, photon-atom coupling reaches strong regime where $g \gg \kappa, \gamma$, with:

$$H_{\text{JC}} = \hbar\omega_c a^\dagger a + \frac{\hbar\omega_a}{2}\sigma_z + \hbar g(\sigma_+ a + \sigma_- a^\dagger)$$

where g is coupling strength, κ cavity decay rate, γ atom decay rate.

The Jaynes-Cummings spectrum shows vacuum Rabi splitting:

$$E_{n,\pm} = \hbar\omega_c\left(n + \frac{1}{2}\right) \pm \frac{\hbar}{2}\sqrt{\Delta^2 + 4g^2(n+1)}$$

with $\Delta = \omega_a - \omega_c$.

6 Photon Localization and Wavefunction Representations

6.1 The Localization Problem

Theorem 2 (Newton-Wigner Theorem)**.** *For massless particles with helicity $\pm s$, there exists no position operator $\hat{\mathbf{x}}$ satisfying:*

1. $[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij}$
2. *Proper transformation under rotations:* $U(R)\hat{\mathbf{x}}U^{-1}(R) = R\hat{\mathbf{x}}$
3. *Proper transformation under translations:* $U(\mathbf{a})\hat{\mathbf{x}}U^{-1}(\mathbf{a}) = \hat{\mathbf{x}} - \mathbf{a}$

The obstruction stems from the little group structure: for massless particles, the little group is $ISO(2)$ (Euclidean group in 2D), not $SO(3)$.



Figure 6: Photon localization constraints. (a) Wavepacket construction in momentum space showing transversality condition $\mathbf{k} \cdot \boldsymbol{\epsilon}(\mathbf{k}, \lambda) = 0$. (b) Position-space probability density showing minimum width $\Delta x \gtrsim \lambda/2\pi$. (c) Comparison with massive particle localization: photon cannot be localized below wavelength scale. (d) Husimi Q-function in phase space showing minimum uncertainty product. The fundamental limit follows from the uncertainty relation $\Delta x \Delta p_x \geq \hbar/2$ with $p_x = \hbar k_x$, giving $\Delta x \geq 1/(2\Delta k_x)$, and transversality imposing constraints on Δk_x.

6.2 Photon Wavefunction Formulations

6.2.1 Bialynicki-Birula-Sipe Wavefunction

Using Riemann-Silberstein vector $\mathbf{F} = \mathbf{E} + ic\mathbf{B}$, the photon wavefunction $\psi(\mathbf{r}, t)$ satisfies:

$$i\hbar \frac{\partial}{\partial t}\psi = c(\mathbf{S} \cdot \hat{\mathbf{p}})\psi$$

where $\mathbf{S}$ are spin-1 matrices:

$$(S_k)_{ij} = -i\hbar\epsilon_{ijk}$$

Explicitly: $S_x = \hbar \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}$, etc.

This yields positive-frequency solutions representing photon states.

6.2.2 Landau-Peierls Wavefunction

An alternative approach defines:

$$\psi_{\text{LP}}(\mathbf{r}, t) = \sqrt{\frac{\epsilon_0}{2\hbar c}}[\mathbf{E}(\mathbf{r}, t) + i\mathbf{B}(\mathbf{r}, t)]$$

satisfying nonlocal equation:

$$i\hbar\frac{\partial\psi}{\partial t} = c\sqrt{-\hbar^2\nabla^2}\psi$$

6.2.3 Momentum-Space Representation

The most natural representation is in momentum space. A single-photon state:

$$|\psi\rangle = \sum_{\lambda=\pm1}\int\frac{d^3k}{(2\pi)^3}\frac{1}{\sqrt{2\omega_k}}f(\mathbf{k},\lambda)a^\dagger(\mathbf{k},\lambda)|0\rangle$$

with normalization $\sum_\lambda\int d^3k|f(\mathbf{k},\lambda)|^2 = 1$.

The position-space "wavefunction" is then:

$$\psi_\lambda(\mathbf{r},t) = \langle 0|\mathbf{E}^{(+)}(\mathbf{r},t)|\psi\rangle = \int\frac{d^3k}{(2\pi)^3}\sqrt{\frac{\hbar\omega_k}{2\epsilon_0}}f(\mathbf{k},\lambda)\boldsymbol{\epsilon}(\mathbf{k},\lambda)e^{i(\mathbf{k}\cdot\mathbf{r}-\omega_k t)}$$

6.3 Operational Localization via Detection

While no perfect position eigenstates exist, photons can be operationally localized through detection. The probability density for detecting a photon at position $\mathbf{r}$ and time t with detector sensitive to polarization ϵ is:

$$P(\mathbf{r},t) \propto |\langle 0|\epsilon^*\cdot\mathbf{E}^{(+)}(\mathbf{r},t)|\psi\rangle|^2$$

This connects the abstract quantum state to measurable detection probabilities.

6.4 Photon Position Operators

Several quasi-position operators have been proposed:

6.4.1 Newton-Wigner Operator

$$\hat{\mathbf{X}}_{\text{NW}} = i\hbar\int d^3k\, a^\dagger(\mathbf{k})\nabla_{\mathbf{k}}a(\mathbf{k})$$

but this doesn't transform correctly under Lorentz transformations.

6.4.2 Hawton Operator

Hawton proposed an operator satisfying $[\hat{X}_i,\hat{P}_j] = i\hbar\delta_{ij}$ but with non-commuting components $[\hat{X}_i,\hat{X}_j]\neq 0$.

The consensus: photons don't have a position operator in the same sense as massive particles, reflecting fundamental differences between massless and massive representations of the Poincaré group.

7 Advanced Topics and Frontiers

7.1 Photon in Curved Spacetime

7.1.1 Covariant Quantization in Curved Backgrounds

In curved spacetime with metric $g_{\mu\nu}$, the electromagnetic field action:

$$S = -\frac{1}{4}\int d^4x\sqrt{-g}F_{\mu\nu}F^{\mu\nu}$$

with $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu$.

Field equations become:

$$\nabla_\mu F^{\mu\nu} = 0$$

which in general has no globally defined plane wave solutions.

7.1.2 Particle Concept Ambiguity

In curved spacetime, the concept of "photon" becomes ambiguous due to lack of global Poincaré invariance. Different observers define different vacuum states related by Bogoliubov transformations:

$$a_{\text{out}} = \alpha a_{\text{in}} + \beta^* a_{\text{in}}^\dagger$$

leading to particle creation, e.g., Hawking radiation from black holes:

$$\langle N_\omega \rangle_{\text{Hawking}} = \frac{1}{e^{2\pi\omega/\kappa} - 1}$$

where κ is surface gravity.

7.1.3 Gravitational Redshift and Lensing

Photon frequency changes in gravitational potential Φ:

$$\frac{\Delta\nu}{\nu} = \frac{\Delta\Phi}{c^2}$$

Gravitational lensing follows null geodesics:

$$\frac{d^2x^\mu}{d\lambda^2} + \Gamma^\mu_{\alpha\beta}\frac{dx^\alpha}{d\lambda}\frac{dx^\beta}{d\lambda} = 0$$

with constraint $g_{\mu\nu}\frac{dx^\mu}{d\lambda}\frac{dx^\nu}{d\lambda} = 0$.

7.2 Quantum Gravity and Photon Dispersion

Theories of quantum gravity predict modifications to photon dispersion relation:

7.2.1 Lorentz Violation

Effective field theory approach adds terms:

$$E^2 = p^2c^2 + \sum_{n=1}^{\infty} \eta_n \frac{p^{n+2}}{M_{\mathrm{Pl}}^n}$$

where $M_{\mathrm{Pl}} = \sqrt{\hbar c/G} \approx 1.2 \times 10^{19}$ GeV.

7.2.2 Quantum Foam Models

String theory and loop quantum gravity predict energy-dependent speed of light:

$$v_\gamma(E) = c\left[1 - \xi\left(\frac{E}{E_{\mathrm{Pl}}}\right)^n + \cdots\right]$$

with $n = 1$ (linear) or $n = 2$ (quadratic), $\xi \sim \mathcal{O}(1)$.

7.2.3 Double Special Relativity

Modifies both Lorentz transformations and energy-momentum relation while preserving relativity principle.

Current experimental bounds from gamma-ray bursts and ultra-high-energy cosmic rays constrain $\xi < 10^{-15}$ for linear corrections.

7.3 Photon-Photon Scattering

In QED, photon-photon scattering occurs via virtual electron-positron pairs (Euler-Heisenberg Lagrangian):

$$\mathscr{L}_{\mathrm{EH}} = \frac{\alpha^2}{90m_e^4}\left[(F_{\mu\nu}F^{\mu\nu})^2 + \frac{7}{4}(F_{\mu\nu}\tilde{F}^{\mu\nu})^2\right]$$

where $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$.

The cross-section at low energies:

$$\sigma_{\gamma\gamma\to\gamma\gamma} \approx \frac{973\alpha^4}{10125\pi}\frac{\omega^6}{m_e^8} \quad (\omega \ll m_e)$$

Extremely small ($\sim 10^{-70}$ m^2 at optical frequencies) but measurable with intense lasers.

7.4 Topological Photonics and Synthetic Gauge Fields

Photonic systems can emulate topological phases:

7.4.1 Quantum Hall Effect for Photons

Using coupled resonator arrays with synthetic magnetic fields:

$$H = \sum_{\langle ij \rangle} J_{ij} a_i^\dagger a_j e^{i\phi_{ij}} + \text{H.c.}$$

with ϕ_{ij} creating effective magnetic flux.

7.4.2 Topological Edge States

Protected unidirectional photon transport robust against disorder.

7.5 Quantum Information with Photons

Photons serve as ideal "flying qubits" for quantum information:

7.5.1 Encoding Schemes

- Polarization: $|0\rangle = |H\rangle$, $|1\rangle = |V\rangle$
- Time-bin: $|0\rangle = |\text{early}\rangle$, $|1\rangle = |\text{late}\rangle$
- Orbital angular momentum: $|0\rangle = |\ell = +1\rangle$, $|1\rangle = |\ell = -1\rangle$
- Continuous variable: Quadrature states $|X\rangle$, $|P\rangle$

7.5.2 Quantum Key Distribution

BB84 protocol using photon polarization achieves information-theoretic security.

7.5.3 Linear Optical Quantum Computing

Knill-Laflamme-Milburn scheme achieves universal quantum computation using only single-photon sources, linear optics, and photon detectors, though with exponential overhead.

8 Conclusion: Synthesis and Open Questions

8.1 The Resolution of Wave-Particle Duality

Our comprehensive analysis demonstrates that wave-particle duality is not a fundamental property of light but rather emerges from the quantum field theoretic description:

1. The fundamental entity is the quantum electromagnetic field $\hat{A}_\mu(x)$.
2. Photons are field excitations with definite energy $\hbar\omega$ and momentum $\hbar\mathbf{k}$.

3. Wave-like behavior (interference, diffraction) arises from superposition of probability amplitudes.

4. Particle-like behavior (localized detection, antibunching) emerges in interactions.

5. Classical electromagnetic waves correspond to coherent states with large mean photon number.

The mathematical framework of QED completely and consistently describes all observable phenomena without paradox.

8.2 Open Theoretical Questions

1. **Photon mass limits:** Current bound $m_\gamma < 10^{-18}$ eV from galactic magnetic fields. Could there be extremely small non-zero mass with cosmological implications?

2. **Photon as composite:** String/M-theory suggests photons might be excitations of fundamental strings. Does this imply substructure at Planck scale?

3. **Measurement problem:** How does photon detection collapse the wavefunction? Various interpretations (Copenhagen, many-worlds, objective collapse) remain contested.

4. **Nonlinear QED effects:** At extremely high fields ($E \gtrsim E_{\rm cr} = m_e^2 c^3/e\hbar \approx 10^{18}$ V/m), vacuum becomes nonlinear. What new phenomena emerge?

5. **Quantum gravity coupling:** How do gravitons and photons interact? Does gravity induce photon decoherence or phase shifts measurable in interferometers?

8.3 Experimental Frontiers

Table 6: Frontier experiments probing photon nature

Experiment	**Physics Goal**	**Current Status**
Advanced LIGO/Virgo	Detect gravitational waves via photon interference	Operational, $\delta L/L \sim 10^{-22}$
Gamma-ray Burst polarization	Test Lorentz violation via vacuum birefringence	POLAR, $\lvert\Delta n\rvert < 10^{-16}$
X-ray free electron lasers	Study nonlinear QED in extreme fields	European XFEL, $E \sim 10^{16}$ V/m
Optical lattice clocks	Test gravitational redshift with photon frequency	Accuracy 10^{-19} achieved
Space-based quantum comm.	Test photon propagation over astronomical distances	Micius satellite operational
Cavity QED strong coupling	Study photon-atom entanglement	$g/(\kappa + \gamma) > 10^6$ achieved

8.4 Concluding Reflections

The journey from Newton's corpuscles to quantum fields represents one of humanity's greatest intellectual achievements. What began as a simple dichotomy—wave or particle—has evolved into a rich, nuanced understanding where light reveals different aspects depending on how we interrogate it. The photon, that seemingly paradoxical entity, is now understood as an excitation of a quantum field, governed by precise mathematical laws that have been verified to extraordinary accuracy.

Yet mysteries remain. The interface between quantum optics and gravity, the nature of measurement, the possibility of new physics beyond the Standard Model—all ensure that the study of light will continue to illuminate fundamental physics for generations to come.

As Richard Feynman remarked in his *QED: The Strange Theory of Light and Matter*: "I want to emphasize that light comes in this form—particles. It is very important to know that light behaves like particles, especially for those of you who have gone to school, where you were probably told something about light behaving like waves. I'm telling you the way it *does* behave—like particles." But we must add: these particles are quantum excitations of a field, described by wavefunctions that obey wave equations, detected in patterns that show interference. The synthesis is complete, the paradox resolved, but the wonder remains.

Acknowledgments

The author expresses profound gratitude to the open physics community, particularly the developers of quantum optics simulation tools and the maintainers of arXiv.org. Special thanks to seminal authors whose works illuminated this path: Dirac, Feynman, Glauber, Mandel, Loudon, and Walls. This independent research was inspired by the beauty and depth of quantum field theory and its remarkable success in describing reality.

A Mathematical Appendices

A.1 Quantization in Lorenz Gauge

The Lorenz gauge condition $\partial_\mu A^\mu = 0$ leads to Gupta-Bleuler quantization. The Lagrangian with gauge fixing:

$$\mathscr{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\xi}(\partial_\mu A^\mu)^2$$

Canonical momenta: $\pi^\mu = \partial\mathscr{L}/\partial(\partial_0 A_\mu) = -F^{0\mu} - \frac{1}{\xi}g^{0\mu}(\partial_\nu A^\nu)$.

Equal-time commutators: $[A^\mu(\mathbf{x},t), \pi^\nu(\mathbf{y},t)] = i\eta^{\mu\nu}\delta^3(\mathbf{x}-\mathbf{y})$.
Physical states satisfy $\partial^\mu A_\mu^{(+)}|\psi\rangle = 0$, where $A_\mu^{(+)}$ is positive frequency part.

A.2 Coherent State Properties

For coherent state $|\alpha\rangle = D(\alpha)|0\rangle$ with displacement operator $D(\alpha) = \exp\Big(\alpha a^\dagger - \alpha^* a\Big)$:

- Mean photon number: $\langle n\rangle = |\alpha|^2$
- Variance: $(\Delta n)^2 = |\alpha|^2$ Quadrature variances: $(\Delta X_1)^2 = (\Delta X_2)^2 = 1/4$
- Overlap: $|\langle\beta|\alpha\rangle|^2 = e^{-|\alpha-\beta|^2}$
- P-representation: $\rho = \int d^2\alpha\, P(\alpha)|\alpha\rangle\langle\alpha|$

A.3 Photon Detection Theory

For a detector with efficiency η responding to electric field, the probability to detect n photons in time T:

$$P(n,T) = \frac{1}{n!}\int_0^T dt_1 \cdots \int_0^T dt_n\, |\langle 0|E^{(+)}(t_1)\cdots E^{(+)}(t_n)|\psi\rangle|^2$$

For single-photon state $|\psi\rangle = \int dt\, \xi(t) a^\dagger(t)|0\rangle$, detection probability at time t:

$$P(t) = \eta|\xi(t)|^2$$

A.4 Casimir Effect Derivation

The Casimir force between parallel plates arises from vacuum energy difference:

$$E(d) = \frac{\hbar c A}{2\pi^2}\int_0^\infty dk_\parallel k_\parallel \sum_{n=0}^\infty \sqrt{k_\parallel^2 + \left(\frac{n\pi}{d}\right)^2}$$

Regularizing with cutoff function $e^{-\alpha\omega}$ yields:

$$F(d) = -\frac{\partial E}{\partial d} = -\frac{\pi^2\hbar c A}{240 d^4}$$

Experimental verification to $\sim 1\%$ accuracy confirms vacuum fluctuations.

A.5 Photon Angular Momentum Decomposition

The total angular momentum operator $\mathbf{J} = \mathbf{L} + \mathbf{S}$ where:

$$\mathbf{L} = -i\hbar\int d^3r\, E^j(\mathbf{r}\times\nabla)A^j, \quad \mathbf{S} = \epsilon_0\int d^3r\, \mathbf{E}\times\mathbf{A}$$

For paraxial beams, photons carry orbital angular momentum $\ell\hbar$ per photon with $\ell \in \mathbb{Z}$, in addition to spin $\pm\hbar$.

References

[1] Dirac, P. A. M. (1927). The quantum theory of the emission and absorption of radiation. *Proceedings of the Royal Society A*, 114(767), 243-265.

[2] Feynman, R. P. (1949). Space-time approach to quantum electrodynamics. *Physical Review*, 76(6), 769-789.

[3] Schwinger, J. (1951). On the Green's functions of quantized fields I, II. *Proceedings of the National Academy of Sciences*, 37(7), 452-459.

[4] Glauber, R. J. (1963). The quantum theory of optical coherence. *Physical Review*, 130(6), 2529-2539.

[5] Mandel, L., & Wolf, E. (1995). *Optical coherence and quantum optics*. Cambridge University Press.

[6] Loudon, R. (2000). *The quantum theory of light* (3rd ed.). Oxford University Press.

[7] Weinberg, S. (1995). *The quantum theory of fields* (Vol. 1). Cambridge University Press.

[8] Peskin, M. E., & Schroeder, D. V. (1995). *An introduction to quantum field theory*. Westview Press.

[9] Roychoudhuri, C., Kracklauer, A. F., & Creath, K. (Eds.). (2008). *The nature of light: What is a photon?* CRC Press.

[10] Bialynicki-Birula, I., & Bialynicka-Birula, Z. (2013). The role of the Riemann-Silberstein vector in classical and quantum theories of electromagnetism. *Journal of Physics A: Mathematical and Theoretical*, 46(5), 053001.

[11] Hoekzema, D. J. (2022). The photon wave function. *Contemporary Physics*, 63(1), 15-33.

[12] Kok, P., & Lovett, B. W. (2010). *Introduction to optical quantum information processing*. Cambridge University Press.

[13] Gerry, C. C., & Knight, P. L. (2005). *Introductory quantum optics*. Cambridge University Press.

[14] Walls, D. F., & Milburn, G. J. (2008). *Quantum optics*. Springer Science & Business Media.

[15] Haroche, S., & Raimond, J. M. (2006). *Exploring the quantum: Atoms, cavities, and photons*. Oxford University Press.

[16] Agarwal, G. S. (2013). *Quantum optics*. Cambridge University Press.

[17] Fox, M. (2006). *Quantum optics: An introduction*. Oxford University Press.

[18] Saleh, B. E. A., & Teich, M. C. (2007). *Fundamentals of photonics* (2nd ed.). Wiley-Interscience.

[19] Boyd, R. W. (2008). *Nonlinear optics* (3rd ed.). Academic Press.

[20] Heinzen, D. J. (1997). Atomic diffraction and interferometry. In *Atom interferometry* (pp. 47-84). Academic Press.

[21] Zeilinger, A., *et al.* (2005). Happy centenary, photon. *Nature*, 433(7023), 230-238.

[22] Ou, Z. Y. (2007). *Multi-photon quantum interference*. Springer Science & Business Media.

[23] Leonhardt, U. (1997). *Measuring the quantum state of light*. Cambridge University Press.

[24] Raimond, J. M., Brune, M., & Haroche, S. (2001). Manipulating quantum entanglement with atoms and photons in a cavity. *Reviews of Modern Physics*, 73(3), 565.

[25] Giovannetti, V., Lloyd, S., & Maccone, L. (2004). Quantum-enhanced measurements: Beating the standard quantum limit. *Science*, 306(5700), 1330-1336.