

Projected shoot area (a.u.) under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




Fv/Fm under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




Fq'/Fm' under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




NPQ under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




qP under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




qN under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




qL under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




qI under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




ΦNO under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




ΦNPQ under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




NPQ(t) under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.




Chlorophyll index under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.



Anthocyanin reflectance index under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.



NDVI under a drought gradient, by Mimulus population group

Mean ± SE per watering regime. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD;
each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05.
Series starts at 144 h, the projected-shoot-area trough in all four groups; earlier timepoints are spline-start noise on very small plants.
Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants.

