

Projected shoot area (a.u.) under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




Fv/Fm under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




Fq'/Fm' under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




NPQ under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




qP under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.



qN under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.



qL under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




qI under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




ΦNO under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




ΦNPQ under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




NPQ(t) under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.




Chlorophyll index under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.



Anthocyanin reflectance index under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.



NDVI under a drought gradient, by Mimulus maternal line

Mean ± SE per watering regime, one panel per maternal line; rows are the population group, columns the lines within it. Lower strip: one-way ANOVA across the five regimes at every timepoint, then Tukey HSD; each line is one regime against 100% of optimal watering, labelled by that regime. Red line = p 0.05. Only 3-5 plants per regime per line, so these tests are much weaker than the pooled-group version. Series starts at 144 h. Grey dashed line at 409.5 h: Exp 20243 (left) and Exp 20244 (right) are separate plant cohorts, so the series is not continued on the same plants. Oregon 2014 has two lines, so its third column is blank.

