

An analytic regularisation scheme on curved space–times with applications to cosmological space–times

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Class. Quantum Grav. 33 095009

(<http://iopscience.iop.org/0264-9381/33/9/095009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.251.60.15

This content was downloaded on 14/04/2016 at 16:52

Please note that [terms and conditions apply](#).

An analytic regularisation scheme on curved space-times with applications to cosmological space-times

Antoine G  re¹, Thomas-Paul Hack^{1,3} and Nicola Pinamonti^{1,2}

¹ Dipartimento di Matematica, Universit   di Genova, Via Dodecaneso 35, I-16146 Genova, Italy

² Istituto Nazionale di Fisica Nucleare, Sezione di Genova, Via Dodecaneso, 33 I-16146 Genova, Italy

E-mail: gere@dima.unige.it, hack@dima.unige.it and pinamont@dima.unige.it

Received 27 September 2015

Accepted for publication 18 January 2016

Published 13 April 2016



CrossMark

Abstract

We develop a renormalisation scheme for time-ordered products in interacting field theories on curved space-times that consists of an analytic regularisation of Feynman amplitudes and a minimal subtraction of the resulting pole parts. This scheme is directly applicable to space-times with Lorentzian signature, manifestly generally covariant, invariant under any space-time isometries present, and constructed to all orders in perturbation theory. Moreover, the scheme correctly captures the nongeometric state-dependent contribution of Feynman amplitudes, and it is well suited for practical computations. To illustrate this last point, we compute explicit examples on a generic curved space-time and demonstrate how momentum space computations in cosmological space-times can be performed in our scheme. In this work, we discuss only scalar fields in four space-time dimensions, but we argue that the renormalisation scheme can be directly generalised to other space-time dimensions and field theories with higher spin as well as to theories with local gauge invariance.

Keywords: quantum field theory on curved space-times, perturbation theory, renormalisation

1. Introduction

In the perturbative construction of models in quantum field theory on curved space-times, one encounters time-ordered products of field polynomials that are *a priori* ill defined due to the

³ Author to whom any correspondence should be addressed.

appearance of UV divergences. Several renormalisation schemes that deal with these divergences in the presence of nontrivial space–time curvature have been discussed in the literature, such as for example, local momentum space methods [Bu81], dimensional regularisation in combination with heat kernel techniques [Lü82, To82], differential renormalisation [CHL95, Pr97], ζ -function renormalisation [BF13], generic Epstein–Glaser renormalisation [BF00, HW01, HW04], and, on cosmological space–times, Mellin–Barnes techniques [Ho10] and dimensional regularisation with respect to the comoving spatial coordinates [BCK10].

Some of these schemes, such as heat–kernel approaches, ζ -function techniques, and local momentum space methods are based on constructions that are initially only well defined for space–times with Euclidean signatures. These constructions can be partly transported to general Lorentzian space–times by the local Wick-rotation techniques developed in [Mo99]. However, whereas, the Feynman propagator is essentially unique on Euclidean space–times, this is not the case on Lorentzian space–times where this propagator has a nonunique contribution depending on the quantum state of the field model. Consequently, the Euclidean renormalisation techniques and the numerous practical computations already performed by means of these methods—see, for example, the monographs [BD82, BOS92, PT09]—are able to capture the correct divergent and geometric parts of Feynman amplitudes but *a priori* not their nongeometric and state-dependent contributions.

A renormalisation scheme that is directly applicable to curved space–times with Lorentzian signature has been developed in [BF00, HW01, HW04] in the framework of algebraic quantum field theory. This scheme implements ideas of [EG73, St71] and is based on microlocal techniques that replace the momentum space methods available in Minkowski space–time and have been introduced to quantum field theory in curved space–time by the seminal work [Ra96]. However, although the generalised Epstein–Glaser scheme developed in [BF00, HW01, HW04] is conceptually clear and mathematically rigorous, it is not easily applicable in practical computations. On the other hand, Lorentzian schemes that are better suited for this purpose have not been developed to all orders in perturbation theory [CHL95, Pr97], are tailored to specific space–times [Ho10], or are not manifestly covariant [BCK10].

Motivated by this, we develop a renormalisation scheme for time-ordered products in interacting field theories on curved space–times that is directly applicable to space–times with Lorentzian signature, manifestly generally covariant, invariant under any space–time isometries present, and constructed to all orders in perturbation theory. Moreover, the scheme correctly captures the nongeometric state-dependent contribution of the Feynman amplitudes, and it is well suited for practical computations. In this work, we discuss only scalar fields in four space–time dimensions, but we will argue that the renormalisation scheme can be directly generalised to other space–time dimensions and field theories with higher spin as well as to theories with local gauge invariance. Our analysis will take place in the framework of perturbative algebraic quantum field theory (pAQFT) [BDF09, BF00, FR12, FR14, HW01, HW04], which is a conceptually clear framework in which fundamental physical properties of perturbative interacting models on curved space–times can be discussed. However, we will make an effort to review how the formulation of pAQFT is related to the more standard formulation of pQFT.

The renormalisation scheme we propose is inspired by the works [DFKR14, Ke10] that deal with pQFT in Minkowski space–time. In these works, the authors introduce an analytic regularisation of the position–space Feynman propagator in Minkowski space–time that is similar to the one discussed in [BG72]. Based on this, time-ordered products are constructed recursively by an Epstein–Glaser type procedure, and it is shown that this recursion can be

resolved by a position–space forest formula similar to the one of Zimmermann used in Bogoliubov–Parasiuk–Hepp–Zimmermann renormalisation in momentum space.

In order to extend the scheme proposed in [DFKR14] to curved space–times and motivated by [BG72] and by the form of Feynman propagators on curved space–times, we introduce an analytic regularisation $\Delta_F^{(\alpha)}$ of a Feynman propagator Δ_F by

$$\Delta_F^{(\alpha)} := \lim_{\epsilon \rightarrow 0^+} \frac{1}{8\pi^2} \left(\frac{u}{(\sigma + i\epsilon)^{1+\alpha}} + \frac{v}{\alpha} \left(1 - \frac{1}{(\sigma + i\epsilon)^\alpha} \right) \right) + w,$$

where u , v , and w are the so-called Hadamard coefficients and σ is $1/2$ times the squared geodesic distance. This analytic regularisation, namely, the construction of certain distributions by means of powers of the geodesic distance, is reminiscent of the use of Riesz distributions to define advanced and retarded Green’s functions on Minkowski space–time. A careful discussion of Riesz distributions and their extension to the curved case is presented in [BGP07]. The regularisation we use is loosely related to dimensional regularisation because the leading singularity of a Feynman propagator in N space–time dimensions is proportional to $(\sigma + i\epsilon)^{1-N/2}$, see, e.g. ([Mo03] appendix A). A regularisation of the Feynman propagator similar to the one above has recently been discussed in [Da15]. In this work, we will combine the analytic regularisation of the Feynman propagator with the minimal subtraction scheme encoded in a forest formula of the kind discussed in [DFKR14, Ho10, Ke10] in order to obtain a time-ordered product that satisfies the causal factorisation property, i.e., a product that is, indeed, time ordered. In order to prove that the analytically regularised amplitudes constructed out of $\Delta_F^{(\alpha)}$ have the meromorphic structure necessary for the application of the forest formula and in order to show how the corresponding Laurent series can be computed explicitly, we will make use of generalised Euler operators. The practical feasibility of the renormalisation scheme will be demonstrated by computing a few examples.

A large part of our analysis is devoted to demonstrating that the scheme we propose is consistent and well defined on general curved space–times. Readers interested in directly applying our scheme may use proposition 3.9 with (3.22), (3.23) in order to compute the Laurent series of the Feynman amplitudes (3.11) regularised by means of (3.12). The correct order of pole subtractions is encoded in the forest formula (3.9), which is explained prior to its display. A number of examples is discussed in section 3.5.

In quantum field theory on cosmological space–times, i.e., Friedmann–Lema  tre–Robertson–Walker (FLRW) space–times, one usually exploits the high symmetry of these space–times in order to evaluate analytical expressions in spatial Fourier space. However, the renormalisation scheme discussed in this work operates on quantities, such as the geodesic distance and the Hadamard coefficients, whose explicit position space and momentum space forms are not even explicitly known in FLRW space–times. Notwithstanding, we will devote a large part of this work in order to develop simple methods to evaluate quantities renormalised in our scheme on FLRW space–times in momentum space, and we will illustrate these methods by explicit examples.

The paper is organised as follows. In the next section, we present a brief introduction to pAQFT and its connection with the more standard formulation of pQFT. Afterwards, we introduce the renormalisation scheme, demonstrate that it is well defined, and analyse its properties in section 3 where we also illustrate the scheme by computing examples. In the fourth section, we demonstrate the applicability of the renormalisation scheme to momentum space computations on cosmological space–times. Finally, a few conclusions are drawn in the

last section of this paper. Conventions regarding the various propagators of a scalar field theory and a few technical computations are collected in the appendix.

2. Introduction to pAQFT

2.1. Basic definitions

Throughout this work, we will consider four-dimensional globally hyperbolic space-times (\mathcal{M}, g) , where g is a Lorentzian metric whose signature is $(-, +, +, +)$, and we use the sign conventions of [Wa84] regarding the definitions of curvature tensors.

We recall the perturbative construction of an interacting quantum field theory on a generic curved space-time in the framework of **pAQFT** recently developed in [BDF09, FR12, FR14] based on earlier work. In this construction, the basic object of the theory is an algebra of observables, which is realised as a suitable set of functionals on field configurations equipped with a suitable product. In order to implement the perturbative constructions following the ideas of Bogoliubov and others, the **field configurations** ϕ are assumed to be off shell. Namely, $\phi \in \mathcal{E}(\mathcal{M}) = C^\infty(\mathcal{M})$ is a smooth function on the globally hyperbolic space-time (\mathcal{M}, g) , and observables are modelled by functionals $F: \mathcal{E}(\mathcal{M}) \rightarrow \mathbb{C}$ satisfying further properties. In particular, all the functional derivatives exist as distributions of compact support where we recall that the functional derivative of a functional F is defined for all $\psi_1, \dots, \psi_n \in \mathcal{D}(\mathcal{M}) = C_0^\infty(\mathcal{M})$ as

$$F^{(n)}(\phi)(\psi_1 \otimes \dots \otimes \psi_n) := \frac{d^n}{d\lambda_1 \dots d\lambda_n} F(\phi + \lambda_1 \psi_1 + \dots + \lambda_n \psi_n)|_{\lambda_1 = \dots = \lambda_n = 0} \in \mathcal{E}'(\mathcal{M}^n).$$

The set of these functionals is indicated by \mathcal{F} . Further regularity properties are assumed for the construction of an algebraic product. In particular, the set of **local functionals** $\mathcal{F}_{\text{loc}} \subset \mathcal{F}$ is formed by the functionals whose n th order functional derivatives are supported on the total diagonal $d_n = \{(x, \dots, x), x \in \mathcal{M}\} \subset \mathcal{M}^n$. Furthermore, their singular directions are required to be orthogonal to d_n , namely, $\text{WF}(F^{(n)}) \subset \{(x, k) \in T^*\mathcal{M}^n, x \in d_n, k \perp Td_n\}$ where WF denotes the wave front set. A generic local functional is a polynomial $P(\phi)(x)$ in ϕ and its derivatives integrated against a smooth and compactly supported tensor. The functionals, whose functional derivatives are compactly supported smooth functions, are instead called **regular functionals** and are indicated by \mathcal{F}_{reg} .

The quantum theory is specified once a product among elements of \mathcal{F}_{loc} and a $*$ -operation (an involution on \mathcal{F}) are given. For the case of free (linear) theories, the product can be explicitly given by a \star -**product**,

$$F \star_H G = \sum_n \frac{\hbar^n}{n!} \langle F^{(n)}, H_+^{\otimes n} G^{(n)} \rangle, \quad (2.1)$$

where H_+ is a Hadamard distribution of the linear theory we are going to quantize, namely, a distribution whose antisymmetric part is proportional to the commutator function $\Delta = \Delta_R - \Delta_A$ and whose WF set satisfies the Hadamard condition, see, e.g. [BFK96, Ra96] for further details and section A.1 for our propagator conventions. Owing to the properties of H_+ , iterated \star_H -products of local functionals $F_1 \star_H \dots \star_H F_n$ are well defined, and \star_H is associative.

In a normal neighbourhood of (\mathcal{M}, g) , a Hadamard distribution H_+ is of the form

$$H_+(x, y) = \frac{1}{8\pi^2} \left(\frac{u(x, y)}{\sigma_+(x, y)} + v(x, y) \log(M^2 \sigma_+(x, y)) \right) + w(x, y), \quad (2.2)$$

where $\sigma_+(x, y) = \sigma(x, y) + i\epsilon(t(x) - t(y)) + \epsilon^2/2$ with t as a time function, i.e., a global time coordinate $2\sigma(x, y)$ is the squared geodesic distance between x and y , and M is an arbitrary mass scale. The Hadamard coefficients u and v are purely geometric and, thus, state independent, whereas, w is smooth and state dependent if $H_+(x, y)$ is the two-point function of a quantum state.

For the perturbative construction of interacting models, we further need a **time-ordered product** \cdot_{T_H} on local functionals. This product is characterised by **symmetry** and the **causal factorisation property**, which requires that

$$F \cdot_{T_H} G = F \star_H G, \quad \text{if } F \succsim G, \quad (2.3)$$

where $F \succsim G$ indicates that F is later than G , i.e., there exists a Cauchy surface Σ of (\mathcal{M}, g) such that $\text{supp}(F) \subset J^+(\Sigma)$ and $\text{supp}(G) \subset J^-(\Sigma)$. However, the causal factorisation uniquely fixes only the time-ordered products among regular functionals in which case,

$$F \cdot_{T_H} G = \sum_n \frac{\hbar^n}{n!} \langle F^{(n)}, H_F^{\otimes n} G^{(n)} \rangle, \quad (2.4)$$

where H_F is the time-ordered (Feynman) version of H_+ , i.e., $H_F = H_+ + i\Delta_A$ with Δ_A as the advanced propagator of the free theory, cf. section A.1. For local functionals, (2.4) is only correct up to the need to employ a nonunique renormalisation procedure, cf. section 3.1. This renormalisation can be performed in such a way that iterated \cdot_{T_H} products of local functionals $F_1 \cdot_{T_H} \dots \cdot_{T_H} F_n$ are well defined with \cdot_{T_H} being associative. Moreover, \star_H products of such time-ordered products of local functionals are well defined as well, cf. [BDF09, FR12, FR14, HW02]. Consequently, we may consider the algebra $\mathcal{A}_0 \star_H$ generated by iterated \cdot_{T_H} products of local functionals. This algebra contains all observables of the free theory that are relevant for perturbation theory.

In the perturbative construction of interacting models, namely, when the free action is perturbed by a nonlinear local functional V , the observables associated with the interacting theory are represented on the free algebra \mathcal{A}_0 by means of the **Bogoliubov formula**. This is given in terms of the local S matrix, i.e., the time-ordered exponential,

$$S(V) = \sum_{n=0}^{\infty} \frac{i^n}{n! \hbar^n} \underbrace{V \cdot_{T_H} \dots \cdot_{T_H} V}_{n \text{ times}}, \quad (2.5)$$

where V is the interacting Lagrangean. In particular, for every interacting observable F , the corresponding representation on the free algebra \mathcal{A}_0 is given by

$$\mathcal{R}_V(F) = S^{-1}(V) \star_H (S(V) \cdot_{T_H} F), \quad (2.6)$$

where $S^{-1}(V)$ is the inverse of $S(V)$ with respect to the \star_H product. The problem in using $\mathcal{R}_V(F)$ as generators of the algebra of interacting observables lies in the construction of the time-ordered product, which *a priori* is an ill-defined operation.

This problem can be solved using ideas that go back to Epstein and Glaser, see, e.g. [BF00], by means of which the time-ordered product among local functionals is constructed recursively. The time-ordered products can be expanded in terms of distributions smeared with compactly supported smooth functions that play the role of coupling constants (multiplied by a space-time cutoff). At each recursion step, the causal factorisation property (2.3)

permits constructing the distributions defining the time-ordered product up to the total diagonal. The extension to the total diagonal can be performed extending the distributions previously obtained without altering the scaling degree towards the diagonal. In this procedure, there is the freedom of adding finite local contributions supported on the total diagonal. This freedom is the well known renormalisation freedom. In addition to the properties already discussed, the renormalised time-ordered product is required to satisfy further physically reasonable conditions. We refer to [HW02, HW04] for details on these properties and the proof that they can be implemented in the recursive Epstein–Glaser construction.

Despite the theoretical clarity of this construction, the Epstein–Glaser renormalisation is quite difficult to implement in practise. The aim of this paper is to discuss a renormalisation scheme that is suitable for practical computations.

2.2. Relation to the standard formulation of pQFT

In this subsection, we outline the relation of the pAQFT framework to the standard formulation of pQFT. As an example, we demonstrate how the two-point (Wightman) function of the interacting field in ϕ^4 theory on a four-dimensional curved space–time is computed where we assume that the quantum state of the interacting field is just the state of free field modified by the interacting dynamics. We further assume that the free field is in a pure and Gaussian Hadamard state.

Let us recall the relevant formulas in perturbative algebraic quantum field theory where we will always try to write expressions both in the pAQFT and in the more standard notation, indicating the latter by a \doteq . Given a local action V , such as $V = \int_{\mathcal{M}} d^4x \sqrt{-g} \frac{\lambda}{4} \phi(x)^4$ in ϕ^4 theory, the corresponding S -matrix, which is, loosely speaking, the ‘ S -matrix in the interaction picture,’ is defined by (2.5) and corresponds to $S(V) \doteq T e^{\frac{i}{\hbar} V}$.

The interacting field, i.e., ‘the field in the interaction picture’ $\phi_I(x)$, is defined by the Bogoliubov formula,

$$\phi_I(x) = \mathcal{R}_V(\phi(x)) = S(V)^{-1} \star_H (S(V) \cdot_{T_H} \phi(x)) \doteq T(e^{\frac{i}{\hbar} V})^{-1} T(e^{\frac{i}{\hbar} V} \phi(x)), \quad (2.7)$$

similar to (2.6) where, by unitarity, $S(V)^{-1} = S(V)^*$. Interacting versions of more complicated expressions in the field, e.g., polynomials at different and coinciding points, are defined analogously. A thorough discussion of the relation between the Bogoliubov formula and the more common formulation of observables in the interaction picture may be found, e.g., in ([Li13] section 3.1). We only remark that, in the Minkowski vacuum state Ω_0 , the expectation value of the Bogoliubov formula can be shown to read (also for more general expressions in the field)

$$\langle \phi_I(x) \rangle_{\Omega_0} \doteq \langle T(e^{\frac{i}{\hbar} V})^{-1} T(e^{\frac{i}{\hbar} V} \phi(x)) \rangle_{\Omega_0} = \frac{\langle T(e^{\frac{i}{\hbar} V} \phi(x)) \rangle_{\Omega_0}}{\langle T(e^{\frac{i}{\hbar} V}) \rangle_{\Omega_0}},$$

which is the theorem of Gell–Mann and Low, see [Du96, DF00] for details.

In the algebraic formulation, one usually cuts off the interaction in order to avoid infrared problems by replacing $\lambda \rightarrow \lambda f(x)$ with a compactly supported smooth function f and considers the adiabatic limit $f \rightarrow 1$ in the end when computing expectation values. As our aim is to compute expectation values in this section, we will write the results in the adiabatic limit keeping in mind that proving the absence of infrared problems, i.e., the convergence of the space–time integrals, is nontrivial and may depend on the state of the free field chosen. Note that the so-called ‘in–in formalism,’ often used in pQFT on cosmological space–times, corresponds to considering a cutoff function f of the form $f(t, \vec{x}) = \Theta(t - t_0)$, i.e., f is a step

function in time and the parameter t_0 corresponds to the time where the interaction is switched on.

Our choice for the quantum state Ω of the interacting field implies that, e.g., the interacting two-point function,

$$\langle \phi_I(x) \star_H \phi_I(y) \rangle_\Omega \doteq \langle \phi_I(x) \phi_I(y) \rangle_\Omega$$

is computed by writing ϕ_I in terms of the free field ϕ and by computing the expectation value of the resulting observable of the free field in the chosen pure Gaussian Hadamard state of the free field that we may, thus, denote by the same symbol Ω . The interacting vacuum state in Minkowski space–time is of this form, whereas, interacting thermal states in flat space–time do not belong to this class as they, roughly speaking, require to take into account both the change in dynamics and the change in spectral properties induced by V [FL13].

The functionals in the functional picture of pAQFT correspond to Wick-ordered quantities of the free field in the sense we will explain now. To this avail, we recall the form of the (quantum) \star_H product and (time-ordered) \cdot_{T_H} product in (2.1), (2.4) that are defined by means of a Hadamard distribution H_+ and its Feynman version $H_F = H_+ + i\Delta_A$. Up to renormalisation of the time-ordered product, these products computed for the special case of the functional $\phi^2(x)$ give

$$\begin{aligned} \phi(x)^2 \star_H \phi(y)^2 &= \phi(x)^2 \phi(y)^2 + 4\hbar \phi(x) \phi(y) H_+(x, y) + 2\hbar^2 H_+^2(x, y), \\ \phi(x)^2 \cdot_{T_H} \phi(y)^2 &= \phi(x)^2 \phi(y)^2 + 4\hbar \phi(x) \phi(y) H_F(x, y) + 2\hbar^2 H_F^2(x, y). \end{aligned}$$

This example shows that the \star_H product (\cdot_{T_H} product) implements the Wick theorem for normal-ordered (time-ordered) fields, and, thus, the previous formulas can be interpreted in more standard notation as

$$\begin{aligned} \phi(x)^2 \cdot_H \phi(y)^2 &= \phi(x)^2 \phi(y)^2 + 4\hbar \phi(x) \phi(y) H_+(x, y) + 2\hbar^2 H_+^2(x, y), \\ T(\phi(x)^2 \cdot_H \phi(y)^2) &= \phi(x)^2 \phi(y)^2 + 4\hbar \phi(x) \phi(y) H_F(x, y) + 2\hbar^2 H_F^2(x, y), \end{aligned}$$

where

$$\begin{aligned} :A;_H &:= \alpha_{-H_+}(A) := e^{-\hbar \langle H_S(x, y), \frac{\delta}{\delta \phi(x)} \otimes \frac{\delta}{\delta \phi(y)} \rangle} A, \\ H_S(x, y) &:= \frac{1}{2}(H_+(x, y) + H_+(y, x)), \end{aligned} \tag{2.8}$$

e.g.,

$$\phi(x)^2 \cdot_H = \lim_{x \rightarrow y} (\phi(x) \phi(y) - H_+(x, y)).$$

Equation (2.8) is a convenient way to encode the combinatorics of normal ordering, whereby the exponential series terminates for polynomial functionals, such as $A = \phi(x)^2$.

The Wick theorem relates (time-ordered) products of Wick-ordered quantities to sums of Wick-ordered versions of contracted products where the definition of Wick ordering and ‘contraction’ are directly related, they both depend on the Hadamard distribution H_+ chosen. Thus, if we choose a particular H_+ to define \star_H and \cdot_{T_H} in pAQFT, we immediately fix the interpretation of all functionals in terms of expressions Wick ordered with respect to H_+ .

For the algebraic formulation, the choice of H_+ is not important, indeed, choosing a different H'_+ with the same properties, one has that $w := H'_+ - H_+ = H'_F - H_F$ because the advanced propagator Δ_A is unique and, thus, universal. Moreover, w is real, smooth, and symmetric, and

$$A \star_{H'} B = \alpha_w(\alpha_{-w}(A) \star_H \alpha_{-w}(B)), \quad A \cdot_{T_{H'}} B = \alpha_w(\alpha_{-w}(A) \cdot_{T_H} \alpha_{-w}(B)),$$

with α defined as in (2.8), and, thus, the algebras associated with \star_H , \cdot_{T_H} and $\star_{H'}$, $\cdot_{T_{H'}}$ are isomorphic via

$$\alpha_w : \mathcal{A}_0 \rightarrow \mathcal{A}'_0,$$

where we recall that \mathcal{A}_0 is algebra \star_H generated by \cdot_{T_H} products of local functionals.

Hence, one may choose a suitable H_+ according to one's needs. However, since $\alpha_d(A) \neq A$ for functionals containing multiple field powers, statements, such as 'the potential is ϕ^4 ', are ambiguous in pAQFT and, in fact, in the standard treatment of QFT. They become nonambiguous only if one says 'the potential is $\phi^4_{:H}$, i.e., ϕ^4 Wick ordered with respect to H_+ .' In pAQFT, the corresponding nonambiguous statement would be 'the potential is the functional ϕ^4 in the algebra \mathcal{A}_0 constructed by means of H_+ .' If one, then, passes to the algebra \mathcal{A}'_0 constructed by means of H'_+ , the potential picks up quadratic and c -number terms as we will compute explicitly below. Alternatively, this ambiguity may be seen to correspond to the renormalisation ambiguity of tadpoles in Feynman diagrams.

Given a Gaussian and Hadamard free field state Ω , a convenient choice or representation of the algebra is to take $H_+ = \Delta_+$, where $\Delta_+(x, y) = \langle \phi(x) \star_\Delta \phi(y) \rangle_\Omega \doteq \langle \phi(x) \phi(y) \rangle_\Omega$ is the two-point function of the free field in state Ω . This corresponds to standard normal ordering, and, consequently, in this representation, the expectation values of all expressions that contain nontrivial powers of the field vanish, i.e.,

$$\langle A \rangle_\Omega = A|_{\phi=0} \doteq \langle :A:_\Delta \rangle_\Omega. \quad (2.9)$$

Keeping a state Ω fixed but passing on to a representation of the algebra with arbitrary H_+ , the expectation value is computed as

$$\langle A \rangle_\Omega = \alpha_w(A)|_{\phi=0} \doteq \langle :A:_H \rangle_\Omega, \quad w = \Delta_+ - H_+,$$

for instance,

$$\langle \phi^2(x) \rangle_\Omega = \alpha_w(\phi^2(x))|_{\phi=0} = (\phi^2(x) + w(x, x))|_{\phi=0} = w(x, x) \doteq \langle :\phi^2(x):_H \rangle_\Omega,$$

which, in more standard terms, would be computed as

$$\langle :\phi^2(x):_H \rangle_\Omega = \lim_{x \rightarrow y} \langle \phi(x) \phi(y) - H_+(x, y) \rangle_\Omega = \lim_{x \rightarrow y} (\Delta_+(x, y) - H_+(x, y)) = w(x, x).$$

In QFT in curved space-times, normal ordering is, in principle, problematic because (pointlike) observables should be defined in a local and generally covariant way, i.e., they should only depend on the space-time in an arbitrarily small neighbourhood of the observable localisation [BFV01, HW01]. This is not satisfied for, e.g., field polynomials Wick ordered with $\Delta_+(x, y)$ because this distribution satisfies the Klein–Gordon equation and, thus, it encodes nonlocal information on the curved space-time [HW01]. It is still possible to compute in the convenient normal-ordered representation in the following way. In the example of ϕ^4 theory, one defines the potential $\frac{\lambda}{4} \phi(x)^4$ as a local and covariant observable by identifying it with the corresponding monomial in a representation of the algebra furnished by a purely geometric H_+ , i.e., a H_+ of the form (2.2) with $w = 0$.

In other words, we set, once and for all, in the H_+ representation,

$$V_H = \int_{\mathcal{M}} d^4x \sqrt{-g} \frac{\lambda}{4} \phi(x)^4 \doteq \int_{\mathcal{M}} d^4x \sqrt{-g} \frac{\lambda}{4} : \phi(x)^4 :_H.$$

This does not fix V uniquely because H depends on the scale M inside of the logarithm but the freedom in defining V_H and, analogously, the free/quadratic part of the Klein–Gordon action as above, corresponds to the usual freedom in choosing the 'bare mass' m , 'bare coupling to

would like to compute in the convenient normal-ordered representation with respect to Ω , leads to the introduction of an effective space–time-dependent and state-dependent (squared) mass term $\mu(x) = 3\lambda w(x, x)$ in the interaction potential, which, of course, leads to additional Feynman graphs in perturbation theory, cf. Figures 1 and 2. The field-independent term $\frac{3\lambda}{4}w(x, x)^2$ plays no role for computations of quantities that do not involve functional derivatives of the extended Einstein–Hilbert–Klein–Gordon action with respect to the metric (an example where it does play a role is the stress–energy tensor), just as the modification of the free action by the change of representation plays no role for the computation of such quantities. A similar phenomenon as in (2.10) occurs in thermal quantum field theory on Minkowski space–time where the effective mass generated by changing from the normal-ordered picture with respect to the free vacuum state to the normal-ordered picture with respect to the free thermal state is termed ‘thermal mass,’ cf. ([Li13] section 2.3.2.) for details.

After these general considerations, we can proceed to compute, as an example, the two-point function of the interacting field ϕ_I in ϕ^4 up to second order in λ , whereby ϕ_I is assumed to be in a state induced by a Gaussian Hadamard state of the free field. To this avail, we will exclusively compute in the associated normal-ordered representation and thus omit the subscripts on the star product and the time-ordered product $\star := \star_\Delta$, $\cdot_T := \cdot_{T_\Delta}$.

We start from the Bogoliubov formula (2.7) and compute (from now on, $\hbar = 1$)

$$\begin{aligned} S(V) &= 1 + iV - \frac{1}{2}V \cdot_T V + O(\lambda^3) \\ S(V)^{\star-1} &= 1 - iV + \frac{1}{2}V \cdot_T V - V \star V + O(\lambda^3) \\ \phi_I &= \phi - iV \star \phi + iV \cdot_T \phi + \frac{1}{2}(V \cdot_T V) \star \phi - V \star V \star \phi \\ &\quad - \frac{1}{2}V \cdot_T V \cdot_T \phi + V \star (V \cdot_T \phi) + O(\lambda^3). \end{aligned}$$

It remains to compute the \star -product of $\phi_I(x)$ and $\phi_I(y)$ and to set $\phi = 0$ in the remaining expression in order to obtain the expectation value in state Ω . The result can, as always, be conveniently expressed in terms of Feynman diagrams where we use the Feynman rules depicted in figure 1.

In the computation of $\langle \phi_I(x)\phi_I(y) \rangle_\Omega$, many expressions can be shortened considerably by using the relation $\Delta_F - \Delta_+ = i\Delta_A$, in particular, this holds for the external legs of the appearing Feynman diagrams. The resulting Feynman diagrams are depicted in figure 2.

3. Analytic regularisation and minimal subtraction on curved space–times

As discussed above, the main problem in using the Bogoliubov formula (2.6),

$$\mathcal{R}_V(F) = \frac{\hbar}{i} \frac{d}{d\lambda} S(V)^{-1} \star_H S(V + \lambda F)|_{\lambda=0}$$

for constructing interacting fields perturbatively is that it is given in terms of the S matrix, which is the time-ordered exponential (2.5). Unfortunately, the time-ordered product defined in terms of a ‘deformation’ (2.4) written by means of a Feynman propagator H_F is well defined only on regular functionals because the singularities present in H_F forbid their application to more general functionals.

In order to proceed, there is the need for employing a renormalisation procedure to construct the time-ordered products. In this work, we discuss the use of certain analytic

methods to solve this problem. The procedure we will pursue is the following. We deform the Feynman propagator by means of complex parameter α with values in the neighbourhood of the origin obtaining a function with distributional values $\alpha \mapsto H_F^{(\alpha)}$. The deformation we are looking for needs to be such that, in the limit $\alpha \rightarrow 0$, we recover the ordinary Feynman propagator. Furthermore, when α is nonvanishing but sufficiently small, pointwise powers of $H_F^{(\alpha)}$ and integral kernels of more complicated loop diagrams should be well defined. If this is the case since the corresponding distributions obtained in the limit $\alpha \rightarrow 0$ are well defined outside of the total diagonal, the poles of $\alpha \mapsto H_F^{(\alpha)}$ and more complicated loop expressions are supported on the total diagonal. The idea, similar to what happens in dimensional regularisation, is that it is possible to renormalise these distributions by simply removing the poles.

3.1. Analytic regularisation of time-ordered products and the minimal subtraction scheme

In order to discuss the analytic regularisation of time-ordered products, we employ the notation used, e.g., in [DFKR14] that efficiently encodes the full combinatorics of Feynman diagrams in a compact form. Namely, the time-ordered product of n local functionals V_1, \dots, V_n can be formally defined in the following way⁴:

$$V_1 \cdot_{T_H} \dots \cdot_{T_H} V_n := \mathcal{T}_n(V_1 \otimes \dots \otimes V_n) := m \circ T_n(V_1 \otimes \dots \otimes V_n), \quad (3.1)$$

where m denotes the pointwise product $m(F_1 \otimes \dots \otimes F_n)(\phi) = F_1(\phi) \dots F_n(\phi)$ and the operator T_n is written in terms of an exponential,

$$T_n = \exp \left(\sum_{1 \leq i < j \leq n} \Delta_{ij} \right) = \prod_{1 \leq i < j \leq n} \sum_{l_{ij} \geq 0} \frac{\Delta_{ij}^{l_{ij}}}{l_{ij}!}, \quad (3.2)$$

with

$$\Delta_{ij} := \left\langle H_F, \frac{\delta^2}{\delta \phi_i \delta \phi_j} \right\rangle. \quad (3.3)$$

Here, the functional derivative $\frac{\delta}{\delta \phi_i}$ acts on the i th element of the tensor product $V_1 \otimes \dots \otimes V_n$, and $H_F = H_+ + i\Delta_A$ is the time-ordered version of the Hadamard distribution H_+ entering the construction of the free algebra \mathcal{A}_0 via \star_H . The exponential (3.2) admits the usual representation in terms of Feynman graphs. More precisely, it can be written as a sum over all graphs Γ in \mathcal{G}_n , the set of all graphs with vertices $V(\Gamma) = \{1, \dots, n\}$, and l_{ij} edges $e \in E(\Gamma)$ joining the vertices i, j . Furthermore, in this construction, there are no tadpoles $l_{ii} = 0$ (cf. section 2.2 for details on why these are absent), and the edges are not oriented $l_{ij} = l_{ji}$. With this in mind,

$$T_n = \sum_{\Gamma \in \mathcal{G}_n} \frac{1}{N(\Gamma)} \left\langle \tau_\Gamma, \frac{\delta^{2|E(\Gamma)|}}{\prod_{i \in V(\Gamma)} \prod_{E(\Gamma) \ni e \supset i} \delta \phi_i(x_i)} \right\rangle, \quad (3.4)$$

where $N(\Gamma) = \prod_{i < j} l_{ij}!$ is a numerical factor counting the possible permutations among the lines joining the same two vertices, the second product $\prod_{e \supset i}$ is over the edges having i as a vertex, and x_i is a point in \mathcal{M} corresponding to the vertex i . Moreover, τ_Γ is a distribution that

⁴ In fact, in view of locality and covariance, a better definition of the time-ordered product is $T_1(V_1) \cdot_{T_H} \dots \cdot_{T_H} T_1(V_n) := \mathcal{T}_n(V_1 \otimes \dots \otimes V_n)$ where $T_1: \mathcal{F}_{\text{loc}} \rightarrow \mathcal{F}_{\text{loc}} \subset \mathcal{A}_0$ plays the role of identifying local and covariant (smeared) Wick polynomials as particular elements of the free algebra \mathcal{A}_0 , cf. [HW04]. As we will not touch upon this point in our renormalisation scheme, we choose to omit T_1 in our formulas for simplicity.

is well defined outside of all partial diagonals, namely, on $\mathcal{M}^n \setminus D_n$, where

$$D_n := \{x_1, \dots, x_n \mid x_i = x_j \text{ for, at least, one pair } (i, j), i \neq j\}, \quad (3.5)$$

and τ_Γ has the form

$$\tau_\Gamma = \prod_{e=(i,j) \in E(\Gamma)} H_F(x_i, x_j) = \prod_{1 \leq i < j \leq n} H_F(x_i, x_j)^{l_{ij}}. \quad (3.6)$$

The *a priori* restricted domain of τ_Γ is the reason why T_n defined as above is not a well-defined operation on $\mathcal{F}_{\text{loc}}^{\otimes n}$. In this context, we recall that the total diagonal $d_n \subset D_n$ is defined as

$$d_n := \{(x, \dots, x), x \in \mathcal{M}\} \subset \mathcal{M}^n. \quad (3.7)$$

In order to complete the construction, we need to extend the obtained distributions to the diagonals D_n . This is not a straightforward limit because the singular structure of the Feynman propagator H_F contains the one of the δ distribution and because pointwise products of the latter distribution are ill defined. Consequently, a renormalisation procedure needs to be implemented in order to extend τ_Γ to the full \mathcal{M}^n . This extension is, in general, not unique but subject to renormalisation freedom.

Here, we will discuss a procedure to extend the distributions τ_Γ to D_n called **minimal subtraction (MS)**, which makes use of an analytic regularisation $\Delta_{ij}^{\alpha_{ij}}$ of Δ_{ij} given in terms of a family of deformations $H_F^{\alpha_{ij}}$ of the Feynman propagator H_F parametrised by complex parameters α_{ij} contained in some neighbourhood of $0 \in \mathbb{C}$. To this end, we follow [DFKR14] and call $t^{(\alpha)}$ an analytic regularisation of a distribution t defined outside of a point $x_0 \in \mathcal{M}$ if for all $f \in \mathcal{D}(\mathcal{M})$ $\langle t^{(\alpha)}, f \rangle$ is a meromorphic function in α for α in some neighbourhood of 0, which is analytic for $\alpha \neq 0$. Moreover, $t^{(\alpha)}$ may be extended to x_0 for $\alpha \neq 0$, whereas, $\lim_{\alpha \rightarrow 0} t^{(\alpha)} = t$ on $\mathcal{M} \setminus \{x_0\}$.

We will introduce an analytic regularisation of the Feynman propagator H_F in the following section, but the basic idea of the MS scheme is independent of the details of the analytic regularisation. Namely, given any analytic regularisation $H_F^{(\alpha)}$ of H_F , we repeat the formal construction of T_n presented above by replacing H_F by $H_F^{(\alpha)}$ in (3.3) and Δ_{ij} by the induced $\Delta_{ij}^{\alpha_{ij}}$ in (3.2). Proceeding in this way, we define

$$T_n^{(\alpha)} := e^{\sum_{i < j} \Delta_{ij}^{\alpha_{ij}}} \text{ with } \alpha := \{\alpha_{ij}\}_{i < j},$$

and the corresponding integral kernels $\tau_\Gamma^{(\alpha)}$ of Feynman graphs Γ in analogy to (3.4). We expect that the distributions $\tau_\Gamma^{(\alpha)}$ are multivariate meromorphic functions that have poles at the origin for some of the α_{ij} . Hence, in order to obtain well-defined distributions in the limit α_{ij} to 0 and, consequently, a renormalised time-ordered product \cdot_{T_H} , all these poles need to be subtracted.

The properties of the analytically regularised Feynman propagator imply that $\tau_\Gamma^{(\alpha)}$ is well defined on $\mathcal{M}^n \setminus D_n$ (3.5), even if all α_{ij} are vanishing. Since $\tau_\Gamma^{(\alpha)}$ is a multivariate meromorphic function in α that is analytic if restricted to $\mathcal{M}^n \setminus D_n$, we may deduce that the principal part of $\tau_\Gamma^{(\alpha)}$ for some α_{ij} must be supported on a partial diagonal of \mathcal{M}^n . In fact, in order for the time-ordered products to fulfil the factorisation property (2.3), the subtraction of the principal parts of $\tau_\Gamma^{(\alpha)}$ needs to be done in such a way that, at each step, only local terms are subtracted. However, the previous discussion only implies that the support of the principal parts is contained in D_n , i.e., the union of all the partial diagonals in \mathcal{M}^n . In order to satisfy the causal factorisation property, the principal parts need to be removed in a recursive way starting from the partial diagonals corresponding to two vertices and

proceeding with the partial diagonals corresponding to an increasing number $m \leq n$ of vertices $\mathfrak{d}_I := \{(x_1, \dots, x_n) \in M^n, x_i = x_j, i, j \in I \subset \{1, \dots, n\}, |I| = m\}$.

The correct recursion procedure is implemented by the so-called Epstein–Glaser forest formula, which is a position–space analogue of the Zimmermann forest formula, see [DFKR14, Ho10, Ke10] for a careful analysis of the subject. Here, we will follow the treatment discussed in [DFKR14]. To this end, we consider the set of indices $\bar{n} := \{1, \dots, n\}$ and define a forest F as

$$F = \{I_1, \dots, I_k\}, \quad I_j \subset \bar{n} \quad \text{and} \quad |I_j| \geq 2,$$

where, for every pair $I_i, I_j \in F$,

$$I_i \cap I_j = \emptyset \quad \text{or} \quad I_i \subset I_j \quad \text{or} \quad I_j \subset I_i.$$

The set of all forests of n indices together with the empty forest $\{\}$ is indicated by $\mathfrak{F}_{\bar{n}}$.

For every subset $I \subset \bar{n}$, we indicate, by R_I , the operator that extracts the principal part with respect to α_I of a multivariate meromorphic function $f(\{\alpha_{ij}\}_{i < j})$, where, for every $i, j \in I$, $\alpha_{ij} = \alpha_I$, and multiply it with -1 ,

$$R_I f := -\text{pp} \lim_{\substack{\alpha_{ij} \rightarrow \alpha_I \\ \forall i, j \in I}} f(\{\alpha_{ij}\}_{i < j}). \quad (3.8)$$

We complement this definition by setting $R_{\{\}}$ to be the identity.

Given all these data, we define the renormalised time-ordered product in the MS scheme as in, e.g. ([DFKR14] theorem 3.1) by

$$\mathcal{T}_n = (\mathcal{T}_n)_{\text{MS}} := \lim_{\alpha \rightarrow 0} m \circ \left(\sum_{F \in \mathfrak{F}_{\bar{n}}} \prod_{I \in F} R_I \right) \circ T_n^{(\alpha)}, \quad (3.9)$$

where, in the product over $I \in F$, R_I appears before R_J if $I \subset J$. Furthermore, for each graph Γ , the limit $\alpha = \{\alpha_{ij}\}_{i < j} \rightarrow 0$ is computed by setting $\alpha_{ij} = \alpha_\Gamma$ for every $i < j$ before taking the sum over the forests and finally by considering the limit α_Γ to 0. In this context, we recall that, for every element of the sum over $\mathfrak{F}_{\bar{n}}$, part of the limit $\alpha_{ij} \rightarrow \alpha_\Gamma$ is already taken by applying R_I , see (3.8).

Given the renormalised \mathcal{T}_n in the MS scheme, the corresponding local S -matrix may be constructed as

$$S(V) = \sum_{n=0}^{\infty} \frac{i^n}{\hbar^n n!} \mathcal{T}_n(V \otimes \dots \otimes V)$$

for any local interaction Lagrangean V .

In order to implement the minimal subtraction scheme as outlined above, we first need to specify an analytic regularisation $H_F^{(\alpha)}$ of the Feynman propagator H_F on generic curved space–times. Afterwards, we have to demonstrate that, for all graphs $\Gamma \in \mathcal{G}_n$, the analytically regularised integral kernels,

$$\tau_\Gamma^{(\alpha)} = \prod_{e=(i,j) \in \Gamma} H_F^{\alpha_{ij}}(x_i, x_j) = \prod_{1 \leq i < j \leq n} (H_F^{\alpha_{ij}}(x_i, x_j))^{l_{ij}}, \quad (3.10)$$

appearing in

$$T_n^{(\alpha)} = \sum_{\Gamma \in \mathcal{G}_n} \frac{1}{N(\Gamma)} \left\langle \tau_\Gamma^{(\alpha)}, \frac{\delta^{2|E(\Gamma)|}}{\prod_{i \in V(\Gamma)} \prod_{e \supset i} \delta \phi_i(x_i)} \right\rangle \quad (3.11)$$

satisfy the properties necessary for the implementation of the MS scheme. In particular, we need to demonstrate that the distribution $\tau_{\Gamma}^{(\alpha)}$, which is *a priori* defined only on $\mathcal{M}^n \setminus D_n$, can be uniquely extended to the full \mathcal{M}^n without renormalisation where the uniqueness of this extension is important in order to obtain a definite renormalisation scheme. Moreover, we need to show that this distribution $\tau_{\Gamma}^{(\alpha)} \in \mathcal{D}'(\mathcal{M}^n)$ is weakly meromorphic in α in a neighbourhood of 0 where, in view of the forest formula, it is only necessary to show that setting $\alpha_{ij} = \alpha_I$ for all $i, j \in I$, $\tau_{\Gamma}^{(\alpha)}$ is weakly meromorphic in α_I . Additionally, we need to prove that, if τ_{Γ} prior to regularisation is well defined outside of the partial diagonal d_I , then, the pole of $\tau_{\Gamma}^{(\alpha)}$ with $\alpha_{ij} = \alpha_I$ for all $i, j \in I$ in α_I is supported on d_I and, thus, local. Finally, we need to prove that our MS scheme satisfies all properties given in [HW02, HW04], which a physically meaningful renormalisation scheme on curved space-times should satisfy, and we need to provide means to explicitly compute the minimal subtraction, which, after all, is the main motivation for this work.

Our plan to construct the mentioned quantities and to prove their required properties is as follows.

1. In section 3.2, we construct an analytic regularisation $H_F^{(\alpha)}$ of the Feynman propagator based on the observation that locally H_F is of the form (2.2) up to considering instead of σ_+ the half squared geodesic with the Feynman ϵ prescription $\sigma_F := \sigma + i\epsilon$. Motivated by the fact that the singular structure of H_F originates from the form in which σ_F appears, we set locally,

$$H_F^{(\alpha)} := \lim_{\epsilon \rightarrow 0^+} \frac{1}{8\pi^2} \left(\frac{u}{M^{2\alpha} \sigma_F^{1+\alpha}} + \frac{v}{\alpha} \left(1 - \frac{1}{M^{2\alpha} \sigma_F^\alpha} \right) \right) + w, \quad (3.12)$$

where we use the (arbitrary but fixed) mass scale M present in (2.2) also for preserving the mass dimension of H_F in the regularisation.

2. In proposition 3.7, we then prove that the relevant distributions,

$$t_{\Gamma}^{(\alpha)} := \prod_{1 \leq i < j \leq n} \frac{1}{\sigma_F^{I_{ij}(1+\alpha_{ij})}} \in \mathcal{D}'(\mathcal{M}^n \setminus D_n) \quad (3.13)$$

are multivariate analytic functions. The distribution (3.13) only displays the most singular contribution of $\tau_{\Gamma}^{(\alpha)}$ (3.10), but the subleading contributions are clearly of the same form up to replacing some of the factors $(1 + \alpha_{ij})$ in the exponents by α_{ij} or 0.

3. In order to show that $t_{\Gamma}^{(\alpha)}$ can be uniquely extended from $\mathcal{M}^n \setminus D_n$ to \mathcal{M}^n in a weakly meromorphic fashion, i.e., that the singularities relevant for the forest formula are poles of finite order, we follow a strategy similar to the one used in [HW02] and consider a scaling expansion with respect to a suitable scaling transformation. We first argue, in proposition 3.8, that an analytically regularised distribution $t^{(\alpha)} \in \mathcal{D}'(\mathcal{M}^n \setminus d_n)$, which can be written as a sum of homogeneous terms with respect to this scaling transformation plus a sufficiently regular remainder, can be extended to \mathcal{M}^n in a weakly meromorphic way where the uniqueness of the extension follows from its weak meromorphicity. In proposition 3.10, we give a sufficient condition for the existence of such a homogeneous expansion, and we demonstrate, in proposition 3.12, that the distributions $t_{\Gamma}^{(\alpha)}$ satisfy this condition.
4. The above-mentioned results are proved by means of generalised Euler operators (see [Da13] for a related concept), which can be written abstractly in terms of a scaling transformation, but also in terms of covariant differential operators whose explicit form can be straightforwardly computed as we argue in section 3.3.1. In proposition 3.9, we

use these operators in order to demonstrate how the full relevant pole structure of $t_{\Gamma}^{(\alpha)}$ can be computed, thus, showing the practical feasibility of the MS scheme. We find that our renormalisation scheme corresponds, in fact, to a particular form of differential renormalisation and expand on this by computing a few examples in section 3.5.

5. Finally, in proposition 3.14, we prove that the MS scheme satisfies the axioms of [HW02, HW04] for time-ordered products and preserves invariance under any space–time isometries present.

Remark 3.1. The local Hadamard expansion (2.2) of H_F and correspondingly the analytically continued $H_F^{(\alpha)}$ defined in (3.12) are only meaningful on normal neighbourhoods \mathcal{N} of (\mathcal{M}, g) . In order to define $H_F^{(\alpha)}$ and the induced distributions $\tau_{\Gamma}^{(\alpha)}$ (3.10) globally, we may employ suitable partitions of unity. Rather than providing general and cumbersome formulas, we prefer to illustrate the idea at the example of the triangular graph,

$$\tau_{\Gamma} = H_{F,13} H_{F,23} H_{F,12}^2 := H_F(x_1, x_3) H_F(x_2, x_3) H_F(x_1, x_2)^2,$$

the renormalisation of which is discussed in detail in section 3.5.3. We recall that a corollary of lemma 10 of chapter 5 in [O’N83] guarantees that there exists a covering \mathcal{C} of \mathcal{M} consisting of open geodesically convex sets such that $\mathcal{N}_i \cap \mathcal{N}_j$ is geodesically convex for every $\mathcal{N}_i, \mathcal{N}_j \in \mathcal{C}$ ⁵. With this \mathcal{C} at our disposal, we define the sets,

$$\mathcal{N}_{12} := \bigcup_{\mathcal{N} \in \mathcal{C}} \mathcal{N} \times \mathcal{N} \subset \mathcal{M}^2, \quad \mathcal{N}_{123} := \bigcup_{\mathcal{N} \in \mathcal{C}} \mathcal{N} \times \mathcal{N} \times \mathcal{N} \subset \mathcal{M}^3.$$

We call sets of the form \mathcal{N}_{12} and \mathcal{N}_{123} a **normal neighbourhood of the total diagonal**. This definition is essentially motivated by the fact that, for every $x \in \mathcal{M}$, we can find a normal neighborhood $\mathcal{N}_x \in \mathcal{C}$ of x in \mathcal{M} . The squared geodesic distance σ then is well defined on \mathcal{N}_{12} , whereas, the same is, in general, not true if we replace \mathcal{C} in the previous formula with a covering of \mathcal{M} formed by all open geodesically convex sets. Setting $\sigma_{ij} := \sigma(x_i, x_j)$, we observe that σ_{12} is well defined on \mathcal{N}_{12} and that σ_{12}, σ_{13} , and σ_{23} are well defined on \mathcal{N}_{123} . We now consider smooth and compactly supported functions $\chi_{12} \in \mathcal{D}(\mathcal{N}_{12})$, $\chi_{123} \in \mathcal{D}(\mathcal{N}_{123})$, which are such that $\chi_{12} = 1$ on $d_2 \subset \mathcal{N}_{12}$ and $\chi_{123} = 1$ on $d_3 \subset \mathcal{N}_{123}$. Note that, by construction, χ_{12} and χ_{123} vanish outside of \mathcal{N}_{12} and \mathcal{N}_{123} , respectively. We may now define the analytically regularised distribution $\tau_{\Gamma}^{(\alpha)}$ by setting

$$\begin{aligned} \tau_{\Gamma}^{(\alpha)} := & H_{F,13}^{(\alpha_{13})} H_{F,23}^{(\alpha_{23})} (H_{F,12}^{(\alpha_{12})})^2 \chi_{12} \chi_{123} + H_{F,13} H_{F,23} H_{F,12}^2 (1 - \chi_{12}) \\ & + H_{F,13} H_{F,23} (H_{F,12}^{(\alpha_{12})})^2 \chi_{12} (1 - \chi_{123}), \end{aligned}$$

where the Feynman propagators are regularised as in (3.12). By construction, $\tau_{\Gamma}^{(\alpha)}$ is globally well defined, and the analysis outlined above and performed in the following sections implies that it can be uniquely extended to a weakly meromorphic distribution on the full \mathcal{M}^3 . Moreover, the local pole contributions corresponding to $\alpha_{12} = \alpha_I$ with $I = \{1, 2\}$ and $\alpha_{12} = \alpha_{13} = \alpha_{23} = \alpha_J$ with $J = \{1-3\}$ are clearly independent of the choice of χ_{12}, χ_{123} , and $\mathcal{N}_{12}, \mathcal{N}_{123}$ such that the MS-regularised amplitude $(\tau_{\Gamma})_{\text{MS}}$ is both globally well-defined and independent of the quantities entering the global definition of the analytic regularisation. Keeping this approach to define global analytically regularised quantities in mind, we will, for simplicity, work only with local quantities in the following.

⁵ We would like to thank V. Moretti for pointing this result out to us.

3.2. Analytic regularisation of the Feynman propagator H_F on curved space-times

Following the plan outlined in section 3.1, we would like to define an analytic regularisation $H_F^{(\alpha)}$ of H_F by (3.12). To this end, we start our analysis by constructing the distribution $1/\sigma_F^{1+\alpha}$ in \mathcal{M}^2 for $\alpha \in \mathbb{C} \setminus \mathbb{N}$. As anticipated in section 3.1, we will make use of scaling properties of $1/\sigma_F^{1+\alpha}$ and the induced quantities $t_\Gamma^{(\alpha)}$ (3.13) with respect to a particular geometric scaling transformation.

For every pair of points x_1, x_i in a normal neighbourhood $\mathcal{N} \subset (\mathcal{M}, g)$, there exists a unique geodesic γ connecting x_1 and x_i . We will assume that $\gamma: \lambda \mapsto x_i(\lambda)$ is affinely parametrised and that $x_i(0) = x_1$, whereas, $x_i(1) = x_i$. For all $\lambda \geq 0$ and all $f \in \mathcal{D}(\mathcal{N}_n)$ with $\mathcal{N}_n \subset \mathcal{M}^n$, a normal neighbourhood of the total diagonal d_n (cf. remark 3.1), the geometric scaling transformation we will consider is

$$f_\lambda := \lambda^{4(n-1)} f(x_1, x_2(\lambda), \dots, x_n(\lambda)) \prod_{i=2}^n \frac{\sqrt{g(x_i(\lambda))}}{\sqrt{g(x_i)}}, \quad (3.14)$$

where $g(x)$ is the absolute value of the determinant of the metric expressed in normal coordinates. For $\lambda > 1$, it may happen that $x_i(\lambda)$ lies outside of \mathcal{N}_n and is, thus, not well defined, in general. In this case, we set $f_\lambda = 0$, which is well defined because $f = 0$ outside of \mathcal{N}_n . For later purposes, we recall that the determinant of the metric computed in normal coordinates centred at x_1 is such that,

$$\sqrt{g(x_i)} = \frac{1}{u^2(x_1, x_i)},$$

where u is the Hadamard coefficient in (2.2) and u^2 is the van Vleck–Morette determinant, see, e.g. ([PPV11] (8.5)).

By means of this transformation, relevant information about the behaviour of a distribution in the neighbourhood of the total diagonal d_n can be obtained.

We recall two definitions and a few results that we will use in the following. To this end, we consider a distribution t in $\mathcal{D}'(\mathcal{M}^n \setminus d_n)$. The restrictions of t to the sets,

$$\mathcal{O}_x = \{(x, y) \in \mathcal{M}^n \mid y \in \mathcal{M}^{n-1}\}$$

for $x \in \mathcal{M}$ are well defined if $\text{WF}(t)$ is disjoint from the **conormal bundle** $N^*\mathcal{O}_x = \{(y, k) \in T^*\mathcal{M} \mid \langle k, \xi \rangle = 0, \forall \xi \in T_y\mathcal{O}_x\}$, see, e.g., Theorem 8.2.4 in the book by Hörmander [Hö90]. By assumption, these restrictions are defined everywhere on \mathcal{O}_x up to the single points $\mathcal{O}_x \cap d_n$. The extension of the restricted distributions to that point can be constructed by means of particular scaling properties of that distribution, see, e.g., theorems 5.2 and 5.3 in [BF00]. However, in order to discuss the extension of t to the full diagonal d_n , it is useful to introduce the scaling degree of t towards d_n in the following way, which is an adaption of the notion of a transversal scaling degree presented in ([BF00] section 6) to our setting.

Definition 3.2. Let $\Gamma \subset T^*\mathcal{M}^n$ be a closed conical set that is equal to the conormal bundle N^*d_n on d_n and is disjoint from $N^*\mathcal{O}_x$ for every x , and let t be a distribution in $\mathcal{D}'_\Gamma(X)$, $X \in \{\mathcal{M}^n, \mathcal{M}^n \setminus d_n\}$, i.e., $t \in \mathcal{D}'(X)$ and $\text{WF}(t) \subset \Gamma$. The **scaling degree** of t towards d_n is defined as

$$\text{sd}_{d_n}(t) := \inf \{w \in \mathbb{R} \mid \lim_{\lambda \rightarrow 0^+} \lambda^w t_\lambda = 0\},$$

where the distributions $t_\lambda(f) := t(f_{1/\lambda})$ are defined in terms of the scaling transformation (3.14) and where the limits are taken in the topology of $\mathcal{D}'_\Gamma(\mathcal{M}^n)$.

If a distribution has scaling degree towards d_n lower than the total dimension of the scaled coordinates $4(n - 1)$, then, it possesses a unique extension towards d_n with the same scaling degree, see, e.g., theorem 6.9 of [BF00]. The proof of the existence of the extensions presented there is constructive and can be obtained as the limit to large n of $t(1 - \theta_n)$, where θ_n are suitable smoothed characteristic functions of a neighbourhood of the diagonal, whose supports become smaller and smaller for large n . In particular, the previous notion of scaling degree permits controlling the WF set of every $t(1 - \theta_n)$ and of the limiting distribution. In fact, the extended distribution can be restricted to every \mathcal{O}_x as well because Γ and $N^*\mathcal{O}_x$ are disjoint sets. The uniqueness of the extension descends from the uniqueness of the extension of the distributions restricted to \mathcal{O}_x and from the observation that the scaling degree of the restricted distributions towards the point $(x, \dots, x) \in \mathcal{O}_x$ is such that $\text{sd}(t|_{\mathcal{O}_x}) \leq \text{sd}_{d_n}(t)$. If not strictly necessary, we will drop the suffix d_n in the following. Having this result at our disposal implies that, in order to construct an extension towards d_n , it is sufficient to construct extensions of the distribution restricted to \mathcal{O}_x .

Remark 3.3. The distributions relevant for our analysis are the ones of the form (3.10), respectively (3.13). As we discuss briefly in the proof of proposition 3.7, these distributions satisfy the microlocal spectrum condition introduced in [BFK96] and, consequently, the WF set condition required in definition 3.2.

The scaling degree towards a partial diagonal may be defined in analogy to definition 3.2. The same geometric transformation (3.14) can be used to introduce relevant homogeneity properties of a distribution.

Definition 3.4. A distribution $t \in \mathcal{D}'(\mathcal{M}^n)$ or $t \in \mathcal{D}'(\mathcal{M}^n \setminus d_n)$, which satisfies the equality,

$$\lambda^\delta \langle t, f_\lambda \rangle = \langle t, f \rangle \quad \forall \lambda > 0,$$

under transformations of the form (3.14) for all $f \in \mathcal{D}(\mathcal{N}_n \setminus d_n)$ and for a $\delta \in \mathbb{C}$, is called **homogeneous of degree δ** .

These definitions imply, by direct inspection, that a distribution that is homogeneous of degree δ and whose WF set restricted to d_n is contained in N^*d_n has scaling degree $-\text{Re}(\delta)$. We further notice that homogeneous distributions $t \in \mathcal{D}(\mathcal{M}^n \setminus d_n)$, whose $\text{WF}(t) \cap T_{d_n}^*\mathcal{M}$ is contained in N^*d_n , possess unique extensions to \mathcal{M}^n with the same degree of homogeneity δ if $-(\delta + 4(n - 1)) \notin \mathbb{N}$. A proof of this claim can be obtained following very closely the proof of theorem 3.2.3 in [Hö90]. In particular, the uniqueness of these extensions descends once again from the uniqueness of the extensions of the distributions restricted to \mathcal{M}_x^n , whereas, the existence of the extension is obtained employing a fundamental solution of the Euler operator E_1 we will introduce below in (3.17).

In the next proposition, we introduce the distributions we will use as building blocks for the construction of regularised Feynman propagators on Lorentzian manifolds. Although not completely analogous, the construction we are going to present is similar to the extension of Riesz distributions to curved spaces presented in section 1.4 of [BGP07]. In particular, here, we will discuss the boundary value of $1/(\sigma + i\epsilon)^\alpha$ for $\epsilon \rightarrow 0$, while ordinary Riesz distributions are related to the antisymmetric part of a different boundary value of the functions $1/\sigma^\alpha$.

Proposition 3.5. Consider a normal neighbourhood $\mathcal{N}_2 \subset \mathcal{M}^2$ of d_2 (cf. remark 3.1) and the following expression for $\alpha \in \mathbb{C}$ and $f \in \mathcal{D}(\mathcal{N}_2)$:

$$\left\langle \frac{1}{\sigma_F^\alpha}, f \right\rangle := \lim_{\epsilon \rightarrow 0^+} \int_{\mathcal{M}^2} \frac{1}{(\sigma(x, y) + i\epsilon)^\alpha} f(x, y) d\mu_g(x) d\mu_g(y).$$

Then, the following statements hold.

- (a) $1/\sigma_F^\alpha$ restricted to $\mathcal{D}(\mathcal{N}_2 \setminus d_2)$ is a distribution that is weakly analytic in α .
- (b) $1/\sigma_F^\alpha$ is homogeneous of degree -2α with respect to transformations of the form (3.14) for $f \in \mathcal{D}(\mathcal{N}_2 \setminus d_2)$.
- (c) $1/\sigma_F^\alpha$ is well defined as a distribution on \mathcal{N}_2 for $2\alpha - 4 \notin \mathbb{N}$. Furthermore, for all $f \in \mathcal{D}(\mathcal{N}_2)$ $\langle 1/\sigma_F^\alpha, f \rangle$ is analytic for $2\alpha - 4 \notin \mathbb{N}$ and meromorphic for $\alpha \in \mathbb{C}$ with simple poles at $2\alpha - 4 \in \mathbb{N}$.

Proof. (a) For every $x \in \mathcal{M}$, we fix a normal coordinate system $\xi_x : y \rightarrow \mathbb{R}^4$ in order to parametrise points y in a normal neighbourhood of x . Consequently, on \mathcal{N}_2 , the squared geodesic distance divided by 2 can easily be expressed as

$$\sigma(x, y) = \frac{1}{2} \eta(\xi_x(y), \xi_x(y)) = \frac{1}{2} \xi_x^a \xi_{xa},$$

where η is the standard Minkowski metric given in Cartesian coordinates. Furthermore,

$$\left\langle \frac{1}{\sigma_F^\alpha}, f \right\rangle = \lim_{\epsilon \rightarrow 0^+} \int_{\mathcal{M}} \int_{\mathbb{R}^4} \frac{2^\alpha}{(\xi_x^a \xi_{xa} + i\epsilon)^\alpha} f(x, \xi_x) \sqrt{g(\xi_x)} d^4 \xi_x d\mu_g(x). \quad (3.15)$$

which is well defined for $f \in \mathcal{D}(\mathcal{N}_2)$. Observe that $1/(\xi_x^a \xi_{xa})^\alpha$, for $\xi_x^a \in \{z \in \mathbb{C}^4 \mid \text{Im}(z) \in V^\pm\}$, where V^\pm is the forward or past light cone with respect to the Minkowski metric, is analytic both in ξ and in α . Furthermore, in the limit $\epsilon \rightarrow 0^+$, $1/(\xi_x^a \xi_{xa} + i\epsilon)^\alpha$ can be seen as the boundary value of that analytic function. Since this function grows, at most, polynomially for large $1/\text{Im}(\xi_x^a \xi_{xa})$, its boundary value defines a distribution, see, e.g. ([Hö90] theorem 3.1.15). The analytic dependence on α is weakly preserved in the limit $\epsilon \rightarrow 0^+$, and, thus, the resulting distribution is weakly analytic. (b) The transformation defined in (3.14) acts on points parametrised by normal coordinates as $\xi \rightarrow \lambda \xi$. Furthermore, $1/(\xi_x^a \xi_{xa})^\alpha$ on $A \subset \mathbb{C}^4$ is homogenous of degree 2α with respect to the transformation $\xi \rightarrow \lambda \xi$. The statement follows from this observation, taking into account (3.14) and (3.15). (c) theorem 3.2.3 in [Hö90] ensures that the distribution $i \doteq 1/(\xi_x^a \xi_{xa})^\alpha \in \mathcal{D}'(\mathbb{R}^4 \setminus 0)$ has a unique extension t to 0 preserving the degree of homogeneity for every $2\alpha - 4 \notin \mathbb{N}$. Hence, $\mathbb{I} \otimes t$ defines a distribution on $\mathcal{M} \times \mathbb{R}^4$. Finally, notice that there exists a neighbourhood \mathcal{O} of the diagonal in \mathcal{M}_2 where $f(x, \xi_x) \sqrt{g(\xi_x)}$ is a smooth and compactly supported function for every $f \in C_0^\infty(\mathcal{O})$. Consequently, the statement follows by choosing a partition of unity adapted to \mathcal{O} . We refer to the discussion after definition 3.4 for general arguments why, in fact, theorem 3.2.3 in [Hö90] can be generalised to extensions of suitable homogeneous distributions to the diagonal rather than a point. \square

The previous proposition guarantees that $1/\sigma_F^\alpha$ is weakly meromorphic in α with simple poles at $2\alpha - 4 \in \mathbb{N}$. This property is preserved under taking linear combinations and multiplication by smooth functions. Consequently, the analytically regularised Feynman

propagator $H_F^{(\alpha)}$ defined by (3.12) is well defined on a normal neighbourhood of the diagonal and weakly meromorphic in α .

Proposition 3.6. *Consider a normal neighbourhood \mathcal{N}_2 of the diagonal $d_2 \in \mathcal{M}^2$. The following statements hold for the analytically continued Feynman propagator $H_F^{(\alpha)} \in \mathcal{D}'(\mathcal{N}_2)$ defined in (3.12).*

- (a) $\lim_{\alpha \rightarrow 0} H_F^{(\alpha)} = H_F$.
- (b) $\text{WF} H_F^{(\alpha)} \in \text{WF} H_F$.
- (c) *The scaling degree of $H_F^{(\alpha)}$ tends to $-\infty$ when the real part of α tends to ∞ .*

Proof. The proof of this proposition follows from the properties of $\sigma_F^{1+\alpha}$ obtained in proposition 3.5. In particular, (a) and (c) can be directly obtained from the weak analyticity, while b) follows from the fact that the distribution $1/\sigma_F^\alpha$ is well defined on $\mathcal{N}_2 \setminus d_2$ where it coincides either with $1/\sigma_+^\alpha$ or with $1/\sigma_-^\alpha$. In order to analyse the WF sets of $1/\sigma_\pm^\alpha$, we pass to a normal coordinate system and obtain $1/\sigma_\pm^\alpha = 2/(\xi^a \xi_a \pm i\epsilon \xi^0)^\alpha$. This distribution can be extended to a tempered distribution for every α , and, thus, its Fourier transform can be directly computed. One finds that, for $1/\sigma_\pm^\alpha$, only the null future/past directed directions do not decay rapidly, consequently, $H_F^{(\alpha)}$ restricted to $\mathcal{N}_2 \setminus d_2$ has $\text{WF}(H_F^{(\alpha)}) \subset \text{WF}(H_F)$. Finally, we observe that the extension of $H_F^{(\alpha)}$ to \mathcal{N}_2 may possess further singularities supported on the diagonal with singular directions orthogonal to d_2 . Hence, $\text{WF}(H_F^{(\alpha)}) \subset \text{WF}(H_F)$ still holds for $H_F^{(\alpha)} \in \mathcal{D}'(\mathcal{N}_2)$. \square

We are now able to discuss the analytical regularisation $\tau_\Gamma^{(\alpha)}$ (3.10) of the distributions τ_Γ given in (3.6) that appear in the graph expansion (3.4) of the time-ordered products \mathcal{T}_n (3.1). As anticipated in section 3.1, owing to the form of $H_F^{(\alpha)}$ given in (3.12), the relevant distributions that need to be discussed are $t_\Gamma^{(\alpha)}$ introduced in (3.13) and are analysed in the following proposition.

Proposition 3.7. *The operation,*

$$\langle t_\Gamma^{(\alpha)}, f \rangle := \int_{\mathcal{M}^n} \prod_{1 \leq i < j \leq n} \frac{1}{\sigma_F(x_i, x_j)^{l_{ij}(1+\alpha_{ij})}} f \, dx_1 \cdots dx_n \quad (3.16)$$

defined for $f \in \mathcal{D}(\mathcal{M}^n \setminus D_n \cap \mathcal{N})$ where \in is a normal neighbourhood of the total diagonal (cf. remark 3.1) has the following properties.

- (a) $t_\Gamma^{(\alpha)}$ is a distribution on $\mathcal{M}^n \setminus D_n \cap \in$.
- (b) $\langle t_\Gamma^{(\alpha)}, f \rangle$ is a continuous function for $\alpha = \{\alpha_{ij}\}_{i < j} \in \mathbb{R}^{n(n-1)/2}$.
- (c) $\langle t_\Gamma^{(\alpha)}, f \rangle$ is analytic for every α_{ij} with $i < j$ and, thus, a multivariate analytic function.

Proof. (a) The domain $\mathcal{M}^n \setminus D_n \cap \mathcal{N}$ is a disjoint union of connected components. On every connected component, \mathcal{C} $\sigma_F(x_i, x_j)$ equals either $\sigma_+(x_i, x_j)$ or $\sigma_+(x_j, x_i)$ depending on the causal relation between x_i and x_j , which is fixed in \mathcal{C} . Hence, on \mathcal{C} , the WF set of $\sigma_F(x_i, x_j)^{-1}$ is contained either in \mathcal{V}_+ or in \mathcal{V}_- , where $\mathcal{V}_{+/-} = \{(x, x', k, k') \in T^*\mathcal{M}^2 \setminus 0, (x, k) \sim (x', -k'), k \lhd/\rhd 0\}$.

Consequently, $\sigma_F(x_i, x_j)^{-1}$ satisfies the Hadamard condition up to a permutation of the arguments. The very same holds for the distributions $\sigma_F(x_i, x_j)^{l_{ij}(1+\alpha_{ij})}$ for every l_{ij} and every α_{ij} that have been discussed in proposition 3.5. Owing to the form of their WF set, the pointwise products of these distributions present in $t_\Gamma^{(\alpha)}$ are well defined because the Hörmander criterion for multiplication of distributions is satisfied. In fact, up to some fixed permutation of the arguments (x_1, \dots, x_n) , $t_\Gamma^{(\alpha)}$ satisfies the microlocal spectrum condition introduced in [BFK96]. Hence, $t_\Gamma^{(\alpha)}$ is a well-defined distribution on every connected component \mathcal{C} of $\mathcal{M}^n \setminus D_n \cap \mathcal{N}$, and, thus, it is well defined on $\mathcal{M}^n \setminus D_n \cap \mathcal{N}$. (b) In order to check continuity for $\alpha = \{\alpha_{ij}\}_{i < j} \in \mathbb{R}^{n(n-1)/2}$ in a fixed point $\bar{\alpha}$, we may analyse the distribution on a fixed connected component \mathcal{C} of the domain of $t_\Gamma^{(\alpha)}$ and may factorize the distribution in two parts. In fact, due to the WF set of $t_\Gamma^{(\alpha)}$ on \mathcal{C} , the factorisation $t_\Gamma^{(\alpha)} = t_\Gamma^{(\bar{\alpha})} \cdot \tau_\Gamma^{(\beta)}$ is unique where the integral kernel of $\tau_\Gamma^{(\beta)}$ is $\prod_{1 \leq i < j \leq n} \frac{1}{\sigma_F(x_i, x_j)^{\beta_{ij}}}$. For β in a sufficiently small neighbourhood of 0, $\tau_\Gamma^{(\beta)}$ is an integrable function that is differentiable for $\beta = 0$ as can be obtained by dominated convergence. Finally, the continuity is preserved by pointwise multiplication with $t_\Gamma^{(\bar{\alpha})}$. c) For an arbitrary but fixed pair of indices i, j , α_{ij} appears in the product displayed in (3.16) as $1/\sigma_F(x_i, x_j)^{l_{ij}(1+\alpha_{ij})}$, and we have already analysed the analyticity property of such a distribution in proposition 3.5. We will, thus, interpret $t_\Gamma^{(\alpha)}$ as a composition of distributions, namely, as $1/\sigma_F^{\alpha_{ij}} \circ z$ where z is an operator that maps $\mathcal{D}(\mathcal{M}^n \setminus D_n \cap \mathcal{N})$ to $\mathcal{D}'(\mathcal{M}^2 \setminus D_2 \cap \mathcal{N}_2)$ for a suitable $\mathcal{N}_2 \supset D_2 = d_2$. The ϵ -regularised integral kernel of z corresponds to the product present in (3.16) with the factor $1/\sigma_F^{\alpha_{ij}}$ removed. Because of the singular structure of z , for every $f \in \mathcal{D}(\mathcal{M}^n \setminus D_n \cap \mathcal{N})$, $\langle z, f \rangle$ is, in fact, a compactly supported smooth function supported on $\mathcal{M}^2 \setminus D_2 \cap \mathcal{N}_2$. Hence, the analysis of its composition with $1/\sigma_F^{\alpha_{ij}}$ is straightforward. These considerations imply separate analyticity of $t_\Gamma^{(\alpha)}$ in each α_{ij} , whereas, joint analyticity follows from the continuity proved in (b). \square

3.3. Generalised Euler operators and principal parts of homogeneous expansions

The next step in the strategy outlined at the end of section 3.1 is to extend the distributions $t_\Gamma^{(\alpha)}$, which are *a priori* defined only outside of the union of all partial diagonals D_n in a normal neighborhood \mathcal{N} of the total diagonal (cf. remark 3.1) to $D_n \cap \mathcal{N}$ and to show that this extension is weakly meromorphic in α_i upon setting $\alpha_{ij} = \alpha_i$ for all $i, j \in I \subset \{1, \dots, n\}$. As anticipated, we will prove this by using particular homogeneity properties of $t_\Gamma^{(\alpha)}$ with respect to the scaling transformations (3.14). Even if $t_\Gamma^{(\alpha)}$ is not homogeneous in the strong sense of definition 3.4, it has weaker homogeneity properties that are still strong enough in order to obtain the wanted results. In this section, we analyse analytically regularised distributions satisfying this weaker homogeneity condition, provide sufficient conditions for this weaker homogeneity to hold, and show how the principal part of a distribution of this type can be efficiently computed.

To this avail, we consider a normal neighbourhood \mathcal{N}_n of the total diagonal d_n (cf. remark 3.1) and define the **generalised Euler operator** $E_p : \mathcal{D}(\mathcal{N}_n) \rightarrow \mathcal{D}(\mathcal{N}_n)$ by

$$E_p f(x_1, \dots, x_n) := (-1)^p \lambda^{p+4(n-1)} \frac{d^p}{d\lambda^p} (\lambda^{-4(n-1)} f_\lambda(x))|_{\lambda=1}, \quad (3.17)$$

where the scaling transformation (3.14) is used. We then consider a family of distributions $t^{(\alpha)} \in \mathcal{D}'_1(\mathcal{N}_n \setminus d_n) \subset \mathcal{D}'(\mathcal{N}_n \setminus d_n)$ (cf. Definition 3.2) defined for α in some neighbourhood \mathcal{O}

of $0 \in \mathbb{C}$ and assume that $t^{(\alpha)}$ can be expanded as

$$t^{(\alpha)} = \sum_{k=0}^m t_k^{(\alpha)} + r^{(\alpha)},$$

where $t_k^{(\alpha)}$ are homogeneous with degree $a_k = -\delta_\alpha + k$, whose real part is smaller than or equal to $-4(n-1)$ and a remainder $r^{(\alpha)} \in \mathcal{D}'(\mathcal{N}_n \setminus d_n)$ that has a scaling degree smaller than $4(n-1)$ and can, thus, be uniquely extended to d_n for every $\alpha \in \mathcal{O}$ by ([BF00] theorem 6.9), see also the discussion after definition 3.2. Owing to its homogeneity, every $t_k^{(\alpha)}$ can be rewritten by means of the generalised Euler operator E_p as

$$\langle t_k^{(\alpha)}, f \rangle = \frac{1}{\prod_{j=0}^{p-1} (a_k + j + 4(n-1))} \langle t_k^{(\alpha)}, E_p f \rangle. \quad (3.18)$$

Note that, $E_p f(x_1, \dots, x_n)$ is smooth and vanishes for $y = (x_1, \dots, x_n) \rightarrow x = (x_1, \dots, x_l)$ as $C|y-x|^p$, i.e., it is in the class $O(|y-x|^p)$. For this reason, if p is chosen sufficiently large as $p > -a_k - 4(n-1)$, $t_k^{(\alpha)} \circ E_p$ possesses a unique extension to d_n . We recall that, in order to renormalise $t^{(\alpha)}$ for $\alpha = 0$ in the MS scheme, we have to subtract its principal part before computing the limit of vanishing α ,

$$\langle (t_k)_{\text{MS}}, f \rangle := \lim_{\alpha \rightarrow 0} (\langle t_k^{(\alpha)}, f \rangle - \text{pp} \langle t_k^{(\alpha)}, f \rangle).$$

However, if we use the representation of $t_k^{(\alpha)}$ provided by the right hand side of equation (3.18), its poles are manifestly exposed and can be subtracted easily. We recall that, since the original distribution $t_k^{(\alpha)}$ is well defined on $\mathcal{N}_n \setminus d_n$ even for $\alpha = 0$, the principal part we are subtracting can only be supported on d_n . We summarise this discussion in the following proposition.

Proposition 3.8. *Consider a normal neighbourhood \mathcal{N}_n of the total diagonal d_n and a distribution $t \in \mathcal{D}'(\mathcal{N}_n \setminus d_n)$. Assume that $t^{(\alpha)} \in \mathcal{D}'(\mathcal{N}_n \setminus d_n)$ is an analytic regularisation of t , i.e., $t^{(\alpha)}$ is weakly analytic for α in a neighbourhood \mathcal{O} of the origin of \mathbb{C} and $\lim_{\alpha \rightarrow 0} t^{(\alpha)} = t$. Moreover, assume that $t^{(\alpha)}$ can be decomposed as*

$$t^{(\alpha)} = \sum_{k=0}^m t_k^{(\alpha)} + r^{(\alpha)},$$

where $t_k^{(\alpha)}$ are weakly analytic distributions that scale homogeneously under transformations of the form (3.14) with degree $a_k = -\delta_\alpha + k$ and $r_k^{(\alpha)}$ is a weakly analytic distribution whose scaling degree towards d_n is strictly smaller than $4(n-1)$. Then, if $\text{WF}(t)$ and $\text{WF}(t^{(\alpha)})$ do not intersect the conormal bundle of the orbits of the scaling transformation (3.14), the following statements hold.

- (a) $t^{(\alpha)}$ can be extended to $i^{(\alpha)} \in \mathcal{D}'(\mathcal{N}_n)$ for every $\alpha \in \mathcal{O} \setminus \{0\}$.
- (b) $i^{(\alpha)}$ is weakly meromorphic for $\alpha \in \mathcal{O}$ with possible poles for $\alpha = 0$, and it is the unique weakly meromorphic extension of $t^{(\alpha)}$.
- (c) The pole of $i^{(\alpha)}$ in 0 is supported on d_n .
- (d) The limit

$\alpha \rightarrow 0$ can be considered after subtracting the pole part, namely,

$$\langle t_{\text{MS}}, f \rangle := \lim_{\alpha \rightarrow 0} (\langle i^{(\alpha)}, f \rangle - \text{pp} \langle i^{(\alpha)}, f \rangle)$$

is well defined for all $f \in \mathcal{D}(\mathcal{N}_n)$, and t_{MS} is an extension of t , which preserves the scaling degree.

Proof. The proof of *a*) and *b*) follows from the discussion after definition 3.4 as an application of ([Hö90] theorem 3.2.3) to every $t_k^{(\alpha)}$ restricted to every orbit of the action of the scaling transformation (3.14). Furthermore, since the scaling degree of $r^{(\alpha)}$ towards d_n is strictly smaller than $4(n-1)$, $r^{(\alpha)}$ possesses a unique extension towards d_n , cf. ([BF00] theorem 6.9). In order to prove *c*), we note that the original distribution $t^{(\alpha)}$ defined on $\mathcal{N}_n \setminus d_n$ is weakly analytic and that an explicit construction of the weakly meromorphic extension $i^{(\alpha)}$ to \mathcal{N}_n is provided by (3.18), choosing for every component $t_k^{(\alpha)}$ a sufficiently large p and using again ([BF00] theorem 6.9). Hence, the poles of $i^{(\alpha)}$ can only be supported on d_n . For this reason, after subtracting the principal part of the distribution, the limit $\alpha \rightarrow 0$ can safely be taken. The such obtained distribution prior to considering the limit $\alpha \rightarrow 0$ coincides with $t^{(\alpha)}$ on $\mathcal{N}_n \setminus d_n$, and the same holds in the limit $\alpha \rightarrow 0$. Consequently, t_{ms} is an extension of t . Finally, $\text{sd}(t_{ms}) = \text{sd}(t)$ because our assumptions and the above analysis imply that $\text{sd}(i^{(\alpha)}) = \text{sd}(t^{(\alpha)}) = \text{Re}(\delta_\alpha)$, $\text{sd}(\text{pp}(i^{(\alpha)})) \leq \text{Re}(\delta_\alpha)$, and $\lim_{\alpha \rightarrow 0} \text{Re}(\delta_\alpha) = \text{sd}(t)$. \square

We now discuss how equation (3.18) can be used in order to regularise the most singular part of a distribution $t^{(\alpha)}$ that is known to be of the form $t^{(\alpha)} = \sum_{k=0}^m t_k^{(\alpha)} + r^{(\alpha)}$ but where the distributions $t_k^{(\alpha)}$ are not explicitly known. To this end, observe that equation (3.18) implies

$$\langle t^{(\alpha)}, E_p f \rangle = \sum_{k=0}^m \left(\prod_{j=0}^{p-1} (a_k + j + 4(n-1)) \right) \langle t_k^{(\alpha)}, f \rangle + \langle r^{(\alpha)}, E_p f \rangle.$$

Moreover, we may assume without loss of generality, as in proposition 3.8, that the homogeneity degrees a_k of $t_k^{(\alpha)}$ are of the form $a_k = -\delta_\alpha + k$ where $\text{Re}(\delta_\alpha)$ is the scaling degree of $t^{(\alpha)}$. Consequently, $t_0^{(\alpha)}$ is the contribution with the highest scaling degree that may be extracted by introducing the coefficients,

$$c_k := \prod_{j=0}^{p-1} (a_k + j + 4(n-1)),$$

and considering

$$\langle t^{(\alpha)}, E_p f \rangle - c_0 \langle t^{(\alpha)}, f \rangle = \sum_{k=1}^m (c_k - c_0) \langle t_k^{(\alpha)}, f \rangle + \langle r^{(\alpha)}, E_p f \rangle - c_0 \langle r^{(\alpha)}, f \rangle, \quad (3.19)$$

where the distribution on the right hand side has a scaling degree smaller than $\text{Re}(\delta_\alpha) = -\text{Re}(a_0)$. Hence, although, in general, the distribution $t^{(\alpha)}$ does not scale homogeneously, equation (3.18) still holds up to distributions with a lower scaling degree. Knowing the decreasing degree of homogeneity of the components in the expansion of $t^{(\alpha)}$, we may use a recursive procedure in order to expose the pole part of this distribution. In fact, the previous discussion straightforwardly implies the validity of the following proposition.

Proposition 3.9. *We consider a distribution $t^{(\alpha)}$ with the properties assumed in proposition 3.8 and set*

$$u_0 := t^{(\alpha)}, \quad u_{k+1} := c_k u_k - u_k \circ E_{p_k}, \quad 0 \leq l < m,$$

where p_k are the smallest natural numbers chosen in such a way that $p_k + \text{Re}(a_k) + 4(n-1) > 0$ and $c_k := \prod_{j=0}^{p_k-1} (a_k + j + 4(n-1))$. Then, in order to expose the poles of $t^{(\alpha)}$, we may invert the recursive definition of u_k obtaining

$$t^{(\alpha)} = \frac{1}{c_0} \left(u_0 \circ E_{p_0} + \frac{1}{c_1} \left(u_1 \circ E_{p_1} + \dots + \frac{1}{c_n} (u_n \circ E_{p_n} + u_{n+1}) \right) \right). \quad (3.20)$$

In order to be able use the previous results for our purposes, we provide, in the next proposition, a criterion that is sufficient to ensure that a generic distribution can be decomposed into the sum of a homogeneous distribution and a remainder with lower scaling degree. We will use this criterion in order to prove that the distributions $t^{(\alpha)}$ defined in proposition 3.7 have the desired property.

Proposition 3.10. *Let \mathcal{N}_n be a normal neighbourhood of the total diagonal d_n , and suppose that $t \in \mathcal{D}'(\mathcal{N}_n)$ has scaling degree s_1 towards d_n under transformations of the form (3.14) and that there exists an α with $-\text{Re}(\alpha) = s_1$ such that $t \circ (E_1 + \alpha + 4(n-1))$ has scaling degree $s_2 < s_1$. Then, t can be decomposed into the sum of a homogeneous distribution with degree α and a remainder with scaling degree smaller than or equal to s_2 .*

Proof. We start by observing that, for every test function $f \in \mathcal{D}(\mathcal{N}_n)$,

$$F(\lambda, f) := \langle t, f_{1/\lambda} \rangle$$

is a continuous linear functional of f that is smooth in λ for $\lambda > 0$. Moreover, since the scaling degree of t is s_1 , $\lambda^a F(\lambda, f)$ vanishes in the limit $\lambda \rightarrow 0$ for every $a > s_1$ and for every $f \in \mathcal{D}(\mathcal{N}_n)$. Let us now consider

$$G(\lambda, f) := \langle (-E_1 + \alpha + 4(n-1))t, f_{1/\lambda} \rangle.$$

$G(\lambda, \cdot)$ is, again, a family of distributions on \mathcal{N}_n that depends smoothly on λ for positive λ . Furthermore, $\lambda^a G(\lambda, f)$ vanishes in the limit $\lambda \rightarrow 0$ for every $a > s_2$ and every $f \in \mathcal{D}(\mathcal{N}_n)$. Hence, $\lambda^{\alpha-1} G(\lambda, \cdot)$ tends to 0 in $\mathcal{D}'(\mathcal{N}_0)$ for $\lambda \rightarrow 0$, and the Banach–Steinhaus theorem implies that

$$|\lambda^a G(\lambda, f)| \leq C \sum_{\alpha \leq k} |\partial^\alpha f| \quad (3.21)$$

for every $a > s_2$, uniformly for f supported in a compact set $K \subset \mathcal{N}_n$ and for suitable C and k that do not depend on λ . After these preparatory considerations, we observe that G and F are related by the generalised Euler operator in the following way:

$$G(\lambda, f) = \lambda^{-\alpha+1} \frac{d}{d\lambda} \lambda^\alpha F(\lambda, f).$$

We can invert this relation to obtain

$$F(\lambda, f) = \frac{C(f)}{\lambda^\alpha} + \frac{1}{\lambda^\alpha} \int_0^\lambda \tilde{\lambda}^{\alpha-1} G(\tilde{\lambda}, f) d\tilde{\lambda},$$

where, $C(f)$ is a suitable constant that depends on f . We want to prove that $C(\cdot)$ is, in fact, a distribution. To this end, we note that, owing to the bound (3.21), the integral in $\tilde{\lambda}$ can be performed and the result of this integration is a distribution for every $\lambda > 0$ because $\text{Re}(\alpha) > s_2$. This implies that

$$C(f) = F(1, f) - \int_0^1 \tilde{\lambda}^{\alpha-1} G(\tilde{\lambda}, f) d\tilde{\lambda}$$

is a distribution because it is a linear combination of distributions. By construction, $F(1, f_{1/\lambda}) = F(\lambda, f)$ and $C \circ (E_1 + \alpha) = 0$, hence, C is a homogeneous distribution of degree α . By means of a direct computation, we also find that the scaling degree of the remainder $F(1, f) - C(f)$ is smaller than or equal to the scaling degree of G , which is s_2 . \square

3.3.1. The differential form of generalised Euler operators and homogeneous expansions of Feynman amplitudes. In order to make the previous discussion operative, we have to analyse the action of the generalised Euler operators E_p appearing in (3.18) on test functions. In fact, we will see that E_p corresponds to a particular geometric partial differential operator. To this end, we observe that $E_p = (E_1 - (p - 1))E_{p-1}$. Hence, knowing the differential form of the generalised Euler operator E_1 , it is possible to construct recursively every E_p .

Regarding the differential form of E_1 , we note that it can be written in terms of the geodesic distance and the van Vleck–Morette determinant⁶ u^2 as

$$E_1 f(x_1, \dots, x_n) = \sum_{j=2}^n (\sigma^a(x_j) \nabla_a^{x_j} - (2\sigma^a(x_j) \nabla_a^{x_j} \log(u(x_j, x_1)))) f(x_1, \dots, x_n),$$

where $\nabla_a^{x_j}$ indicates the a th component of the covariant derivative computed in x_j and $\sigma^a(x_j) := \nabla^{x_j a} \sigma(x_1, x_j)$. Considering the adjoint E_p^\dagger of E_p , we have $t \circ E_p = E_p^\dagger t$ where, using the relation $\square \sigma + 2\sigma^a \nabla_a \log(u) = 4$, we find for $p = 1$,

$$\begin{aligned} E_1^\dagger t(x_1, \dots, x_n) &= \sum_{j=2}^n (-\nabla_a^{x_j} \sigma^a(x_j) - 2\sigma^a(x_j) (\nabla_a^{x_j} \log(u(x_j, x_1)))) t(x_1, \dots, x_n) \\ &= - \left(4(n-1) + \sum_{j=2}^n \sigma^a(x_j) \nabla_a^{x_j} \right) t(x_1, \dots, x_n). \end{aligned} \quad (3.22)$$

We finally observe that the recursive identity for E_p also implies that E_p^\dagger can be constructed recursively starting from E_1^\dagger as

$$E_p^\dagger = E_{p-1}^\dagger (E_1^\dagger - (p-1)). \quad (3.23)$$

We proceed by showing that, upon applying E_1^\dagger introduced in (3.22) to a distribution $t_\Gamma^{(\alpha)}$ of the form

$$t_\Gamma^{(\alpha)} = \prod_{1 \leq i < j \leq n} \frac{1}{\sigma_F(x_i, x_j)^{l_{ij}(1+\alpha_{ij})}},$$

that has scaling degree $\text{sd}(t_\Gamma^{(\alpha)}) = \sum_{i < j} 2l_{ij}(1 + \text{Re}(\alpha_{ij}))$ towards the thin diagonal d_n , the result is a term proportional to $t_\Gamma^{(\alpha)}$ plus a remainder that has a lower scaling degree as foreseen in (3.19). Hence, proposition 3.10 implies that $t_\Gamma^{(\alpha)}$ can be written as a homogeneous distribution plus a remainder with a lower scaling degree. If the scaling degree of the remainder is not sufficiently low, we reiterate the procedure in order to obtain a full almost homogeneous expansion of the desired form.

In order to analyse this issue, we will only consider the relevant differential operator on \mathcal{M}^n appearing in E_1^\dagger , namely,

⁶ Recall that the square root of the van Vleck–Morette determinant coincides with the Hadamard coefficient u appearing in (2.2).

$$\rho := - \sum_{j=2}^n \sigma^a(x_j) \nabla_a^j. \quad (3.24)$$

We start by analysing the action of ρ on $\sigma(x_2, x_3)$ for x_2, x_3 in a normal neighbourhood of the point x_1 .

Lemma 3.11. *Let \mathcal{N}_{x_1} be a normal neighbourhood of the point x_1 , and let $x_2, x_3 \in \mathcal{N}_{x_1}$. Then,*

$$\rho\sigma(x_2, x_3) = 2\sigma(x_2, x_3) + G(x_1, x_2, x_3),$$

where G is a smooth function that vanishes in the limit $x_2, x_3 \rightarrow x_1$ as a monomial of order 4 in the normal coordinates of x_2 and x_3 centred in x_1 .

Proof. Using the notation in the proof of proposition 3.5, we write the action of ρ on $\sigma_{23} := \sigma(x_2, x_3)$ as

$$\rho\sigma_{23} = \xi_a(x_2)\sigma_{23}^a + \xi_b'(x_3)\sigma_{23}^{b'}.$$

Recall that σ_{23}^a is the covector in $T_{x_2}^*M$ cotangent to the unique geodesic joining x_2 and x_3 , that $-\sigma_{23}^{b'}$ is equal to the parallel transport of σ_{23}^a from x_2 to x_3 along the geodesic γ joining the two points, and that $\xi^c(x_i) := \sigma^c(x_1, x_i)$. Let us parametrise the image of γ with an affine parameter λ such that $x(0) = x_2$ and $x(1) = x_3$. In order to simplify the notation, we indicate, by $t(\lambda)$, the tangent vector of the geodesic in $x(\lambda)$. As argued before, we have

$$t^a(0) = \sigma_{23}^a \quad \text{and} \quad t^{b'}(1) = -\sigma_{23}^{b'}.$$

Consequently,

$$\begin{aligned} \rho\sigma_{23} &= \xi^a t_a(0) - \xi^b t_b(1) = - \int_0^1 \frac{d}{d\lambda} (\xi^a t_a)(\lambda) d\lambda \\ &= - \int_0^1 t^a \nabla_t \xi_a d\lambda = \int_0^1 t^a t^b \sigma_{ab}(x(\lambda), x_1) d\lambda, \end{aligned}$$

where $\sigma_{ab} := \nabla_a \nabla_b \sigma$. If we now consider the covariant Taylor expansion of $\sigma_{ab}(x(\lambda), x_1)$ around $x(\lambda)$ (see, e.g. [PPV11]), we find that $E_{ab}(x, x_1) := \sigma_{ab}(x, x_1) - g_{ab}(x)$ is a smooth function that vanishes for $x \rightarrow x_1$ as $O(\sigma(x, x_1))$, hence,

$$\rho\sigma_{23} = \int_0^1 t^a t^b g_{ab}(x(\lambda)) d\lambda + \int_0^1 t^a t^b E_{ab}(x(\lambda), x_1) d\lambda = 2\sigma(x_2, x_3) + G(x_1, x_2, x_3),$$

where the remainder is smooth because of the smoothness of the metric g and can be further expanded as

$$\begin{aligned} G(x_1, x_2, x_3) &= \int_0^1 t^a(\lambda) t^b(\lambda) (\sigma_{ab}(x(\lambda), x_1) - g_{ab}(x(\lambda))) d\lambda \\ &= \int_0^1 t^a(\lambda) t^b(\lambda) t^c(\lambda) t^d(\lambda) R_{acbd}(x(\lambda)) d\lambda + \dots \\ &= O(|\xi(x_2)|^4 + |\xi(x_3)|^4), \end{aligned} \quad (3.25)$$

where the absolute value of the normal coordinates $|\xi(x_i)|$ of x_i , $i = 2, 3$ is intended in the Euclidean sense. \square

We are now in position to analyse the action of ρ on the distribution $t_\Gamma^{(\alpha)}$ introduced in (3.16).

Proposition 3.12. The distribution $t_\Gamma^{(\alpha)}$ introduced in (3.16) can be written as a sum of homogeneous distributions with respect to scaling towards the total diagonal d_n plus a remainder. The degrees of homogeneity of these homogeneous distributions are contained in the following set:

$$\left\{ k - \sum_{1 \leq i < j \leq n} 2l_{ij}(1 + \alpha_{ij}), k \in \mathbb{N} \cup \{0\} \right\}.$$

Proof. We perform this analysis with ϵ in σ_F taken to be strictly positive. We start by applying ρ given in (3.24) to $t_\Gamma^{(\alpha)}$. Thanks to the results stated in lemma 3.11, we have

$$\rho t_\Gamma^{(\alpha)} = C t_\Gamma^{(\alpha)} + r_\Gamma^{(\alpha)},$$

where the constant C is

$$C = - \sum_{1 \leq i < j \leq n} 2l_{ij}(1 + \alpha_{ij}).$$

Furthermore, lemma 3.11 and, in particular, (3.25) imply that the remainder $r_\Gamma^{(\alpha)}$ has a scaling degree towards d_n , which is lower than the one of $t_\Gamma^{(\alpha)}$ by at least 2,

$$\text{sd}(r_\Gamma^{(\alpha)}) \leq \text{sd}(t_\Gamma^{(\alpha)}) - 2 = \sum_{1 \leq i < j \leq n} 2l_{ij}(1 + \text{Re}(\alpha_{ij})) - 2. \quad (3.26)$$

Proposition 3.10, then, implies that the distribution $t_\Gamma^{(\alpha)}$ can be written as a homogeneous distribution of degree C plus a remainder with lower scaling degree. In order to finalise the proof, we need to control the recursive application of ρ , therefore, we discuss the application of ρ on $\rho^n t_\Gamma^{(\alpha)}$ for an arbitrary n . Let us start with $n = 1$. In this case, we observe that the relevant contribution is the one given by the remainder $\rho r_\Gamma^{(\alpha)}$, which reads

$$r_\Gamma^{(\alpha)} = \sum_{1 \leq i < j \leq n} l_{ij}(1 + \alpha_{ij}) \frac{G(x_1, x_i, x_j)}{\sigma_F(x_i, x_j)} t_\Gamma^{(\alpha)}.$$

Note that, for every $i < j$, $\sigma_F(x_i, x_j) t_\Gamma^{(\alpha)}$ has the same structure, such as $t_\Gamma^{(\alpha)}$ but the scaling degree $\text{sd}(t_\Gamma^{(\alpha)}) + 2$, whereas $G(x_1, x_i, x_j)$ defined in (3.25) is a smooth function, whose Taylor expansion for x_i, x_j around x_1 starts with components of order 4. Hence, if we apply ρ to $r_\Gamma^{(\alpha)}$, we obtain a constant multiple of $r_\Gamma^{(\alpha)}$ plus a remainder that has a scaling degree lower or equal to $\text{sd}(r_\Gamma^{(\alpha)}) - 1$ where the difference with respect to (3.26) stems from the fact that G can be expanded as a polynomial in $\sigma_a(x_i)$, whose lowest components are monomials of degree 4 multiplied by curvature tensors. These monomials are homogeneous and, thus, contribute to the degree of homogeneity of $\rho r_\Gamma^{(\alpha)}$, while the contributions in G with degrees higher or equal to 5 influence the scaling degree of the remainder. Repeating this analysis for a generic n , we find that similar results hold when ρ is applied recursively to the remainder. Consequently, an iterated application of proposition 3.10 implies that the distribution $t_\Gamma^{(\alpha)}$ can be written as a finite sum of homogeneous distributions plus a remainder. Furthermore, since

the scaling degree of these distributions is always finite, the degree of homogeneity of these components is finite as well. \square

As outlined at the end of section 3.1, we can use proposition 3.12 in conjunction with propositions 3.8, 3.9 in order to extend the distributions $t_\Gamma^{(\alpha)}$ in a unique and weakly meromorphic fashion to the union of all partial diagonal D_n in a normal neighborhood of the total diagonal (cf. remark 3.1) and in order to compute the relevant pole part of this extension as used in the forest formula, cf. (3.9), (3.10), (3.12). To this avail, we stress that proposition 3.12 holds, in particular, for any subgraph Γ_I , $I \subset \{1, \dots, n\}$ of Γ and the corresponding distribution $t_{\Gamma_I}^{(\alpha)}$, which is obtained by omitting all factors in $t_\Gamma^{(\alpha)}$ that correspond to edges not contained in Γ_I . Finally, the recursive structure of the forest formula (3.9) implies that we are dealing not only with expressions of the form $t_{\Gamma_I}^{(\alpha)}|_{\alpha_{ij}=\alpha_I \forall i,j \in I}$, but also with expressions that are of this form up to a subtraction of their principal part. However, our above analysis and, in particular, the discussion in the proof of proposition 3.8 implies that propositions 3.12, 3.9 also hold in this case.

Remark 3.13. Proposition 3.9 and the above analysis imply that our renormalisation scheme is, in fact, a particular form of differential renormalisation. Notwithstanding, the advantage of formulating this scheme in terms of analytic regularisation and minimal subtraction is the ability to define the renormalisation scheme in a closed form at all orders by means of the forest formula (3.9).

3.4. Properties of the minimal subtraction scheme

We conclude the general analysis of the renormalisation scheme introduced in this work by demonstrating that this scheme satisfies—up to one property we will mention at the end of this section—all axioms of [HW02, HW01, HW04], which, as argued in these works, any physically meaningful scheme to renormalise time-ordered products should satisfy. We refer to these works for a detailed formulation and discussion of these axioms. In addition to showing these properties of the scheme, we argue that it preserves invariance under any space–time isometries present.

Proposition 3.14. *The time-ordered product \mathcal{T}_n defined by means of (3.9) where the quantities appearing in this formula are defined by means of (3.8), (3.11), (3.10), (3.12) and where we recall remark 3.1 have the following properties.*

- (a) \mathcal{T}_n is symmetric and satisfies the causal factorisation condition.
- (b) \mathcal{T}_n is unitary.
- (c) \mathcal{T}_n is local and covariant.
- (d) \mathcal{T}_n satisfies the microlocal spectrum condition.
- (e) \mathcal{T}_n is ϕ independent.
- (f) \mathcal{T}_n satisfies the Leibniz rule.
- (g) \mathcal{T}_n satisfies the principle of perturbative agreement (PPA) for perturbations of the generalised mass term μ in the free Klein–Gordon equation $P\phi := (-\square + \mu)\phi = 0$.
- (h) If the space–time (\mathcal{M}, g) has nontrivial isometries and if the Feynman propagator H_F is chosen such as to be invariant under these isometries, then \mathcal{T}_n is invariant under these isometries as well.

Proof. (a) holds because we constructed the renormalised time-ordered product by means of the forest formula (3.9) and because, as implied by proposition 3.8, all counterterms subtracted in the forest formula are local. (b) Unitarity holds because the operation of extracting the relevant principal part of a regularised amplitude $\tau_{\Gamma}^{(\alpha)}$ commutes with complex conjugation (even if α is not real). (c) The regularised amplitudes $\tau_{\Gamma}^{(\alpha)}$ satisfy locality and covariance. Upon setting $\alpha_{ij} = \alpha_I$ for $i, j \in I \subset \{1, \dots, n\}$, $\tau_{\Gamma}^{(\alpha)}$ is weakly meromorphic in α_I . Thus, locality and covariance hold for each term in the corresponding Laurent series and, consequently, after subtracting the principal part of this series. (d) As argued in the proof of proposition 3.7, the distributions $t_{\Gamma}^{(\alpha)}$ defined in (3.16) satisfy the microlocal spectrum condition, i.e., they have the correct WF set. Consequently, the regularised amplitudes $\tau_{\Gamma}^{(\alpha)}$ have the correct WF set as well. As $\tau_{\Gamma}^{(\alpha)}$ is weakly meromorphic in the sense recalled in the proof of (c), each term in the corresponding Laurent series has a WF set bounded by the WF set of $\tau_{\Gamma}^{(\alpha)}$. Consequently, the microlocal spectrum condition holds after subtracting the principal part and considering the limit of vanishing regularisation parameters. (e) This property follows directly from the construction. In particular, the subtraction of counterterms is defined in terms of numerical distributions and is independent of the field ϕ . (f) In analogy to (b), the Leibniz rule holds because the operation of extracting the relevant principal part of a regularised amplitude $\tau_{\Gamma}^{(\alpha)}$ commutes with all partial differential operators. (g) The PPA for perturbations of the generalised mass term μ demands essentially that, upon setting $\mu = \mu_0 + \mu_1$, the renormalisation of \mathcal{T}_n commutes with the operation of perturbatively expanding quantities in μ_1 around μ_0 . A Feynman propagator H_F depends on μ only via the Hadamard coefficients v and w in (2.2). However, in the definition of the analytically regularised H_F^{α} in (3.12) and the corresponding regularised amplitudes $\tau_{\Gamma}^{(\alpha)}$ defined in (3.10), these coefficients are not altered, but only the σ -dependent terms multiplying these coefficients are modified. Consequently, the analytic regularisation and minimal subtraction scheme we consider commutes with a perturbative expansion in μ_1 around μ_0 . (h) As recalled in (g), all operations in our analytic regularisation and minimal subtraction scheme act directly on quantities defined entirely in terms of the geometric quantity σ . As σ is invariant under any space–time isometries present, the renormalisation scheme preserves this invariance. \square

Remark 3.15. Note that the PPA, as introduced in [HW04], also poses conditions on \mathcal{T}_1 , i.e., the renormalisation of local and covariant Wick polynomials, which we omitted in our analysis, cf. Footnote 1 on p. 10. However, given \mathcal{T}_n for $n > 1$, \mathcal{T}_1 can be adjusted in order to satisfy the PPA for changes of μ by using, e.g. ([DHP15] theorem 3.3). Moreover, the PPA, as introduced in [HW04], further demands that, setting $g = g_0 + g_1$, the renormalisation also commutes with perturbatively expanding quantities in g_1 around an arbitrary but fixed background metric g_0 . Since σ depends on g , it is not easy to check whether a perturbative expansion in g_1 commutes with our analytic regularisation and minimal subtraction scheme and, thus, it might well be that the renormalisation scheme discussed in the present work fails to satisfy this part of the PPA. However, if this is the case, the scheme can be modified according to the construction in [HW04] in order to also satisfy this condition while preserving the other properties in proposition 3.14, including the invariance under any space–time isometries present.

Remark 3.16. We have omitted the explicit dependence of renormalised quantities on the mass scale M appearing in the analytically regularised Feynman propagator $H_F^{(\alpha)}$ (3.12), but our analysis implies that the dependence of these quantities on M is such that all renormalised

quantities are polynomials of (derivatives of) $\log(M^2\sigma_F(x_i, x_j))$, see also the examples in the next section. Thus, the renormalisation group flow with respect to changes in M may easily be computed.

3.5. Examples

In this section, we illustrate the method developed in section 3.3 to explicitly compute renormalised quantities in our scheme by considering, first, the example of the fish graph and the sunset graph, i.e., Δ_F^n for $n = 2, 3$. These pointwise powers of the Feynman propagator are the only ones occurring in renormalisable scalar field theories in four space–time dimensions. Afterwards, we will consider a triangular graph in section 3.5.3 in order to illustrate the method in the case of more than two vertices. Recalling remark 3.1, we will work only on subsets of the space–time where the geodesic distance is well defined without loss of generality.

In the special case of Δ_F^n , we are dealing with distributions that are already defined on $\mathcal{M}^2 \setminus d_2$ and have to be extended to \mathcal{M}^2 . In order to accomplish this task, we will use (3.18) in order to expose the poles before subtracting them. In this context, we note that E_1^\dagger given in (3.22) applied to a distribution t whose integral kernel $t(\sigma_F)$ depends on x, y only via $\sigma_F(x, y)$ can be further simplified. In particular, introducing $t_1(\sigma_F)$ such that $\nabla^a t_1(\sigma_F) = \sigma^a t(\sigma_F)$, we have

$$\begin{aligned} E_1^\dagger t(\sigma_F) &= -(4 + \sigma^a \nabla_a) t(\sigma) = -\nabla_a \sigma^a t - 2\sigma^a (\nabla_a \log(u)) t(\sigma_F) \\ &= -\square t_1(\sigma_F) - 2 \frac{\nabla_a u}{u} \nabla^a t_1(\sigma_F), \end{aligned} \quad (3.27)$$

where x is considered to be arbitrary but fixed and all the covariant derivatives are taken with respect to y .

3.5.1. Computation of the renormalised fish and sunset graphs in our scheme. We recall that the Feynman propagator $\Delta_F(x, y) := \langle \phi(x) \cdot_{T_\Delta} \phi(y) \rangle_\Omega$ in any Hadamard state Ω is locally of the form

$$\Delta_F(x, y) = \frac{1}{8\pi^2} \left(\frac{u(x, y)}{\sigma_F(x, y)} + v(x, y) \log(M^2 \sigma_F(x, y)) \right) + w(x, y), \quad \sigma_F := \sigma + i\epsilon. \quad (3.28)$$

From (3.28), we can infer that, in order to renormalise Δ_F^2 and Δ_F^3 , i.e., in order to extend them from $\mathcal{M}^2 \setminus d_2$ to \mathcal{M}^2 , we need to renormalise the three distributions,

$$\frac{1}{\sigma_F^2}, \quad \frac{\log(M^2 \sigma_F)}{\sigma_F^2}, \quad \frac{1}{\sigma_F^3}, \quad (3.29)$$

because all other occurring powers of σ_F , i.e., $\sigma_F^{-m} \log^n(\sigma_F)$ for $m \in \{0, 1\}$ and $n \in \{0-3\}$ have a scaling degree for $y \rightarrow x$ smaller than 4 and, thus, can be uniquely extended to the diagonal. To this avail, we define

$$\sigma_{a_1 \dots a_n} := \nabla_{a_n} \dots \nabla_{a_1} \sigma, \quad [B](x) := B(x, x),$$

where the covariant derivatives are taken with respect to x and B is a general bitensor and recall the following basic identities satisfied by σ :

$$\sigma_a \sigma^a = 2\sigma, \quad \sigma_{ab} \sigma^b = \sigma_a, \quad \square \sigma = 4 - 2 \frac{\sigma^a \nabla_a u}{u}. \quad (3.30)$$

For our purposes, it will prove useful to use the last identity in the form

$$\square \sigma_F = 4 + f \sigma_F \quad \text{with} \quad f := -2 \frac{\sigma^a \nabla_a u}{u \sigma_F}, \quad (3.31)$$

where f is a distribution, which, considered as a distribution in y for fixed x , has scaling degree zero for $y \rightarrow x$ as can be seen from the covariant Taylor expansion $u = [u] + ([\nabla_a u] - \nabla_a[u])\sigma^a + \mathcal{R}_u = 1 + \mathcal{R}_u$ where the remainder \mathcal{R}_u vanishes towards the diagonal faster than σ_a (see, e.g. ([PPV11] section 5)).

Remark 3.17. As f has a vanishing scaling degree for $y \rightarrow x$, the pointwise product $f(x, y)t(x, y)$ with any bidistribution t of scaling degree for $y \rightarrow x$ lower than 4 may be uniquely extended to the diagonal. However, we will also encounter expressions that are naively of the form $f(x, y)\delta(x, y)$ and that are *a priori* ill defined because f is, in general, divergent for x and y lightlike related and, thus, not continuous on the diagonal. Notwithstanding, the distribution $f(x, y)\delta(x, y)$, which is well defined and identically vanishing outside of the diagonal $x = y$, may be extended to the diagonal. In fact, our scheme in which expressions of the form $f(x, y)\delta(x, y)$ appear as $\alpha \rightarrow 0$ limits of particular weakly analytic expressions provides a unique and nonvanishing extension of $f(x, y)\delta(x, y)$ to the diagonal by the very analyticity of the aforementioned expressions. In particular, our scheme implies the following unique and well-defined definitions of distributions on \mathcal{M}^2 .

$$f \square \frac{\log^n(M^2 \sigma_F)}{\sigma_F} := \lim_{\alpha \rightarrow 0} f \square \frac{\log^n(M^2 \sigma_F)}{\sigma_F^{1+\alpha}}, \quad n \geq 0. \quad (3.32)$$

Hereby, uniqueness and weak analyticity of $f \square (\log^n(M^2 \sigma_F)/\sigma_F^{1+\alpha})$ follow from arguments used throughout section 3.3.

From proposition 3.5, we know that $1/\sigma_F^{n+\alpha}$ is weakly meromorphic in α . In order to compute the Laurent series, we use the above-mentioned identities for σ and obtain

$$\frac{1}{\sigma_F^{n+1+\alpha}} = \frac{1}{2(n+\alpha)(n-1+\alpha)} (\square + (n+\alpha)f) \frac{1}{\sigma_F^{n+\alpha}},$$

in accordance with (3.18) and (3.27).

Using this and recalling remark 3.17, we may compute the following Laurent series where we recall that, in $\Delta_F^{(\alpha)}$ (3.12), we use the same (arbitrary) constant M present in the logarithmic term of (3.28) to correct for the change in dimension and a sufficiently regular function k for later purposes,

$$\begin{aligned} \frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} &= \frac{1}{2} (\square + f) \left(\frac{1}{\alpha \sigma_F} - \frac{\log(M^2 \sigma_F)}{\sigma_F} \right) \\ &\quad - \frac{\log(k^2)}{2} (\square + f) \frac{1}{\sigma_F} - \square \frac{1}{2\sigma_F} + O(\alpha), \\ \frac{d}{d\alpha} \frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} &= \frac{1}{2} (\square + f) \left(-\frac{1}{\alpha^2 \sigma_F} + \frac{\log^2(M^2 \sigma_F)}{2\sigma_F} \right) \end{aligned}$$

$$\begin{aligned}
& + \square \frac{\log(M^2 \sigma_F) + 1}{2\sigma_F} + \log^2(k^2)(\square + f) \frac{1}{4\sigma_F} \\
& + \log(k^2) \left(\square \frac{1}{2\sigma_F} + (\square + f) \frac{\log(M^2 \sigma_F)}{2\sigma_F} \right) + O(\alpha), \\
\frac{1}{(Mh)^{2\alpha}} \frac{1}{\sigma_F^{3+\alpha}} &= \frac{1}{8} (\square + 2f)(\square + f) \left(\frac{1}{\alpha \sigma_F} - \frac{\log(M^2 \sigma_F)}{\sigma_F} \right) \\
& - \frac{\log(k^2)}{8} (\square + 2f)(\square + f) \frac{1}{\sigma_F} \\
& - \frac{1}{16} ((5\square + 8f)(\square + f) - 2(\square + 2f)f) \frac{1}{\sigma_F} + O(\alpha).
\end{aligned} \tag{3.33}$$

Note that, by means of lemma 3.18 b), one may explicitly check that the pole terms in these Laurent series are local expressions as expected.

Using the Laurent series, the lowest renormalised powers of σ_F may be defined and may be computed as⁷, where we recall, once again, remark 3.17,

$$\begin{aligned}
\left(\frac{1}{\sigma_F^2} \right)_{\text{MS}} &:= \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} - \text{pp} \frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} \right) \\
&= -\frac{1}{2} (\square + f) \frac{\log(M^2 \sigma_F)}{\sigma_F} - \square \frac{1}{2\sigma_F}, \\
\left(\frac{\log(M^2 \sigma_F)}{\sigma_F^2} \right)_{\text{MS}} &:= -\lim_{\alpha \rightarrow 0} \left(\frac{d}{d\alpha} \frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} - \text{pp} \frac{d}{d\alpha} \frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} \right) \\
&= -\frac{1}{4} (\square + f) \frac{\log^2(M^2 \sigma_F)}{\sigma_F} - \square \frac{\log(M^2 \sigma_F) + 1}{2\sigma_F}, \\
\left(\frac{1}{\sigma_F^3} \right)_{\text{MS}} &:= \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{3+\alpha}} - \text{pp} \frac{1}{M^{2\alpha}} \frac{1}{\sigma_F^{3+\alpha}} \right) \\
&= -\frac{1}{8} (\square + 2f)(\square + f) \frac{\log(M^2 \sigma_F)}{\sigma_F} \\
&\quad - \frac{1}{16} ((5\square + 8f)(\square + f) - 2(\square + 2f)f) \frac{1}{\sigma_F}.
\end{aligned} \tag{3.34}$$

Finally, $(\Delta_F^2)_{\text{MS}}$ and $(\Delta_F^3)_{\text{MS}}$ are defined and are computed by expanding the unrenormalised powers Δ_F^2 and Δ_F^3 and by replacing the three problematic expressions (3.29) by their renormalised versions (3.34).

3.5.2. Alternative computation of the renormalised fish and sunset graphs. As a preparation towards the application of our renormalisation scheme to QFT in cosmological space-times, we will now derive an alternative way to compute $(\Delta_F^2)_{\text{ms}}$ and $(\Delta_F^3)_{\text{ms}}$, which is better suited for practical computations. We start by stating and by proving a few distributional identities.

⁷ Note that here, we use a definition of the analytic regularisation of the logarithm in terms of a direct derivative rather than a limit of differences, such as in (3.12). While the two definitions differ up to a constant factor in the principal part, they coincide in the constant regular part and thus give the same $(\sigma_F^{-2} \log(M^2 \sigma_F))_{\text{MS}}$.

Lemma 3.18. *The following distributional identities hold.*

(a) *For any continuous F_0 and any twice continuously differentiable F_2 ,*

$$\begin{aligned}\sigma F_0 \delta &= 0, & \sigma_a F_0 \delta &= 0, & F_0 \nabla_{\nabla \sigma} \delta &= -[F_0 \square \sigma] \delta, \\ F_2 \square \delta &= [\square F_2] \delta + \square[F_2] \delta - 2 \nabla^a [\nabla_a F_2] \delta.\end{aligned}$$

(b)

$$\begin{aligned}(\square + f) \frac{1}{\sigma_F} &= 8\pi^2 i \delta, \\ (\square + 2f)(\square + f) \frac{1}{\sigma_F} &:= \lim_{\alpha \rightarrow 0} (\square + 2f)(\square + f) \frac{1}{\sigma_F^{1+\alpha}} = 8\pi^2 i \left(\square - \frac{R}{3} \right) \delta\end{aligned}$$

(c) *For all $n_1, n_2, n_3 \in \mathbb{N}_0$ and $n_4, n_5, n_6 \in \{0, 1\}$ with $n_2 - n_3 + n_4 \geq -1$,*

$$\begin{aligned}\log^{n_1}(\sigma_F) (\sigma_F^{n_4}) \sigma_F^{n_2} \left(\frac{1}{\sigma_F^{n_3}} \right)_{\text{MS}} &= \log^{n_1}(\sigma_F) (\sigma_F^{n_4}) \sigma_F^{n_2 - n_3}, \\ \square \log(\sigma_F) &= \frac{\square \sigma - 2}{\sigma_F}, & \nabla_a \frac{\log^{n_5}(\sigma_F)}{\sigma_F^{n_6}} &= \frac{(n_5 - n_6 \log^{n_5}(\sigma_F)) \nabla_a \sigma}{\sigma_F^{n_6+1}}.\end{aligned}$$

(d)

$$\sigma_F \left(\frac{1}{\sigma_F^3} \right)_{\text{MS}} = \left(\frac{1}{\sigma_F^2} \right)_{\text{MS}}.$$

Proof. (a) These identities follow from $B\delta = [B]\delta$ for any continuous bitensor B , $[\sigma] = 0$, $[\sigma_a] = 0$ and the definition of weak derivatives. (b) The first identity holds in Minkowski space-time because $1/(8\pi^2 \sigma_F)$ is the Feynman propagator of the massless vacuum state. In curved space-times, (3.30) imply that $(\square + f)1/\sigma_F$ vanishes outside of the origin and, thus, must be a sum of derivatives of δ distributions. Because σ depends smoothly on the metric, the coefficients in this sum must be smooth functions of the metric with appropriate mass dimension, and, thus, $(\square + f)1/\sigma_F = c\delta$ with a constant c that can be fixed in Minkowski space-time. In order to prove the second identity, we recall remark 3.17 and observe that it is sufficient to compute

$$t := \lim_{\alpha \rightarrow 0} f(\square + f) \frac{1}{\sigma_F^{1+\alpha}}.$$

This expression has, for $y \rightarrow x$, a scaling degree ≤ 4 that vanishes outside of $x = y$, depends smoothly on the metric, is covariant, and has mass dimension 6. Consequently, $t = cR\delta$ where the dimensionless constant c can be computed on any space-time with $R \neq 0$. Moreover, in any space-time where f is actually continuous in a neighbourhood of the diagonal, we have $t(x, y) = 8\pi i f(x, x) \delta(x, y)$. A space-time that satisfies both properties is (the patch of) a de Sitter space-time defined in conformal coordinates by the metric line element,

$$ds^2 = \frac{1}{H^2 \tau^2} (-d\tau^2 + dx^2),$$

on $(-\infty, 0) \times \mathbb{R}^3$, where H is a constant. On this space-time, we have $R = 12H^2$ and

$$\mu^2 := 2H^2 \sigma(\tau_1, \vec{x}_1, \tau_2, \vec{x}_2) = \cos^{-1} \left(\frac{\tau_1^2 + \tau_2^2 - (\vec{x}_1 - \vec{x}_2)^2}{2\tau_1 \tau_2} \right)$$

see, e.g. [A185], where analytic continuation of \cos^{-1} is understood for timelike separations. From this, one can infer

$$\square \sigma = 1 + 3\mu \cot(\mu) \quad \Rightarrow \quad f = \frac{\square \sigma - 4}{\sigma} = 6H^2 \frac{\mu \cot(\mu) - 1}{\mu^2} = -\frac{R}{6} + O(\mu^2),$$

which demonstrates that, on de Sitter space-time, f is continuous in a neighbourhood of the diagonal with $f(x, x) = -R/6$. (c) The distributions on both sides of each equation, considered as distributions in y for fixed x , have the same scaling degree < 4 for $y \rightarrow x$ and agree outside of the diagonal. Thus, they agree also on the diagonal as unique extensions. (d) As in the proof of (a), we observe that the potential local correction term on the right hand side must be a sum of derivatives of δ with coefficients that depend smoothly on the metric because σ does. Thus, the correction term must be of the form $c\delta$ with a constant c that can be computed in Minkowski space-time. This computation may be performed by using (3.30), the previous statements of this lemma, and the following identities, which are valid in Minkowski space-time for any function F such that $F(\sigma_F)$ is a distribution,

$$\begin{aligned} \sigma_F \square F(\sigma_F) &= \square \sigma_F F(\sigma_F) - 4F(\sigma_F) - 2\nabla_{\nabla \sigma_F} F(\sigma_F), \\ \sigma_F \square^2 F(\sigma_F) &= \square^2 \sigma_F F(\sigma_F) - 4\square F(\sigma_F) - 4\square \nabla_{\nabla \sigma_F} F(\sigma_F), \end{aligned}$$

whereby, one finds that $c = 0$. □

These identities can be used to compute $(\Delta_F^2)_{\text{MS}}$ and $(\Delta_F^3)_{\text{MS}}$ in an alternative way under certain conditions.

Proposition 3.19. *Let (\mathcal{M}, g) be such that \mathcal{M} is a normal neighbourhood and let Δ_F be a distribution on \mathcal{M}^2 of Feynman–Hadamard form (3.28). Then, the following identities hold.*

(a) *If Δ_F^α is a well-defined distribution that is weakly meromorphic in α , then,*

$$(\Delta_F^2)_{\text{MS}} = \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} \Delta_F^{2+\alpha} - \text{pp} \frac{1}{M^{2\alpha}} \Delta_F^{2+\alpha} \right) + \frac{i \log(8\pi^2)}{16\pi^2} \delta.$$

(b) *If Δ_F^α is a well-defined distribution that is weakly meromorphic in α , then,*

$$(\Delta_F^2 \log(M^{-2} \Delta_F))_{\text{MS}} = \lim_{\alpha \rightarrow 0} \left(\frac{d}{d\alpha} \frac{1}{M^{2\alpha}} (\Delta_F)^{2+\alpha} - \text{pp} \frac{d}{d\alpha} \frac{1}{M^{2\alpha}} \Delta_F^{2+\alpha} \right) - \frac{i \log^2(8\pi^2)}{32\pi^2} \delta.$$

(c) *If Δ_F^α is a well-defined distribution that is weakly meromorphic in α and $[v] = 0$, then,*

$$(\Delta_F^3)_{\text{MS}} = \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} \Delta_F^{3+\alpha} - \text{pp} \frac{1}{M^{2\alpha}} \Delta_F^{3+\alpha} \right) + \frac{i((1 + 2 \log(8\pi^2))R + 192\pi^2[w])}{48(8\pi^2)^2} \delta.$$

Proof. (a) Setting $h = 8\pi^2\sigma_F\Delta_F$ and $k = \sqrt{8\pi^2/h}$, we obtain

$$\frac{1}{M^{2\alpha}}\Delta_F^{2+\alpha} = \frac{h^2}{(8\pi^2)^2} \frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}}.$$

Using (3.33), $[h^2] = [u^2] = 1$, and lemma 3.18 a), b), c), we may compute

$$\begin{aligned} & \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}}\Delta_F^{2+\alpha} - \text{pp} \frac{1}{M^{2\alpha}}\Delta_F^{2+\alpha} \right) \\ &= \frac{h^2}{(8\pi^2)^2} \lim_{\alpha \rightarrow 0} \left(\frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} - \text{pp} \frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{2+\alpha}} \right) \\ &= \frac{h^2}{(8\pi^2)^2} \left(\left(\frac{1}{\sigma_F^2} \right)_{\text{MS}} - \frac{\log(k^2)}{2} (\square + f) \frac{1}{\sigma_F} \right) = (\Delta_F^2)_{\text{MS}} - \frac{i \log(8\pi^2)}{16\pi^2} \delta. \end{aligned}$$

(b) In analogy to (a), we may compute

$$\begin{aligned} & \lim_{\alpha \rightarrow 0} \left(\frac{d}{d\alpha} \frac{1}{M^{2\alpha}}\Delta_F^{2+\alpha} - \text{pp} \frac{d}{d\alpha} \frac{1}{M^{2\alpha}}\Delta_F^{2+\alpha} \right) \\ &= \frac{h^2}{(8\pi^2)^2} \left(- \left(\frac{\log(M^2\sigma_F)}{\sigma_F^2} \right)_{\text{MS}} - \log\left(\frac{8\pi^2}{h^2}\right) \left(\frac{1}{\sigma_F^2} \right)_{\text{MS}} + \frac{\log^2\left(\frac{8\pi^2}{h^2}\right)}{4} (\square + f) \frac{1}{\sigma_F} \right) \\ &= (\Delta_F^2 \log(M^{-2}\Delta_F))_{\text{MS}} + \frac{i \log^2(8\pi^2)}{32\pi^2} \delta. \end{aligned}$$

(c) This can be proven in analogy to (a), (b), whereby one also needs lemma 3.18 (d) and the fact that $[v] = 0$ implies by means of the covariant expansion of bitensors near the diagonal (see, e.g. ([PPV11] section 5)) that

$$v = [v] + ([\nabla_a v] - \nabla_a[v])\sigma^a + \mathcal{R}_v = [\nabla_a v]\sigma^a + \mathcal{R}_v,$$

where the remainder term \mathcal{R}_v vanishes towards the diagonal faster than σ_a . Thus, the assumption $[v] = 0$ implies that the term in Δ_F^3 proportional to $\sigma_F^{-2} \log M^2\sigma_F$ does not need to be renormalised, which is crucial for the present proof. The correction term arises from the $\log h/(8\pi^2)$ term in the expansion of

$$\frac{1}{(Mk)^{2\alpha}} \frac{1}{\sigma_F^{3+\alpha}},$$

whose contribution may be computed as

$$\begin{aligned} & \frac{h^3 \log\left(\frac{h}{8\pi^2}\right)}{8(8\pi^2)^3} (\square + 2f)(\square + f) \frac{1}{\sigma_F} = - \frac{i \left(-\log(8\pi^2)^{\frac{R}{3}} - [\square h^3 \log(h)] \right) \delta}{8(8\pi^2)^2} \\ &= \frac{i \left(-\log(8\pi^2)^{\frac{R}{3}} + [\square u + 8\pi^2 w \square \sigma] \right) \delta}{8(8\pi^2)^2} = \frac{i((1 + 2 \log(8\pi^2))R + 192\pi^2[w])}{48(8\pi^2)^2} \delta, \end{aligned}$$

where, again, lemma 3.18 (a), (b) prove to be useful. \square

3.5.3. A more complicated graph. In order to show how the proposed renormalisation scheme works for graphs that have more than two vertices, we discuss the renormalisation of the following triangular graph:

$$\tau_T := \Delta_{F,13} \Delta_{F,23} \Delta_{F,12}^2,$$

where $\Delta_{F,ij} := \Delta_F(x_i, x_j)$. In order to apply the forest formula (3.9) to renormalise this graph, we note that the forests that correspond to divergent contributions are

$$\{12\}, \quad \{123\}, \quad \{12, 123\}.$$

The renormalisation of τ_T , thus, reads

$$(\tau_T)_{\text{MS}} = (1 + R_{12} + R_{123} + R_{123}R_{12})\tau_T^{(\alpha)} = (1 + R_{123})(1 + R_{12})\tau_T^{(\alpha)}.$$

In order to illustrate the explicit form of the R , we consider only the most singular contribution to $\tau_T^{(\alpha)}$, namely,

$$t_{T,0}^{(\alpha)} := \frac{1}{\sigma_{13}^{1+\alpha_{13}}} \frac{1}{\sigma_{12}^{2(1+\alpha_{12})}} \frac{1}{\sigma_{23}^{1+\alpha_{23}}},$$

where $\sigma_{ij} := \sigma_F(x_i, x_j)$. Note that, with obvious notation, $(8\pi^2)^{-4} u_{13} u_{12}^2 u_{23} t_{T,0}$ is, in fact, the only contribution to τ_T that needs to be renormalised. The application of $1 + R_{12}$ to $t_{T,0}^{(\alpha)}$ has already been discussed in the preceding sections and corresponds to the renormalisation of the fish graph. Indeed, after setting α_{12} , α_{23} and α_{13} to $\alpha = \alpha_I$ for $I = \{1-3\}$, we obtain

$$t_{T,1}^{(\alpha)} := \lim_{\alpha_{ij} \rightarrow \alpha} (1 + R_{12})t_{T,0}^{(\alpha)} = \left(\left(\frac{1}{\sigma_{12}^2} \right)_{\text{MS}} + O(\alpha) \right) \frac{1}{(\sigma_{13})^{1+\alpha}} \frac{1}{(\sigma_{23})^{1+\alpha}}.$$

The distribution $(1/\sigma_{12}^2)_{\text{MS}}$ is a homogeneous distribution of degree $\delta = -4$ under scaling of x_2 towards x_1 , consequently, $t_{T,1}^{(\alpha)}$ has scaling degree $8 + 4\alpha$.

Owing to proposition 3.12, we know that $t_{T,1}^{(\alpha)}$ can be decomposed into the sum of a homogeneous distribution of degree $-8 - 4\alpha$ and a remainder. Hence, in order to expose the poles of $t_{T,1}^{(\alpha)}$, we can directly apply proposition 3.9 with $m = 1$ and $c_0 = -4\alpha$. To this end, we set $u_0 := t_{T,1}^{(\alpha)}$ and find

$$u_1 := -4\alpha u_0 - E_1^\dagger u_0 = \left(\left(\frac{1}{\sigma_{12}^2} \right)_{\text{MS}} + O(\alpha) \right) \frac{1}{(\sigma_{13})^{1+\alpha}} \frac{1}{(\sigma_{23})^{2+\alpha}} G,$$

where $G = G(x_1, x_2, x_3)$ is the smooth function introduced in lemma 3.11. From (3.20), we can infer that the principal part of $t_{T,1}^{(\alpha)}$ is

$$\text{pp } t_{T,1}^{(\alpha)} = -\frac{1}{4\alpha} \left(E_1^\dagger + \frac{G}{\sigma_{23}} \right) \left(\left(\frac{1}{\sigma_{12}^2} \right)_{\text{MS}} \frac{1}{\sigma_{13}} \frac{1}{\sigma_{23}} \right),$$

whereas the constant regular part can be easily computed as well. Consequently, the renormalised distribution,

$$(t_{T,0})_{\text{MS}} = \lim_{\alpha \rightarrow 0} (t_{T,1}^{(\alpha)} - \text{pp } t_{T,1}^{(\alpha)})$$

can be straightforwardly computed in explicit terms.

4. Explicit computations in cosmological space-times

The aim of this section is provide *pr  t-  -porter* formulas for doing perturbative computations in the renormalisation scheme devised in the previous sections for the special case of FLRW space-times. We, thus, consider space-times (\mathcal{M}, g) of the form $\mathcal{M} = I \times \mathbb{R}^3 \subset \mathbb{R}^4$ and, in comoving coordinates,

$$g = -dt^2 + a(t)^2 dx^2 = a(\tau)^2 (-d\tau^2 + dx^2).$$

Here, t is cosmological time, and τ is conformal time related to t by $dt = a d\tau$ and

$$H := \partial_t \log(a) = \frac{\partial_\tau a}{a^2} =: \frac{\mathcal{H}}{a}, \quad R = 6(\partial_t H + 2H^2) = \frac{\partial_\tau^2 a}{a^3}. \quad (4.1)$$

We consider here the spatially flat FLRW case for simplicity. Note that these space-times are normal neighbourhoods so that (3.28) can be considered as a global expression and all Feynman amplitudes can be analytically regularised without the need of introducing partitions of unity, such as in remark 3.1.

4.1. Propagators in Fourier space

In comoving coordinates with conformal time, the Klein–Gordon operator reads

$$P = -\square + \xi R + m^2 = \frac{1}{a(\tau)^3} \left(\partial_\tau^2 - \vec{\nabla}^2 + \left(\xi - \frac{1}{6} \right) R a^2 + m^2 a^2 \right) a(\tau).$$

It is convenient to employ Fourier transformations with respect to the spatial coordinates in order to expand quantities in QFT on FLRW space-times in terms of mode solutions of the free Klein–Gordon equation,

$$\phi_{\vec{k}}(\tau, \vec{x}) = \frac{\chi_k(\tau) e^{i\vec{k}\vec{x}}}{(2\pi)^{\frac{3}{2}} a(\tau)},$$

where the temporal modes $\chi_k(\tau)$ satisfy

$$\left(\partial_\tau^2 + k^2 + m^2 a^2 + \left(\xi - \frac{1}{6} \right) R a^2 \right) \chi_k(\tau) = 0, \quad (4.2)$$

and the normalisation condition,

$$\chi_k \partial_\tau \overline{\chi_k} - \overline{\chi_k} \partial_\tau \chi_k = i. \quad (4.3)$$

Here, $k := |\vec{k}|$, and $\bar{\cdot}$ denotes complex conjugation.

In particular, we can use the mode expansion in order to give explicit expressions for the various propagators of the free Klein–Gordon quantum field in a pure, Gaussian, homogeneous, and isotropic state Ω (see [LR90, Pi10, Zs13] for associated technical conditions on the mode functions). To this avail, we define

$$\Delta_{\#}(x_1, x_2) =: \lim_{\epsilon \downarrow 0} \frac{1}{8\pi^3 a(\tau_1) a(\tau_2)} \int_{\mathbb{R}^3} d^3k \widehat{\Delta}_{\#}(\tau_1, \tau_2, k) e^{ik(x_1 - x_2) - \epsilon k}, \quad (4.4)$$

where $\Delta_{\#}$ stands for Δ_+ (two-point function), $\Delta_{R/A}$ (retarded/advanced propagator) or Δ_F (Feynman propagator). See section A.1 for our conventions for these propagators and their relations. Recall that our renormalisation scheme preserves invariance under space-time isometries and, thus, we know that renormalised powers of the Feynman propagator may also be written in the form (4.4).

The Fourier versions of the single propagators read

$$\begin{aligned}\widehat{\Delta}_+(\tau_1, \tau_2, k) &= \chi_k(\tau_1) \overline{\chi_k(\tau_2)}, & \widehat{\Delta}_-(\tau_1, \tau_2, k) &= \overline{\widehat{\Delta}_+(\tau_1, \tau_2, k)}, \\ \widehat{\Delta}_F(\tau_1, \tau_2, k) &= \Theta(\tau_1 - \tau_2) \widehat{\Delta}_+(\tau_1, \tau_2, k) + \Theta(\tau_2 - \tau_1) \widehat{\Delta}_-(\tau_1, \tau_2, k), \\ \widehat{\Delta}_{R/A}(\tau_1, \tau_2, k) &= \mp i\Theta(\pm(\tau_1 - \tau_2))(\widehat{\Delta}_+(\tau_1, \tau_2, k) - \widehat{\Delta}_-(\tau_1, \tau_2, k)),\end{aligned}\quad (4.5)$$

whereas, by the convolution theorem, we have the following Fourier versions of products and convolutions of multiple propagators, provided those products and convolutions are well defined. Defining

$$\begin{aligned}[\Delta_{\#1} *_4 \Delta_{\#2}](x, y) &:= \int_{\mathcal{M}} d^4x \sqrt{-g} \Delta_{\#1}(x_1, x) \Delta_{\#2}(x, x_2), \\ [\widehat{\Delta}_{\#1} *_1 \widehat{\Delta}_{\#2}](\tau_1, \tau_2, k) &:= \int_I d\tau a(\tau)^2 \widehat{\Delta}_{\#1}(\tau_1, \tau, k) \widehat{\Delta}_{\#2}(\tau, \tau_2, k), \\ [\widehat{\Delta}_{\#1} *_3 \widehat{\Delta}_{\#2}](\tau_1, \tau_2, k) &:= \int_{\mathbb{R}^3} d^3p \widehat{\Delta}_{\#1}(\tau_1, \tau_2, p) \widehat{\Delta}_{\#2}(\tau_1, \tau_2, |\vec{k} - \vec{p}|),\end{aligned}\quad (4.6)$$

we have

$$\begin{aligned}\prod_{i=1}^n \widehat{\Delta}_{\#i}(\tau_1, \tau_2, k) &= \frac{1}{((2\pi)^3 a(\tau_1)^2 a(\tau_2)^2)^{n-1}} [\widehat{\Delta}_{\#1} *_3 \dots *_3 \widehat{\Delta}_{\#n}](\tau_1, \tau_2, k), \\ \overline{\Delta_{\#1} *_4 \dots *_4 \Delta_{\#n}} &= \widehat{\Delta}_{\#1} *_1 \dots *_1 \widehat{\Delta}_{\#n}.\end{aligned}\quad (4.7)$$

Choosing a pure, Gaussian, homogeneous, and isotropic state Ω of the quantized free Klein–Gordon field on a spatially flat FLRW space–time amounts to choosing a solution of (4.2), (4.3) for each k . In order for Ω to be a Hadamard state, the temporal modes χ_k have to satisfy certain conditions in the limit of large k that are difficult to formulate precisely. Heuristically, a necessary but not sufficient condition is that the dominant part of χ_k for large k , when the mass and curvature terms in (4.2) are dominated by k^2 , is $\frac{1}{\sqrt{2k}} e^{-ik\tau}$, i.e., a positive frequency solution. Note that the retarded and advanced propagators are state independent and, thus, $\widehat{\Delta}_{R/A}(\tau_1, \tau_2, k)$ is independent of the particular χ_k chosen for each k .

4.2. The renormalised fish and sunset graphs in Fourier space

In perturbative calculations at low orders, we encounter (pointwise) powers of Δ_{\pm} and Δ_F . While the powers of Δ_{\pm} are well defined if Ω is a Hadamard state on account of the WF set properties of these distributions, we need to renormalise the powers of Δ_F by means of the scheme developed in the previous sections. In order to be useful for explicit computations in FLRW space–times, we have to develop a spatial Fourier-space version of this scheme. Having in mind the application to ϕ^4 theory, we will compute $(\widehat{\Delta}_F)_{\text{MS}}^n(\tau_1, \tau_2, k)$ for $n = 2, 3$. The difficulty in achieving this is that, to our knowledge, despite the large symmetry of flat FLRW space–times, neither σ nor the Hadamard coefficients u , v , and w may be written in a tractable form that can easily be Fourier transformed. Our strategy to circumvent this problem is the following.

Computational strategy

1. For a general mass m , coupling to the scalar curvature ξ , and a general homogeneous, isotropic, pure, and Gaussian Hadamard state Ω , split Δ_F as

$$\Delta_F = \Delta_{F,0} + d, \quad d := \Delta_F - \Delta_{F,0}, \quad (4.8)$$

where $\Delta_{F,0}$ must satisfy the following conditions.

- $\Delta_{F,0}$ is explicitly known in position space and Fourier space.
- $\Delta_{F,0}$ is of the form

$$\Delta_{F,0} = \frac{1}{8\pi^2} \left(\frac{u_0}{\sigma_F} + v_0 \log(M^2 \sigma_F) \right) + w_0,$$

with $u_0 = u$, i.e., it agrees with Δ_F in the most singular term but not necessarily in the subleading singularities.

- $[v_0] = 0$ and $\Delta_{F,0}^\alpha$ is weakly meromorphic in α such that $(\Delta_{F,0}^2)_{\text{MS}}$, $(\Delta_{F,0}^2 \log(M^{-2} \Delta_{F,0}))_{\text{MS}}$, and $(\Delta_{F,0}^3)_{\text{MS}}$ may be computed with proposition 3.19. This is crucial for preserving the explicit knowledge of $\Delta_{F,0}$ in position space in the renormalisation procedure so that one may hope to compute the Fourier transforms of the renormalised powers.
2. With these assumptions on $\Delta_{F,0}$, it follows that the renormalised fish and sunset graphs may be computed as

$$\begin{aligned} (\Delta_F)_{\text{MS}}^2 &= (\Delta_{F,0})_{\text{MS}}^2 + 2\Delta_{F,0}d + d^2, \\ (\Delta_F)_{\text{MS}}^3 &= (\Delta_{F,0})_{\text{MS}}^3 + 3(\Delta_{F,0}^2 d)_{\text{MS}} + 3\Delta_{F,0}d^2 + d^3, \end{aligned} \quad (4.9)$$

because the nonrenormalised terms in the above formulas are distributions with scaling degree < 4 for $y \rightarrow x$ and, thus, can be directly and uniquely extended to the diagonal.

3. $(\Delta_{F,0}^2)_{\text{MS}}$ and $(\Delta_{F,0}^3)_{\text{MS}}$ may be computed with proposition 3.19 as anticipated. In order to compute $(\Delta_{F,0}^2 d)_{\text{MS}}$, we further split d as

$$d = d_1 + d_2, \quad d_1 := -\frac{[v] \log(M^{-2} \Delta_{F,0})}{8\pi^2}, \quad d_2 := d - d_1. \quad (4.10)$$

Because $v = [v] + O(\sigma_a)$, d_1 contains the leading logarithmic singularity in d (and, thus, Δ_F), which is the only logarithmic singularity relevant for the renormalisation of the sunset graph. Consequently,

$$(\Delta_{F,0}^2 d)_{\text{MS}} = -\frac{[v]}{8\pi^2} (\Delta_{F,0}^2 \log(M^{-2} \Delta_{F,0}))_{\text{MS}} + d_2 (\Delta_{F,0}^2)_{\text{MS}}, \quad (4.11)$$

and, thus, proposition 3.19 can be applied again.

4. Due to the symmetry of FLRW space-times and the assumption that the pure and Gaussian Hadamard state Ω is invariant under this symmetry, $[v]$ and $[w]$ do not depend on the spatial coordinates. Given that one succeeds to compute the spatial Fourier transforms of $\log(M^{-2} \Delta_{F,0})$, $(\Delta_{F,0}^2)_{\text{MS}}$, $(\Delta_{F,0}^2 \log(M^{-2} \Delta_{F,0}))_{\text{MS}}$, and $(\Delta_{F,0}^3)_{\text{MS}}$, $(\widehat{\Delta_F})_{\text{MS}}^n(\tau_1, \tau_2, k)$ for $n \in \{2, 3\}$ may be computed by means of the convolution identities (4.7) since the Fourier transforms of $\Delta_{F,0}$, d_1 , and d_2 are known by construction.

In order to follow the computational strategy outlined above, we first compute $[v]$ and $[w]$. Indeed, the coinciding point limit of the Hadamard coefficient v reads (see, e.g. ([Ha10] section III.1.2) for details)

$$[v] = \frac{m^2 + \left(\xi - \frac{1}{6}\right)R}{2}. \quad (4.12)$$

Moreover, using the method of [Sc10] to compute a spatial Fourier representation of the Hadamard parametrix H_F —here considered as (3.28) with $w = 0$ —in FLRW space-times,

one can compute (see the review in [De13] and a related method in [Pi10] for the conformally coupled case)

$$\begin{aligned}
[w] &= \lim_{x \rightarrow y} (\Delta_F(x, y) - H_F(x, y)) \\
&= \frac{1}{(2\pi)^3 a^2} \int_{\mathbb{R}^3} d^3k |\chi_k(\tau)|^2 - \frac{1}{2\sqrt{k^2 + a^2 m^2 + a^2 \left(\xi - \frac{1}{6}\right) R}} \\
&\quad + \frac{1}{16\pi^2} \left(m^2 + \left(\xi - \frac{1}{6}\right) R \right) \left(2\gamma - 1 + \log \left(\frac{m^2 + \left(\xi - \frac{1}{6}\right) R}{2M^2} \right) \right) - \frac{R}{36(8\pi^2)},
\end{aligned} \tag{4.13}$$

where γ is the Euler–Mascheroni constant and H_F is taken with the mass scale M inside of the logarithm of σ ⁸.

As anticipated, we see that $[v]$ and $[w]$ are functions of time only (recall (4.1)). Moreover, we see that $[v] = 0$ for a conformally coupled ($\xi = \frac{1}{6}$) massless scalar field. Thus, in order to pursue our computational strategy, we should look for a candidate for $\Delta_{F,0}$ among the Feynman propagators in suitable states of this theory. In fact, choosing the conformal vacuum state of the massless conformally coupled scalar field does the job. The conformal vacuum is given by choosing the modes $\chi_k(\tau) = e^{-ik\tau}/\sqrt{2k}$, and thus the Feynman propagator $\Delta_{F,0}$ in this state is of the form

$$\Delta_{F,0}(x_1, x_2) = \frac{1}{8\pi^2 a(\tau_1) a(\tau_2)} \frac{1}{\sigma_{F,\mathbb{M}}(x_1, x_2)}, \quad \widehat{\Delta_{F,0}}(\tau_1, \tau_2, k) = \frac{e^{-ik|\tau_1 - \tau_2|}}{2k}. \tag{4.14}$$

Here, and in the following, the index \mathbb{M} indicates quantities in Minkowski space–time, in particular, $\sigma_{\mathbb{M}}(x_1, x_2) = \frac{1}{2}(\vec{x}_1 - \vec{x}_2)^2 - \frac{1}{2}(\tau_1 - \tau_2)^2$. $\Delta_{F,0}^\alpha$ is weakly meromorphic in α because the massless vacuum Feynman propagator in Minkowski space–time has this property and the conformal rescaling by a does not violate it. Thus, we may follow our computational strategy and compute $(\Delta_{F,0}^2)_{\text{MS}}$, $(\Delta_{F,0}^2 \log(M^{-2} \Delta_{F,0}))_{\text{MS}}$, and $(\Delta_{F,0}^3)_{\text{MS}}$ by means of proposition 3.19. This is easily done using (3.33) for $\sigma_{F,\mathbb{M}}$ rather than σ_F and $h = \sqrt{8\pi^2 a(\tau_1) a(\tau_2)} = \sqrt{8\pi^2 a \otimes a}$. The results are

$$\begin{aligned}
(\Delta_{F,0}^2)_{\text{MS}} &= \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} (\Delta_{F,0})^{2+\alpha} - \text{pp} \frac{1}{M^{2\alpha}} (\Delta_{F,0})^{2+\alpha} \right) + \frac{i \log(8\pi^2)}{16\pi^2} \delta \\
&= \lim_{\alpha \rightarrow 0} \frac{1}{(8\pi^2)^2 a^2 \otimes a^2} \left(\frac{1}{(M\sqrt{8\pi^2 a \otimes a})^{2\alpha}} \frac{1}{\sigma_{F,\mathbb{M}}^{2+\alpha}} \right. \\
&\quad \left. - \text{pp} \frac{1}{(M\sqrt{8\pi^2 a \otimes a})^{2\alpha}} \frac{1}{\sigma_{F,\mathbb{M}}^{2+\alpha}} \right) + \frac{i \log(8\pi^2)}{16\pi^2} \delta \\
&= - \frac{1 + 2 \log(a)}{16\pi^2 a^4} i \delta_{\mathbb{M}} - \frac{1}{2(8\pi^2)^2 a^2 \otimes a^2} \square_{\mathbb{M}} \frac{\log(M^2 \sigma_{F,\mathbb{M}})}{\sigma_{F,\mathbb{M}}},
\end{aligned}$$

⁸ Note that one may take, instead of the function $F(k) = 1/\left(2\sqrt{k^2 + a^2 m^2 + a^2 \left(\xi - \frac{1}{6}\right) R}\right)$ in (4.13), any distribution $F'(k)$ such that $F'(k) - F(k)$ is $O(k^{-5})$ for large k and integrable. By taking, e.g., $F'(k) = 1/(2k) - \Theta(k - am) \left(a^2 m^2 + a^2 \left(\xi - \frac{1}{6}\right) R\right)/(4k^3)$, one may cancel the $\log R$ term outside of the integral.

$$\begin{aligned}
(\Delta_{F,0}^2 \log(M^{-2} \Delta_{F,0}))_{\text{MS}} &= \lim_{\alpha \rightarrow 0} \left(\frac{d}{d\alpha} \frac{1}{M^{2\alpha}} (\Delta_{F,0})^{2+\alpha} - \text{pp} \frac{d}{d\alpha} \frac{1}{M^{2\alpha}} (\Delta_{F,0})^{2+\alpha} \right) \\
&\quad - \frac{i \log^2(8\pi^2)}{32\pi^2} \delta \\
&= \frac{2 + 2 \log(a^2 8\pi^2) + \log^2(a^2)}{32\pi^2 a^4} i \delta_{\mathbb{M}} \\
&\quad + \frac{1}{4(8\pi^2)^2 a^2 \otimes a^2} \square_{\mathbb{M}} \frac{\log^2(M^2 \sigma_{F,\mathbb{M}})}{\sigma_{F,\mathbb{M}}} \\
&\quad + \frac{1 + \log(8\pi^2 a \otimes a)}{2(8\pi^2)^2 a^2 \otimes a^2} \square_{\mathbb{M}} \frac{\log(M^2 \sigma_{F,\mathbb{M}})}{\sigma_{F,\mathbb{M}}},
\end{aligned}$$

and

$$\begin{aligned}
(\Delta_{F,0})_{\text{MS}}^3 &= \lim_{\alpha \rightarrow 0} \left(\frac{1}{M^{2\alpha}} (\Delta_{F,0})^{3+\alpha} - \text{pp} \frac{1}{M^{2\alpha}} (\Delta_{F,0})^{3+\alpha} \right) \\
&\quad + \frac{i((1 + 2 \log(8\pi^2))R + 192\pi^2[w])}{48(8\pi^2)^2} \delta \\
&= - \frac{(15 + 12 \log(a)) \square_{\mathbb{M}} + 6(\square_{\mathbb{M}} \log(a)) + 2(\partial_{\tau^2}^2 a)/a}{48(8\pi^2)^2 a^6} i \delta_{\mathbb{M}} \\
&\quad - \frac{1}{8(8\pi^2)^3 a^3 \otimes a^3} \square_{\mathbb{M}}^2 \frac{\log(M^2 \sigma_{F,\mathbb{M}})}{\sigma_{F,\mathbb{M}}}.
\end{aligned}$$

where we have used $\delta = \delta_{\mathbb{M}}/a^4$, $f_{\mathbb{M}} = 0$ and the fact that, by (4.13), $8\pi^2[w_0] = -R/36$ for the conformal vacuum state of the massless conformally coupled scalar field.

Using these results as well as the Fourier representation of $1/\sigma_{F,\mathbb{M}}$ (4.14), $\log(M^2 \sigma_{F,\mathbb{M}})$ (A.2), and convolution identities, we can finally obtain the Fourier versions of the renormalised powers of $\Delta_{F,0}$. For instance, we find for $(\Delta_{F,0}^2)_{\text{MS}}$,

$$\begin{aligned}
\widehat{(\Delta_{F,0}^2)_{\text{MS}}}(\tau_1, \tau_2, k) &= - \frac{1 + 2 \log(a(\tau_1))}{16a(\tau_1)^2 \pi^2} \delta(\tau_1 - \tau_2) \\
&\quad - \frac{1}{16\pi^3 a(\tau_1) a(\tau_2)} (\partial_{\tau_1}^2 + k^2) \int_{\mathbb{R}^3} d^3 p \left(\frac{1}{2} \left(\frac{1}{p^3} \right)_{\text{ren}, M} \right. \\
&\quad \left. + \frac{i|\tau_1 - \tau_2|}{2p^2} \right) \frac{1}{2|\vec{k} - \vec{p}|} e^{-i(p+|\vec{k}-\vec{p}|)|\tau_1-\tau_2|},
\end{aligned} \tag{4.15}$$

where the appearing renormalisation of $1/p^3$ is defined in (A.1). Note that the \vec{p} -integral has no convergence problems for large p because one may write the potentially dangerous $-i|\tau_1 - \tau_2|e^{-2ip|\tau_1-\tau_2|}/p$ contribution as $\partial_p(e^{-2ip|\tau_1-\tau_2|}/(2p^2))$ plus an $O(p^{-3})$ term. Regarding the convergence for small p , we observe that the integral is manifestly convergent if $k \neq 0$, thus, yielding a well-defined distribution in \vec{k} on $\mathbb{R}^3 \setminus \{0\}$. The scaling degree of this distribution is easily seen to be $1 < 3$, and, thus, a unique extension towards the origin exists. In practical terms, this means that the integral for $k = 0$ may be computed as a limit $k \rightarrow 0$ of the integral with nonvanishing k without any renormalisation.

4.3. Example: the two-point function for a quartic potential up to second order

In order to compute the analytic expressions corresponding to the graphs in figure 2, we may use the Fourier versions of the appearing propagators (4.5), (4.15) and the analogous expressions for $(\Delta_{F,0}^2 \log(\widehat{M^{-2} \Delta_{F,0}^2}))_{\text{MS}}(\tau_1, \tau_2, k)$ and $(\Delta_{F,0}^3)_{\text{MS}}(\tau_1, \tau_2, k)$, the explicit form of $\mu(x) = 3\lambda w(x, x)$ in (4.13), as well as (4.9), (4.11) and the identities for products and convolutions (4.6), (4.7). Note that $\mu(x)$ is, in fact, only time dependent because Ω was chosen homogeneous and isotropic. Thus, the integrals with μ -vertices can be computed partly with the above-mentioned identities by means of

$$(\widehat{1 \otimes \mu})\Delta_{\#}(\tau_1, \tau_2, k) = \mu(\tau_2)\widehat{\Delta_{\#}}(\tau_1, \tau_2, k), \quad (\widehat{\mu \otimes 1})\Delta_{\#}(\tau_1, \tau_2, k) = \mu(\tau_1)\widehat{\Delta_{\#}}(\tau_1, \tau_2, k).$$

Similarly, the bubbles in the third line of figure 2 contribute only time-dependent vertex factors, which can be computed as

$$h_{\#}(\tau) := \int_{\mathcal{M}} d\tau_1 d^3x_1 a(\tau_1)^4 \mu(\tau_1) \Delta_{\#}(\tau, \tau_1, \vec{x} - \vec{x}_1) = \frac{1}{a(\tau)} \int_I d\tau_1 a(\tau_1)^3 \mu(\tau_1) \widehat{\Delta_{\#}}(\tau, \tau_1, 0),$$

where $\Delta_{\#}$ is either Δ_+^2 or $(\Delta_F^2)_{\text{MS}}$.

With these preparations, we can compute, e.g., the first graphs of the fourth and fifth lines in figure 2 in Fourier space as

$$\begin{aligned} \Delta_R^* \widehat{(h_F \otimes 1)} \Delta_+ &= \widehat{\Delta_R}^* \widehat{(h_F \otimes 1)} \Delta_+ \\ &= \int_{I^2} d\tau_3 d\tau_4 a(\tau_3) a(\tau_4)^3 \mu(\tau_4) \widehat{\Delta_R}(\tau_1, \tau_3, k) \widehat{\Delta_+}(\tau_3, \tau_2, k) (\widehat{\Delta_F^2})_{\text{MS}}(\tau_3, \tau_4, 0), \end{aligned}$$

and

$$\begin{aligned} \Delta_R^* \widehat{(\Delta_F^3)_{\text{MS}}} \Delta_+ &= \widehat{\Delta_R}^* \widehat{(\Delta_F^3)_{\text{MS}}} \Delta_+ \\ &= \int_{I^2} d\tau_3 d\tau_4 a(\tau_3)^2 a(\tau_4)^2 \widehat{\Delta_R}(\tau_1, \tau_3, k) (\widehat{\Delta_F^3})_{\text{MS}}(\tau_3, \tau_4, k) \widehat{\Delta_+}(\tau_4, \tau_2, k). \end{aligned}$$

4.4. More complicated graphs on cosmological space-times

In order to compute the Fourier transforms of more complicated graphs on FLRW space-times, one can use a strategy generalising the one employed in section 4.2. Namely, one again decomposes the Feynman propagator Δ_F into several pieces that capture the relevant singularities and can be expressed in terms of the conformal vacuum Feynman propagator $\Delta_{F,0}$, whose explicit form in position and Fourier space is well known in contrast to the form of σ itself. The corresponding decomposition of general Feynman amplitudes τ_{Γ} is straightforward. The only nontrivial step is to generalise proposition 3.19 to the case of general amplitudes, i.e., to compute the difference between the minimal subtraction schemes used in conjunction with either analytically regularising powers of σ directly or analytically regularising powers of the full propagator $\Delta_{F,0}$. However, we do not foresee any problems in obtaining such a generalisation by proving versions of lemma 3.11 and proposition 3.12 for $\Delta_{F,0}$ rather than σ .

In fact, one can also skip this last step by taking a rather pragmatic approach and by working directly with the renormalisation scheme consisting of decomposition in $\Delta_{F,0}$, analytic regularisation of powers of this propagator, and minimal subtraction of the principal parts. This scheme, clearly applicable only to conformally flat space-times, satisfies all properties proved in proposition 3.14 with two exceptions. It is not obvious whether the PPA with respect to generalised mass perturbations holds for this scheme, whereas, locality and

covariance, of course, only hold in the sense restricted to conformally flat space–times. In this respect, it is essential that the Feynman propagator of the conformal vacuum $\Delta_{F,0}$ on conformally flat space–times is manifestly geometric because the corresponding propagator of the massless Minkowski vacuum has this property.

5. Summary and outlook

In this work, we have introduced a renormalisation scheme on curved space–times consisting of a particular analytic regularisation of the Feynman propagator and, thus, of all Feynman diagrams and a minimal subtraction of the principal (pole) part of the resulting meromorphic expressions. We have argued that this scheme has all properties that a physically meaningful renormalisation scheme on curved space–times should have and that it is, in fact, a particular form of differential renormalisation. The renormalisation scheme discussed in this work has the advantage that it is

- (1) directly applicable to space–times with Lorentzian signature,
- (2) manifestly (local and) covariant,
- (3) manifestly invariant under any space–time isometries present,
- (4) capturing correctly the nongeometric and nonunique state–dependent contributions of Feynman amplitudes and not only the geometric divergent part, which is unique up to finite renormalisations,
- (5) well suited for practical computations, e.g., in cosmological space–times,
- (6) constructed to all orders in perturbation theory,
- (7) and mathematically rigorous.

To the best of our knowledge, other renormalisation schemes on curved space–times discussed in the literature, such as dimensional regularisation, local momentum space methods, ζ -function regularisation, heat-kernel techniques, generic Epstein–Glaser renormalisation, and, on cosmological space–times, dimensional regularisation only with respect to spatial variables, lack, at least, one of the features listed above.

In order to demonstrate the practical applicability of the scheme, we have computed several examples on generic curved space–times. Moreover, we have shown how explicit computations in cosmological space–times can be done, in particular, how the renormalisation scheme initially defined in position space can be interpreted in terms of quantities Fourier transformed with respect to comoving spatial coordinates.

We have discussed the renormalisation scheme only for scalar fields in four space–time dimensions, however, the extension to other space–time dimensions is straightforward. Moreover, as the analytic regularisation discussed in this work consists of regularising only inverse powers of the squared geodesic distance, it can be straightforwardly generalised to field theories with higher spin with and without gauge invariance. In particular, spinorial quantities can be directly regularised without the need to worry about their dependence on the dimension, such as in dimensional regularisation. Finally, we expect that a generalisation of the scheme introduced in this work to gauge theories yields a scheme that preserves or can be modified to preserve the local gauge symmetry in theories that are free of anomalies.

Acknowledgments

The authors would like to thank K. Fredenhagen and M. Fröb for interesting discussions. The work of T.-P.H. has been supported by a Research Fellowship of Deutsche Forschungsgemeinschaft (DFG).

Appendix. Conventions and computational details

A.1. Propagators of the free Klein–Gordon field and their relations

Here, we have

$$\Delta_+(x, y) = \langle \phi(x) \phi(y) \rangle_\Omega = \langle \phi(x) \star \phi(y) \rangle_\Omega = \overline{\langle \phi(y) \phi(x) \rangle_\Omega} = \overline{\Delta_-(x, y)},$$

$$\Delta(x, y) = \frac{1}{i}(\Delta_+(x, y) - \Delta_-(x, y)) = \Delta_R(x, y) - \Delta_A(x, y),$$

$$\Delta_{R/A} = \pm \Theta(\pm(t_x - t_y)) \Delta(x, y),$$

$$\begin{aligned} \Delta_F(x, y) &= \langle T(\phi(x) \phi(y)) \rangle_\Omega = \langle \phi(x) \cdot_T \phi(y) \rangle_\Omega \\ &= \Theta(t_x - t_y) \Delta_+(x, y) + \Theta(t_y - t_x) \Delta_-(x, y) \end{aligned}$$

$$\begin{aligned} \Rightarrow \quad \Delta_F(x, y) &= \frac{1}{2}(\Delta_+(x, y) + \Delta_-(x, y)) + \frac{i}{2}(\Delta_R(x, y) + \Delta_A(x, y)) \\ &= \Delta_+(x, y) + i\Delta_A(x, y) \\ &= \Delta_-(x, y) + i\Delta_R(x, y). \end{aligned}$$

A.2. Fourier transform of the logarithmic term on Minkowski space–time

In order to compute the Fourier transform of $\log(M^2 \sigma_{F, \mathbb{M}})$, we recall that the Feynman propagator of the Klein–Gordon field with mass m in the Minkowski vacuum is given by

$$\begin{aligned} \Delta_{F, m, \mathbb{M}} &= \lim_{\epsilon \downarrow 0} \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} d^3k \frac{e^{-i\sqrt{k^2 + m^2} |\tau_1 - \tau_2|} e^{i\vec{k} \cdot (\vec{x} - \vec{y})} e^{-\epsilon k}}{2\sqrt{k^2 + m^2}} \\ &= \frac{1}{8\pi^2} \sqrt{\frac{2m^2}{\sigma_{F, \mathbb{M}}}} K_1(\sqrt{2m^2 \sigma_{F, \mathbb{M}}}) \\ &= \frac{1}{8\pi^2} \left(\frac{1}{\sigma_{F, \mathbb{M}}} + \frac{m^2}{2} \left(1 + \frac{m^2 \sigma_{F, \mathbb{M}}}{4} \right) \log \left(\frac{e^{2\gamma} m^2 \sigma_{F, \mathbb{M}}}{2} \right) \right. \\ &\quad \left. - \frac{m^2}{2} \left(1 + \frac{5m^2 \sigma_{F, \mathbb{M}}}{8} \right) \right) + O(m^4), \end{aligned}$$

where K_1 is a modified Bessel function and γ is the Euler–Mascheroni constant. Using this, we find

$$\begin{aligned}
& \log(M^2 \sigma_{F, \mathbb{M}}) \\
&= \lim_{m \rightarrow 0} \left(16\pi^2 \frac{d\Delta_{F, m, \mathbb{M}}}{dm^2} - \log\left(\frac{e^{2\gamma} m^2}{2M^2}\right) \right) \\
&= - \lim_{m \rightarrow 0} \left(\lim_{\epsilon \downarrow 0} \frac{1}{\pi} \int_{\mathbb{R}^3} d^3k \frac{1 + i\sqrt{k^2 + m^2} |\tau_1 - \tau_2|}{2(k^2 + m^2)^{\frac{3}{2}}} e^{-i\sqrt{k^2 + m^2} |\tau_1 - \tau_2|} e^{i\vec{k} \cdot (\vec{x} - \vec{y})} e^{-\epsilon k} \right. \\
&\quad \left. + \log\left(\frac{e^{2\gamma} m^2}{2M^2}\right) \right) \\
&= - \lim_{\epsilon \downarrow 0} \frac{1}{\pi} \int_{\mathbb{R}^3} d^3k \left(\lim_{m \rightarrow 0} \left(\frac{1}{2(k^2 + m^2)^{\frac{3}{2}}} + \pi \log\left(\frac{e^{2\gamma} m^2}{2M^2}\right) \delta(\vec{k}) \right) \right. \\
&\quad \left. + \frac{i|\tau_1 - \tau_2|}{2k^2} \right) e^{-ik|\tau_1 - \tau_2|} e^{i\vec{k} \cdot (\vec{x} - \vec{y})} e^{-\epsilon k} \\
&= - \lim_{\epsilon \downarrow 0} \frac{1}{\pi} \int_{\mathbb{R}^3} d^3k \left(\frac{1}{2} \left(\frac{1}{k^3} \right)_{\text{ren}, M} + \frac{i|\tau_1 - \tau_2|}{2k^2} \right) e^{-ik|\tau_1 - \tau_2|} e^{i\vec{k} \cdot (\vec{x} - \vec{y})} e^{-\epsilon k} \\
&= \frac{1}{(2\pi)^{\frac{3}{2}}} \lim_{\epsilon \downarrow 0} \int_{\mathbb{R}^3} d^3k \text{flog}(\tau_1 - \tau_2, k) e^{i\vec{k} \cdot (\vec{x} - \vec{y})} e^{-\epsilon k},
\end{aligned}$$

where the appearing renormalisation of the (tempered) distribution $1/k^3$ is

$$\left(\frac{1}{k^3} \right)_{\text{ren}, M} := \lim_{m \rightarrow 0} \left(\frac{1}{(k^2 + m^2)^{\frac{3}{2}}} + \pi \log\left(\frac{e^{4\gamma} m^4}{4M^4}\right) \delta(\vec{k}) \right), \quad (\text{A.1})$$

and

$$\text{flog}(\tau_1 - \tau_2, k) := -\sqrt{8\pi} \left(\frac{1}{2} \left(\frac{1}{k^3} \right)_{\text{ren}, M} + \frac{i|\tau_1 - \tau_2|}{2k^2} \right) e^{-ik|\tau_1 - \tau_2|} \quad (\text{A.2})$$

is the sought-for spatial Fourier transform of $\log(M^2 \sigma_{F, \mathbb{M}})$.

References

- [Al85] Allen B 1985 Vacuum States in de Sitter Space *Phys. Rev. D* **32** 3136
- [BCK10] Baacke J, Covi L and Kevlishvili N 2010 Coupled scalar fields in a flat FRW universe: renormalisation *JCAP* **1008** 026
- [BGP07] B  r C, Ginoux N and Pf  ffle F 2007 *Wave Equations on Lorentzian Manifolds and Quantization* 1st edn (Z  rich: Eur. Math. Soc.) [arXiv:0806.1036[math.DG]].
- [BF13] Bilal A and Ferrari F 2013 Multi-Loop Zeta Function Regularization and Spectral Cutoff in Curved Spacetime *Nucl. Phys. B* **877** 956
- [BD82] Birrell N D and Davies P C W 1982 *Quantum Fields in Curved Space* (Cambridge: Cambridge University Press)
- [BG72] Bollini C G and Giambiagi J J 1972 Dimensional Renormalization: The Number of Dimensions as a Regularizing Parameter *Nuovo Cimento B* **12** 20
- [BDF09] Brunetti R, Duetsch M and Fredenhagen K 2009 Perturbative Algebraic Quantum Field Theory and the Renormalization Groups *Adv. Theor. Math. Phys.* **13** 1541
- [BF00] Brunetti R and Fredenhagen K 2000 Microlocal analysis and interacting quantum field theories: renormalization on physical backgrounds *Commun. Math. Phys.* **208** 623
- [BFK96] Brunetti R, Fredenhagen K and Kohler M 1996 The Microlocal spectrum condition and Wick polynomials of free fields on curved space-times *Commun. Math. Phys.* **180** 633

- [BFV01] Brunetti R, Fredenhagen K and Verch R 2003 The generally covariant locality principle: a new paradigm for local quantum field theory *Commun. Math. Phys.* **237** 31
- [BOS92] Buchbinder I L, Odintsov S D and Shapiro I L 1992 *Effective Action in Quantum Gravity* (Bristol, UK: IOP) 413
- [Bu81] Bunch T S 1981 Bphz Renormalization of $\Lambda\phi^4$ Field Theory in Curved Space-time *Ann. Phys.* **131** 118
- [Da13] Dang N V 2013 Renormalization of quantum field theory on curved space-times, a causal approach *PhD Thesis* Université Paris Diderot arXiv:1312.5674[math-ph]
- [Da15] Dang N V *Complex powers of analytic functions and meromorphic renormalization in QFT* arXiv:1503.00995[math-ph]
- [De13] Degner A 2013 Properties of states of low energy on cosmological spacetimes, DESY-THESIS-2013-002 *PhD thesis* University of Hamburg
- [DHP15] Drago N, Hack T-P and Pinamonti N *The Generalised Principle of Perturbative Agreement and the Thermal Mass* arXiv:1502.02705 [math-ph]
- [Du96] Duetsch M 1997 Slavnov-Taylor identities from the causal point of view *Int. J. Mod. Phys. A* **12** 3205
- [DF00] Duetsch M and Fredenhagen K 2001 Algebraic quantum field theory, perturbation theory, and the loop expansion *Commun. Math. Phys.* **219** 5
- [DFKR14] Duetsch M, Fredenhagen K, Keller K J and Rejzner K 2014 Dimensional Regularization in Position Space, and a Forest Formula for Epstein-Glaser Renormalization *J. Math. Phys.* **55** 122303
- [EG73] Epstein H and Glaser V 1973 The role of locality in perturbation theory *Ann. Inst. Henri Poincaré A* **19.3** 211–95
- [FL13] Fredenhagen K and Lindner F 2014 Construction of KMS States in Perturbative QFT and Renormalized Hamiltonian Dynamics *Commun. Math. Phys.* **332** 895
- [FR12] Fredenhagen K and Rejzner K in *Mathematical Aspects of Quantum Field Theory Perturbative algebraic quantum field theory* (Berlin: Springer)
- [FR13] Fredenhagen K and Rejzner K 2013 Batalin-Vilkovisky formalism in perturbative algebraic quantum field theory *Commun. Math. Phys.* **317** 697
- [FR14] Fredenhagen K and Rejzner K *QFT on curved spacetimes: axiomatic framework and examples* arXiv:1412.5125 [math-ph]
- [Ha10] Hack T P 2010 On the backreaction of scalar and spinor quantum fields in curved spacetimes *PhD Thesis* University of Hamburg arXiv:1008.1776 [gr-qc]
- [Ho10] Hollands S 2013 Correlators, Feynman diagrams, and quantum no-hair in deSitter spacetime *Commun. Math. Phys.* **319** 1
- [HW01] Hollands S and Wald R M 2001 Local Wick polynomials and time ordered products of quantum fields in curved space-time *Commun. Math. Phys.* **223** 289
- [HW02] Hollands S and Wald R M 2002 Existence of local covariant time ordered products of quantum fields in curved space-time *Commun. Math. Phys.* **231** 309
- [HW04] Hollands S and Wald R M 2005 Conservation of the stress tensor in interacting quantum field theory in curved spacetimes *Rev. Math. Phys.* **17** 227
- [Hö90] Hörmander L 1990 *The Analysis of Linear Partial Differential Operators I* 2nd edn (Berlin: Springer)
- [CHL95] Comellas J, Haagensen P E and Latorre J I 1995 A Generic renormalization method in curved spaces and at finite temperature *Int. J. Mod. Phys. A* **10** 2819
- [Ke10] Keller K J 2010 Dimensional Regularization in Position Space and a Forest Formula for Regularized Epstein-Glaser Renormalization *PhD Thesis* University of Hamburg arXiv:1006.2148 [math-ph]
- [Li13] Lindner F 2013 Perturbative algebraic quantum field theory at finite temperature *PhD Thesis* University of Hamburg DESY-THESIS-2013-029.
- [LR90] Lueders C and Roberts J E 1990 Local quasiequivalence and adiabatic vacuum states *Commun. Math. Phys.* **134** 29
- [Lü82] Luscher M 1982 Dimensional regularization in the presence of large background fields *Ann. Phys.* **142** 359
- [Mo99] Moretti V 2000 Proof of the symmetry of the off diagonal Hadamard/Seeley-deWitt's coefficients in C infinity Lorentzian manifolds by a 'local Wick rotation' *Commun. Math. Phys.* **212** 165

- [Mo03] Moretti V 2003 Comments on the stress energy tensor operator in curved space-time *Commun. Math. Phys.* **232** 189
- [O’N83] O’Neill B 1983 *Semi-Riemannian Geometry* (San Diego: Academic)
- [PT09] Parker L E and Toms D 2009 *Quantum Field Theory in Curved Spacetime: Quantized Field and Gravity* (Cambridge University Press)
- [Pi10] Pinamonti N 2011 On the initial conditions and solutions of the semiclassical Einstein equations in a cosmological scenario *Commun. Math. Phys.* (Cambridge: Cambridge Univ. Press) **305** 563
- [PPV11] Poisson E, Pound A and Vega I 2011 The Motion of point particles in curved spacetime *Living Rev. Relativ.* **14** 7
- [Pr97] Prange D 1999 Epstein-Glaser renormalization and differential renormalization *J. Phys. A* **32** 2225
- [Ra96] Radzikowski M J 1996 Micro-local approach to the Hadamard condition in quantum field theory on curved space-time *Commun. Math. Phys.* **179** 529
- [Sc10] Schlemmer J 2010 Local Thermal Equilibrium on Cosmological Spacetimes *Ph.D. thesis* University of Leipzig
- [St71] Steinmann O 1971 *Perturbation Expansions in Axiomatic Field Theory Lecture Notes in Physics* 11 (Berlin: Springer-Verlag)
- [To82] Toms D J 1982 Renormalization of Interacting Scalar Field Theories in Curved Space-time *Phys. Rev. D* **26** 2713
- [Wa84] Wald R M 1984 *General Relativity* (Chicago University Press)
- [Zs13] Zschoche J 2014 The Chaplygin Gas Equation of State for the Quantized Free Scalar Field on Cosmological Spacetimes *Ann. Henri Poincaré* (Chicago, IL: Chicago Univ. Press) **15** 1285