

The Relational Spin-Closure Hypothesis

Mathematical baseline, open problems, and falsification criteria

Editor: Jan Seeck
AI Assistant: ChatGPT 5.6 Sol
AI Lean Agent: Aristotle

Revised after the companion mathematical paper — 13 September 2026

Abstract

This document formulates a hypothesis and a falsification programme. Its mathematical baseline is the companion paper *A Conditional Construction of Lorentzian Geometry from Euclidean Three-Space: Clifford Double-Cover Data, Gluing, and a Soldering Isomorphism* [1]. Starting from the positive-definite Euclidean space

$$V = \mathbb{R}^3,$$

that paper constructs

$$\mathcal{S} = \mathbb{R} \oplus V, \quad N(a, v) = a^2 - \langle v, v \rangle, \quad \text{Cl}_3 = \text{Cl}(V, q),$$

the subgroup G_{Spin} defined in Paper I by the Clifford norm-one condition, a proper orthochronous Lorentz group $G_{\text{Lor}} \subset O(B)$, and a central two-fold covering

$$1 \longrightarrow \{\pm 1\} \longrightarrow G_{\text{Spin}} \xrightarrow{\rho} G_{\text{Lor}} \longrightarrow 1.$$

It also constructs the orthogonal Lie algebra

$$\mathfrak{so}(B) \cong \Lambda^2 \mathcal{S}, \quad \dim \mathfrak{so}(B) = 6.$$

Global Lorentzian geometry requires additional data. A gluing datum yields a smooth four-manifold only under explicit topological and smoothness hypotheses. Independently, a G_{Spin} -valued Čech 1-cocycle $\tilde{g}_{ij}$ projects to $g_{ij} = \rho(\tilde{g}_{ij})$. Local soldering data determine a vector-bundle isomorphism

$$s : E_g \xrightarrow{\sim} TM$$

and hence the Lorentzian metric

$$g_x(X, Y) = B(s_x^{-1}X, s_x^{-1}Y).$$

Neither the global gluing datum nor the soldering isomorphism follows from the local Euclidean input alone.

The explicit periodic one-parameter family studied in Paper I is used here as a deliberately symmetric reference family: its members are gauge equivalent under G_{Spin} -valued Čech 0-cochains and corresponding soldering data induce the same Lorentzian metric. The hypothesis therefore does not treat that family as an attempted construction of curvature. The local spin-factor, Clifford and Lorentz structures are to remain fixed. A later finite transport model may instead use one global scalar λ to scale an edge-dependent generator assignment,

$$U_{xy}(\lambda) = \exp(\lambda K_{xy}), \quad K_{yx} = -K_{xy}, \quad K_{xy} \in \mathfrak{g},$$

without introducing independent local parameters. This is a comparison family for nontrivial discrete holonomy; it is not a dynamical law.

Each local spin factor contains the distinguished unit $e_x = (1, 0)$ with $N(e_x) = 1$. The future unit hyperboloid $N(u) = 1$ therefore supplies an intrinsic dimensionless mass-shell normalization. No dimensionful mass or energy scale is thereby generated. If a later scale $m > 0$ is independently supplied or derived, the standard dimensionless variables are

$$\varepsilon = E/(mc^2), \quad \boldsymbol{\pi} = \mathbf{p}/(mc), \quad \varepsilon^2 - \|\boldsymbol{\pi}\|^2 = 1.$$

The same local Lorentz-boost subgroup is to be compared in three settings: special-relativistic kinematics, a future comparison with Levi-Civita parallel transport and timelike geodesics, and the composition of kinematic and transport contributions. The distinguishing geometric information is not a deformation of the local Minkowski model but the path dependence, holonomy and, in a continuum formulation, curvature of the transport law.

Subsequent formal work has now constructed the (-1) -eigenspace $\mathfrak{s} \subset \text{Cl}_3$ of Clifford conjugation, the bracket-preserving real-linear isomorphism $D_B : \mathfrak{s} \xrightarrow{\sim} \mathfrak{so}(B)$, and, for every $X \in \mathfrak{s}$, the algebraic one-parameter subgroup $t \mapsto \exp(tX) \in G_{\text{Spin}}$. It also proves that the derivative at $t = 0$ of the projected action on $\mathcal{S}$ is exactly $D_B(X)$. The immediate formal verification is the explicit grade-one rapidity relation

$$\rho\left(\exp\left(\frac{\eta}{2} \iota(v)\right)\right) e = (\cosh \eta, \sinh \eta v) \quad (\langle v, v \rangle = 1),$$

including the factor $\eta = 2t$ between the Clifford exponential parameter and Lorentz rapidity. The smooth Lie-group identification $\text{Lie}(G_{\text{Spin}}) \cong \mathfrak{s}$, the differential $d\rho_e$, connections, parallel transport, holonomy, curvature and torsion remain open in the present development.

The Relational Spin-Closure Hypothesis begins only after these distinctions. It proposes that a future classical configuration space carries a gauge-equivariant constraint whose solution set couples state and geometric variables nontrivially. Quantization and particle interpretations are later tests, not premises.

Contents

1	Status convention	4
2	Mathematical results used from Paper I	4
2.1	Local algebraic construction	4
2.2	Global quotient and tangent transitions	5
2.3	Soldering isomorphism and induced Lorentzian metric	5
2.4	Lifting obstruction on the chosen cover	6
2.5	Examples and nonuniqueness	6
2.6	One-parameter Čech 1-cocycle family	6
3	The Relational Spin-Closure Hypothesis	7
3.1	Configuration space and gauge symmetry	7
3.2	Symmetric reference configuration	8
3.3	One-parameter transport families and constrained dynamics	9
3.4	Linearized coupling between state and geometric variables	12
3.5	Classical formulation prior to quantization	12
4	Current infinitesimal results and the open connection problem	13
4.1	The (-1) -eigenspace of Clifford conjugation and the induced Lie-algebra map	13
4.2	Algebraic one-parameter subgroups and projected derivatives	13
4.3	Remaining smooth Lie-group identification	14
4.4	Connection, curvature and torsion	14
5	Discrete connection on a graph	14

6	Falsification criteria and later benchmarks	15
6.1	Mathematical falsification criteria	15
6.2	Benchmark hierarchy	16
7	Research order	16
8	Object and status table	17
9	Relation to established mathematics	18
10	Conclusion	19

1 Status convention

The labels used below have the following meanings:

Paper I	proved or constructed in Paper I;
Formalized	proved in the subsequent formal Lean development;
Standard	standard external mathematics;
Open	open mathematical problem;
Hypothesis	hypothesis introduced here;
Benchmark	later mathematical or empirical benchmark.

No statement labelled Open, Hypothesis or Benchmark is asserted as a theorem of Paper I.

2 Mathematical results used from Paper I

2.1 Local algebraic construction

The primitive local datum is

$$V = \mathbb{R}^3, \quad \langle u, v \rangle > 0 \quad (u \neq 0). \quad (2.1)$$

No manifold, tangent bundle, Lorentzian metric, orientation, principal bundle or connection is part of this datum.

Paper I constructs the real spin factor

$$\mathcal{S} = \mathbb{R} \oplus V, \quad N(a, v) = a^2 - \langle v, v \rangle, \quad (2.2)$$

with polarized bilinear form B of signature $(1, 3)$, together with the real Clifford algebra

$$\text{Cl}_3 = \text{Cl}(V, q), \quad q(v) = \langle v, v \rangle. \quad (2.3)$$

The Clifford norm-one group G_{Spin} acts on the paravector model and yields

$$1 \longrightarrow \{\pm 1\} \longrightarrow G_{\text{Spin}} \xrightarrow{\rho} G_{\text{Lor}} \longrightarrow 1, \quad (2.4)$$

where $G_{\text{Lor}} \subset O(B)$ is the proper orthochronous Lorentz group defined by the determinant and future-cone conditions used in Paper I. The homomorphism ρ is continuous and admits local continuous sections. No smooth Lie-group structure on G_{Spin} is asserted there.

The B -skew-adjoint endomorphisms form

$$\mathfrak{so}(B) = \{A \in \text{End}_{\mathbb{R}}(\mathcal{S}) : B(Ax, y) + B(x, Ay) = 0\}, \quad (2.5)$$

and Paper I proves a linear isomorphism

$$\Lambda^2 \mathcal{S} \xrightarrow{\sim} \mathfrak{so}(B), \quad \dim_{\mathbb{R}} \mathfrak{so}(B) = 6. \quad (2.6)$$

The identification

$$\text{Lie}(G_{\text{Spin}}) \cong \mathfrak{so}(B) \quad (2.7)$$

is *not* a result of Paper I.

2.2 Global quotient and tangent transitions

Let $\{D_i\}_{i \in I}$ be local four-dimensional pieces. A gluing datum specifies overlap domains and homeomorphisms

$$\phi_{ij} : W_{ij} \longrightarrow W_{ji} \quad (2.8)$$

satisfying the stated identity, inverse and cocycle conditions. Under the closed-gluing-relation and countability hypotheses of Paper I, the quotient is Hausdorff and second countable. If the transition maps are smooth, the resulting atlas defines a smooth four-manifold M .

The local pieces alone do not determine a unique quotient:

$$\{D_i\}_{i \in I} \not\Rightarrow \text{a unique } M. \quad (2.9)$$

The tangent transition cocycle is

$$T_{ij}(y) = D\phi_{ij}(y). \quad (2.10)$$

Independently, let

$$\tilde{g}_{ij} : U_i \cap U_j \longrightarrow G_{\text{Spin}} \quad (2.11)$$

be a G_{Spin} -valued Čech 1-cocycle, and set

$$g_{ij} := \rho \circ \tilde{g}_{ij} : U_i \cap U_j \longrightarrow G_{\text{Lor}}. \quad (2.12)$$

In general,

$$D\phi_{ij} \neq g_{ij}. \quad (2.13)$$

Thus the tangent and Lorentz-valued transition cocycles are distinct data before an additional compatibility condition is imposed.

2.3 Soldering isomorphism and induced Lorentzian metric

Let $E_g \rightarrow M$ be the Lorentz vector bundle defined by the cocycle g_{ij} . Local soldering data are smooth maps

$$A_i : D_i \longrightarrow \text{GL}(\mathcal{S}) \quad (2.14)$$

satisfying

$$A_j(\phi_{ij}(y)) = D\phi_{ij}(y) A_i(y) g_{ij}(x_y). \quad (2.15)$$

They determine a vector-bundle isomorphism

$$s : E_g \xrightarrow{\sim} TM. \quad (2.16)$$

The inverse

$$s^{-1} : TM \xrightarrow{\sim} E_g \quad (2.17)$$

has the conventional direction of an E_g -valued solder form.

The induced metric

$$g_x(X, Y) := B\left(s_x^{-1}X, s_x^{-1}Y\right) \quad (2.18)$$

is a smooth Lorentzian metric on M .

The logical dependence is therefore

$$\boxed{(V, \langle \cdot, \cdot \rangle) \Longrightarrow (\mathcal{S}, B, \text{Cl}_3, G_{\text{Spin}}, \rho, \mathfrak{so}(B))} \quad (2.19)$$

locally, but globally

$$\boxed{\text{gluing datum} + \tilde{g} + \text{soldering data} \Longrightarrow (M, TM, g)}. \quad (2.20)$$

Neither additional term on the left of (2.20) is derived solely from (2.19).

2.4 Lifting obstruction on the chosen cover

For a Lorentz-valued Čech 1-cocycle g_{ij} , assume continuous lifts

$$h_{ij} : U_i \cap U_j \longrightarrow G_{\text{Spin}}, \quad \rho(h_{ij}) = g_{ij} \quad (2.21)$$

have been chosen on the double overlaps. On triple overlaps define

$$\epsilon_{ijk} := h_{jk} h_{ij} h_{ik}^{-1} \in \{\pm 1\}. \quad (2.22)$$

This defines the $\mathbb{Z}/2$ -valued Čech 2-cocycle class used in Paper I. The paper does not identify this class with the classical characteristic class $w_2(TM)$. Likewise the determinant condition obtained from (2.15) is not identified there with $w_1(TM)$.

In particular, the following identifications are not part of the mathematical baseline:

$$[\epsilon] = w_2(TM), \quad [\det D\phi] = w_1(TM). \quad (2.23)$$

2.5 Examples and nonuniqueness

Paper I establishes the following limitations:

$$\text{topological gluing} \not\Rightarrow \text{smooth gluing}, \quad (2.24a)$$

$$\text{pointwise fibre isomorphisms} \not\Rightarrow \text{continuous soldering}, \quad (2.24b)$$

$$\text{orientation-reversing two-chart gluing} \Rightarrow \text{no soldering data}, \quad (2.24c)$$

$$\text{combinatorial labels} \not\Rightarrow \text{existence of soldering data}. \quad (2.24d)$$

On the chosen periodic cover, distinct lifts may have the same Lorentz projection; this is a statement about the chosen cover and the specified equivalence relation, not a computation of $H^1(M; \mathbb{Z}/2)$.

2.6 One-parameter Čech 1-cocycle family

For the explicit periodic family of Paper I, let $\tilde{g}^{(\lambda)}$ denote the G_{Spin} -valued Čech 1-cocycle and let

$$\mathcal{E}_\lambda := \{\text{local soldering data for } \tilde{g}^{(\lambda)}\}. \quad (2.25)$$

Then

$$\forall \lambda \in \mathbb{R}, \quad \mathcal{E}_\lambda \neq \emptyset. \quad (2.26)$$

Moreover, for all $\lambda_1, \lambda_2 \in \mathbb{R}$, there are continuous G_{Spin} -valued Čech 0-cochains u_i such that

$$\tilde{g}_{ij}^{(\lambda_2)} = u_i \tilde{g}_{ij}^{(\lambda_1)} u_j^{-1}, \quad (2.27)$$

and there is an explicit bijection

$$\mathcal{T}_{\lambda_1, \lambda_2} : \mathcal{E}_{\lambda_1} \xrightarrow{\sim} \mathcal{E}_{\lambda_2} \quad (2.28)$$

with

$$g_{\mathcal{T}_{\lambda_1, \lambda_2}}(A) = g_A. \quad (2.29)$$

No statement quantified over arbitrary covers, arbitrary cocycle families or arbitrary one-parameter subgroups follows from (2.26)–(2.29).

The selected one-parameter subgroup used in Paper I is explicitly a Lorentz-boost subgroup,

$$G(\lambda) = \cosh\left(\frac{\lambda}{2}\right) + \sinh\left(\frac{\lambda}{2}\right) \gamma_0,$$

whose Lorentz projection acts on the distinguished timelike–spacelike plane by

$$\begin{pmatrix} \cosh \lambda & \sinh \lambda \\ \sinh \lambda & \cosh \lambda \end{pmatrix}.$$

The direction γ_0 is selected, not canonical. In the present hypothesis this local one-parameter Lorentz-boost subgroup is not assigned exclusively to either special-relativistic momentum or gravitation. The intended distinction lies in the transport structure: special-relativistic kinematics may coexist with path-independent transport and trivial loop products, whereas a future connection-theoretic gravitational comparison would require path-dependent transport with nontrivial holonomy or curvature after those objects have been constructed.

3 The Relational Spin-Closure Hypothesis

The constructions above determine the present mathematical boundary. The hypothesis begins beyond that boundary.

3.1 Configuration space and gauge symmetry

The future classical variables have not yet been fully constructed. Accordingly, their final product structure is not assumed. Let

$$\tilde{\mathcal{Q}} \tag{3.1}$$

denote a future space of admissible classical configurations extending the geometric structures already constructed, and let $\mathcal{G}$ be its gauge group. If a quotient is mathematically well behaved, write

$$\mathcal{Q} := \tilde{\mathcal{Q}}/\mathcal{G}. \tag{3.2}$$

No manifold structure on $\mathcal{Q}$ is assumed at this stage.

Hypothesis 3.1 (Relational Spin-Closure Hypothesis). *There exist a classical configuration space $\tilde{\mathcal{Q}}$, a gauge group $\mathcal{G}$, a target space $\mathcal{Y}$, and a map*

$$\mathfrak{C} : \tilde{\mathcal{Q}} \longrightarrow \mathcal{Y} \tag{3.3}$$

such that:

(i) $\mathfrak{C}$ is gauge equivariant and $0 \in \mathcal{Y}$ is fixed by the gauge action;

(ii) the physical solution set

$$\mathcal{Z} := \mathfrak{C}^{-1}(0)/\mathcal{G} \tag{3.4}$$

is nonempty and contains nontrivial gauge classes;

(iii) *the geometric variables entering $\tilde{\mathcal{Q}}$ are solved jointly with the state variables rather than prescribed as fixed external functions;*

(iv) *the local Euclidean/Clifford/Lorentz construction of Paper I is not altered in order to obtain a desired macroscopic equation.*

The notation $\mathfrak{C}$ names the proposed equation only. It is not identified with torsion, curvature, a Čech cocycle, a connection, or any previously constructed object.

3.2 Symmetric reference configuration

The flat and homogeneous family studied in Paper I is an intentional reference configuration. It is not evidence that every admissible configuration must be flat. The local algebraic model is held fixed:

$$(V_x, \langle \cdot, \cdot \rangle_x, \mathcal{S}_x, B_x, \text{Cl}_x) \cong (\mathbb{R}^3, \langle \cdot, \cdot \rangle, \mathcal{S}, B, \text{Cl}_3)$$

for every local copy under consideration. No gravitational degree of freedom is introduced by deforming this local Clifford/Lorentz model.

The previously studied one-parameter family is homogeneous at the level of the transition data: the same parameter changes the transition cocycle coherently, and

$$\tilde{g}^{(\lambda_1)} \sim_{\mathcal{G}} \tilde{g}^{(\lambda_2)}, \quad g_{\lambda_1} = g_{\lambda_2}.$$

Its gauge equivalence therefore characterizes that particular family; it is not a theorem that all inhomogeneous or path-dependent transport assignments are gauge equivalent.

Later extensions are required to preserve the local Clifford/Spin and Lorentz structures while allowing transport data between local frames to vary subject to explicitly stated compatibility and evolution equations. In particular, a non-flat geometry must not be obtained merely by prescribing

$$s = s(x), \quad \lambda = \lambda(x), \quad \omega = \omega(x)$$

as independent external functions and then computing the resulting curvature. Such fields may enter only after they have been constructed or constrained by the mathematical variables of the model.

For a finite graph model with identical local spin-factor/Clifford structures and edge transport variables

$$U_{xy} \in G_{\text{Spin}}, \quad U_{yx} = U_{xy}^{-1},$$

the reference configuration is characterized by trivial ordered products around every closed path,

$$H_C = 1 \quad \text{for every closed path } C.$$

A nontrivial conjugacy class

$$[H_C] \neq [1]$$

is therefore a gauge-invariant indication of path dependence in the graph transport data.

In a continuum theory based on a principal connection ω , the corresponding standard notions must be distinguished carefully. The curvature two-form is

$$F_\omega = d\omega + \frac{1}{2}[\omega \wedge \omega].$$

The condition $F_\omega = 0$ means that the connection is flat, whereas $F_\omega \neq 0$ is local curvature of that connection. Nontrivial holonomy can nevertheless occur for a flat connection on a topologically nontrivial base. Consequently, a nontrivial graph loop product is not identified either with continuum curvature or with continuum holonomy until an explicit comparison theorem has been proved.

The mathematical question is therefore whether the coupled compatibility and evolution equations for several locally identical spin-factor/Clifford structures admit or require gauge-inequivalent transport data while preserving the local Lorentzian model.

3.3 One-parameter transport families and constrained dynamics

A one-parameter family used for comparison may be written

$$U_{xy}(\lambda) = \exp(\lambda K_{xy}), \quad K_{xy} \in \mathfrak{s}, \quad K_{yx} = -K_{xy}, \quad (3.5)$$

with one global amplitude $\lambda \in \mathbb{R}$ and a fixed edge-dependent generator assignment K_{xy} . Hence

$$U_{xy}(0) = 1, \quad \left. \frac{d}{d\lambda} U_{xy}(\lambda) \right|_{\lambda=0} = K_{xy}.$$

The scalar λ scales a prescribed family of transport variables; it does not create independent local degrees of freedom. Equation (3.5) is therefore a comparison family, not an explanation of dynamics.

Intrinsic unit normalization and the dual role of Lorentz boosts

Every local spin factor has the distinguished unit

$$e_x = (1, 0), \quad N(e_x) = 1. \quad (3.6)$$

Define the future unit hyperboloid

$$\mathcal{H}_x^+ := \{u = (\varepsilon, \boldsymbol{\pi}) \in \mathcal{S}_x : N(u) = 1, \varepsilon > 0\}. \quad (3.7)$$

The equation $N(u) = 1$ is an intrinsic dimensionless normalization. It is not a dimensional energy law and does not generate a mass scale. If a positive mass scale m is later independently supplied or dynamically derived, the standard dimensionless comparison is

$$\varepsilon = \frac{E}{mc^2}, \quad \boldsymbol{\pi} = \frac{\mathbf{p}}{mc}, \quad \varepsilon^2 - \|\boldsymbol{\pi}\|^2 = 1. \quad (3.8)$$

Thus the local rest state is represented by $u = e_x$, for which the dimensionless rest-energy ratio under this later physical identification is $E_{\text{rest}}/(mc^2) = 1$. Equation (3.8) is a later physical interpretation of the already present Lorentzian unit normalization, not a derivation of m , E , or $\mathbf{p}$.

For a unit spatial direction n , let $L_n(\eta)$ denote the standard proper orthochronous Lorentz boost of rapidity η in direction n . Its action on the rest vector is

$$L_n(\eta)e_x = (\cosh \eta, \sinh \eta n), \quad N(L_n(\eta)e_x) = 1. \quad (3.9)$$

Hence, for an oriented one-dimensional boost subgroup,

$$\varepsilon = \cosh \eta, \quad \pi_{\parallel} = \sinh \eta, \quad \eta = \operatorname{arcosh} \varepsilon = \operatorname{arsinh} \pi_{\parallel}. \quad (3.10)$$

This gives an inverse kinematic reconstruction: a normalized state on $\mathcal{H}_x^+$ determines the rapidity of the corresponding one-dimensional Lorentz boost, subject to the stated orientation convention.

The formal Clifford development has already proved, for $X \in \mathfrak{s}$, that $\exp(tX) \in G_{\text{Spin}}$ is an algebraic one-parameter subgroup and that the derivative at $t = 0$ of its projected action is $D_B(X)$. For a unit vector $v \in V$, it also proves

$$\exp(t \iota(v)) = \cosh t \, 1 + \sinh t \, \iota(v). \quad (3.11)$$

The immediate formal verification is to identify this curve with the standard boost normalization by proving

$$\rho\left(\exp\left(\frac{\eta}{2} \iota(v)\right)\right) e_x = L_v(\eta)e_x = (\cosh \eta, \sinh \eta v), \quad \langle v, v \rangle = 1. \quad (3.12)$$

Thus the expected relation between the Clifford exponential parameter t and Lorentz rapidity is

$$\eta = 2t. \quad (3.13)$$

Equations (3.12)–(3.13) are stated here as the next formal bridge to be verified; they are not yet counted as formal results of the current development.

The same local Lorentz-boost subgroup then has two distinct standard roles in the comparison programme:

Comparison	Mathematical form	Interpretation
Special-relativistic kinematics	$u = L_n(\eta_{\text{kin}})e_x$, trivial discrete holonomy	relative velocity, rapidity and momentum
General-relativistic free fall	$u = \dot{\gamma}$, $\nabla_u^g u = 0$ for the Levi-Civita connection	boost component of orthonormal-frame parallel transport
Combined comparison	$L_{\text{obs}} = L_{\text{tr}}L_{\text{kin}}$	composition of kinematic and transport contributions

The second and third rows are comparison targets for a future connection theory, not present theorems of the hypothesis.

At the finite level, for an oriented closed path C , define the ordered product

$$H_C(\lambda) = \prod_{e \in C} U_e(\lambda). \quad (3.14)$$

The condition $H_C = 1$ for every closed path is the path-independent reference case for this finite model. A nontrivial conjugacy class $[H_C] \neq [1]$ is a gauge-invariant obstruction to such path independence. A nonzero special-relativistic rapidity therefore does not by itself imply curvature or nontrivial holonomy: one may have $\eta_{\text{kin}} \neq 0$ while all loop products remain trivial.

For comparison with general relativity, a future connection constructed by the model must first be related to the Levi-Civita connection ∇^g of the induced Lorentzian metric g , or any torsion and non-metricity must be stated explicitly. Standard timelike free fall in general relativity is then described by a geodesic γ with four-velocity $u = \dot{\gamma}$ satisfying

$$\nabla_u^g u = 0. \quad (3.15)$$

In a local orthonormal frame, the Levi-Civita connection has the standard component form

$$\frac{du^a}{d\tau} + \omega^a_{b\mu} \dot{x}^\mu u^b = 0. \quad (3.16)$$

In a one-generator boost reduction $u(\tau) = L_n(\eta(\tau))e_x$, the boost component would reduce, after fixing the sign and generator conventions, to an equation of the form

$$\frac{d\eta}{d\tau} = -\omega_K(\dot{\gamma}). \quad (3.17)$$

Equations (3.15)–(3.17) are standard comparison equations. They are not yet derived from the hypothesis, and no identification of a future G_{Spin} -connection with the Levi-Civita connection is assumed.

Composition of kinematic and transport contributions

Write a local comparison between an initially boosted state and a transported frame as

$$L_{\text{obs}} = L_{\text{tr}}L_{\text{kin}}. \quad (3.18)$$

For collinear boosts along the same unit direction n ,

$$L_{\text{kin}} = L_n(\eta_{\text{kin}}), \quad L_{\text{tr}} = L_n(\eta_{\text{tr}}), \quad L_{\text{tr}}L_{\text{kin}} = L_n(\eta_{\text{kin}} + \eta_{\text{tr}}), \quad (3.19)$$

so that $\eta_{\text{obs}} = \eta_{\text{kin}} + \eta_{\text{tr}}$. Consequently,

$$\eta_{\text{tr}} = \text{arcosh}(\varepsilon_{\text{obs}}) - \eta_{\text{kin}} \quad (3.20)$$

under the same oriented one-dimensional convention. This is the simplest inverse reconstruction relevant to the combined comparison: once the kinematic rapidity is specified, the normalized observed state fixes the collinear transport rapidity required by the model.

For noncollinear $K_1, K_2 \in \mathfrak{s}$, set $A_i := D_B K_i \in \mathfrak{so}(B)$. If $[A_1, A_2] \neq 0$, scalar rapidity addition is invalid. On the Lorentz-algebra side the Baker–Campbell–Hausdorff expansion gives

$$\log(e^{\xi A_2} e^{\eta A_1}) = \eta A_1 + \xi A_2 + \frac{1}{2} \eta \xi [A_2, A_1] + \cdots. \quad (3.21)$$

The commutator term contains the additional Lorentz-algebra component determined by the chosen generators; for two noncollinear boost generators this includes the familiar rotational contribution. Such terms are consequences of the noncommutative composition law, not additional variables introduced by definition.

If a differentiable trajectory satisfies the invariant normalization $N(u(\tau)) = 1$, then differentiation gives

$$B(u, \dot{u}) = 0, \quad \dot{u} \in T_u \mathcal{H}_x^+. \quad (3.22)$$

Thus any future dynamical mechanism compatible with the local normalization must move the normalized state tangentially to the unit hyperboloid or modify the transport relation between such states; it must not change the defining Lorentzian normalization of the local spin factor.

A one-parameter group-commutator comparison family may use $K_1, K_2 \in \mathfrak{s}$:

$$H(\lambda) = e^{\lambda K_1} e^{\lambda K_2} e^{-\lambda K_1} e^{-\lambda K_2}. \quad (3.23)$$

Formally near $\lambda = 0$,

$$H(0) = 1, \quad H(\lambda) = 1 + \lambda^2 [K_1, K_2] + O(\lambda^3). \quad (3.24)$$

The first nontrivial term is therefore produced by the noncommutativity of the transport generators while the local spin-factor/Clifford structures remain unchanged. This is a prescribed family of group elements; it does not define an evolution equation.

The dynamical hypothesis requires more than a prescribed one-parameter family. Suppose a future formulation provides a $\mathcal{G}$ -equivariant evolution vector field $\mathcal{V}$ on the constraint surface $\mathfrak{C}^{-1}(0)$, so that an admissible trajectory satisfies

$$\mathfrak{C}(q) = 0, \quad \dot{q} = \mathcal{V}(q). \quad (3.25)$$

At a configuration q , evolution tangent to the gauge orbit,

$$\mathcal{V}(q) \in T_q(\mathcal{G} \cdot q), \quad (3.26)$$

represents no change of the corresponding gauge-equivalence class. Accordingly, a configuration is stationary modulo gauge when (3.26) holds.

The hypothesis requires the existence of admissible configurations for which gauge-invariant transport data differ from the reference configuration and for which the evolution is not tangent to the gauge orbit,

$$\mathcal{V}(q) \notin T_q(\mathcal{G} \cdot q). \quad (3.27)$$

The vector field $\mathcal{V}$ and its dependence on the transport and geometric variables must be determined by the coupled equations of the model; they may not be supplied by prescribing the time dependence of λ , U_{xy} , ω , or the metric.

In the heuristic discussion below, the phrase *non-equilibrium* refers only to this dynamical-systems notion of a nonstationary gauge-equivalence class. No thermodynamic interpretation, entropy functional, statistical ensemble, or irreversible dynamics is assumed.

Remark 3.2 (Heuristic picture). One may picture each local copy as carrying the same Lorentzian unit hyperboloid and the same local spin-factor/Clifford structure. Special-relativistic motion changes the normalized state by a Lorentz boost. A gravitational comparison, if a later connection theory realizes it, would instead involve parallel transport between local orthonormal frames and the possible path dependence of that transport. The hypothesis is that the coupled equations can admit nonstationary gauge-equivalence classes while leaving the local Minkowski model unchanged. This picture is explanatory only; the mathematical content is expressed by Lorentz transformations, transport variables, gauge transformations, the constraint surface, and the evolution vector field defined above.

3.4 Linearized coupling between state and geometric variables

Suppose a future differentiable formulation admits a local splitting

$$T_q \tilde{\mathcal{Q}} = X_\Phi \oplus X_G, \quad (3.28)$$

where X_Φ denotes variations of the state variables and X_G denotes variations of the geometric variables. At a solution $q \in \mathfrak{C}^{-1}(0)$, the linearized constraint is

$$D\mathfrak{C}_q(\delta\Phi, \delta G) = 0. \quad (3.29)$$

A minimal form of linearized coupling is the existence of a tangent vector in the linearized solution space with nonzero components in both summands:

$$\ker D\mathfrak{C}_q \cap ((X_\Phi \setminus \{0\}) \times (X_G \setminus \{0\})) \neq \emptyset. \quad (3.30)$$

This states only that some infinitesimal solution variations change both sets of variables; it does not assert a constitutive law or a dynamical equation.

A stronger local statement can be formulated on specified subspaces when the partial derivatives

$$C_\Phi := D_\Phi \mathfrak{C}_q, \quad C_G := D_G \mathfrak{C}_q \quad (3.31)$$

are invertible there. Then the linearized constraint can be solved in either direction,

$$\delta G = -C_G^{-1} C_\Phi \delta\Phi, \quad \delta\Phi = -C_\Phi^{-1} C_G \delta G. \quad (3.32)$$

This is an implicit-function-type regularity condition and is not part of the present mathematical baseline.

3.5 Classical formulation prior to quantization

The hypothesis is classical at its defining stage. No Hilbert-space state axiom, Born rule, canonical commutation or anticommutation relation, Fock-space construction, path integral, or quantum-statistical postulate is used to define $\mathfrak{C}$.

No clock scalar, lapse, preferred foliation or simultaneity convention belongs to the primitive data. If such an object is introduced later, it requires an independent definition and status.

A variational realization, if it exists, would be a stronger statement:

$$\mathcal{A} : \tilde{\mathcal{Q}} \longrightarrow \mathbb{R}, \quad D\mathcal{A} = 0 \implies \mathfrak{C} = 0. \quad (3.33)$$

No particular action functional is assumed.

4 Current infinitesimal results and the open connection problem

4.1 The (-1) -eigenspace of Clifford conjugation and the induced Lie-algebra map

Subsequent formal work defines the (-1) -eigenspace of Clifford conjugation

$$\mathfrak{s} := \{X \in \text{Cl}_3 : c(X) = -X\} = \ker(\text{id} + c). \quad (4.1)$$

It proves

$$\mathfrak{s} = \text{Cl}_3^{(1)} \oplus \text{Cl}_3^{(2)}, \quad \dim_{\mathbb{R}} \mathfrak{s} = 6, \quad [\mathfrak{s}, \mathfrak{s}] \subseteq \mathfrak{s}. \quad (4.2)$$

For the paravector embedding $A : \mathcal{S} \rightarrow \text{Cl}_3$, the associated real-linear map is characterized by

$$A(D_X x) = XA(x) + A(x)\tilde{X}, \quad (4.3)$$

where $\tilde{X}$ denotes reversion. The resulting map is a bracket-preserving real-linear isomorphism

$$D_B : \mathfrak{s} \xrightarrow{\sim} \mathfrak{so}(B). \quad (4.4)$$

These are algebraic results. They do not by themselves identify $\mathfrak{s}$ with the tangent Lie algebra of G_{Spin} .

4.2 Algebraic one-parameter subgroups and projected derivatives

For every $X \in \mathfrak{s}$, the formal development proves

$$\gamma_X(t) := \exp(tX) \in G_{\text{Spin}}, \quad (4.5)$$

with

$$\gamma_X(0) = 1, \quad \gamma_X(t+s) = \gamma_X(t)\gamma_X(s), \quad \gamma_X(-t) = \gamma_X(t)^{-1}. \quad (4.6)$$

It also proves

$$c(\exp(tX)) = \exp(-tX) \quad (4.7)$$

and the exact projected paravector action

$$A(\rho(\exp(tX))x) = \exp(tX)A(x)\exp(t\tilde{X}). \quad (4.8)$$

The derivative at the origin of this explicit projected curve is the previously constructed bracket-preserving linear map:

$$\left. \frac{d}{dt} \rho(\exp(tX))x \right|_{t=0} = D_B(X)x. \quad (4.9)$$

This is a derivative of an explicit curve acting on the finite-dimensional Lorentz vector space $\mathcal{S}$. It is not yet the differential $d\rho_e$ of a smooth Lie-group homomorphism.

For a unit vector $v \in V$, the grade-one exponential is proved in closed form:

$$\langle v, v \rangle = 1 \implies \exp(t\iota(v)) = \cosh t \, 1 + \sinh t \, \iota(v). \quad (4.10)$$

This result provides the Clifford-algebra side of the standard boost parametrization. The immediate formal verification is equation (3.12), including the relation $\eta = 2t$, and the corresponding proof of $N = 1$ for the projected rest state.

The same development computes the directional derivative at the identity of the norm-one constraint

$$F(g) = g c(g) - 1$$

as

$$DF_1(X) = X + c(X), \quad \ker DF_1 = \mathfrak{s}. \quad (4.11)$$

The $+1$ -eigenspace appearing as the image of $\text{id} + c$ is the fixed subspace of *Clifford conjugation*; it must not be confused with the fixed subspace of reversion. No regular-value or submanifold theorem has yet been applied.

4.3 Remaining smooth Lie-group identification

The next structural statements are

$$\mathrm{Lie}(G_{\mathrm{Spin}}) \stackrel{?}{\cong} \mathfrak{s} \xrightarrow{D_B} \mathfrak{so}(B), \quad (4.12)$$

and

$$d\rho_e \stackrel{?}{=} D_B \quad \text{under this identification.} \quad (4.13)$$

A smooth manifold structure compatible with the group operations on G_{Spin} must be constructed before (4.12)–(4.13) can be interpreted literally in the standard Lie-theoretic sense. The explicit one-parameter subgroups and equation (4.9) provide strong compatibility data but do not substitute for that construction.

4.4 Connection, curvature and torsion

Only after the required smooth principal-bundle and infinitesimal representation structures are available can the standard theory of principal connections be applied. If $P \rightarrow M$ is a principal G_{Spin} -bundle with a principal connection ω , then standard differential geometry gives the curvature

$$F_\omega = d\omega + \frac{1}{2}[\omega \wedge \omega]. \quad (4.14)$$

With a solder form $e : TM \rightarrow E_g$, torsion is

$$T = D_\omega e. \quad (4.15)$$

These are standard definitions [8, 3]; neither object has yet been constructed in the present formal development.

A group homomorphism cannot be applied directly to a Lie-algebra-valued one-form. In particular,

$$\rho \circ \omega \quad (4.16)$$

is not a definition of a Lorentz connection. Projection of a connection requires $d\rho_e$ or an explicitly proved equivalent infinitesimal representation.

The intended flat reference case is

$$F_\omega = 0. \quad (4.17)$$

Any nonzero curvature in the hypothesis must be produced by the coupled constraint and evolution equations, not prescribed as input.

5 Discrete connection on a graph

A discrete connection on a graph may be studied before the continuum connection theory is complete, provided that no continuum-limit identification is assumed without proof.

Let

$$\Gamma = (\mathcal{V}, \mathcal{E}) \quad (5.1)$$

be an oriented graph. Assign

$$U_{xy} \in G_{\mathrm{Spin}}, \quad U_{yx} = U_{xy}^{-1} \quad (5.2)$$

to each oriented edge; these group elements play the role of link variables. A vertex gauge transformation $h_x \in G_{\mathrm{Spin}}$ acts by

$$U_{xy} \mapsto h_y U_{xy} h_x^{-1}. \quad (5.3)$$

For a closed path $C = (x_0, \dots, x_n = x_0)$, define the ordered product (discrete holonomy)

$$H_C := U_{x_{n-1}x_n} \cdots U_{x_0x_1}. \quad (5.4)$$

Then

$$H_C \mapsto h_{x_0} H_C h_{x_0}^{-1}, \quad (5.5)$$

so its conjugacy class $[H_C]$ is gauge invariant, as expected for based holonomy under a change of local frame.

The reference configuration has trivial discrete holonomy,

$$H_C = 1 \quad \text{for every closed path } C. \quad (5.6)$$

Nontrivial discrete holonomy is detected by

$$[H_C] \neq [1] \quad (5.7)$$

for at least one closed path. This condition leaves the local Clifford/Lorentz structure unchanged and concerns only the discrete connection.

No identification

$$[H_C] = [\text{Hol}_\omega(C)] \quad (5.8)$$

is asserted before a continuum limit and an explicit comparison theorem have been constructed.

6 Falsification criteria and later benchmarks

6.1 Mathematical falsification criteria

The following implications are excluded by the current mathematics:

$$V = \mathbb{R}^3 \not\Rightarrow M, \quad (6.1a)$$

$$(B, \tilde{g}) \not\Rightarrow \exists s : E_g \xrightarrow{\sim} TM, \quad (6.1b)$$

$$\text{group homomorphism } \rho \not\Rightarrow d\rho_e, \quad (6.1c)$$

$$\prod_{e \in C} U_e \not\equiv \text{Hol}_\omega(C) \quad \text{without a comparison theorem.} \quad (6.1d)$$

Criterion 6.1 (Nontriviality modulo gauge). *A proposed constraint variable fails if every member of the tested family belongs to one gauge orbit and all observables used by the proposal are constant on that orbit.*

The one-parameter family (2.25)–(2.29) is an explicit example of such a negative result at the transition-cocycle level. It does not imply the same conclusion for connection-level families.

Criterion 6.2 (Independence from benchmark field equations). *A candidate $\mathfrak{C}$ does not establish the hypothesis if it is defined by inserting the Einstein equation, a prescribed Schwarzschild field, a geodesic equation, Maxwell/Yang–Mills equations, an externally prescribed redshift/lapse profile, or a desired particle mass as a primitive constitutive law.*

Criterion 6.3 (No prescribed geometric field as explanatory input). *A proposed non-flat geometric configuration does not establish the hypothesis if it is obtained merely by prescribing a position-dependent soldering field, scalar field, connection, or curvature and then computing the desired geometry from that prescription. A global scalar λ may parameterize a fixed edge-dependent transport assignment K_{xy} , but such a family is only a comparison family. Any physical departure from the symmetric reference configuration must instead be determined by the transport variables together with the coupled constraint and evolution equations.*

Criterion 6.4 (No prescribed time dependence as a dynamical explanation). *A nonzero comparison parameter or a nontrivial loop observable is not by itself a dynamical explanation. The hypothesis requires an evolution equation on the admissible configuration space for which some configuration with nontrivial gauge-invariant transport data has evolution not tangent to its gauge orbit. If the same behavior is obtained solely by prescribing the time dependence of λ , U_{xy} , ω , or the metric, no intrinsic dynamical mechanism has been established.*

Criterion 6.5 (Origin of dimensionful scales). *Every dimensionful quantity must have an explicit mathematical source: boundary or initial data, a dimensionful parameter, a normalization of a classical solution, or a later quantum/renormalization mechanism after quantization has independently been justified.*

6.2 Benchmark hierarchy

Required mathematical structure	Later comparison	Status
Classical state space and evolution law	Lorentz-covariant dynamics on a flat Lorentzian background	Benchmark
Connection and a mathematically defined flat reference configuration	special-relativistic observables	Benchmark
Coupled non-flat classical solutions	weak-field gravitational phenomenology	Benchmark
Stable localized finite-energy solutions	energy, momentum and invariant mass	Benchmark
Reduced phase space / Hamiltonian or symplectic structure	possible quantization	Benchmark
Consistent quantized physical modes	particle/statistics/radiation interpretation	Benchmark
Independent internal non-Abelian gauge sector	confinement / QCD comparisons	Benchmark

No row may be used as an assumption to prove an earlier row.

7 Research order

The logical order of the programme is stated without using later structures as premises for earlier ones.

Established baseline from Paper I. The current geometric construction is

$$(V, \langle \cdot, \cdot \rangle) \implies (\mathcal{S}, B, \text{Cl}_3, G_{\text{Spin}}, \rho, \mathfrak{so}(B)), \quad (7.1)$$

$$\text{gluing datum} + \tilde{g} + \text{soldering data} \implies (M, TM, g).$$

Subsequent formalized algebraic results. The Lean development has established

$$\mathfrak{s} = \{X : c(X) = -X\}, \quad D_B : \mathfrak{s} \xrightarrow{\sim} \mathfrak{so}(B), \quad (7.2)$$

$$\gamma_X(t) = \exp(tX) \in G_{\text{Spin}}, \quad \left. \frac{d}{dt} \rho(\gamma_X(t))x \right|_{t=0} = D_B(X)x.$$

For unit $v \in V$, the closed formula

$$\exp(t \iota(v)) = \cosh t \, 1 + \sinh t \, \iota(v)$$

is also formalized. The immediate next verification is the finite-dimensional rapidity identity

$$\rho\left(\exp\left(\frac{\eta}{2}\iota(v)\right)\right)e \stackrel{?}{=} (\cosh \eta, \sinh \eta v), \quad N = 1. \quad (7.3)$$

This comparison does not require a smooth Lie-group structure on G_{Spin} .

Open Lie-group step. The next structural problem is to construct a compatible smooth Lie-group structure and prove

$$\text{Lie}(G_{\text{Spin}}) \stackrel{?}{\cong} \mathfrak{s}, \quad d\rho_e \stackrel{?}{=} D_B. \quad (7.4)$$

Open connection-theoretic step. Only after the preceding identification can the programme introduce a principal connection ω , parallel transport, holonomy, curvature F_ω , and torsion. In the finite graph model the symmetric reference configuration has

$$H_C = 1 \quad \text{for every closed path } C. \quad (7.5)$$

For a continuum principal connection, flatness is the separate condition

$$F_\omega = 0. \quad (7.6)$$

These statements must not be identified without a comparison theorem; in particular, a flat connection may have nontrivial global holonomy.

A one-parameter comparison family may retain one global scalar parameter while assigning different generators to different oriented edges,

$$U_{xy}(\lambda) = e^{\lambda K_{xy}}, \quad K_{xy} \in \mathfrak{s}.$$

The same local Lorentz-boost subgroup can then be compared in special-relativistic kinematics, in a future connection-theoretic description of frame transport, and in the composition of the two. Throughout these comparisons the local normalization $N(u) = 1$ remains fixed.

Hypothesized coupled classical dynamics. Only after the required variables have been constructed does the main hypothesis become a concrete mathematical problem: construct a configuration space $\tilde{\mathcal{Q}}$, a gauge group $\mathcal{G}$, a gauge-equivariant constraint map $\mathfrak{C}$, and an evolution law on the constraint surface such that the quotient contains nontrivial solutions and some admissible configurations are nonstationary modulo gauge.

Physical benchmarks. Special-relativistic observables, gravitational phenomenology, localized finite-energy solutions, quantization, and strong-interaction comparisons are subsequent tests. None of them is used as a definition of the preceding mathematical structures.

The logical dependence is therefore

local and global geometric construction $\longrightarrow$ Clifford–Lorentz algebraic structure $\longrightarrow$ smooth Lie-group structure $\longrightarrow$ principal connection and transport $\longrightarrow$ constrained dynamics $\longrightarrow$ physical comparisons.	(7.7)
---	-------

8 Object and status table

Object or statement	Mathematical form	Status
Euclidean input	$V = \mathbb{R}^3$ with positive-definite inner product	Paper I
Real spin factor	$\mathcal{S} = \mathbb{R} \oplus V$, $N(a, v) = a^2 - \langle v, v \rangle$	Paper I
Distinguished unit and unit hyperboloid	$e = (1, 0)$, $N(e) = 1$, $\mathcal{H}^+ = \{u : N(u) = 1, u_0 > 0\}$	Paper I
Dimensionless mass-shell interpretation	$\varepsilon = E/(mc^2)$, $\boldsymbol{\pi} = \mathbf{p}/(mc)$ if a scale m is later supplied	Benchmark
Real Clifford algebra	$\text{Cl}_3 = \text{Cl}(V, q)$	Paper I
Subgroup defined by the Clifford norm-one condition	G_{Spin} as defined in Paper I	Paper I
Central two-fold covering	$1 \rightarrow \{\pm 1\} \rightarrow G_{\text{Spin}} \xrightarrow{\rho} G_{\text{Lor}} \rightarrow 1$	Paper I
Orthogonal Lie algebra	$\mathfrak{so}(B) \cong \Lambda^2 \mathcal{S}$, dimension 6	Paper I
(-1) -eigenspace of Clifford conjugation	$\mathfrak{s} = \{X : c(X) = -X\}$, a six-dimensional Lie algebra under the commutator	Formalized
Bracket-preserving linear isomorphism	$D_B : \mathfrak{s} \xrightarrow{\sim} \mathfrak{so}(B)$	Formalized
Algebraic one-parameter subgroup	$\gamma_X(t) = \exp(tX) \in G_{\text{Spin}}$ for $X \in \mathfrak{s}$	Formalized
Derivative of projected exponential action	$\frac{d}{dt} \rho(\exp(tX))x _0 = D_B(X)x$	Formalized
Smooth four-manifold	obtained from additional gluing data and smoothness hypotheses	Paper I
Tangent transition cocycle	$D\phi_{ij}$	Paper I
Group-valued Čech 1-cocycle	$\tilde{g}_{ij} : U_i \cap U_j \rightarrow G_{\text{Spin}}$	Paper I
Lorentz-valued cocycle	$g_{ij} = \rho(\tilde{g}_{ij})$	Paper I
Soldering isomorphism	$s : E_g \xrightarrow{\sim} TM$, when local soldering data exist	Paper I
Lorentzian metric	$g_x(X, Y) = B(s_x^{-1}X, s_x^{-1}Y)$	Paper I
Čech lifting class on $\mathcal{U}$	$[\epsilon] \in \check{H}^2(\mathcal{U}; \mathbb{Z}/2)$ for chosen lifts	Paper I
Identification with $w_2(TM)$	not proved in Paper I	Open
Identification with $w_1(TM)$	not proved in Paper I	Open
Lie algebra of G_{Spin}	$\text{Lie}(G_{\text{Spin}})$	Open
Differential at the identity	$d\rho_e$	Open
Principal connection	ω	Open
Curvature and torsion	$F_\omega, T = D_\omega e$	Open
Classical configuration space	$(\tilde{\mathcal{Q}}, \mathcal{G})$	Hypothesis
Gauge-equivariant constraint	$\mathfrak{C} : \tilde{\mathcal{Q}} \rightarrow \mathcal{Y}$, $\mathfrak{C} = 0$	Hypothesis
Linearized state–geometry coupling	mixed state/geometric directions in $\ker D\mathfrak{C}$	Hypothesis
Quantum theory	conditional later extension	Benchmark

9 Relation to established mathematics

The algebraic terminology follows standard Clifford, Jordan-algebra, and spin-geometry usage [4, 5]. Smooth manifolds, vector bundles, principal bundles, gauge transformations, connections, parallel transport, holonomy, solder forms, curvature, torsion, geodesics, and constrained dynamical systems are used in their standard mathematical meanings [6, 7, 8, 3]. No claim of novelty is attached to those structures or terms.

The title “Relational Spin-Closure Hypothesis” is a name for the proposal, not the introduc-

tion of a new mathematical operation called “spin closure.” The actual hypothesis is formulated only in terms of standard objects: configuration spaces, gauge group actions, equivariant constraint maps, evolution vector fields, discrete connections, holonomy, and, if later constructed, principal connections and curvature. Heuristic passages are explicitly marked as such and do not define additional mathematical structures.

The specific new proposal is restricted to the existence and physical adequacy of a future coupled classical system with the properties stated in Section 3. General relativity, gauge theory, soliton models, and non-Abelian strong-interaction theory are comparison targets only [9, 10, 11, 12, 13].

10 Conclusion

The current construction contains the chain

$$\begin{aligned}
 V = \mathbb{R}^3 &\implies (\mathcal{S}, B, \text{Cl}_3, G_{\text{Spin}}, \rho, \mathfrak{so}(B)), \\
 &+ \text{gluing datum} \implies M, TM, \\
 &+ \tilde{g}_{ij}, A_i \implies s : E_g \xrightarrow{\sim} TM \implies (TM, g), \\
 \mathfrak{s} = \{X : c(X) = -X\} &\xrightarrow{D_B} \mathfrak{so}(B), \quad D_B \text{ an algebraic isomorphism,} \\
 X \in \mathfrak{s} \implies \gamma_X(t) = \exp(tX) &\in G_{\text{Spin}}, \quad \left. \frac{d}{dt} \rho(\gamma_X(t))x \right|_{t=0} = D_B(X)x.
 \end{aligned} \tag{10.1}$$

For a unit vector v , the formal development additionally proves

$$\exp(t \iota(v)) = \cosh t \, 1 + \sinh t \, \iota(v). \tag{10.2}$$

The immediate formal verification is the explicit comparison with standard Lorentz rapidity,

$$\rho\left(\exp\left(\frac{\eta}{2} \iota(v)\right)\right) e \stackrel{?}{=} (\cosh \eta, \sinh \eta v), \quad N = 1, \tag{10.3}$$

which makes the expected relation $\eta = 2t$ explicit. The next structural questions are

$$\text{Lie}(G_{\text{Spin}}) \stackrel{?}{\cong} \mathfrak{s}, \quad d\rho_e \stackrel{?}{=} D_B, \tag{10.4}$$

followed, if successful, by connection-level geometry.

The symmetric geometry studied so far is an intentional reference configuration. The local spin-factor, Clifford, and Lorentz structures are held fixed, and the graph model begins with trivial discrete holonomy. The hypothesis does not obtain gravitation by deforming the local Minkowski model or by freely prescribing a position-dependent soldering field, scalar field, connection, or curvature. Each local spin factor retains the normalization $e_x = (1, 0)$, $N(e_x) = 1$; the future unit hyperboloid is therefore a dimensionless mass-shell normalization, not a generated mass scale.

A single global scalar may parameterize the comparison family

$$U_{xy}(\lambda) = e^{\lambda K_{xy}},$$

with an edge-dependent generator assignment K_{xy} . The same local Lorentz-boost subgroup can then be compared with special-relativistic kinematics, with a future connection-theoretic description of orthonormal-frame transport, and with the composition of the two. Collinear rapidities add, whereas noncollinear boosts are governed by the noncommutative Lorentz composition law and its commutator terms. These are standard kinematic and group-theoretic facts; they do not constitute an evolution equation.

The dynamical hypothesis requires a coupled constrained evolution law on $\mathfrak{C}^{-1}(0)$. Some admissible configurations with nontrivial gauge-invariant transport data must have an evolution vector field that is not tangent to the gauge orbit, while the local Lorentzian normalization remains fixed. The time dependence of λ , U_{xy} , a connection, or the metric may not be prescribed merely to obtain this behavior.

In the graph model the distinction between trivial and nontrivial discrete holonomy is

$$H_C = 1 \quad \text{for every closed path } C, \quad \text{versus} \quad [H_C] \neq [1] \quad \text{for some } C. \quad (10.6)$$

For a future continuum principal connection, flatness is instead the local condition

$$F_\omega = 0, \quad (10.7)$$

and nonzero F_ω is curvature of that connection. Flatness does not exclude nontrivial global holonomy. Consequently, the discrete and continuum statements remain distinct until an explicit continuum-limit comparison has been proved.

The later hypothesis is the existence of

$$(\tilde{\mathcal{Q}}, \mathcal{G}, \mathcal{Y}, \mathfrak{C}), \quad \mathfrak{C} : \tilde{\mathcal{Q}} \rightarrow \mathcal{Y}, \quad \mathfrak{C}^{-1}(0)/\mathcal{G} \neq \emptyset, \quad (10.8)$$

with nontrivial gauge classes and nontrivial coupling between state and geometric variables in the linearized solution space. Failure to derive nontrivial gauge-invariant transport data, failure to construct such a classical constraint and evolution law, or failure of the later physical benchmarks falsifies the corresponding claim. Nothing in Paper I by itself establishes (10.8).

References

- [1] J. Seeck (ed.), *A Conditional Construction of Lorentzian Geometry from Euclidean Three-Space: Clifford Double-Cover Data, Gluing, and a Soldering Isomorphism*, companion mathematical paper, 13 September 2026. https://raw.githubusercontent.com/antaris82/The_Relational_Spin-Closure_Hypothesis/main/docs/mathematics/A_Conditional_Construction_of_Lorentzian_Geometry.pdf.
- [2] *The Relational Spin-Closure Hypothesis: formal Lean development*. https://github.com/antaris82/The_Relational_Spin-Closure_Hypothesis.
- [3] M. Nakahara, *Geometry, Topology and Physics*, 2nd ed., Institute of Physics Publishing, Bristol and Philadelphia, 2003. ISBN-13: 978-0-7503-0606-5.
- [4] C. Chevalley, *The Algebraic Theory of Spinors*, Columbia University Press, New York, 1954.
- [5] H. B. Lawson Jr. and M.-L. Michelsohn, *Spin Geometry*, Princeton University Press, Princeton, 1989.
- [6] J. M. Lee, *Introduction to Smooth Manifolds*, 2nd ed., Springer, 2013.
- [7] D. Husemoller, *Fibre Bundles*, 3rd ed., Springer, 1994.
- [8] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry, Vol. I*, Interscience, New York, 1963.
- [9] A. Einstein, *Die Grundlage der allgemeinen Relativitätstheorie*, Annalen der Physik 354(7) (1916), 769–822. DOI: [10.1002/andp.19163540702](https://doi.org/10.1002/andp.19163540702).

- [10] T. W. B. Kibble, *Lorentz Invariance and the Gravitational Field*, Journal of Mathematical Physics 2(2) (1961), 212–221. DOI: [10.1063/1.1703702](https://doi.org/10.1063/1.1703702).
- [11] T. H. R. Skyrme, *A Non-Linear Field Theory*, Proceedings of the Royal Society A 260 (1961), 127–138. DOI: [10.1098/rspa.1961.0018](https://doi.org/10.1098/rspa.1961.0018).
- [12] C. N. Yang and R. L. Mills, *Conservation of Isotopic Spin and Isotopic Gauge Invariance*, Physical Review 96 (1954), 191–195. DOI: [10.1103/PhysRev.96.191](https://doi.org/10.1103/PhysRev.96.191).
- [13] K. G. Wilson, *Confinement of Quarks*, Physical Review D 10 (1974), 2445–2459. DOI: [10.1103/PhysRevD.10.2445](https://doi.org/10.1103/PhysRevD.10.2445).