

**ERC Starting Grant 2025
Research proposal [Part B1]**

A data-driven approach to constructions of abstract polytopes and related objects.

PolyData

Cover Page:

- Name of the Principal Investigator (PI): José Antonio Montero Aguilar.
- Name of the PI's Host Institution for the project: University of Ljubljana.
- Proposal duration in months: 60.

Proposal Summary:

There are fundamental differences between the standard scientific method and the method of mathematical research. Broadly speaking, the standard scientific method is founded upon observations; these observations are eventually used to formulate hypotheses and those are then confirmed or rejected based on carefully design experiments.

In contrast to the natural sciences, in many mathematical disciplines, there is frequently a paucity of initial data upon which hypotheses can be formulated. Quite often experimental methodology is absent, making it challenging to validate conjectures through empirical evidence. Mathematicians are so used to this way of conducting research that the possibility of using experimental data is often overlooked even when available or not too difficult to obtain. This project aims to change this reality in the realm of abstract polytopes.

Abstract polytopes (APs) are combinatorial objects that generalise polygons, polyhedra and other geometrical objects. The study of APs comes hand in hand with the study of their symmetries. When working with symmetric objects, a natural question is: how symmetrical can an object be? More precisely, given certain symmetry conditions, is there such an object that satisfies these conditions? What structural conditions are imposed on such an object by the symmetry constraints? In this proposal we address these questions for APs and related objects.

We propose a data-driven approach with a two-way flow of knowledge: with the aim of generating new datasets of highly symmetric APs and related objects, we will develop new theoretical results and methods, enabling us to devise new algorithms and combinatorial representations of APs. The algorithms will then be carefully implemented and executed. The obtained datasets will then be analysed with the aim of finding interesting patterns, suggesting new conjectures and proposing new directions for further research.

Section a: *Extended Synopsis of the scientific proposal*

Background

There are fundamental differences between the standard scientific method and the method of mathematical research. In essence, the scientific method is founded upon the observation of natural phenomena and the measurement of physical quantities. These are then used to pose hypotheses, which are then tested against cleverly designed experiments. In this way, many false hypotheses can be quickly rejected, while those that survive these tests are eventually recognised as laws of nature. Conversely, in theoretical mathematics, there is frequently a paucity of initial data upon which hypotheses can be formulated. The process of mathematical research thus typically commences with an abstract concept, which may be based on intuition and a limited number of illustrative examples. These ideas eventually evolve to conjectures. However, in contrast to the natural sciences, experimental methodology is absent in many mathematical disciplines, making it challenging to validate conjectures through empirical evidence. As we all know too well, this frequently results in significant time investment in attempts to disprove erroneous conjectures, hours that could be spent on proving theorems instead.

Mathematicians are so used to this way of conducting research that the possibility of using experimental data is often overlooked even when available or not too difficult to obtain. This is why we consider the construction of explicit and manageable interesting examples to be one of the most important lines of research in mathematics. In recent years, with the development of computers, the ability to run experiments and compute large datasets of mathematical objects has opened up new possibilities in mathematical research.

Of our particular interest is to build objects with a large degree of mathematical symmetry. While each science focus on the consequences of the presence or absence of symmetries (see Noether's theorem [39] for an example in Physics, or Jaffe, Altman and Merryman [24] for an example in Chemistry and Medicine), in mathematics we study the phenomenon in arguably its purest essence. This is usually controlled by the action of a certain group of transformations that respect the global structure, often called *automorphisms*. An object is often considered symmetric if it has a *rich* automorphism group; this notion of *richness* might be present in the form of having a large automorphism group or admitting such a group acting in a specific way (for example, transitively) on the atomic parts of the mathematical object.

Sometimes the presence of symmetry impose conditions on the structure of the mathematical object. There are many examples of this phenomenon: the classification of Platonic solids, the Uniformization Theorem [1] and Thurston's geometrization conjecture [51], or the polycirculant conjecture in graph theory [30]; just to mention some. On the other hand, sometimes structural properties of the objects impose restrictions on the presence of symmetry. Hurwitz bound [23] and Tutte's theorem [52] are classical examples of this phenomenon. In general, when studying symmetries from a mathematical perspective we are interested in the following questions

Problem 1. How symmetrical a given mathematical object can be?; what are the structural properties imposed by high degree of symmetry? what are the possible *symmetry types* of a family of objects? are all those symmetry types achievable?

The core ideas of this proposal pose the questions presented in [Problem 1](#) in the context of highly symmetric abstract polytopes. Our research approach is inspired by the first paragraphs of this introduction and relies on the construction of datasets of such objects (explained in detail below). We propose a data-driven approach with a two-way flow of knowledge: with the aim of generating new datasets of highly symmetric APs and related objects, we will develop new theoretical results and methods, enabling us to devise new algorithms and combinatorial representations of APs. The algorithms will then be carefully implemented and executed. The obtained datasets will then be analysed with the aim of finding interesting patterns, suggesting new conjectures and proposing new directions for further research.

Highly symmetric polytopes and related objects

Abstract polytopes are combinatorial objects that generalise (the face lattice) of convex polytopes. Enumeration and classification of highly symmetric convex polytopes goes all the way to the Greeks and the classical problem of classifying what today we know as *Platonic Solids*, to the beginning of last century when the classification of higher dimensional convex polytopes was achieved. By considering combinatorial objects and hence removing the geometric constraints, often imposed by the ambient space, the possibilities for constructions of highly symmetric abstract polytopes are now wide open, and a complete classification problem becomes unattainable and rather turns into a series of construction and enumeration problems of families with particular characteristics.

A *flag* of an abstract polytope of rank n is a n -tuple of mutually incident faces, one of each rank (dimension). The degree of symmetry of a polytope can be measured by the number of flag-orbits of its automorphism group.

Regular polytopes are those that are flag-transitive, meaning they have exactly 1 flag-orbit. This *symmetry type* of polytopes has been traditionally the most studied one and includes classical examples as Platonic solids and regular tilings of Euclidean and Hyperbolic spaces. A slightly less symmetric class of polytopes is that of *2-orbit polytopes* and among those, *chiral polytopes* are the most studied. The notion of chirality is a classical one in other natural sciences, notably in Chemistry. However, in the context of abstract polytopes, chirality has a very specific meaning. Chiral polytopes were formally introduced in [49] by Schulte and Weiss as a combinatorial generalisation of Coxeter's twisted honeycombs [11]. An abstract polytope is chiral if it has 2 orbits on flags such that adjacent flags belong to different orbits. Informally speaking, a chiral polytope is one that admits maximal degree of combinatorial rotations but that do not admit mirror reflections.

Chiral polytopes are, without a doubt, the second most studied symmetry type of abstract polytopes. Even so, the amount of theory developed around this symmetry type is very limited, in particular when compared with the regular case. One potential reason behind this is the building manageable examples of chiral polytopes has proved to be a extremely difficult problem. Moreover the mere existence of regular polytopes in *higher ranks* (every rank larger than 3) was established fairly recently [41]. However, the examples provided by Pellicer are very quickly impractical due to their size. The existence of chiral polytopes in higher ranks can be obtained also a consequence of recent results by Conder, Hubard and O'Reilly [10]. In a joint work with Toledo we used a similar general approach to obtain some results that also prove the existence of chiral polytopes in higher ranks (see [38]).

When we study abstract regular polytopes with less symmetry, that is, with several flag orbits, the number of such orbits gives us only partial information. Here is where the notion of the *symmetry type graph* (STG) comes in handy. The *flag-graph* of a polytopes the (edge-coloured) graph whose nodes are the flags of the polytope in such a way that two of them are connected by an edge of colour i ($i \in \{0, \dots, n-1\}$) if the corresponding flags differ exactly on the face of rank i . It can be proved that the automorphism group of a polytope acts semiregularly as colour-preserving graph automorphisms of the flag-graph. The *symmetry type graph* (STG) of a polytope is the quotient of its flag-graph by its automorphism group [12]. This graph has one vertex for each flag-orbit and its connections show the local arrangement of the flag orbits.

Maniplexes arise naturally as graph-theoretical generalisations of abstract polytopes. A *maniplex of rank n* (or *n -maniplex*) is a properly-edge-colour n -valent graph that satisfies some properties (that can be made precise) that resemble those of the flag-graph of a polytope. *Premaniplexes* (see [19, 20]) are properly edge coloured graphs that satisfy certain (natural) conditions so that, when connected, become the most natural candidate to be the symmetry graph of a polytope, in the sense that every STG of a k -orbit polytope is a connected premaniplex with k vertices.

Determining whether or not there exist polytopes or maniplexes with a given symmetry type is one of the most relevant and long lasting problems on the theory; it has shown to be extremely difficult even for the simplest cases. However, we pose the following conjecture as theoretical driving goal of our research.

Conjecture 1 (Symmetry-type conjecture). *For every n -premaniplex \mathcal{T} there exist an abstract polytope \mathcal{P} such that the symmetry type graph of \mathcal{P} is isomorphic to \mathcal{T} .*

This conjecture has been presented before in the form of a problem (see [14, Problem 12]). In [43], Pellicer, Potočník and Toledo build examples of 2-orbit n -maniplexes for $n \geq 4$ and every symmetry type with 2 vertices. Very recently Mochán [31] proved that several of them are actually polytopes. However, it is still unknown if [Conjecture 1](#) holds for most of the 2-orbit symmetry types. [Conjecture 1](#) is completely solved when \mathcal{T} has 3 vertices, that is, for $n \geq 3$ there exist n -polytopes with each of the possible connected premaniplexes with 3 vertices as their symmetry type graph (see [14, Theorem 4.3])

[Conjecture 1](#) defines our research objective.

Research Objective 1. Prove the Symmetry-type conjecture ([Conjecture 1](#)) in full generality.

We essentially offer two theoretical (independent but not necessarily disconnected) approaches towards proving [Conjecture 1](#): the theory of extensions of abstract polytopes and the theory of graph coverings.

Extensions of abstract polytopes

An $(n + 1)$ -polytope \mathcal{P} is an *extension* of an n -polytope \mathcal{K} if all the facets (maximal faces) of \mathcal{P} are isomorphic to \mathcal{K} . Building extensions is a natural way of building polytopes and related objects, after all is using known examples to build new ones.

The extension problem becomes challenging when we impose symmetry constraints on \mathcal{P} . If \mathcal{P} is regular, then \mathcal{K} must also be regular. This situation has been explored deeply and several papers have appeared on the topic (see [15, 40, 42, 46–48], for example). If \mathcal{P} is chiral, then \mathcal{K} must be either regular or chiral with regular facets. In the latter situation universal [50] and finite [13, 37] constructions are known. However, if \mathcal{K} is regular, little is known on the problem of determining whether or not \mathcal{K} admits a chiral extension. In a recent paper [10] Conder, Hubbard and O'Reilly proved that for every $n \geq 4$, the n -simplex admits a chiral extension. In a recently finished paper with Toledo [38] it is proved that all but finitely many regular tessellations of the n -dimensional torus admit a chiral extension.

However, the problem of finding chiral extensions of regular polytopes is mostly open and due to its importance we pose it as one of our driving questions:

Problem 2. Determine which regular polytopes admit a chiral extension.

The Schläfli symbol of an abstract n -polytope \mathcal{K} is a sequence $\{p_1, \dots, p_{n-1}\}$ that describes the local combinatorics of a \mathcal{K} . In particular, if \mathcal{K} has Schläfli symbol $\{p_1, \dots, p_{n-1}\}$, then the number of facets of \mathcal{K} around each $(n - 3)$ -face is p_{n-1} . Not every abstract polytope has a well-defined Schläfli symbol, however, regular or chiral ones do. If \mathcal{P} is a regular or chiral extension of \mathcal{K} and \mathcal{K} has Schläfli symbol $\{p_1, \dots, p_{n-1}\}$, then \mathcal{P} must have Schläfli symbol $\{p_1, \dots, p_{n-1}, q\}$ for some (possibly infinity) q .

Problem 3. Let \mathcal{K} be a regular n -polytope or a chiral polytope with regular facets and assume that \mathcal{K} has Schläfli symbol $\{p_1, \dots, p_{n-1}\}$. Determine the possible integers q so that \mathcal{K} admits a chiral extension of type $\{p_1, \dots, p_{n-1}, q\}$.

Chiral polytopes are just one of $2^n - 1$ possible symmetry types of 2-orbit n -polytopes. Each symmetry type are in correspondence with a proper subset I of $\{0, \dots, n-1\}$ defined by the property that $i \in I$ if and only if any two i -adjacent flags are on the same orbit. If $I \neq \{0, \dots, n-2\}$, then a polytope in class 2_I is facet-transitive hence it makes explore the extension problem: given \mathcal{K} an n -polytope, does \mathcal{K} admit an extension in class 2_I for $I \neq \{0, \dots, n-1\}$ (note the shift on the rank). The case $I = \{0, \dots, n-1\}$ is in correspondence with the problem of constructing *semiregular alternating polytopes* (see [32–35]). These polytopes have two different types of regular facets occurring in an alternating way around each $(n-3)$ -face.

The theory of extensions of APs in class 2_I is mostly unexplored and we consider it a good first step towards achieving [Research objective 1](#).

Research Objective 2. Develop the theory of 2-orbit non chiral extensions of abstract polytopes and, as a consequence, push forward the results about alternating semiregular polytopes.

Abstract polytopes and graphs coverings

For a graph G with the dart-set $D(G)$ and an abstract group Γ , we say that a function $\zeta : D(G) \rightarrow \Gamma$ is a *voltage assignment* on G provided that it maps inverse darts to inverse group elements. Given such a voltage assignment, one can construct the *derived covering graph* $\text{Cov}(G, \zeta)$ and the *derived regular covering projection* $\wp_\zeta : \text{Cov}(G, \zeta) \rightarrow G$ with Γ acting regularly on each fibre as a subgroup of the covering transformations of \wp . Conversely, every regular covering projection is isomorphic (in a sense that can be made precise) to one derived from a voltage assignment.

The automorphism group of a maniplex acts semiregularly on the flags. Hence, every maniplex may be regarded simply as a regular cover of its symmetry type graph. Equivalently, there exists a voltage assignment ζ such that if \mathcal{T} is the symmetry type of a maniplex \mathcal{M} , then $\mathcal{M} = \text{Cov}(\mathcal{T}, \zeta)$. This means that [Conjecture 1](#) can be formulated as a problem in the language of coverings of graphs or rather, of premaniplexes.

A constant problem when using this approach is that quite often the graph $\text{Cov}(\mathcal{T}, \zeta)$ has more than the desired symmetry. This phenomenon can occur essentially for two reasons:

- i) there might be automorphisms of \mathcal{T} that lift to (undesired) automorphisms of $\text{Cov}(\mathcal{T}, \zeta)$ (e.g. a regular maniplex covering the symmetry type graph of chiral maniplexes);
- ii) or there might be automorphisms of $\text{Cov}(\mathcal{T}, \zeta)$ that do not project to automorphisms of \mathcal{T} (e.g. a regular maniplex covering a 3-vertex premaniplex)

Explicit examples of both situations are known (and they are not uncommon). The theory of coverings of graphs offers a way to understanding both phenomena via the study of the groups $\text{MaxLift}(\wp)$ and $\text{MaxProj}(\wp)$ that, for a graph covering $\wp : \overline{G} \rightarrow G$, describe the automorphisms G that lift to \overline{G} and the automorphisms of \overline{G} that project to G , respectively. (see [26–29] for details).

A graph covering $\wp : \overline{G} \rightarrow G$ is called *stable* if $\text{Aut}(\overline{G}) = \text{MaxProj}(\wp)$. For the purposes of solving [Conjecture 1](#), given a premaniplex \mathcal{T} we shall find a stable covering $\wp : \text{Cov}(\mathcal{T}, \zeta) \rightarrow \mathcal{T}$ that satisfies $\text{MaxLift}(\wp) = 1$. This is not a new problem, and in the more general theory of coverings of graphs, the following conjecture is believed to be true (see [44]):

Conjecture 2. *For every graph G and $\Gamma \leq \text{Aut}(G)$ (satisfying some mild conditions that can be made precise), there exists a stable non-identity regular covering projection $\wp : \overline{G} \rightarrow G$ such that $\Gamma = \text{MaxLift}(\wp)$.*

As part of the implementation of our research we shall explore the validity of [Conjecture 2](#) for coverings of (pre)maniplexes. This shall not only provide with tools to solve [Conjecture 1](#) but it will also give us insight into the more general problem of proving [Conjecture 2](#).

Dataset	Rank 3	Rank 4	Rank ≥ 5
Hartley - Regular [17]	64.55%	31.61%	3.84%
Hartley - Chiral [16]	85.71%	14.29%	0.00%
Conder - Regular [7]	61.51%	34.70%	3.79%
Conder - Chiral [5]	87.01%	12.87%	0.12%
Leemans - Regular [25]	95.35%	4.37%	0.30%
Leemans - Chiral [18]	87.82%	11.97%	0.21%

Table 1: Percentages of examples according to rank

Datasets of abstract polytopes

The research objectives presented above are classical and natural in the theory of abstract polytopes. Our approach would not be innovative in any way if it was not connected with and experimental and data-oriented complement.

Theoretical constructions of abstract polytopes have lead to the constructions of some datasets of highly symmetric polytopes and related objects (see [3, 4, 6–9, 16–18, 25, 45]). However, the existing datasets of polytopes suffer of the following restrictions

- (i) They are mainly focused on regular or chiral polytopes.
- (ii) The size of the examples is very restrictive.
- (iii) They often exhibit numerous examples of rank 3 but the amount of examples of rank higher than 4 drops dramatically.
- (iv) They are not very user-friendly, either because they exist only as raw data or because they are specific-programming language oriented.

In Table 1 we show the proportion of examples according to ranks on the existing datasets of polytopes with examples of rank higher than 3. The points addressed above clearly show a need to push forward the existing datasets. Of course, this leads to our last research objective.

Research Objective 3. Extend the existing and build new datasets of abstract polytopes and related structures with particular focus on

- (i) Building examples on ranks higher than 3;
- (ii) Exploring different symmetry types.

General implementation and project management

I strongly believe in team collaboration and as a junior researcher, I am committed to support my fellow young colleagues. While I plan to carry most of the research, I plan to form a research team consisting mostly of junior researchers. My current research network already counts with young experts in relevant areas for our research: Micael Toledo and Primož Potočnik are leading experts on covers of graphs and maniplexes while Rhys Evans is not only a strong theoretical mathematician but an expert in the development of software packages. Katja Berčič is a leading expert in mathematical knowledge management and her input will be crucial towards completing Research objective 3. Moreover, a number of well established researchers have expressed an interest to spend part of the time collaborating in this research. This list include Marston Conder (University of Auckland), Isabel Hubbard (UNAM - Mexico) and Gabe Cunningham (Wentworth Institute of Technology).

References

- [1] W. Abikoff. “The Uniformization Theorem”. In: *The American Mathematical Monthly* 88.8 (Oct. 1981), pp. 574–592. DOI: [10.1080/00029890.1981.11995320](https://doi.org/10.1080/00029890.1981.11995320).
- [2] J. Collins and A. Montero. “Equivelar Toroids with Few Flag-Orbits”. en. In: *Discrete & Computational Geometry* 65.2 (Jan. 2021), pp. 305–330. DOI: [10.1007/s00454-020-00230-y](https://doi.org/10.1007/s00454-020-00230-y).
- [3] M. Conder. *Chiral orientably-regular maps of genus 2 to 101*. Total number of maps in list below: 594. 2006. URL: <https://www.math.auckland.ac.nz/~conder/ChiralMaps101.txt>.
- [4] M. Conder. *Regular orientable maps of genus 2 to 301*. Total number of maps in list below: 15824. 2011. URL: <https://www.math.auckland.ac.nz/~conder/RegularOrientableMaps301.txt>.
- [5] M. Conder. *Chiral polytopes with up to 2000 flags*. 2012. URL: <https://www.math.auckland.ac.nz/~conder/ChiralPolytopesWithFewFlags-ByOrder.txt>.
- [6] M. Conder. *Regular non-orientable maps of genus 2 to 602*. Total number of maps in list below: 3260. 2012. URL: <https://www.math.auckland.ac.nz/~conder/RegularNonorientableMaps602.txt>.
- [7] M. Conder. *Regular polytopes with up to 2000 flags*. 2012. URL: <https://www.math.auckland.ac.nz/~conder/RegularPolytopesWithFewFlags-ByOrder.txt>.
- [8] M. Conder. *Rotary maps (on orientable or non-orientable surfaces) with up to 1000 edges*. 2012. URL: <https://www.math.auckland.ac.nz/~conder/RotaryMapsWithUpTo1000Edges.txt>.
- [9] M. Conder. *Chiral rotary maps of genus 2 to 301*. Total number of maps in list below: 3870. 2013. URL: <https://www.math.auckland.ac.nz/~conder/ChiralMaps301.txt>.
- [10] M. Conder, I. Hubard, and E. O’Reilly-Regueiro. “Construction of chiral polytopes of large rank with alternating or symmetric automorphism group”. In: *Advances in Mathematics* 452 (Aug. 2024), p. 109819. DOI: [10.1016/j.aim.2024.109819](https://doi.org/10.1016/j.aim.2024.109819).
- [11] H. S. M. Coxeter. *Twisted honeycombs*. Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 4. American Mathematical Society, Providence, R.I., 1970, pp. iv+47. DOI: [10.1090/cbms/004](https://doi.org/10.1090/cbms/004).
- [12] G. Cunningham, M. Del Río-Francos, I. Hubard, and M. Toledo. “Symmetry type graphs of polytopes and maniplaxes”. In: *Ann. Comb.* 19.2 (2015), pp. 243–268. DOI: [10.1007/s00026-015-0263-z](https://doi.org/10.1007/s00026-015-0263-z).
- [13] G. Cunningham and D. Pellicer. “Chiral extensions of chiral polytopes”. In: *Discrete Math.* 330 (2014), pp. 51–60. DOI: [10.1016/j.disc.2014.04.014](https://doi.org/10.1016/j.disc.2014.04.014).
- [14] G. Cunningham and D. Pellicer. “Open problems on k -orbit polytopes”. In: *Discrete Math.* 341.6 (2018), pp. 1645–1661. DOI: [10.1016/j.disc.2018.03.004](https://doi.org/10.1016/j.disc.2018.03.004).
- [15] L. Danzer. “Regular incidence-complexes and dimensionally unbounded sequences of such. I”. In: *Convexity and graph theory (Jerusalem, 1981)*. Vol. 87. North-Holland Math. Stud. North-Holland, Amsterdam, 1984, pp. 115–127. DOI: [10.1016/S0304-0208\(08\)72815-9](https://doi.org/10.1016/S0304-0208(08)72815-9).
- [16] M. Hartley. *An atlas of small chiral polytopes*. 2006. URL: <https://www.abstract-polytopes.com/chiral/>.
- [17] M. Hartley. *The Atlas of Small Regular Polytopes*. 2006. URL: <https://www.abstract-polytopes.com/atlas/>.
- [18] M. Hartley, I. Hubard, and D. Leemans. *An Atlas of Chiral Polytopes for Small Almost Simple Groups*. URL: <https://leemans.dimitri.web.ulb.be/CHIRAL/index.html>.

- [19] I. Hubard and E. Mochán. “All polytopes are coset geometries: Characterizing automorphism groups of k -orbit abstract polytopes”. In: *European Journal of Combinatorics* 113 (Oct. 2023), p. 103746. DOI: [10.1016/j.ejc.2023.103746](https://doi.org/10.1016/j.ejc.2023.103746).
- [20] I. Hubard, E. Mochán, and A. Montero. “Voltage Operations on Maniplexes, Polytopes and Maps”. en. In: *Combinatorica* (May 2023). DOI: [10.1007/s00493-023-00018-7](https://doi.org/10.1007/s00493-023-00018-7).
- [21] I. Hubard, E. Mochán, and A. Montero. “Symmetries of voltage operations on polytopes, maps and maniplexes”. In preparation.
- [22] I. Hubard, A. Orbaníć, D. Pellicer, and A. I. Weiss. “Symmetries of equivelar 4-toroids”. In: *Discrete Comput. Geom.* 48.4 (2012), pp. 1110–1136. DOI: [10.1007/s00454-012-9444-2](https://doi.org/10.1007/s00454-012-9444-2).
- [23] A. Hurwitz. “Ueber algebraische Gebilde mit eindeutigen Transformationen in sich”. de. In: *Mathematische Annalen* 41.3 (Sept. 1892), pp. 403–442. DOI: [10.1007/BF01443420](https://doi.org/10.1007/BF01443420).
- [24] I. A. Jaffe, K. Altman, and P. Merryman. “The Antipyridoxine Effect of Penicillamine in Man”. en. In: *The Journal of Clinical Investigation* 43.10 (Oct. 1964), pp. 1869–1873. DOI: [10.1172/JCI105060](https://doi.org/10.1172/JCI105060).
- [25] D. Leemans, V. Laurence, T. Connor, M. Mixer, and J. Mulpas. *An Atlas of Polytopes for Small Almost Simple Groups*. URL: <https://leemans.dimitri.web.ulb.be/polytopes/index.html>.
- [26] A. Malnič and R. Požar. “On the split structure of lifted groups”. In: *Ars Math. Contemp* 10 (2016), pp. 113–134.
- [27] A. Malnič and R. Požar. “On split liftings with sectional complements”. In: *Math. Comp* 88.316 (2019), pp. 983–1005.
- [28] A. Malnič, D. Marušič, and P. Potočnik. “Elementary Abelian Covers of Graphs”. en. In: *Journal of Algebraic Combinatorics* 20.1 (July 2004), pp. 71–97. DOI: [10.1023/B:JACO.0000047294.42633.25](https://doi.org/10.1023/B:JACO.0000047294.42633.25).
- [29] A. Malnič, R. Nedela, and M. Škoviera. “Lifting Graph Automorphisms by Voltage Assignments”. In: *European Journal of Combinatorics* 21.7 (Oct. 2000), pp. 927–947. DOI: [10.1006/eujc.2000.0390](https://doi.org/10.1006/eujc.2000.0390).
- [30] D. Marušič. “On vertex symmetric digraphs”. In: *Discrete Math.* 36.1 (1981), pp. 69–81. DOI: [10.1016/0012-365X\(81\)90174-6](https://doi.org/10.1016/0012-365X(81)90174-6).
- [31] E. Mochán. “Polytopality of 2-orbit maniplexes”. In: *Discrete Mathematics* 347.9 (Sept. 2024), p. 114055. DOI: [10.1016/j.disc.2024.114055](https://doi.org/10.1016/j.disc.2024.114055).
- [32] B. Monson and E. Schulte. “Semiregular polytopes and amalgamated C-groups”. In: *Advances in Mathematics* 229.5 (Mar. 2012), pp. 2767–2791. DOI: [10.1016/j.aim.2011.12.027](https://doi.org/10.1016/j.aim.2011.12.027).
- [33] B. Monson and E. Schulte. “The Assembly Problem for Alternating Semiregular Polytopes”. In: *Discrete & Computational Geometry* (Aug. 2019). DOI: [10.1007/s00454-019-00118-6](https://doi.org/10.1007/s00454-019-00118-6).
- [34] B. Monson and E. Schulte. “Universal Alternating Semiregular Polytopes”. In: *Canadian Journal of Mathematics* (Feb. 2020), pp. 1–28. DOI: [10.4153/s0008414x20000085](https://doi.org/10.4153/s0008414x20000085).
- [35] B. Monson and E. Schulte. “The interlacing number for alternating semiregular polytopes”. en. In: *The Art of Discrete and Applied Mathematics* 5.3 (Aug. 2022), #P3.11–24 pp. DOI: [10.26493/2590-9770.1406.ec5](https://doi.org/10.26493/2590-9770.1406.ec5).
- [36] A. Montero. “Regular polyhedra in the 3-torus”. In: *Advances in Geometry* 18.4 (2018), pp. 431–450. DOI: [10.1515/advgeom-2018-0017](https://doi.org/10.1515/advgeom-2018-0017).
- [37] A. Montero. “On the Schläfli symbol of chiral extensions of polytopes”. en. In: *Discrete Mathematics* 344.11 (Nov. 2021), p. 112507. DOI: [10.1016/j.disc.2021.112507](https://doi.org/10.1016/j.disc.2021.112507).

- [38] A. Montero and M. Toledo. “Chiral extensions of regular toroids”. Under review.
- [39] E. Noether. “Invariante Variationsprobleme”. deu. In: *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* 1918 (1918), pp. 235–257.
- [40] D. Pellicer. “Extensions of regular polytopes with preassigned Schläfli symbol”. In: *J. Combin. Theory Ser. A* 116.2 (2009), pp. 303–313. DOI: [10.1016/j.jcta.2008.06.004](https://doi.org/10.1016/j.jcta.2008.06.004).
- [41] D. Pellicer. “A construction of higher rank chiral polytopes”. In: *Discrete Math.* 310.6-7 (2010), pp. 1222–1237. DOI: [10.1016/j.disc.2009.11.034](https://doi.org/10.1016/j.disc.2009.11.034).
- [42] D. Pellicer. “Extensions of dually bipartite regular polytopes”. In: *Discrete Math.* 310.12 (2010), pp. 1702–1707. DOI: [10.1016/j.disc.2009.11.023](https://doi.org/10.1016/j.disc.2009.11.023).
- [43] D. Pellicer, P. Potočník, and M. Toledo. “An existence result on two-orbit maniplaxes”. In: *Journal of Combinatorial Theory, Series A* 166 (Aug. 2019), pp. 226–253. DOI: [10.1016/j.jcta.2019.02.014](https://doi.org/10.1016/j.jcta.2019.02.014).
- [44] P. Potočník and P. Spiga. “Lifting a prescribed group of automorphisms of graphs”. In: *Proc. Amer. Math. Soc* 147 (2019), pp. 3787–3796.
- [45] P. Potočník. *Census of chiral maps*. 2014. URL: <https://www.fmf.uni-lj.si/~potocnik/work.htm>.
- [46] E. Schulte. “On arranging regular incidence-complexes as faces of higher-dimensional ones”. In: *European J. Combin.* 4.4 (1983), pp. 375–384. DOI: [10.1016/S0195-6698\(83\)80035-3](https://doi.org/10.1016/S0195-6698(83)80035-3).
- [47] E. Schulte. “Extensions of regular complexes”. In: *Finite geometries (Winnipeg, Man., 1984)*. Vol. 103. Lecture Notes in Pure and Appl. Math. Dekker, New York, 1985, pp. 289–305.
- [48] E. Schulte. “On a class of abstract polytopes constructed from binary codes”. In: *Discrete Mathematics* 84.3 (1990), pp. 295–301. DOI: [10.1016/0012-365X\(90\)90134-4](https://doi.org/10.1016/0012-365X(90)90134-4).
- [49] E. Schulte and A. I. Weiss. “Chiral polytopes”. In: *Applied geometry and discrete mathematics*. Vol. 4. DIMACS Ser. Discrete Math. Theoret. Comput. Sci. Amer. Math. Soc., Providence, RI, 1991, pp. 493–516.
- [50] E. Schulte and A. I. Weiss. “Free extensions of chiral polytopes”. In: *Canad. J. Math.* 47.3 (1995), pp. 641–654. DOI: [10.4153/CJM-1995-033-7](https://doi.org/10.4153/CJM-1995-033-7).
- [51] W. P. Thurston. “Three dimensional manifolds, Kleinian groups and hyperbolic geometry”. en. In: *Bulletin of the American Mathematical Society* 6.3 (1982), pp. 357–381. DOI: [10.1090/S0273-0979-1982-15003-0](https://doi.org/10.1090/S0273-0979-1982-15003-0).
- [52] W. T. Tutte. “On the Symmetry of Cubic Graphs”. In: *Canadian Journal of Mathematics* 11 (1959), pp. 621–624. DOI: [10.4153/CJM-1959-057-2](https://doi.org/10.4153/CJM-1959-057-2).

Section b: Curriculum vitae and Track Record

PERSONAL DETAILS

- **Family name, First name:** Montero, Antonio.
- **Researcher unique identifier(s):**
 - ORCID: [0000-0002-3293-8517](https://orcid.org/0000-0002-3293-8517)
 - Google Scholar: [GEJHkg0AAAAJ](https://scholar.google.com/citations?user=GEJHkg0AAAAJ)
 - MathSciNet Author ID: [1293226](https://mathscinet.ams.org/mathscinet/authorID/1293226)
- **URL for web site:** anteromontonio.github.io

Education and key qualifications

- | | |
|------------|---|
| 19/08/2019 | Ph.D. (Mathematics) , <i>Centro de Ciencias Matematicas, National Autonomous University of Mexico (UNAM)</i> , Mexico.
Under the supervision of Dr. Daniel Pellicer, with the thesis “Chiral extensions of toroids”. |
| 2015 | M. Cs. (Mathematics) , <i>Centro de Ciencias Matematicas, National Autonomous University of Mexico (UNAM)</i> , Morelia, Mexico.
Thesis: “Realizaciones regulares de poliedros en el 3-toro (Regular realizations of polyhedra in the 3-torus)” |
| 2013 | B. Cs. (Mathematics and Physics) , <i>University of Michoacan</i> , Morelia, Mexico.
Thesis: “Poliedros regulares en el 3-toro (Regular polyhedra in the 3-torus)” |

Current position(s)

- | | |
|-------------|--|
| 2022 – 2025 | Znanstveni sodelavec (Research associate, postdoctoral) , <i>Faculty of Mathematics and Physics, University of Ljubljana</i> , Ljubljana, Slovenia. |
| 2024 – | Asistent IX-1 (Teaching assistant) , <i>Faculty of Education, University of Ljubljana</i> , Ljubljana, Slovenia. |

Previous position(s)

- | | |
|-------------|---|
| 2020 – 2022 | Becario postdoctoral (Postdoctoral fellow) , <i>Institute of Mathematics, National Autonomous University of Mexico (UNAM)</i> , Mexico City, Mexico. |
| 2019 – 2020 | Postdoctoral fellow , <i>Department of Mathematics and Statistics, York University</i> , Toronto ON, Canada. |

RESEARCH ACHIEVEMENTS AND PEER RECOGNITION

Research achievements

1. **Constructions of chiral extensions of regular toroids** [38]. In a joint work with Toledo we proved that all but infinitely many tessellations of the torus admit a chiral extension. This results has a consequence the existence of infinitely many chiral polytopes in every rank higher than 4. The research in this paper is closely related to the results in [10] by Conder, Hubard and O'Reilly.

2. **Combinatorial description of classical operations on polytopes** [20, 21]. In joint work with Hubard and Mochán we developed a technique to describe classical geometrical operations of polyhedra, maps, polytopes and similar objects in a combinatorial way. This work lead to the publication of a paper in *Combinatorica* in 2023 [20] and a second paper [21] on the topic is currently under review. I am confident that our work has potential to become a breakthrough in the theory of highly symmetric objects and it has already caught the attention of the community.
3. **Classification of highly symmetric tilings of the n -dimensional torus** [2]. Highly symmetric tessellations of the 2-dimensional torus were described by Coxeter in the 70's. Cubic tilings (regardless of their symmetry degree) of the 3-dimensional torus were classified in 2012 by Hubard, et. al. [22] and they conjectured the non-existence of tilings of a particular symmetry class on higher dimensions. In a paper with Collins [2] we disproved this conjecture and completely classified the highly symmetric cubic tilings of the n -dimensional torus for arbitrary n .
4. **Classification of toroidal regular polyhedra** [36]. Regular polyhedra are classical objects whose study lies in the interplay of combinatorics and geometry. The modern definition of regular polyhedron and a large list of examples were provided by Grünbaum in the 70's and in the 80's Dress achieved a complete classification of the regular polyhedra in the Euclidean space. A natural family of problems arise when we change the ambient space: the analogous problem for the projective space was attacked and eventually solved in the early 2000's while the hyperbolic analogous remains mostly open. In my first paper [36] I complete the classification of regular polyhedra in the 3-dimensional torus.

Peer recognition

January 2021	Candidato a investigador nacional (National research candidate) , <i>National Council of Humanities, Science and Technology (CONAHCyT)</i> , Mexico.. As part of the program <i>Sistema nacional de investigadores (SNI) (National researchers system)</i> .
July 2018	Best student talk award , <i>UP FAMNIT - UP IAM - IMFM</i> , Slovenia.. Award given to the best talk presented by a student during the 8th PhD Summer School in Discrete Mathematics.
October 2012	“Padre de la Patria” , <i>Faculty of Physics and Mathematics, UMSNH</i> , Mexico.. Award given to the best student of each year.
October 2011	“Padre de la Patria” , <i>Faculty of Physics and Mathematics, UMSNH</i> , Mexico.. Award given to the best student of each year.

Other contributions to the research community

Journal Reviewer

- *Ars Mathematica Contemporanea*
- *Discrete Mathematics*
- *Contributions to Discrete Mathematics*
- *The Art of Discrete and Applied Mathematics*
- *Journal of the Australian Mathematical Society*

Organising committees

- June 2023 **10th Slovenian Conference on Graph Theory**, Kranjska Gora, Slovenia..
Member of the organising committee.
- May 2021 **Mexican Mathematical Olympiad in Michoacan**, Morelia, Mich., Mexico..
Member of the local committee.
- February 2018 **Degustaciones Matemáticas 2018**, *Joint Program of Graduate Studies in Mathematics UMSNH-UNAM*, Morelia, Mich., Mexico..
Cycle of talks in the Faculty of Physics and Mathematics, UMSNH.
- February 2017 **Degustaciones Matemáticas 2017**, *Joint Program of Graduate Studies in Mathematics UMSNH-UNAM*, Morelia, Mich., Mexico..
Cycle of talks in the Faculty of Physics and Mathematics, UMSNH.
- February 2016 **Degustaciones Matemáticas 2016**, *Joint Program of Graduate Studies in Mathematics UMSNH-UNAM*, Morelia, Mich., Mexico..
Cycle of talks in the Faculty of Physics and Mathematics, UMSNH.
- February 2015 **Degustaciones Matemáticas 2015**, *Joint Program of Graduate Studies in Mathematics UMSNH-UNAM*, Morelia, Mich., Mexico..
Cycle of talks in the Faculty of Physics and Mathematics, UMSNH.
- Aug. 2014 **Graduate students seminar**, *Joint Program of Graduate Studies in Mathematics UMSNH-UNAM*, Morelia, Mich., Mexico. .
- March 2013 **XXVIII Colloquium “Victor Neumann-Lara” of graph theory, combinatorics and its applications**, Morelia, Mich., Mexico..
Member of the local committee.