

**Problem 1.**

*Solution:* Note that  $f$  is continuous at every point in  $\mathbb{R}^3$ . This implies that Jacobian exists. Let  $f_1 : \mathbb{R}^3 \rightarrow \mathbb{R}$ ,  $f_1(x_1, x_2, x_3) = x_1x_2 + \sin(x_3) + x_1^2$  and  $f_2 : \mathbb{R}^3 \rightarrow \mathbb{R}^1$ ,  $f_2(x_1, x_2, x_3) = 7 + e^{x_2}$ . Therefore

$$\nabla f_1 = [x_2 + 2x_1 \quad x_2 \quad \cos(x_3)] \quad \nabla f_2 = [0 \quad e^{x_2} \quad 0]$$

This implies that

$$J_x = \begin{bmatrix} x_2 + 2x_1 & x_2 & \cos(x_3) \\ 0 & e^{x_2} & 0 \end{bmatrix}$$

We now aim to show what induced one norm on a matrix. For any  $x \in \mathbb{R}^n$  and  $A \in \mathbb{R}^{m \times n}$ , we can see that:

$$\begin{aligned} Ax &= \sum_{j=1}^n a_{ij}x_j \\ \|Ax\|_1 &= \sum_{i=1}^m \left| \sum_{j=1}^n a_{ij}x_j \right| \\ &\leq \sum_{i=1}^m \sum_{j=1}^n |a_{ij}| \cdot |x_j| \\ &\leq \sum_{j=1}^n |x_j| \sum_{i=1}^m |a_{ij}| \\ &\leq \sum_{j=1}^n |x_j| \max_j |c_j| \\ &\leq \max_j |c_j| \end{aligned}$$

where  $c_j$  denotes the sum of the  $j$ th column. To prove the reverse direction, we can see that if we let  $x = e_j$ , where it is the maximum column sum, we can see that

$$\|Ax\|_1 = \sup_{\|x\|_1=1} \|Ax\|_1 \geq \max_j |c_j|$$

which implies that  $\|A\|_1 = \max_j |c_j|$ . Therefore, we see that

$$k_{abs} = \max\{|x_2 + 2x_1|, |x_1 + e^{x_2}|, |\cos(x_3)|\}$$

Therefore, since  $k_{rel} = k_{abs} \cdot \frac{\|x\|_1}{\|f(x)\|_1}$ , we see that:

$$k_{abs} = \max\{|x_2 + 2x_1|, |x_1 + e^{x_2}|, |\cos(x_3)|\} \cdot \frac{|x_1| + |x_2| + |x_3|}{|x_1x_2 + \sin(x_3) + x_1^2 + 7 + e^{x_2}|}$$

□

**Problem 2.**

*Solution:* Let  $x, X, y, Y \in \mathbb{R}$ , the following are derived from the statements given.

$$\begin{aligned}x\|\cdot\|_c &\leq \|\cdot\|_a \leq X\|\cdot\|_c \\y\|\cdot\|_b &\leq \|\cdot\|_c \leq Y\|\cdot\|_b\end{aligned}$$

We can combine these inequalities to find that:

$$xy\|\cdot\|_b \leq x\|\cdot\|_c \leq \|\cdot\|_a \leq X\|\cdot\|_c \leq XY\|\cdot\|_b$$

Thus, showing that  $\|\cdot\|_a$  and  $\|\cdot\|_b$  are indeed equivalent.

□

**Problem 3.**

*Solution:* If  $p = q$ , the statement is obvious. We aim to prove that the 1 norm is equivalent to the  $p$  norm and use transitivity of norms to prove the statement. First, consider  $p, q \in \mathbb{R}, p > q \geq 1$ . We can consider the following for any  $x \in \mathbb{R}^n$  (or  $\mathbb{C}^n$ ):

$$\|x\|_q^q = \sum_i^n |x_i|^q = \sum_{i=1}^n 1 \cdot |x_i|^q$$

By holder's inequality, we know that:

$$\sum_{i=1}^n 1 \cdot (x_i)^q \leq \left( \sum_{i=1}^n 1^a \right)^{\frac{1}{a}} \left( \sum_{i=1}^n (|x_i|^q)^b \right)^{\frac{1}{b}}$$

such that  $\frac{1}{a} + \frac{1}{b} = 1$ . Since the choice of  $a$  and  $b$  are arbitrary, we can let  $b$  a value such that  $b^{-1} = \frac{q}{p}$ . From here, we can perform the following algebra to see that:

$$a = \frac{b}{b-1} = \frac{p}{p-q}$$

and thus:

$$\begin{aligned} \|x\|_q^q &\leq \left( \sum_{i=1}^n 1^a \right)^{\frac{1}{a}} \left( \sum_{i=1}^n (|x_i|^q)^b \right)^{\frac{1}{b}} \\ \|x\|_q^q &\leq \left( \sum_{i=1}^n 1 \right)^{\frac{p-q}{p}} \left( \sum_{i=1}^n |x_i|^p \right)^{\frac{q}{p}} \\ \|x\|_q^q &\leq n^{\frac{p-q}{p}} \|x\|_p^q \\ \|x\|_q &\leq n^{\frac{p-q}{pq}} \|x\|_p \end{aligned}$$

Let  $q = 1$  and  $p > 1$ . we know aim to bound  $\|x\|_1$  from below. Let  $\{e_1, e_2, \dots, e_n\}$  denote the standard basis vectors of  $\mathbb{R}^n$ . Let  $x = (x_1, x_2, \dots, x_n)$ . and we can see that:

$$x = \sum_{i=1}^n e_i x_i$$

Thus, we can see that:

$$\|x\|_p = \left\| \sum_{i=1}^n x_i e_i \right\|_p$$

By the triangle inequality and the properties of the norm, we know that:

$$\left\| \sum_{i=1}^n x_i e_i \right\|_p \leq \sum_{i=1}^n \|x_i e_i\|_p = \sum_{i=1}^n |x_i| \cdot \|e_i\|_p$$

Let  $M = \max_{i \in 1, 2, \dots, n} \|e_i\|_p$ . Thus, it follows that:

$$\|x\|_p \leq \sum_{i=1}^n |x_i| \cdot \|e_i\|_p \leq M \sum_{i=1}^n |x_i| = M \|x\|_1$$

Thus, we can see that:

$$\frac{1}{M} \|x\|_p \leq \|x\|_1 \leq n^{\frac{p-1}{p}} \|x\|_p$$

Thus, we have proved that 1 norm and the  $p$  norm is equivalent. Thus, by transitive of norms, we can see that 1 and  $q$  are equivalent, which implies that  $p$  and  $q$  are equivalent norms.  $\square$

**Problem 4.**

*Solution:* Consider the following: We know that by the definition of the induced norm that:

$$\|Ax\|_a \leq \|A\|_{a \leftarrow c} \cdot \|x\|_c$$

as

$$\|A\|_{a \leftarrow c} := \sup \frac{\|Ax\|_a}{\|x\|_c}, \forall x \in \mathbb{R}^n$$

Let  $y = Bx$ , we see that:

$$\|Ay\|_a \leq \|A\|_{a \leftarrow c} \cdot \|Bx\|_c$$

But, since we know that:

$$\|B\|_{c \leftarrow b} := \sup \frac{\|Bx\|_c}{\|x\|_b}, \forall x \in \mathbb{R}^n$$

we can see that:

$$\|Bx\|_c \leq \|B\|_{c \leftarrow b} \cdot \|x\|_b$$

Thus, we can see that, if we were to combine these two inequalities, we get that:

$$\|Ay\|_a \leq \|A\|_{a \leftarrow c} \cdot \|B\|_{c \leftarrow b} \cdot \|x\|_b$$

We can see that

$$\begin{aligned} \|ABx\|_a &\leq \|A\|_{a \leftarrow c} \cdot \|B\|_{c \leftarrow b} \cdot \|x\|_b \\ \frac{\|ABx\|_a}{\|x\|_b} &\leq \|A\|_{a \leftarrow c} \cdot \|B\|_{c \leftarrow b} \end{aligned}$$

We can take the supremum of  $\frac{\|ABx\|_a}{\|x\|_b}$ , and we can see that:

$$\|AB\|_{a \leftarrow b} \leq \|A\|_{a \leftarrow c} \|B\|_{c \leftarrow b}$$

□

**Problem 5.**

*Solution:* Consider the subspace of  $V_1$  and  $V_2$  spanned by the basis  $\{\cos(n\pi x), \sin(n\pi x)\}$ . We can  $f \in \{\cos(n\pi x), \sin(n\pi x)\}$  is a linear combination, denoted as follows:

$$f = a_n \sin(n\pi x) + b_n \cos(n\pi x)$$

Thus,

$$\frac{d}{dx}f = n\pi(a_n \cos(n\pi x) - b_n \sin(n\pi x))$$

Thus, we can see that:

$$T_n f = \begin{bmatrix} 0 & -n\pi \\ n\pi & 0 \end{bmatrix} \begin{bmatrix} a_n \sin(n\pi x) \\ b_n \cos(n\pi x) \end{bmatrix} = \begin{bmatrix} a_n \cos(n\pi x) \\ -b_n \sin(n\pi x) \end{bmatrix}$$

Which implies that

$$T_n = \begin{bmatrix} 0 & n\pi \\ -n\pi & 0 \end{bmatrix}$$

Which implies that the  $T$  is:

$$T = \begin{bmatrix} T_1 & 0 & 0 & \dots \\ 0 & T_2 & 0 & \dots \\ \vdots & \vdots & \ddots & \dots \\ 0 & 0 & 0 & T_n \end{bmatrix}$$

where  $T_n$  is defined as above. Note that any function in  $V_1$  can be defined as follows:

$$f = \sum_{i=1}^k a_n \sin(n\pi x) + b_n \cos(n\pi x)$$

We now compute  $\langle f, f \rangle$ .

$$\begin{aligned} \langle f, f \rangle &= \int_0^1 \left( \sum_{i=1}^k a_n \sin(n\pi x) + b_n \cos(n\pi x) \right)^2 \\ &= \int_0^1 \sum_{n,m}^k a_n a_m \sin(n\pi x) \sin(m\pi x) + b_n b_m \cos(n\pi x) \cos(m\pi x) + 2a_n b_m \sin(n\pi x) \cos(m\pi x) \end{aligned}$$

Note that:

$$\begin{aligned} \int_0^1 \sin(n\pi x) \sin(m\pi x) &= \begin{cases} 0.5 & n = m \\ 0 & n \neq m \end{cases} \\ \int_0^1 \cos(n\pi x) \cos(m\pi x) &= \begin{cases} 0.5 & n = m \\ 0 & n \neq m \end{cases} \end{aligned}$$

$$\int_0^1 \sin(n\pi x) \cos(n\pi x) = 0, \forall n, m$$

Thus, this implies that:

$$\langle f, f \rangle = \frac{1}{2} \sum_{i=1}^k (a_i^2 + b_i^2)$$

By a similar logic, we can find that:

$$\langle Tf, Tf \rangle = \frac{1}{2} \sum_{n=1}^k (n\pi)^2 (a_n^2 + b_n^2)$$

Therefore, we can see that:

$$\left( \frac{\|Tf\|}{\|f\|} \right)^2 = \frac{\sum_{n=1}^k (n^2 \pi^2) (a_n^2 + b_n^2)}{\sum_{n=1}^k (a_n^2 + b_n^2)}$$

Note that  $\sup_f \left( \frac{\|Tf\|}{\|f\|} \right) = k\pi$ , where this maximum is obtained when  $a_n, b_n = 0$  when  $n \neq k$ , and  $a_k, b_k = 1$ , as this is the sum of all weighted averages, and the maximum is achieved when we put all the weight when  $n = k$ . This implies that  $\|T\| = k\pi$ . We first also wish to prove the existence of a  $T^{-1}$ . Note that the derivative is a bijective function, which implies that  $T^{-1}$  exists. In this case, we can see that  $T^{-1}$  can be defined as follows:

$$T_n^{-1} = \begin{bmatrix} 0 & \frac{-1}{n\pi} \\ \frac{1}{n\pi} & 0 \end{bmatrix}$$

where:

$$T^{-1} = \begin{bmatrix} T_1^{-1} & 0 & 0 & \dots \\ 0 & T_2^{-1} & 0 & \dots \\ \vdots & \vdots & \ddots & \dots \\ 0 & 0 & 0 & T_n^{-1} \end{bmatrix}$$

Let  $Tf = g$ , where  $g, f \in V_1$ . We can see that:

$$\sup_g \left( \frac{\|T^{-1}g\|}{\|g\|} \right)^2 = \sup_f \left( \frac{\|f\|}{\|Tf\|} \right)^2 = \sup_n \left( \frac{1}{(n\pi)^2} \right) = \frac{1}{\pi^2}$$

Thus, this implies that  $\|T^{-1}\| = \frac{1}{\pi}$ . Thus, we see that:

$$\max k_{rel} = \|T\| \|T^{-1}\| = k$$

Thus, this implies that  $k_{rel} = O(k)$ , as the bounded function as follows:




Figure 1: Showing that the relative condition is  $O(k)$

□



**Problem 6.**

*Solution:* Let  $a_1 = 2\sin(\pi x) + \sin(\pi x)$  and  $a_2 = -3\sin(\pi x) + \sin(2\pi x)$ . Let  $q_1 = \frac{a_1}{\|a_1\|}$ . Thus, we can proceed with the following computation:

$$\begin{aligned}\langle a_1, a_1 \rangle &= \langle 2\sin(\pi x) + \sin(\pi x), 2\sin(\pi x) + \sin(\pi x) \rangle \\ &= \int_0^1 (2\sin(2\pi x) + \sin(\pi x))^2 dx \\ &= \int_0^1 4\sin^2(2\pi x) + 4\sin(\pi x)\sin(2\pi x) + \sin^2(\pi x) dx\end{aligned}$$

Note that  $\int_0^1 \sin(\pi x)\sin(2\pi x) dx = 0$ . Thus, we can see that:

$$\begin{aligned}\langle a_1, a_1 \rangle &= \int_0^1 4\sin^2(2\pi x) + 4\sin(\pi x)\sin(2\pi x) + \sin^2(\pi x) dx \\ &= \int_0^1 4\sin^2(2\pi x) + \sin^2(\pi x) dx\end{aligned}$$

Note  $\int_0^1 \sin^2(n\pi x) = 0.5, \forall n \in \mathbb{N}$ . Thus, we can see that:

$$\langle a_1, a_1 \rangle = \int_0^1 4\sin^2(2\pi x) + \sin^2(\pi x) dx = 2 + 0.5 = \frac{5}{2}$$

This implies that:

$$q_1 = \sqrt{\frac{2}{5}}(2\sin(2\pi x) + \sin(\pi x))$$

How, we must consider solving the following  $v_2 = a_2 - \text{proj}_{a_2} q_1 = a_2 - \langle a_2, q_1 \rangle q_1$ . We now find the following integral;

$$\begin{aligned}\sqrt{\frac{2}{5}} \int_0^1 (2\sin(2\pi x) + \sin(\pi x))(\sin(2\pi x) - 3\sin(\pi x)) dx &= \sqrt{\frac{2}{5}} \int_0^1 2\sin^2(2\pi x) - 3\sin^2(\pi x) dx \\ &= \sqrt{\frac{2}{5}} * \frac{-1}{2} \\ &= -\sqrt{\frac{1}{10}}\end{aligned}$$

Thus, we see that  $v_2 = a_2 - \text{proj}_{a_2} q_1 = a_2 - \langle a_2, q_1 \rangle q_1 = a_2 + \left(\sqrt{\frac{1}{10}}\right) \left(\sqrt{\frac{2}{5}}\right) a_1 = a_2 + 0.2a_1$ . Simplifying the vectors, we can see that we can simplify to:

$$-3\sin(\pi x) + \sin(2\pi x) + 0.2(2\sin(2\pi x) + \sin(\pi x)) = \frac{-14}{5}\sin(\pi x) + \frac{7}{5}\sin(2\pi x)$$

We now proceed with the following calculation:

$$\begin{aligned}\int_0^1 \left( \frac{7}{5}(-2\sin(\pi x) + \sin(2\pi x)) \right)^2 dx &= \frac{49}{25} \int_0^1 4\sin^2(\pi x) + \sin^2(2\pi x) dx \\ &= \frac{49}{25} \left( 2 + \frac{1}{2} \right) \\ &= \frac{49}{10}\end{aligned}$$

We see that  $\|v_2\| = \frac{7}{\sqrt{10}}$ . Therefore, we can see that

$$q_1 = \sqrt{\frac{2}{5}}(2\sin(2\pi x) + \sin(\pi x)), q_2 = \frac{\sqrt{10}}{5}(\sin(2\pi x) - 2\sin(\pi x))$$

Note that  $r_{11} = \sqrt{\frac{5}{2}}$ ,  $r_{12} = \frac{\langle a_2, q_1 \rangle}{\|q_1\|} = -\frac{1}{\sqrt{10}}$ ,  $r_{22} = \frac{7}{\sqrt{10}} = \|v_2\|$ . Thus, this implies that:

$$R = \begin{bmatrix} \sqrt{\frac{5}{2}} & -\sqrt{\frac{1}{10}} \\ 0 & \frac{7}{\sqrt{10}} \end{bmatrix}$$

□