

PSET 3

Anthony Yoon

1/29/2025

1

a

We are interested in the following optimization problem:

$$\max \quad px_1^{\frac{1}{3}}x_2^{\frac{1}{3}} - \omega_1x_1 - \omega_2x_2$$

We see that the FOCs are

$$\begin{aligned} [x_1] \quad & \frac{1}{3}px_1^{-\frac{2}{3}}x_2^{\frac{1}{3}} - \omega_1 \leq 0 \quad \text{for } x_1 \geq 0 \\ [x_2] \quad & \frac{1}{3}px_1^{\frac{1}{3}}x_2^{-\frac{2}{3}} - \omega_2 \leq 0 \quad \text{for } x_2 \geq 0 \end{aligned}$$

We can see that $x_1, x_2 \neq 0$ as this would cause the FOCs to become undefined. From here, divide the FOCs to get the relation $\omega_1x_1 = \omega_2x_2$. Using this expression, we can substitute this into the FOCs to get that

$$x_1^* = \frac{p^3}{27\omega_1^2\omega_2} \quad x_2^* = \frac{p^3}{27\omega_1\omega_2^2}$$

Therefore, we see that:

$$y^* = \left(\frac{p^6}{3^6\omega_1^3\omega_2^3} \right)^{\frac{1}{3}} = \frac{p^2}{9\omega_1\omega_2}$$

Thus, we see that

$$PF = py^* - \omega_1x_1^* - \omega_2x_2^* = p \left(\frac{p^2}{9\omega_1\omega_2} \right) - \omega_1 \frac{p^3}{27\omega_1^2\omega_2} - \omega_2 \frac{p^3}{27\omega_1\omega_2^2} = \frac{p^3}{27\omega_1\omega_2}$$

b

We can see that for the IDFs

$$\frac{\partial x_1^*}{\partial \omega_1} = -2 \left(\frac{p^3}{27\omega_2\omega_1^3} \right)$$

and

$$\frac{\partial x_2^*}{\partial \omega_2} = -2 \left(\frac{p^3}{27\omega_1\omega_2^3} \right)$$

Note that both quantities are bounded above by 0, as p, ω are strictly positive. For the ODF, we see that

$$\frac{\partial y^*}{\partial p} = \frac{2p}{9\omega_1\omega_2}$$

which is always positive for the same reasons. For the PF, note that

$$\frac{\partial \pi(\omega, y)}{\partial p} = \frac{p^2}{9\omega_1\omega_2} > 0 \quad \frac{\partial \pi(\omega, p)}{\partial \omega_1} = \frac{-p^3}{27\omega_1^2\omega_2} < 0 \quad \frac{\partial \pi(\omega, p)}{\partial \omega_2} = \frac{-p^3}{27\omega_1\omega_2^2} < 0$$

c

Proof that IDF is homogenous in degree 0, let $t > 0$, we see that

$$x_1^*(t\omega, tp) = \frac{(tp)^3}{27(t\omega_1)^2 t\omega_2} = \frac{t^3 p^3}{27t^3 \omega_1^2 \omega_2} = \frac{p^3}{27\omega_1^2 \omega_2} = x_1^*(\omega, p)$$

and similarly

$$x_2^*(t\omega, tp) = \frac{(tp)^3}{27t\omega_1 (t\omega_2)^2} = \frac{t^3 p^3}{27t^3 \omega_1 \omega_2^2} = \frac{p^3}{27\omega_1 \omega_2^2} = x_2^*(\omega, p)$$

Proof that OSF is homogenous in degree 0, let $t > 0$, we see that

$$y^*(t\omega, tp) = \frac{t^2 p^2}{9t^2 \omega_1 \omega_2} = \frac{p^2}{9\omega_1 \omega_2} = y^*(\omega, p)$$

Proof that PF is homogenous in degree 1, let $t > 0$, we see that

$$\pi(t\omega, pt) = \frac{t^3 p^3}{27t^2 \omega_1 \omega_2} = \frac{tp^3}{27\omega_1 \omega_2} = t\pi(\omega, p)$$

d