



# Analysis of D meson decays at LHCb

Anthony Lestone  
P-3, High Energy Nuclear Physics

8/3/2021

**LA-UR-21-27161**

# Research Objective

To study the quark-level decays of the  $D^+$  and  $D_s^+$  mesons to  
search for new particles


# Quarks: the building blocks of hadronic matter

- Quarks are not allowed to exist freely, so they are always confined within hadrons.
- Most commonly, quarks form clusters of three, called baryons, or clusters of two, called mesons.
- For every quark, there is a corresponding antiquark with opposite charge.
  - Antiquarks are denoted with a bar.

$+\frac{2}{3}$	u	c	t
$-\frac{1}{3}$	d	s	b


mass  $\longrightarrow$  mass

The proton




$$u^{+\frac{2}{3}} + u^{+\frac{2}{3}} + d^{-\frac{1}{3}} = \text{net charge } +1$$

The neutron



$$d^{-\frac{1}{3}} + d^{-\frac{1}{3}} + u^{+\frac{2}{3}} = \text{net charge } 0$$


The pion ( $\pi^+$ )



$$u^{+\frac{2}{3}} + \bar{d}^{+\frac{1}{3}} = \text{net charge } +1$$


# Choosing a dataset: on the hunt for D mesons

- This project looks to compare the  $D^+$  and  $D_s^+$  mesons, so we want a data set that contains both of these particles.
- $D^+$  and  $D_s^+$  both decay far too quickly to be seen directly, so we look for their decay products.
- Both are able to decay into two kaons and a pion.




# The Large Hadron Collider

- 16.8 mile long loop located 300 feet underground at Geneva, Switzerland.
- The most powerful particle accelerator in the world
- Accelerates two proton beams in opposite directions
  - Proton beams are kept separate, experiments are housed at beam crossing locations



# The LHCb Experiment


- Experiment located at the LHC
- As high energy protons collide, their energy is converted into new particles via  $E = mc^2$
- Able to determine energy and momentum of detected particles, as well as particle identity




# Placing selection criteria on the data

- If we took every single  $K^+ K^- \pi^+$  event, we would be left with random, uncorrelated noise.
- We place criteria such as:
  - All 3 particles came from the same location
  - Particle ID criteria from Cherenkov counters
  - Their energies and momenta add up to a mass close to that of the target particle

The particles appear to have come from the same event





The particles *do not* appear to have come from the same event



# The Data


- The mass of the primary particle is recreated from the energies and momenta of the  $K^+$ ,  $K^-$ , and  $\pi^+$ . This graph is called an invariant mass spectrum.
- Black dots are data points, the blue curve is the model
  - The data is modeled by the sum of two Gaussian curves and a linear background






# The Data

- The mass of the primary particle is recreated from the energies and momenta of the  $K^+$ ,  $K^-$ , and  $\pi^+$ . This graph is called an invariant mass spectrum.
- Black dots are data points, the blue curve is the model
  - The data is modeled by the sum of two Gaussian curves and a linear background




# What about intermediate decay states?

- Do the  $D^+$  and  $D_s^+$  really decay directly to  $K^+ K^- \pi^+$ , or is there some sort of “stepping stone” in between?




# What about intermediate decay states?

- Do the  $D^+$  and  $D_s^+$  really decay directly to  $K^+ K^- \pi^+$ , or is there some sort of “stepping stone” in between?




# Two-body invariant mass spectra

- We can create invariant mass spectra of  $K^+ K^-$  and  $K^- \pi^+$  to search for intermediate states.



# Two-body invariant mass spectra



- We can create invariant mass spectra of  $K^+ K^-$  and  $K^- \pi^+$  to search for intermediate states.
- Let's have a closer look at these leftmost peaks...





# Leftmost peaks

- Similar resonances appear in both the  $D^+$  and  $D_s^+$  spectra.
- We can search the literature for particles that satisfy the mass and decay criteria:
  - 1020 MeV and decays to  $K^+ K^-$
  - 896 MeV and decays to  $K^- \pi^+$

$D^+$







$D_s^+$



# The culprits


- We find two particles that fit these descriptions exactly.
- The phi meson is appearing in the  $K^+ K^-$  spectrum while the  $K^*(892)$  is appearing in the  $K^- \pi^+$  spectrum.



## But what about those other peaks?

- We still have many peaks yet unidentified!  
Let's have a look at the middle peak.
- There are no known mesons with mass  $\sim 1120$  MeV which decay to  $K^- \pi^+$ .
  - Clearly I've discovered a new particle!  
Where's my Nobel prize?
  - Or maybe not...



$K^-$  and  $\pi^+$  Invariant Mass for  $D^+$






# This is where Dalitz plot analysis comes in handy

- Named after Richard Dalitz
- Widely used in the study of three-body decays of D mesons, like in our case of  $K^+ K^- \pi^+$ .




# Resonances appear on a Dalitz plot as a band

- The  $\phi$  appears as a stripe at 1020 MeV on the  $K^+ K^-$  axis, while the  $K^*$  appears at 896 MeV on the  $K^- \pi^+$  axis.
- The keen-eyed observer might be able to see where this is going...





# Resonances can manifest themselves on both axes

- The unidentified peaks are not new particles, but rather the  $\phi$  and  $K^*$  creating signals on both axes.
- Particles are correlated due to the decay process, and the Dalitz plot offers a method to study this correlation




# Conclusions


- The decays of the  $D^+$  and  $D_s^+$  mesons to  $K^+ K^- \pi^+$  proceed via two distinct pathways: one with a  $\psi$  meson and one with a  $K^*(892)$  meson.



## The $\phi \pi^+$ pathway





## The $K^* K^+$ pathway



# Future steps

- Several other analyses quote the presence of several additional particle resonances so the inclusion of these resonances in my model might improve the fit to data.
- The asymmetric phi lobes in the  $K^- \pi^+$  spectrum for the  $D^+$  implies a decay pathway exclusive to the  $D^+$ . Further study could reveal the particle that mediates this pathway.



**Thank you for your time! Any questions?**

# Backup Slides

# The starring cast

## Hadrons

$D^+$



Mass: 1869.7 MeV  
Charge: +1  
Spin: 0

$D_s^+$



Mass: 1968.4 MeV  
Charge: +1  
Spin: 0

$\pi^+$



Mass: 139.6 MeV  
Charge: +1  
Spin: 0

$K^+$



Mass: 493.7 MeV  
Charge: +1  
Spin: 0

$K^-$



Mass: 493.7 MeV  
Charge: -1  
Spin: 0

$K^*(892)$



Mass: 895.6 MeV  
Charge: 0  
Spin: 1

$\psi$



Mass: 1019.5 MeV  
Charge: 0  
Spin: 1

## Quarks

up



Mass: ~2.2 MeV  
Charge:  $+\frac{2}{3}$   
Spin:  $\pm\frac{1}{2}$

charm



Mass: ~1270 MeV  
Charge:  $+\frac{2}{3}$   
Spin:  $\pm\frac{1}{2}$

down



Mass: ~4.7 MeV  
Charge:  $-\frac{1}{3}$   
Spin:  $\pm\frac{1}{2}$

strange





Mass: ~93 MeV  
Charge:  $-\frac{1}{3}$   
Spin:  $\pm\frac{1}{2}$





# Leftmost peaks

- Similar resonances appear in both the  $D^+$  and  $D_s^+$  spectra.

$D^+$



$D_s^+$



# Analysis Tools

- ROOT
  - Built on C++
  - A powerful tool used to analyze large datasets
  - Created by CERN, tailored specifically for particle physics applications
    - All histograms in this presentation were created using ROOT



# Welcome to the LANL slide template!

## Please take a moment to read this information.

- How to use the Lab's PowerPoint templates: The LANL PowerPoint templates have been developed to help users easily create professional slide presentations that meet the Lab's new brand standards, effective January 2021. This template is for a widescreen display presentation. Please use the default font (Arial) and default font sizes as much as possible.
- Classification: Unclassified LANL presentations do NOT require marking.
- LA-UR #: All LANL presentations intended for external use must have an LA-UR number. For more information on the LA-UR process, visit <http://int.lanl.gov/library/help/rassti/index.shtml>. A placeholder text box is located in the bottom left corner of the title slide for an LA-UR number if necessary.
- **Date:** The date is set for your presentation in the footer of each slide. It changes to the current date automatically. If you wish to enter a different date, you can do so by entering the slide master and changing it. Here's how:
  - Open the slide master by clicking the View tab and selecting Slide Master.
  - In the left navigation pane, scroll to the top and find the first slide layout. (It's larger than the others and has a "1" next to the top left corner.) Click on this slide.
  - Click in the date text box in the lower right hand corner and change the date.
  - This will change the date on the most of the slides, **but not all of them.**
  - **Scroll to the first blue slide, Title Slide, and manually change the date.**
  - In the left navigation pane, scroll to the second master slide, named Custom, with the blue background (It's larger than the others and has a "2" next to the top left corner.) Click on this slide.
  - Click in the date text box and change the date, scroll through each master template to make sure all the dates have been revised on the slides you are using.
  - Click Close Master View.