## Discrete Mathematics Set 4

Math 435: Complete 7 parts of the following exercises.

Math 530: Exercises 1, 2, and any two of the remaining exercises.

- **1.** Let  $A_n$  be the set of paths in  $\mathbb{R}^2$  which start at (0,0), end at (n,n), and only use steps of the form (1,0) or (0,1).
  - a. Show that there are  $\binom{2n}{n}$  elements in  $A_n$ .
  - b. Denote the number of times the path  $p \in A_n$  touches the line y = x by touch(p). Let

$$A(x,t) = \sum_{n=0}^{\infty} \left( \sum_{p \in A_n} t^{\mathsf{touch}(p)} \right) x^n$$

and let  $c_n$  be the  $n^{th}$  Catalan number. Show that

$$\sum_{p \in A_{n+1}} t^{\mathsf{touch}(p)} = 2t \sum_{k=0}^n c_k \left( \sum_{p \in A_{n-k}} t^{\mathsf{touch}(p)} \right).$$

- c. Show that  $A(x,t) = \frac{t}{1 t + 2t\sqrt{\frac{1}{4} x}}$ .
- d. Find an asymptotic formula for the average number of times a path in  $C_n$  touches the line y = x.
- **2.** Let  $M(x) = \sum_{n=0}^{\infty} m_n x^n$  be the generating function for  $m_n$ , the number of Motzkin paths of length n, as defined in Set 1 Exercise 6.
  - a. Find  $\lim_{x\to 1/3} (1/3-x)^{1/2} x M'(x)$  and use this result to find an asymptotic formula for  $m_n$ . (As always, calculations can be done using a computer algebra system if desired.)
  - b. Let  $a_n$  be as defined in Set 1 Exercise 6b. Find an asymptotic formula for  $a_n$  and use it to find an asymptotic formula for the probability that a random path in the plane that from (0,0) to (n,n) using steps of the form (1,1),(1,-1) and (1,0) is a Motzkin path.
- **3.** Find an asymptotic formula for the average number of sets in an ordered set partition of n (see Set 2 Exercise 3, Set 3 Exercise 1b, and Set 3 Exercise 1d).
- **4.** Find an asymptotic formula for the average number of cycles in a permutation of n with ordered cycles (see Set 3 Exercise 1c and Set 3 Exercise 1d.)
- **5.** Find an asymptotic formula for the number of permutations of n that do not have any cycles of length 1, 2 or 3.
- 6. Define
  - 1. the Chebyshev polynomial of the first kind  $T_n(y)$  by  $\sum_{n=0}^{\infty} T_n(y) x^n = \frac{1-yx}{1-2yx+x^2}$ ,
  - 2. the Chebyshev polynomial of the second kind  $U_n(y)$  by  $\sum_{n=0}^{\infty} U_n(y) x^n = \frac{1}{1-2yx+x^2}$ , and

3. the Legendre polynomial 
$$P_n(y)$$
 by  $\sum_{n=0}^{\infty} P_n(y) x^n = \frac{1}{\sqrt{1-2yx+x^2}}$ .

Find asymptotic formulas for  $T_n(5/4)$ ,  $U_n(5/4)$ , and  $P_n(5/4)$ .