
# Mathematics 2263 Exercises


**1.** Match each of the following functions with the corresponding contour plot and plot in $\mathbb{R}^3$:


$$a(x,y) = \cos x \cos y \, e^{-\sqrt{x^2+y^2}/4} \qquad b(x,y) = -\frac{xy^2}{x^2+y^2} \qquad c(x,y) = \frac{1}{4x^2+y^2+1}$$


$$d(x,y) = e^{-y/10}\cos x \qquad e(x,y) = xy\frac{x^2-y^2}{x^2+y^2} \qquad f(x,y) = y^4 - x^4$$


Contour plots:


1. 

2. 


3. 


4. 


5. 


6. 


Plots in $\mathbb{R}^3$:


i. 

ii. 

iii. 

iv. 

v. 

vi. 

**2.** Sketch the level curves $f(x,y) = c$ on the same set of axes for the given values of $c$:

a. $f(x,y) = x + y - 1, c = -3, -2, -1, 0, 1, 2, 3$

b. $f(x,y) = x^2 + y^2, c = 0, 1, 4, 9, 16, 25$

c. $f(x,y) = xy, c = -9, -4, -1, 0, 1, 4, 9$

d. $f(x,y) = \sqrt{25 - x^2 - y^2}, c = 0, 1, 2, 3, 4$

**3.** Sketch sample level curves and sketch the following functions in $\mathbb{R}^3$:

a. $f(x,y) = y^2$

b. $f(x,y) = x^2 + y^2$

c. $f(x,y) = 4 - x^2 - y^2$

d. $f(x,y) = 4 - |x| - |y|$

**4.** Find these limits or explain why they do not exist:

a. $\lim\limits_{(x,y)\to(0,0)} \dfrac{3x^2 - y^2 + 5}{x^2 + y^2 + 2}$

b. $\lim\limits_{(x,y)\to(1,\pi/6)} \dfrac{x \sin y}{x^2 + 1}$

c. $\lim\limits_{(x,y)\to(1,1), x\neq y} \dfrac{x^2 - 2xy + y^2}{x - y}$

d. $\lim\limits_{\substack{(x,y)\to(0,0)\\ x\neq y, x\geq 0, y\geq 0}} \dfrac{x - y + 2\sqrt{x} - 2\sqrt{y}}{\sqrt{x} - \sqrt{y}}$

e. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x^4}{x^4 + y^2}$

f. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x}{\sqrt{x^2 + y^2}}$

g. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x^2 y}{\sqrt{x^4 + y^2}}$

h. $\lim\limits_{\substack{(x,y)\to(1,1)\\ x\neq y}} \dfrac{x^2 - y^2}{x - y}$

i. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x^3 - xy^2}{x^2 + y^2}$

j. $\lim\limits_{(x,y)\to(0,0)} \dfrac{\sin(x^2 + y^2)}{x^2 + y^2}$

k. $\lim\limits_{(x,y)\to(0,0)} \dfrac{x^2 - y^2}{x^2 + y^2}$

**5.** At what points $(x,y)$ in $\mathbb{R}^2$ are the following functions continuous?

a. $\sin(x + y)$

b. $\ln(x^2 + y^2)$

c. $\dfrac{x^2 + y^2}{x^2 - 3x + 2}$

d. $\dfrac{1}{x^2 - y}$

**6.** Find $\partial f/\partial x$, find $\partial f/\partial y$, and (in the cases where $f$ is a function of $z$) find $\partial f/\partial z$:

a. $f(x,y) = \ln(x + y)$

b. $f(x,y) = e^{xy} \ln y$

c. $f(x,y) = \displaystyle\int_x^y g(t)\, dt$ where $g$ is continuous

d. $f(x,y,z) = xy + xz + yz$

e. $f(x,y,z) = (x^2 + y^2 + z^2)^{-1/2}$

f. $f(x,y,z) = \ln(x + 2y + 3z)$

**7.** Find all second order partial derivatives:

a. $f(x,y) = x + y + xy$

b. $f(x,y) = \sin(xy)$

c. $f(x,y) = xy^2 + x^2y^3 + x^3y^4$

**8.** Find the value of $\partial z/\partial x$ at the point $(1,1,1)$ if the equation

$$xy + z^3x - 2yz = 0$$

defines $z$ as a function of the independent variables $x$ and $y$.

**9.** Show that each of the following functions satisfy the equation $\dfrac{\partial^2 f}{\partial x^2} + \dfrac{\partial^2 f}{\partial y^2} + \dfrac{\partial^2 f}{\partial z^2} = 0$:

a. $f(x,y,z) = x^2 + y^2 - 2z^2$

b. $f(x,y,z) = \ln\sqrt{x^2 + y^2}$

c. $f(x,y,z) = \arctan(x/y)$

**10.** Show that each of the following functions satisfy the equation $\dfrac{\partial^2 f}{\partial t^2} = c^2 \dfrac{\partial^2 f}{\partial x^2}$:

a. $f(x,t) = \sin(x + ct)$

b. $f(x,t) = \tan(2x - 2ct)$

**11.** Assume that $w = f(s^3 + t^2)$ and $f'(x) = e^x$. Find $\partial w/\partial t$ and $\partial w/\partial s$.

**12.** Assume that $w = f(ts^2, s/t)$, $\partial f(x,y)/\partial x = xy$, and $\partial f(x,y)/\partial y = x^2/2$. Find $\partial w/\partial t$ and $\partial w/\partial s$.

**13.** Suppose that $f$ is a function of $u$, $v$, and $w$ where $u = x - y$, $v = y - z$, and $w = z - x$. Show that

$$\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} + \frac{\partial f}{\partial z} = 0.$$

**14.** Suppose that we substitute the polar equations $x = r\cos\vartheta$ $y = r\sin\vartheta$ into $w = f(x,y)$.

a. Show that

$$\frac{\partial w}{\partial r} = f_x \cos\vartheta + f_y \sin\vartheta$$

and

$$\frac{1}{r}\frac{\partial w}{\partial \vartheta} = -f_x \sin\vartheta + f_y \cos\vartheta.$$

b. Solve the equations in part a. to express $f_x$ and $f_y$ in terms of $\partial w/\partial r$ and $\partial w/\partial \vartheta$.

c. Show that

$$(f_x)^2 + (f_y)^2 = \left(\frac{\partial w}{\partial r}\right)^2 + \frac{1}{r^2}\left(\frac{\partial w}{\partial \vartheta}\right)^2.$$

**15.** Show that if $w = f(u,v)$ satisfies

$$f_{uu} + f_{vv} = 0$$

and if $u = (x^2 - y^2)/2$ and $v = xy$, then $w$ satisfies

$$w_{xx} + w_{yy} = 0.$$

**16.** Find the gradient of the function at the given point. Sketch the gradient and the level curve that passes through the point:

a. $f(x,y) = y - x$ at $(2,1)$

b. $f(x,y) = \ln(x^2 + y^2)$ at $(1,1)$

c. $f(x,y) = x^2/2 - y^2/2$ at $(\sqrt{2},1)$

**17.** Find the derivative of the function at $P$ in the direction of $\mathbf{u}$:

a. $f(x,y) = 2x^2 + y^2$ with $P = (-1,1)$ and $\mathbf{u} = \langle 3,-4\rangle$

b. $f(x,y) = \dfrac{x-y}{xy+2}$ with $P = (1,-1)$ and $\mathbf{u} = \langle 12,5\rangle$

c. $f(x,y,z) = x^2 + 2y^2 - 3z^2$ with $P = (1,1,1)$ and $\mathbf{u} = \langle 1,1,1\rangle$

**18.** Find the directions in which the functions increase and decrease most rapidly at $P$ Then find the derivatives of the functions in these directions.

a. $f(x,y) = x^2y + e^{xy}\sin y$ with $P = (1,0)$

b. $f(x,y,z) = xe^y + z^2$ with $P = (1,\ln 2, 1/2)$

c. $f(x,y,z) = \ln(xy) + \ln(yz) + \ln(xy)$ with $P = (1,1,1)$

**19.** Is there a direction $\mathbf{u}$ in which the rate of change of $f(x,y) = x^2 - 3xy + 4y^2$ at $(1,2)$ equals 14? Why?

**20.** Find the tangent plane and the normal line for the surface at the point $P$:

a. $x^2 + y^2 + z^2 = 3$ at $P = (1,1,1)$

b. $x^2 + y^2 - z^2 = 18$ at $P = (3,5,-4)$

c. $2z - x^2 = 0$ at $P = (2,0,2)$

d. $x^2 + y^2 - 2xy - x + 3y - z = -4$ at $P = (2,-3,18)$

e. $z = \ln(x^2 + y^2)$ at $P = (1,0,0)$

f. $z = 4x^2 + y^2$ at $P = (1,1,5)$

**21.** Find all local maxima, local minima, and saddle points for these functions

a. $f(x,y) = 2xy - 5x^2 - 2y^2 + 4x + 4y - 4$

b. $f(x,y) = 5xy - 7x^2 + 3x - 6y + 2$

c. $f(x,y) = 2x^2 + 3xy + 4y^2 - 5x + 2y$

d. $f(x,y) = x^3 + 3xy + y^3$

e. $f(x,y) = 4xy - x^4 - y^4$

f. $f(x,y) = \ln(x+y) + x^2 - y$

**22.** Find the absolute maxima and absolute minima of the function on the given domain:

a. $f(x,y) = 2x^2 - 4x + y^2 - 4y + 1$ on the triangular region bounded by the lines $x = 0$, $y = 2$, $y = 2x$ in the first quadrant

b. $f(x,y) = x^2 + xy + y^2 - 6x$ on $[0,5] \times [-3,3]$

c. $f(x,y) = (4x - x^2)\cos y$ on $[1,3] \times [-\frac{\pi}{4}, \frac{\pi}{4}]$

**23.** Find two numbers $a$ and $b$ with $a \leq b$ that maximizes

$$\int_a^b (6 - x - x^2)\, dx.$$

**24.** Find the maximum value of $xy + yz + xz$ where $x + y + z = 6$.

**25.** Find the minimum distance from $z = \sqrt{x^2 + y^2}$ to the point $(-6, 4, 0)$.

**26.** Find the points on the ellipse $x^2 + 2y^2 = 1$ where $f(x,y) = xy$ has its extreme values.

**27.** Find the extreme values of $f(x,y) = xy$ subject to the constraint $g(x,y) = x^2 + y^2 - 10 = 0$.

**28.** Find the points on $x^2 y = 2$ closest to the origin.

**29.** Find the points on the curve $x^2 + xy + y^2 = 1$ in the $xy$-plane that are nearest and farthest from the origin.

**30.** Find the dimensions of the rectangle of largest perimeter that can be inscribed in the ellipse $x^2/a^2 + y^2/b^2 = 1$ with sides parallel to the coordinate axes. What is this perimeter?

**31.** Find the maximum and minimum values of $3x - y + 6$ subject to the constraint $x^2 + y^2 = 4$.

**32.** Maximize and minimize $xyz^2$ on $x^2 + y^2 + z^2 = 1$.

**33.** Evaluate the following integrals:

a. $\displaystyle\int_0^2 \int_{-1}^1 (x - y)\, dx\, dy$

b. $\displaystyle\int_0^1 \int_0^1 \left(1 - \frac{x^2 + y^2}{2}\right) dx\, dy$

c. $\displaystyle\int_0^1 \int_1^2 xye^x\, dx\, dy$

d. $\displaystyle\iint_R xy\cos y\, dA$ where $R = [-1,1] \times [0,\pi]$

e. $\displaystyle\iint_R y\sin(x + y)\, dA$ where $R = [-\pi, 0] \times [0,\pi]$

f. $\displaystyle\iint_R \frac{y}{x^2y^2 + 1}\, dA$ where $R = [0,1] \times [0,1]$

g. $\displaystyle\iint_R \frac{1}{xy}\, dA$ where $R = [1,2] \times [1,2]$

**34.** Find the volume of the region bounded above by the paraboloid $z = x^2 + y^2$ and below $[-1,1] \times [-1,1]$.

**35.** Sketch the described regions of integration in $\mathbb{R}^2$:

a. $-2 \leq y \leq 2, y^2 \leq x \leq 4$

b. $0 \leq y \leq 1, y \leq x \leq 2y$

c. $1 \leq x \leq e^2, 0 \leq y \leq \ln x$.

**36.** Write an iterated integral for $\iint_R dA$ over the region $R$ using both vertical cross sections and horizontal cross sections:

a. The triangle in the first quadrant of $\mathbb{R}^2$ bounded by the graphs of $x = 3$ and $y = 2x$.

b. The region in the first quadrant of $\mathbb{R}^2$ bounded by the lines $x = 2$, $y = 1$, and the graph of the function $y = e^x$.

c. The region bounded by $y = 3 - 2x$, $y = x$, and $x = 0$.

d. The region bounded by $y = x^2$ and $y = x + 2$.

**37.** Sketch the region and then evaluate the integral:

a. $\displaystyle\int_0^\pi \int_0^x x\sin y\, dy\, dx$

b. $\displaystyle\int_1^2 \int_y^{y^2} dx\, dy$

c. $\displaystyle\int_0^1 \int_0^{\sqrt{1-x^2}} 8y\, dy\, dx$

**38.** Reverse the order of integration and then evaluate the integral:

a. $\displaystyle\int_0^\pi \int_x^\pi \frac{\sin y}{y}\, dy\, dx$

b. $\displaystyle\int_0^1 \int_y^1 x^2 e^{xy}\, dx\, dy$

c. $\displaystyle\int_0^{2\sqrt{\ln 3}} \int_{y/2}^{\sqrt{\ln 3}} e^{x^2}\, dx\, dy$

d. $\displaystyle\iint_R (y - 2x^2)\, dA$ where $R$ is the region bounded by $|x| + |y| = 1$ in $\mathbb{R}^2$.

**39.** Find the volume of the solid in the first octant bounded by the coordinate planes, the plane $x = 3$, and the parabolic cylinder $z = 4 - y^2$.

**40.** Find the area of the following regions using a double integral:

a. The coordinate axes and the line $x + y = 2$.

b. The parabola $x = -y^2$ and $y = 4$.

c. The curve $y = e^x$ and the lines $y = 0$, $x = 0$, and $x = \ln 2$.

d. The lines $y = 2x$, $y = x/2$, and $y = 3 - x$.

**41.** Change from Cartesian into polar and evaluate:

a. $\displaystyle\int_{-1}^{1}\int_{0}^{\sqrt{1-x^2}} dy\,dx$

b. $\displaystyle\int_{0}^{1}\int_{0}^{\sqrt{1-y^2}} (x^2+y^2)\,dx\,dy$

c. $\displaystyle\int_{-a}^{a}\int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} dy\,dx$

d. $\displaystyle\int_{\sqrt{2}}^{2}\int_{0}^{x} y\,dy\,dx$

e. $\displaystyle\int_{-1}^{1}\int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \frac{2}{(1+x^2+y^2)^2}\,dy\,dx$

f. $\displaystyle\int_{0}^{1}\int_{x}^{\sqrt{2-x^2}} (x+2y)\,dy\,dx$

g. $\displaystyle\int_{0}^{\infty}\int_{0}^{\infty} \frac{1}{(1+x^2+y^2)^2}\,dy\,dx$

**42.** The average value of a function $f(x,y)$ over the region $R$ in $\mathbb{R}^2$ is given by

$$\frac{1}{\text{area}(R)}\iint_R f(x,y)\,dx\,dy.$$

Find the average values of $f(x,y) = \sqrt{a^2-x^2-y^2}$ over the region described by $x^2+y^2 \le a^2$.

**43.** Find the average distance from a point on the disk $x^2+y^2 \le a^2$ to the origin.

**44.** Let $I = \int_0^\infty e^{-x^2}\,dx$. Evaluate $I^2$ by noticing that

$$\begin{aligned} I^2 &= \left(\int_0^\infty e^{-x^2}\,dx\right)\left(\int_0^\infty e^{-y^2}\,dy\right) \\ &= \int_0^\infty\int_0^\infty e^{-(x^2+y^2)}\,dx\,dy \end{aligned}$$


and switching into polar.

**45.** Write six different iterated triple integrals for the volume of the following regions in $\mathbb{R}^3$:

a. The first octant enclosed by the cylinder $x^2+z^2 = 4$ and the plane $y = 3$

b. The region bounded by $z = 8 - x^2 - y^2$ and $z = x^2 + y^2$.

c. The region bounded in the first octant that satisfies $z + y \le 1$ and $x^2 \le y$:




**46.** Evaluate the integrals:

a. $\displaystyle\int_0^1\int_0^1\int_0^1 (x^2+y^2+z^2)\,dz\,dy\,dx$

b. $\displaystyle\int_0^e\int_0^{e^2}\int_0^{e^3} \frac{1}{xyz}\,dx\,dy\,dz$

c. $\displaystyle\int_0^1\int_0^{3-3x}\int_{x^2+3y^2}^{8-x^2-y^2} dz\,dy\,dx$


d. $\displaystyle\int_0^3\int_0^{\sqrt{9-x^2}}\int_0^{\sqrt{9-x^2}} dz\,dy\,dx$

e. $\displaystyle\int_0^1\int_0^{2-x}\int_0^{2-x-y} dz\,dy\,dx$

f. $\displaystyle\int_0^\pi\int_0^\pi\int_0^\pi \cos(x+y+z)\,dz\,dy\,dx$


**47.** Find the volumes of the following regions in $\mathbb{R}^3$:

a. The region in the first octant bounded by the coordinate planes, the plane $y + z = 2$, and the cylinder $x = 4 - y^2$:




b. The tetrahedron in the first octant bounded by the coordinate planes and the plane passing through $(1,0,0)$, $(0,2,0)$ and $(0,0,3)$.

c. The region in the first octant bounded the by the coordinate planes, the plane $y = 1 - x$ and the surface $z = \cos(\pi x/2)$ for $0 \leq x \leq 1$:



d. The region cut from the cylinder $x^2 + y^2 = 4$ by the plane $z = 0$ and the plane $x + z = 3$:




**48.** Find the center of mass of a thin plate of uniform density bounded by the lines $x = 0$, $y = x$, and the parabola $y = 2 - x^2$ in the first quadrant.

**49.** Find the center of mass of a thin plate of nonuniform density in the shape of a triangle bounded by the $y$-axis and the lines $y = x$ and $y = 2 - x$ if the density at the point $(x, y)$ is $6x + 3y + 3$.

**50.** Find the mass of the solid and find the center of mass for the following regions in $\mathbb{R}^3$:

a. The solid region in the first octant bounded by the coordinate planes and the plane $x + y + z = 2$ where the solid has density given by $\delta(x, y, z) = 2x$.

b. The solid region in the first octant bounded by the planes $y = 0$ and $z = 0$ and the surfaces $z = 4 - x^2 4$ and $x = y^2$ where the solid has density given by $\delta(x, y, z) = xy$:



c. A solid cube in the first octant bounded by the planes $x = 1, y = 1$, and $z = 1$ where the density of the cube at $(x, y, z)$ is given by $x + y + z + 1$.

**51.** Evaluate the integrals in cylindrical coordinates:

a. $\displaystyle\int_0^{2\pi}\int_0^{1}\int_r^{\sqrt{2-r^2}} dz r\, dr\, d\vartheta$

b. $\displaystyle\int_0^{2\pi}\int_0^{\vartheta/(2\pi)}\int_0^{3+24r^2} dz r\, dr\, d\vartheta$


c. $\displaystyle\int_0^{2\pi}\int_0^{3}\int_0^{z/3} r^3\, dr\, dz\, d\vartheta$

**52.** Convert

$$\int_{-1}^{1}\int_{0}^{\sqrt{1-y^2}}\int_{0}^{x}(x^2+y^2)\, dz\, dx\, dy$$

into cylindrical coordinates and evaluate.

**53.** Let $R$ be the region bounded below by the cone $z = \sqrt{x^2 + y^2}$ and above by the paraboloid $z = 2 - x^2 - y^2$:




Set up the triple integrals that give the volume of $R$ in these three orders: $dz\,dr\,d\vartheta$, $dr\,dz\,d\vartheta$ and $d\vartheta\,dz\,dr$.

**54.** Evaluate the integrals in spherical coordinates (some orders of integration may be easier than others):


a. $\displaystyle\int_0^{\pi}\int_0^{\pi}\int_0^{2\sin\varphi} \rho^2 \sin\varphi\, d\rho\, d\varphi\, d\vartheta$

b. $\displaystyle\int_0^{2\pi}\int_0^{\pi/4}\int_0^{2} \rho^3 \cos\varphi \sin\varphi\, d\rho\, d\varphi\, d\vartheta$

c. $\displaystyle\int_0^{2}\int_{-\pi}^{0}\int_{\pi/4}^{\pi/2} \rho^3 \sin(2\varphi)\, d\varphi\, d\vartheta\, d\rho$

d. $\displaystyle\int_{\pi/6}^{\pi/2}\int_{-\pi/2}^{\pi/2}\int_{\csc\varphi}^{2} 5\rho^4 \sin^3\varphi\, d\rho\, d\vartheta\, d\varphi$

**55.** Find the volumes of the region above $z = (x^2 + y^2)^2 - 1$ and below $z = 4 - 4(x^2 + y^2)$:




**56.** Find the average value of the function $f(\rho, \varphi, \vartheta) = \rho$ over the solid ball described by $\rho \le 1$ in spherical coordinates.

**57.** Let $u = x - y$ and $v = 2x + y$.

a. Solve for $x$ and $y$ in terms of $u$ and $v$.

b. Find the Jacobian $\begin{vmatrix} \dfrac{\partial x}{\partial u} & \dfrac{\partial x}{\partial v} \\ \dfrac{\partial y}{\partial u} & \dfrac{\partial y}{\partial v} \end{vmatrix}$.

c. Sketch the image of the triangle with corners $(0, 0)$, $(1, 1)$, and $(1, -2)$ in the $x, y$ plane after the transformation into the $u, v$ plane given by $u = x - y$ and $v = 2x + y$.

d. Use this transformation to evaluate the integral

$$\iint_R (2x^2 - xy - y^2)\, dx\, dy$$

where $R$ is the region in the first quadrant bounded by the lines $y = -2x + 4$, $y = -2x + 7$, $y = x - 2$, and $y = x + 1$.

**58.** Let $u = 3x + 2y$ and $v = x + 4y$.

a. Solve for $x$ and $y$ in terms of $u$ and $v$.

b. Find the Jacobian $\begin{vmatrix} \dfrac{\partial x}{\partial u} & \dfrac{\partial x}{\partial v} \\ \dfrac{\partial y}{\partial u} & \dfrac{\partial y}{\partial v} \end{vmatrix}$.

c. Sketch the image of the triangle bounded by the $x$-axis, the $y$-axis, and the line $x + y = 1$ in the $x, y$ plane after the transformation into the $u, v$ plane given by $u = 3x + 2y$ and $v = x + 4y$.

d. Use this transformation to evaluate the integral

$$\iint_R (3x^2 + 14xy + 8y^2)\, dx\, dy$$

where $R$ is the region in the first quadrant bounded by the lines $y = -(3/2)x + 1$, $y = -(3/2)x + 3$, $y = -(1/4)x$, and $y = -(1/4)x + 1$.

**59.** Find the area of the ellipse $x^2/a^2 + y^2/b^2 = 1$ in the $x, y$-plane by using the transformation $x = au, y = bv$.

**60.** Use the transformation $x = u^2 - v^2$, $y = 2uv$ to evaluate

$$\int_0^1 \int_0^{2\sqrt{1-x}} \sqrt{x^2 + y^2}\, dy\, dx.$$

**61.** Evaluate the following line integrals:

a. $\displaystyle\int_C (x + y)\, ds$ where $C$ is the straight line segment from $(0, 1, 0)$ to $(1, 0, 0)$.

b. $\int_C (x - y + z - 2)\, ds$ where $C$ is the straight line segment from $(0,1,1)$ to $(1,0,1)$.

c. $\int_C (x + \sqrt{y} - z^2)\, ds$ where $C$ is the path that starts at $(0,0,0)$ and then ends at $(1,1,1)$ by moving along $\langle t, t^2, 0\rangle$ for $t \in [0,1]$ and then moving along $\langle 1,1,s\rangle$ for $s \in [0,1]$.

d. $\int_C x\, ds$ where $C$ is the straight line segment from $(0,0)$ to $(4,2)$ in $\mathbb{R}^2$.

e. $\int_C x\, ds$ where $C$ follows the parabolic curve $y = x^2$ from $(0,0)$ to $(2,4)$ in $\mathbb{R}^2$.

f. $\int_C x^2/(y^{4/3})\, ds$ where $C$ follows the path described by the parametric equations $x = t^2, y = t^3$ for $t \in [1,2]$.

g. $\int_C \dfrac{1}{x^2+y^2+1}\, ds$ where $C$ travels counterclockwise along the perimeter of the square in $\mathbb{R}^2$ with corners $(0,0)$, $(1,0)$, $(1,1)$, and $(0,1)$.

**62.** Find the gradient fields for the following functions:

a. $f(x,y,z) = (x^2+y^2+z^2)^{-1/2}$

b. $f(x,y,z) = xy + yz + xz$

**63.** Find work done by $\mathbf{F}$ along the paths $C_1$ and $C_2$ where $C_1$ is the straight line segment from $(0,0,0)$ to $(1,1,1)$ and $C_2$ is the path described by $\langle t, t^2, t^3\rangle$ for $t \in [0,1]$:

a. $\mathbf{F} = \langle 3y, 3x, 4z\rangle$

b. $\mathbf{F} = \langle yz, xz, xy\rangle$

**64.** Find the work done by $\mathbf{F}$ over the following curves:

a. $\mathbf{F} = \langle xy, y, -yz\rangle$ over $\mathbf{r}(t) = \langle t, t^2, t\rangle$ for $t \in [0,1]$

b. $\mathbf{F} = \langle 2y, 3x, x+y\rangle$ over $\mathbf{r}(t) = \langle \cos t, \sin t, t/6\rangle$ for $t \in [0, 2\pi]$

c. $\mathbf{F} = \langle xy, y - x\rangle$ over the straight line in $\mathbb{R}^2$ from $(1,1)$ to $(2,3)$

d. $\mathbf{F}$ is the gradient of $f(x,y) = (x+y)^2$ over the path that travels counterclockwise once around the circle $x^2 + y^2 = 4$ in $\mathbb{R}^2$ that starts and ends at $(2,0)$.

**65.** State whether or not the vector field is conservative:

a. $\mathbf{F} = \langle yz, xz, xy\rangle$

b. $\mathbf{F} = \langle y, x+z, -y\rangle$

c. $\mathbf{F} = \langle -y, x\rangle$

**66.** Find a potential function $f$ for the field $\mathbf{F}$:

a. $\mathbf{F} = \langle 2x, 3y, 4z\rangle$

b. $\mathbf{F} = \langle y+z, x+z, x+y\rangle$

**67.** Show that the values of the follow integrals do not depend on the path from $A$ to $B$:

a. $\int_A^B z^2\, dx + 2y\, dy + 2xz\, dz$

b. $\int_A^B \dfrac{x\, dx + y\, dy + z\, dz}{\sqrt{x^2+y^2+z^2}}$

**68.** Find a potential function for the gravitational field

$$\mathbf{F} = -GmM\frac{\langle x,y,z\rangle}{(x^2+y^2+z^2)^{3/2}}$$

where $G$, $m$, and $M$ are constants. Then show that the word done by the gravitational field in moving a particle from a distance $a$ to a distance $b$ away from the origin is

$$GmM\left(\frac{1}{b} - \frac{1}{a}\right).$$

**69.** Find the work done by $\mathbf{F}$ in moving a particle once counterclockwise around the given curve $C$:

a. $\mathbf{F} = \langle 2xy^3, 4x^2y^2\rangle$ with $C$ the boundary of the region in the first quadrant enclosed by the $x$-axis, the line $x = 1$, and the curve $y = x^3$

b. $\mathbf{F} = \langle 4x - 2y, 2x - 4y\rangle$ with $C$ the boundary of the circle $(x-2)^2 + (y-2)^2 = 4$

**70.** We sometimes use the notation $\oint_C (P\, dx + Q\, dy)$ to denote the work done by $F = \langle P, Q\rangle$ along $C$. With this notation, apply Green's theorem to evaluate the integrals:

a. $\oint_C (y^2\, dx + x^2\, dy)$ with $C$ the triangle bounded by $x = 0$, $x + y = 1$, and $y = 0$.

b. $\oint_C (3y\, dx + 2x\, dy)$ with $C$ the boundary of $0 \le x \le \pi, 0 \le y \le \sin x$

**71.** Green's area formula says that the area of a region $R$ enclosed by a simple closed curve $C$ in $\mathbb{R}^2$ is given by

$$\frac{1}{2}\oint_C x\,dy - y\,dx.$$

Use this formula to find the areas of the regions enclosed by these curves:

a. The circle $\mathbf{r}(t) = \langle a\cos t, a\sin t\rangle$ for $t \in [0, 2\pi]$

b. The ellipse $\mathbf{r}(t) = \langle a\cos t, b\sin t\rangle$ for $t \in [0, 2\pi]$

c. The asteroid $\mathbf{r}(t) = \langle \cos^3 t, \sin^3 t\rangle$ for $t \in [0, 2\pi]$

**72.** Find the surface area of the surface parameterized by $r(u, v) = \langle uv, v + u^2, v - u^2\rangle$ for $u, v \in [0, 1]$.

**73.** Find the surface integrals:

a. Integrate $xyz$ over the triangular surface with vertices $(1, 0, 0)$, $(0, 2, 0)$, and $(0, 1, 1)$.

b. Integrate $yz$ over the part of the sphere $x^2 + y^2 + z^2 = 4$ that lies above the cone $z = \sqrt{x^2 + y^2}$.

c. Integrate $z - x$ over the cone $z = \sqrt{x^2 + y^2}$ for $0 \le z \le 1$.

**74.** Find the flux of $\mathbf{F} = \langle -y^2x, xy\rangle$ along the curve in $\mathbb{R}^2$ parameterized by $\mathbf{r}(t) = \langle t^2, t - t^3\rangle$ for $t \in [0, 1]$.

**75.** Find the divergence of the gravitational field

$$\mathbf{F}(x, y, z) = -GmM\frac{\langle x, y, z\rangle}{(x^2 + y^2 + z^2)^{3/2}}$$

where $G, m, M$ are constants.

**76.** Use the divergence theorem the calculate the flux of the following vector fields $\mathbf{F}$ over the closed surface $S$:

a. $\mathbf{F} = \langle y - z, z - y, y - x\rangle$ and $S$ is the cube bounded by the planes $x = \pm 1$, $y = \pm 1$, and $z = \pm 1$.

b. $\mathbf{F} = \langle x^2, xz, 3z\rangle$ and $S$ is the unit sphere with center the origin.

c. $\mathbf{F} = \langle 6x^2 + 2xy, 2y + x^2z, 4x^2y^3\rangle$ and $S$ is the cylinder above $z = 0$, below $z = 3$, and contained within $x^2 + y^2 = 4$.

**77.** Use Stokes' theorem to find the flux of the curl for $\mathbf{F} = \langle x^2 - y, 4z, x^2\rangle$ around the surface of the cone $z = \sqrt{x^2 + y^2}$ for $z \in [0, 1]$.

**78.** Use Stokes' theorem to calculate the flux of the curl for $\mathbf{F} = \langle y, -xz, xz^2\rangle$ around the surface $z = x^2 + 4y^2$ beneath the plane $z = 1$.