

**UNIVERSITY OF TORONTO MISSISSAUGA
DECEMBER 2015 FINAL EXAMINATION**

MAT102H5F - Introduction to Mathematical Proofs

Shay Fuchs, Ali Mousavidehshikh, Marina Tvalavadze

Duration: 2 hours

Aids: None

NAME (PRINT):

Tan

Last / Surname

Anthony

First / Given Name

STUDENT #:

1002583402

SIGNATURE:

Anthony Tan

The University of Toronto Mississauga and you, as a student, share a commitment to academic integrity. You are reminded that you may be charged with an academic offence for possessing any unauthorized aids during the writing of an exam. Clear, sealable, plastic bags have been provided for all electronic devices with storage, including but not limited to: cell phones, SMART devices, tablets, laptops, calculators, and MP3 players. Please turn off all devices, seal them in the bag provided, and place the bag under your desk for the duration of the examination. You will not be able to touch the bag or its contents until the exam is over. If, during an exam, any of these items are found on your person or in the area of your desk other than in the clear, sealable, plastic bag, you may be charged with an academic offence. A typical penalty for an academic offence may cause you to fail the course.

*Please note, you **CANNOT** petition to re-write an examination once the exam has begun.*

INSTRUCTIONS

ISSUED TO STUDENT

- There are two parts to this examination:
PART I (40 marks): Ten short questions. Each question is worth 4 marks.
PART II (60 marks): Six written questions. Each question is worth 10 marks.
- This examination has 9 different pages including this page. Make sure your copy of the examination has 9 different pages and **sign** at the top of this page. You can use the back of page 9 for rough work.

Good Luck!

Question	Part I	Part II Q#1	Part II Q#2	Part II Q#3	Part II Q#4	Part II Q#5	Part II Q#6	TOTAL
Marks	<i>20</i> /40	<i>0</i> /10	<i>4</i> /10	<i>5</i> /10	<i>6</i> /10	<i>5</i> /10	<i>10</i> /10	<i>50</i> /100

PART I (40 marks)

Answer the following short questions. Each question is worth **4 marks**.


No explanation is needed.

1. Let D be the following subset of \mathbb{R} :

$$D = [1, 3] \cup \{4\}$$

Draw, on the provided grid, the set $D \times D$.

Use **solid** and **dotted** lines to indicate whether the boundary is or is not part of the set.



2. Complete the following sentence:

The **image** of the function $f: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{R}$, $f(n, m) = (-1)^n + (-1)^m$ is ~~(0, \infty)~~.

3. Complete the addition table on the right for a field with four elements $F = \{0, 1, a, b\}$.

+	0	1	a	b
0	0	1	a	b
1	1	0	b	a
a	a	b	0	1
b	b	a	1	0

4. Consider the following statement:

$P =$ "There exists an integer M , such that $x^2 \leq M$ for all real numbers x ."

- (a) Write the statement P using the **logic symbols**.

~~$(\exists M \in \mathbb{Z})(\forall x \in \mathbb{R})(x^2 \leq M)$~~ $(\exists M \in \mathbb{Z})(\forall x \in \mathbb{R})(x^2 \leq M)$

- (b) Write the **negation** of P using the **logic symbols** (but without using the symbol ' \neg ').

$(\forall M \in \mathbb{Z})(\exists x \in \mathbb{R})(x^2 > M)$

- (c) Which statement is true: P or $\neg P$? **Circle the correct answer.**

P is TRUE

$\neg P$ is TRUE

5. Complete with a single number:

$$\gcd(24, 54 + 24^7) = \underline{3}$$

6. Describe (in words) the set of all integers m , for which the equation $3x + m \cdot y = 1$ has integer solutions.

m must be an integer such that $\gcd(3, m) = 1$. This means 3 and m must be relatively prime.

In Questions 7-10, there is only ONE correct answer. Circle it!

7. Let P and Q be two statements.

If P is true and Q is false, then which of the following statements is TRUE?

• $P \wedge Q$

• $P \Rightarrow Q$

• $Q \Rightarrow P$

• $(\neg P) \vee Q$

8. The function $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \begin{cases} x & \text{if } x \leq 0 \\ \frac{1}{x} & \text{if } x > 0 \end{cases}$ is...

• a surjection but not an injection.

• an injection but not a surjection.

• a bijection.

• neither a surjection nor an injection.

This is neither surjective or injective.

9. Which of the following sets is countable?

• $\mathbb{R} - \mathbb{Q}$

• $(0, \infty)$

• $P(\mathbb{Q})$

• $\mathbb{Z} - \mathbb{N}$

10. Consider the following equivalence relation on \mathbb{Z} :

$$a \sim b \quad \text{if and only if} \quad a - b \text{ is divisible by } 5.$$

Then the set $\{\dots, -7, -2, 3, 8, 13, 18, \dots\}$ is the equivalence class of...

• 22

• 33

• -11

• -15

PART II (60 marks)

Answer the following questions. Each question is worth **10 marks**.

Provide **complete solutions and justify your arguments**.

1. Is the following statement **TRUE** or **FALSE**?

"For any two sets A and B : if $A - B = B - A$, then $A = B$."

If the statement is **true**, prove it! If the statement is **false**, provide a counter-example.

a) $A - B \subseteq B - A$

~~Let $x \in A - B$~~

If $A = B$,

$x \in A$

$x \in B$

$A - B = \emptyset$

$A - B \subseteq B - A$

If $A = B$, x would be both an element of A and B . This means if A and B are subtracted, an empty set is formed.

If the sets are equal, the order of subtraction can be changed to achieve the same answer.

b) $A - B \supseteq B - A$

If $A = B$

$y \in B$

$y \in A$

$B - A = \emptyset$

$B - A \subseteq A - B$

Following the same idea,

$B - A$ can be written as

$A - B$, meaning $A = B$

\emptyset

∴ The statement holds true, $A - B = B - A$ if $A = B$

not what the question is asking?

2. Prove, by induction, the following identity for all $n \in \mathbb{N}$:

$$1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{4n^3 - n}{3}$$

Base case:

$$n=1$$

$$(2(1)-1)^2 = \frac{4(1)^3 - 1}{3}$$

$$1 = \frac{3}{3}$$

$$1=1 \text{ so True}$$

Induction.

Assume $\frac{4k^3 - k}{3}$ holds true for $k \in \mathbb{N}$. Prove this holds true for $k+1$

$$1^2 + 3^2 + 5^2 + \dots + (2k-1)^2 + (2(k+1)-1)^2 = \frac{4(k+1)^3 - (k+1)}{3}$$

$$\text{I.H.} \rightarrow \frac{4k^3 - k}{3} + (2(k+1)-1)^2 = \frac{4(k+1)^3 - (k+1)}{3}$$

$$4k^3 - k + 3(2(k+1)-1)^2 = 4(k+1)^3 - (k+1)$$

$$4k^3 - k + 3(2k+1)^2 = 4(k+1)^3 - (k+1)$$

$$4k^3 - k + 3(4k^2 + 4k + 1) = 4(k^3 + 3k^2 + 3k + 1) - (k+1)$$

$$4k^3 + 12k^2 + 12k + 3 = 4k^3 + 12k^2 + 12k + 4 - k - 1$$

$$4k^3 + 12k^2 + 12k + 3$$

$$4k^3 - k + 12k^2 + 12k + 3 = (4k^2 + 8k + 4)(k+1) - k - 1$$

$$4k^3 + 12k^2 + 11k + 3 = 4k^3 + 4k^2 + 8k^2 + 8k + 4k + 4 - k - 1$$

$$4k^3 + 12k^2 + 11k + 3 = 4k^3 + 12k^2 + 11k + 3$$

$$\text{so L.S.} = \text{R.S.}$$

By PMI, the statement holds true for $n \in \mathbb{N}$

3. Prove, that for any **odd integer** k , the number $\sqrt{2k}$ is **irrational**.

(Hint: Use contradiction.)

~~Proof by contradiction. Let's show~~

lets assume $\sqrt{2k}$ is rational for odd values of k . let ~~there~~ $m \in \mathbb{Z}$

$$\frac{p}{q} = \sqrt{2(2m-1)} \quad p, q \in \mathbb{Z}$$

$$\frac{p^2}{q^2} = 2(2m-1)$$

$$p^2 = q^2(4m-2)$$

WHY?

p is a multiple of $(4m-2)$

$$k(4m-2)^2 = q^2(4m-2), \quad k \in \mathbb{Z}$$

q is also a multiple of $(4m-2)$

There is a contradiction since the fraction $\frac{p}{q}$ is not in lowest terms
 $\& \sqrt{2k}$ is irrational when k is an odd number.

4. Define a relation R on the set of all real number as follows:

$$(x, y) \in R \quad \text{if and only if} \quad |x + y| = |x| + |y|.$$

Is this relation reflexive? Is it symmetric? Is it transitive?

Is this an equivalence relation? Explain.

a)

Reflexive: let $x=y$.

$$|x + x| = |x| + |x|$$

$$|2x| \neq 2|x|$$

∴ The relation is not reflexive.

3

Symmetric: $|x+y| = |x|+|y|$

$$|y+x| = |x|+|y|$$

$$|x+y| = |y+x|$$

addition laws allow us to switch the position of 2 numbers and have the same answer

∴ Symmetric.

3

Transitive:

$$|x+y| = |x|+|y| \Rightarrow |x+y| - |y| = |x|$$

$$|y+z| = |y|+|z| \Rightarrow |y+z| - |y| = |z|$$

$$|x+z| \neq |x|+|z|$$

$$|x+z| = |x+y| - |y| + |y+z| - |y|$$

$$|x+z| \neq |x+y| + |y+z| - 2|y|$$

∴ The relation is not transitive.

0

∴ The relation is only symmetric

b) This is not an equivalence relation since it does not satisfy reflexivity and transitivity. An equivalence relation requires all 3.

5. Consider the function $f: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, $f(a, b) = 12^a \cdot 18^b$.

(a) Show that f is an **injection** (one-to-one).

$$\text{let } a=1$$

$$\text{let } b=1$$

$$f(1, b_1) = f(1, b_2)$$

$$f(a_1, 1) = f(a_2, 1)$$

$$12 \cdot 18^{b_1} = 12 \cdot 18^{b_2}$$

$$12^{a_1} \cdot 18 = 12^{a_2} \cdot 18$$

$$18^{b_1} = 18^{b_2}$$

$$12^{a_1} = 12^{a_2}$$

$$\text{so } b_1 = b_2$$

$$\text{so } a_1 = a_2$$

∴ The function f is an injection.

X

(-5)

(b) Is f a **surjection** (onto)? Explain.

No. by substituting $a, b = 1$ (the smallest case) we get $f(1, 1) = 12 \cdot 18$


This means that $f(a, b) \geq 12 \cdot 18$. All natural numbers less than $12 \cdot 18$ will not be mapped. ∴ The function is not a surjection.

✓

6. Consider the function $g: \mathbb{R} \rightarrow \mathbb{R}$, $g(x) = 2x + 1$.

Prove, by induction, that for any $n \in \mathbb{N}$, $g^n(x) = 2^n \cdot x + 2^n - 1$.

Note: g^n is the function obtained by composing n copies of g : $g^n = \underbrace{g \circ g \circ \dots \circ g}_{n \text{ times}}$.

base case: 

$n=1$

$$\begin{aligned} g^1(x) &= 2^1 \cdot x + 2^1 - 1 \\ &= 2x + 1 \end{aligned}$$



is True

Induction:

assume $g^k(x) = 2^k \cdot x + 2^k - 1$, for $k \in \mathbb{N}$. Prove this for $k+1$

$$g^k(2x+1) = g^{k+1}(x)$$

$$\text{I.H.} \rightarrow 2^k \cdot (2x+1) + 2^k - 1 = 2^{k+1} \cdot x + 2^{k+1} - 1$$

$$2^k \cdot (2x+1) + 2^k = 2^{k+1} \cdot x + 2^{k+1}$$

$$2^k 2x + 2^k + 2^k = 2^{k+1} x + 2^{k+1}$$

$$2^k (2x+1+1) = 2^{k+1} (x+1)$$

$$2^k (2x+2) = 2^{k+1} (x+1)$$

$$2^k 2(x+1) = 2^{k+1} (x+1)$$

$$2^{k+1} (x+1) = 2^{k+1} (x+1)$$



By PMI, the statement holds true for $n \in \mathbb{N}$

