

# MATH1M03 Test #2, Review Session

TA: Anthony Tan (tana4@mcmaster.ca)

Office hours: 2:30p - 6:30p Fri in Math  
Help Centre

① Substitution (or some articles/  
books say "u-substitution")

$$\int f(x) dx \longrightarrow \int g(u) du$$

> We are replacing  $x$ 's  
with  $u$ 's and (hopefully)  
things cancel nicely

> There is something in my  
expression that looks like a  
derivative of  $u$

## ② Integration by Parts

$$\int u dv = uv - \int v du$$

↳ We declare one function  
to be  $u$  and another to  
be  $dv$

Eventually you'll a set-up  
like  $\left\{ \begin{array}{l} u = ? , \quad du = ? \\ dv = ? , \quad v = ? \end{array} \right\}$


Alternatively, if you prefer  
 $\left\{ \begin{array}{l} u = ? , \quad u' = ? \\ v' = ? , \quad v = ? \end{array} \right\}$

③ For a closed interval  
 $[a, b]$  (or alternatively  
 $a \leq x \leq b$ )


the average value is  
given by

$$\frac{1}{b-a} \int_a^b f(x) dx$$

# ④ Area between curves



How do I find the area of this region?



After cutting "a bunch of times"  $g(x)$

Area between  $f(x)$  and  $g(x)$  on  $[a, b]$  = "The continuous sum of rectangles between  $a$  and  $b$ "

$$= \int_a^b [f(x) - g(x)] dx$$

## 2nd Sample Test #2

2. The slope of a graph at any  $x$  is given by  $f'(x) = e^2 - e^x$  and the function passes  $(0, 1)$ .

Find  $f(x)$ .

Thoughts:

- Want  $f(x)$  given  $f'(x) \dots$

$$f'(x) \xrightarrow{\int dx} f(x)$$

-  $f(0) = 1$  (same as  $(0, 1)$ )

Soln

$$\int e^2 - e^x dx = \underbrace{e^2 x - e^x}_{f(x)} + C$$

$$f(0) = 1$$

$$1 = \cancel{e^2(0)} - e^0 + C$$

$$C = 2$$

$$\therefore f(x) = e^2 x - e^x + 2 \quad \boxed{E}$$

$$7. \int \frac{e^{2x}}{\sqrt{3+e^{2x}}} dx = \int \frac{\cancel{e^{2x}}}{\sqrt{u}} \cdot \frac{du}{\cancel{2e^{2x}}}$$

$$\left\{ \begin{array}{l} \underline{u} = 3 + e^{2x} \\ \underline{\frac{du}{dx}} = \frac{2e^{2x}}{1} \\ \underline{dx} = \frac{du}{2e^{2x}} \end{array} \right\} = \int \frac{1}{2} \cdot u^{-1/2} du$$

$$= \frac{1}{2} \cdot \frac{1}{-1/2+1} u^{-1/2+1} + C$$

$$= \frac{1}{2} \cdot 2 u^{1/2} + C$$

$$= (3 + e^{2x})^{1/2} + C$$

C



Ex (Not in sample test, but it's a tricky problem)

$$\int_1^3 t^5 \sqrt{t^3 + 4} \, dt$$

Pause and compute  $\int t^5 \sqrt{t^3 + 4} \, dt$

Set-up:  $u = t^3 + 4$   
 $\frac{du}{dt} = 3t^2$

modify set-up:

$$t^3 = u - 4$$

$$\frac{du}{3} = t^2 dt$$

$$\int t^5 \sqrt{t^3 + 4} \, dt$$

$$= \int t^2 t^3 \sqrt{t^3 + 4} \, dt$$

$$= \int \underbrace{t^3}_{u-4} \underbrace{\sqrt{t^3+4}}_u \underbrace{t^2 dt}_{du/3}$$

$$= \int (u-4) u^{1/2} \frac{du}{3}$$

$$= \frac{1}{3} \int (u^{3/2} - 4u^{1/2}) \, du$$

$$= \frac{1}{3} \left[ \frac{1}{1+3/2} u^{3/2+1} - \frac{4}{1+1/2} u^{1/2+1} \right] + C$$

$$= \frac{1}{3} \left[ \frac{2}{5} u^{5/2} - 4 \cdot \frac{2}{3} u^{3/2} \right] + C$$

$$= \frac{1}{3} \left[ \frac{2}{5} (t^3+4)^{5/2} - \frac{8}{3} (t^3+4)^{3/2} \right] + C$$

$F(t)$

$$\int_1^3 t^5 \sqrt{t^3+4} dt = F(3) - F(1)$$

---


$$\text{Ans: } \frac{2}{15} (31)^{5/2} - \frac{2}{15} (5)^{5/2} - \frac{8}{9} (31)^{3/2} + \frac{8}{9} (5)^{3/2}$$

18. Evaluate  $\int_1^2 x^2 e^x dx$

For now, forget about  $\int_1^2$  and focus on  $\int dx$  (indefinite int)

$$\int x^2 e^x dx \quad \overset{\text{OLD}}{\left\{ \begin{array}{l} u = x^2, du = 2x \\ dv = e^x, v = e^x \end{array} \right\}}$$

$$= uv - \int v du$$

$$\overset{\text{OLD}}{=} x^2 e^x - \int 2x e^x dx \quad \overset{\text{NEW}}{\left\{ \begin{array}{l} u = x, du = 1 \\ dv = e^x, v = e^x \end{array} \right\}}$$

$$= x^2 e^x - 2 \left( \int x e^x dx \right) \longrightarrow$$

$$= x^2 e^x - 2 \left( uv - \int v du \right)$$

NEW

$$= x^2 e^x - 2 \left( x e^x - \int e^x dx \right)$$

$$= x^2 e^x - 2 \left( x e^x - e^x \right) + C$$

$F(x)$

$$\begin{aligned} \text{Then } \int_1^2 x^2 e^x dx &= F(2) - F(1) \\ &= 2e^2 - e \quad \boxed{C} \end{aligned}$$

19. A manufacturer determines that when  $x$  hundred units of a particular commodity are produced, the profit generated is  $P(x)$  thousand dollars, where

$$P(x) = \frac{500 \ln(x+1)}{(x+1)^2}$$

What is the average profit over the production range  $0 \leq x \leq 10$ ? (★)

$a$

$b$

$$\begin{aligned}
 \textcircled{\star} &= \frac{1}{b-a} \int_a^b P(x) dx \\
 &= \frac{1}{10} \int_0^{10} \frac{500 \ln(x+1)}{(x+1)^2} dx \\
 &= 50 \int_0^{10} \frac{\ln(x+1)}{(x+1)^2} dx
 \end{aligned}$$

---

 PAUSE

$$\begin{aligned}
 &50 \int \frac{\ln(x+1)}{(x+1)^2} dx \\
 &= 50 \int \frac{\ln u}{u^2} du \quad \left\{ \begin{array}{l} u = x+1 \\ \frac{du}{dx} = 1 \\ \text{or } du = dx \end{array} \right\}
 \end{aligned}$$

$$\underbrace{\ln u}_v \cdot \underbrace{\frac{1}{u^2}}_{dw}$$

$$\left\{ \begin{array}{l} v = \ln u, \quad dv = 1/u \\ dw = 1/u^2, \quad w = -1/u \end{array} \right\}$$

$$= 50 \left( vw - \int w \, dv \right)$$

$$= 50 \left( -\frac{\ln u}{u} - \int -\frac{1}{u} \cdot \frac{1}{u} \, du \right)$$



$$= 50 \left( -\frac{\ln u}{u} + \int \frac{1}{u^2} du \right)$$

$$= 50 \left( -\frac{\ln u}{u} - \frac{1}{u} \right) + C$$

$$= 50 \left( -\frac{\ln(x+1)}{x+1} - \frac{1}{x+1} \right) + C$$

$$= \frac{-50}{x+1} (\ln(x+1) + 1) + C$$

$$\underbrace{\hspace{10em}}_{\hat{P}(x)} \quad \swarrow [C]$$


$$\text{Ans: } \hat{P}(10) - \hat{P}(0) \approx 34555.02$$

after  
(mult by  
1000)

15. Find the area of the region bounded by  $f(x) = x^2 + 9x - 28$

$$g(x) = -x^2 + 5x + 2$$

Soln Expect a picture like



- $g(x) > f(x)$
- Points of Intersection

$$f(x) = g(x)$$

$$\Leftrightarrow x^2 + 9x - 28 = -x^2 + 5x + 2$$

$$\Leftrightarrow 0 = x^2 + x^2 + 9x - 5x - 28 - 2$$

$$\Leftrightarrow \frac{0}{2} = \frac{2x^2}{2} + \frac{4x}{2} - \frac{30}{2}$$

$$\Leftrightarrow 0 = x^2 + 2x - 15$$

$$\Leftrightarrow 0 = (x-3)(x+5)$$

$$\Leftrightarrow x = \underbrace{-5}_a \text{ or } \underbrace{3}_b$$

Evaluate:

Area between

$f(x)$  and  $g(x)$

$$= \int_{-5}^3 \text{height} \cdot \text{width}$$

$$= \int_{-5}^3 [g(x) - f(x)] dx$$

$$= \int_{-5}^3 [-2x^2 - 4x + 30] dx$$

$$= -\frac{2x^3}{3} - 2x^2 + 30x \Big|_{-5}^3$$

$$= \left[ -\frac{2(3)^3}{3} - 2(3)^2 + 30(3) \right] - \left[ -\frac{2(-5)^3}{3} - 2(-5)^2 + 30(-5) \right]$$

$$\approx 170.67 \quad \boxed{1E}$$

Ex (Not from sample) Determine if the following integrals converge or diverge.

$$(a) \int_4^{\infty} \frac{2}{x} dx = \lim_{t \rightarrow \infty} \int_4^t \frac{2}{x} dx$$

$$= 2 \lim_{t \rightarrow \infty} \int_4^t \frac{1}{x} dx$$

$$= 2 \lim_{t \rightarrow \infty} \left[ \ln|x| \right]_4^t$$

$$= 2 \lim_{t \rightarrow \infty} \left[ \cancel{\ln |t|}^{\infty \text{ (slowly)}} - \ln |4| \right]$$

$$= \infty \Rightarrow \text{DIV}$$

$$(b) \int_9^{\infty} \frac{5}{x^{3/2}} dx$$

$$= 5 \lim_{t \rightarrow \infty} \int_9^t x^{-3/2} dx$$

$$= 5 \lim_{t \rightarrow \infty} \left[ \frac{1}{1+(-3/2)} x^{-3/2+1} \Big|_9^t \right]$$


$$= 5 \lim_{t \rightarrow \infty} \left[ -2x^{-1/2} \middle| t \atop q \right]$$

$$= -10 \lim_{t \rightarrow \infty} \left[ \frac{1}{x^{1/2}} \middle| t \atop q \right]$$

$$= -10 \lim_{t \rightarrow \infty} \left[ \frac{1}{\cancel{t^{1/2}}} - \frac{1}{3} \right]$$

$$= -10 \left[ 0 - \frac{1}{3} \right]$$

$$= \frac{10}{3}$$



$$\Rightarrow \lim = a \neq$$

$$\Rightarrow \text{CONV}$$

Further,

$$\int_{-\infty}^a f(x) dx = \lim_{t \rightarrow -\infty} \int_t^a f(x) dx$$

CONV  
if limit  
exists

DIV  
if limit  
DNE

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^{\infty} f(x) dx \rightarrow$$



$$\int_{-\infty}^a f(x) dx + \int_a^{\infty} f(x) dx$$

$$= \lim_{t \rightarrow -\infty} \underbrace{\int_t^a f(x) dx}_{(A)} + \lim_{s \rightarrow \infty} \underbrace{\int_a^s f(x) dx}_{(B)}$$

$$\text{If } \int_{-\infty}^{\infty} f(x) dx \text{ CONV,}$$

Then (A) and (B) CONV

