

A geometric approach to clustering

Identity of indiscernibles	$d(x,y) = 0 \Leftrightarrow x = y$
Non-negativity	$d(x,y) \ge 0$
Symmetry	d(x,y) = d(y,x)
Triangle inequality	$d(x,y) \le d(x,z) + d(z,y)$


Properties of any metric space


Simple random geometric graph

- 1. Sprinkle N nodes uniformly on a disk of radius R.
- 2. Connect any nodes separated by a distance less than r.
- ✓ high clustering

binomial/Poisson degree distribution

Assume that the nodes are embedded in a metric space and that any two nodes are connected with a probability that is a decreasing function of the distance between them. For further info, see Phys. Rep. 499, 1-101 (2011)





A geometric approach to clustering

Assume that the nodes are embedded in a metric space and that any two nodes are connected with a probability that is a decreasing function of the distance between them.

Properties of any metric space

Identity of indiscernibles $d(x,y) = 0 \Leftrightarrow x = y$


Non-negativity $d(x,y) \ge 0$

Symmetry d(x,y) = d(y,x)

Triangle inequality $d(x,y) \le d(x,z) + d(z,y)$

Simple random geometric graph

- 1. Sprinkle N nodes uniformly on a disk of radius R.
- 2. Connect any nodes separated by a distance less than r.
- ✓ high clustering
- binomial/Poisson degree distribution





For further info, see Phys. Rep. 499, 1–101 (2011)

A geometric approach to clustering: Hyperbolic geometry

Simple random geometric graph

- 1. Sprinkle N nodes uniformly on the hyperbolic disk of radius R.
- 2. Connect any nodes separated by a distance less than r = R.





- ✓ high clustering
- ✓ power-law degree distribution with exponent -3

Phys. Rev. E 82, 036106 (2010)