

Extra credit problems

Math 427

0. Find a mistake or misprint in the book. (The score depends on the type of mistake.)

1. Describe all the motions of the Manhattan plane.

2. Construct a metric space \mathcal{X} and a distance preserving map $f: \mathcal{X} \rightarrow \mathcal{X}$ that is not a motion of \mathcal{X} .

3. Note that the following quantity

$$\tilde{\angle}ABC = \begin{cases} \pi & \text{if } \angle ABC = \pi \\ -\angle ABC & \text{if } \angle ABC < \pi \end{cases}$$

can serve as the angle measure; that is, the axioms hold if one changes everywhere \angle to $\tilde{\angle}$.

(a). Show that \angle and $\tilde{\angle}$ are the only possible angle measures on the plane.

(b). Show that without Axiom IIIc, this is not longer true.

4. Let M be the midpoint of the side $[AB]$ of $\triangle ABC$ and M' be the midpoint of the side $[A'B']$ of $\triangle A'B'C'$. Assume $C'A' = CA$, $C'B' = CB$, and $C'M' = CM$. Prove that $\triangle A'B'C' \cong \triangle ABC$.

5. (solved) Show that a composition of three reflections in the sides of a nondegenerate triangle does not have a fixed point.

6. Consider triangle $\triangle ABC$ with $D \in (AC)$ such that $(BD) \perp (AC)$, and points N and M such that $AN = DC$, $CM = AD$, $(AN) \perp (AB)$ and $(CM) \perp (BC)$.

Prove that M and N are equidistant from B .

7. Lines ℓ and m are tangent to two circles of radii r and R on such a way the circles are on one side of ℓ and on different sides of m . Let A and B be tangential points of ℓ and Q be the point of intersection ℓ and m . Show that

$$QA \cdot QB = R \cdot r.$$

8. Given two parallel lines ℓ and m and a point P , use only ruler to construct the line through P parallel to ℓ and m . (You can play with the java applet “Third parallel line” on anton-petrinin.github.io/birkhoff/car/.)

9. Let ABC be a nondegenerate triangle and $A' \in (BC)$, $B' \in (CA)$, $C' \in (AB)$ be the points such that

$$2 \cdot \angle AA'B \equiv 2 \cdot \angle BB'C \equiv 2 \cdot \angle CC'A \equiv \frac{2}{3}\pi.$$

Show that the triangle formed by the lines (BB') , (CC') , and (AA') , is congruent to $\triangle ABC$.

10. Construct a triangle with the given perimeter, base, and the opposite angle (You can play with the java applet “Triangle with given base, perimeter and angle.html” on anton-petrinin.github.io/birkhoff/car/.)