Alexandrov geometry

preliminary version no. 1

S. Alexander, V. Kapovitch, A. Petrunin

Contents

I Preliminaries **1**

1 Model plane **3**
1.1 Trigonometry 3
1.2 Hemisphere lemma 6

2 Metric spaces **9**
2.1 Metrics and their relatives 9
2.2 Notations 11
2.3 Length spaces 13
2.4 Convex sets 16
2.5 Quotient spaces 17
2.6 Gluing and doubling 19
2.7 Kuratowsky embedding 20

3 Ultralimits **21**
3.1 Ultrafilters 21
3.2 Ultralimits of points 22
3.3 Ultralimits of spaces 24
3.4 Ultratangent space 26
3.5 Gromov–Hausdorff convergence and ultralimits 27
3.6 Ultralimits of sets 29
3.7 Ultralimits of functions 29

4 Maps and functions **31**
4.1 Submaps 31
4.2 Lipschitz conditions 32
4.3 Isometries and sumbetries 33
4.4 Speed of curves 35
4.5 Convex real-to-real functions 36
4.6 Convex functions on a metric space 38

5 The ghost of Euclid **41**
5.1 Geodesics, triangles and hinges 41
5.2 Model angles and triangles 42
5.3 Angles and the first variation 44
5.4 Constructions 46
5.5 Space of directions and tangent space 47
5.6 Velocity of curves 47
5.7 Differential 49
5.8 Remarks 50

6 Dimension theory **51**
6.1 Definitions 51
6.2 Linear dimension 53

II Definitions and constructions **55**

7 Definitions of curvature bounded below **57**
7.1 Four-point comparison 57
7.2 Geodesics 60
7.3 More comparisons 62
7.4 Function comparison 65
7.5 Development 66
7.6 Local definitions and globalization 70
7.7 Properties of geodesics and angles 76
7.8 On positive lower bound 81
7.9 Remarks and open problems 84

8 Definitions of curvature bounded above **85**
8.1 Four-point comparison. 85
8.2 Geodesics 86
8.3 More comparisons 88
8.4 Thin triangles 91
8.5 Function comparison 94
8.6 Convex sets 95
8.7 Development 98
8.8 Patchwork globalization 99
8.9 Reshetnyak's gluing theorem 101
8.10 Space of geodesics 103
8.11 Lifting globalization 107
8.12 Reshetnyak majorization 111
8.13 Hadamard–Cartan theorem 118
8.14 Angles 122
8.15 Remarks and open problems 124

9 Kirszbraun revisited **125**
9.1 Short map extension definitions. 125
9.2 $(1+n)$-point comparison 129
9.3 Helly's theorem 131
9.4 Kirszbraun's theorem 132
9.5 Curvature free 138
9.6 Remarks and open problems 139

10 Warped products **141**
10.1 Definition 141
10.2 Extended definition 144
10.3 Examples 144
10.4 1-dimensional base 145
10.5 General case 147

III Structure and tools **149**

11 First order differentiation **151**
11.1 Intrinsic property of tangent space 151
11.2 Differential 154
11.3 Ultratangent space 156
11.4 Gradient 156
11.5 Polar vectors 162
11.6 Linear subspace of tangent space 163
11.7 Comments 165

12 Dimension of CAT spaces **167**
12.1 The case of complete geodesic spaces 167
12.2 The case of CAT spaces 172
12.3 Dimension of CBA spaces 176
12.4 Finite-dimensional CAT spaces 179
12.5 Remarks and open problems 181

13 Dimension of CBB spaces **183**
13.1 Struts and rank 183
13.2 Right-inverse theorem 184
13.3 Dimension theorem for CBB 187
13.4 Finite dimensional CBB spaces 191
13.5 One-dimensional CBB spaces 194

14 Gradient flow **197**
14.1 Gradient-like curves 197
14.2 Gradient curves 200
14.3 Gradient flow 207
14.4 Line splitting theorem 208
14.5 Radial curves 210
14.6 Radial comparisons 212
14.7 Gradient exponent 217
14.8 Remarks 218

Bibliography **229**

Part I

Preliminaries

Chapter 1

Model plane

1.1 Trigonometry

Given a real number κ, the *model κ-plane* will be a complete simply connected 2-dimensional Riemannian manifold of constant curvature κ.

The model κ-plane will be denoted by $\mathbb{M}^2(\kappa)$.

- ◇ If $\kappa > 0$, $\mathbb{M}^2(\kappa)$ is isometric to a sphere of radius $\frac{1}{\sqrt{\kappa}}$; the unit sphere $\mathbb{M}^2(1)$ will be also denoted by $\mathbb{S}^2$.
- ◇ If $\kappa = 0$, $\mathbb{M}^2(\kappa)$ is the Euclidean plane, which is also denoted by $\mathbb{E}^2$
- ◇ If $\kappa < 0$, $\mathbb{M}^2(\kappa)$ is the Lobachevsky plane with curvature κ.

Set $\varpi^\kappa = \operatorname{diam} \mathbb{M}^2(\kappa)$, so $\varpi^\kappa = \infty$ if $\kappa \leqslant 0$ and $\varpi^\kappa = \pi/\sqrt{\kappa}$ if $\kappa > 0$.

The distance between points $x, y \in \mathbb{M}^2(\kappa)$ will be denoted by $|x-y|$, and $[xy]$ will denote the segment connecting x and y. The segment $[xy]$ is uniquely defined for $\kappa \leqslant 0$ and for $\kappa > 0$ it is defined uniquely if $|x-y| < \varpi^\kappa = \pi/\sqrt{\kappa}$.

A triangle in $\mathbb{M}^2(\kappa)$, with vertices x, y, z will be denoted by $[xyz]$. Formally, a triangle is an ordered set of its sides, so $[xyz]$ is just a short notation for a triple $([yz], [zx], [xy])$.

The angle of $[xyz]$ at x will be denoted by $\measuredangle[x\,{}^{y}_{z}]$.

By $\tilde{\triangle}^\kappa\{a, b, c\}$ we denote a triangle in $\mathbb{M}^2(\kappa)$ with sidelengths a, b, c, so $[xyz] = \tilde{\triangle}^\kappa\{a, b, c\}$ means that $x, y, z \in \mathbb{M}^2(\kappa)$ are such that

$$|x-y| = c, \quad |y-z| = a, \quad |z-x| = b.$$

For $\tilde{\triangle}^\kappa\{a, b, c\}$ to be defined, the sides a, b, c must satisfy the triangle

inequality. If $\kappa > 0$, we require in addition that $a + b + c < 2\cdot\varpi^\kappa$; otherwise $\tilde\triangle^\kappa\{a,b,c\}$ is considered to be undefined.

Trigonometric functions. We will need three "trigonometric functions" in $\mathbb{M}^2(\kappa)$: cs^κ, sn^κ and md^κ; cs stands for *cosine*, sn stands for *sine* and md, for *modified distance*.

They are defined as the solutions of the following initial value problems respectively:

$$\begin{cases} x'' + \kappa\cdot x = 0, \\ x(0) = 1, \\ x'(0) = 0. \end{cases} \qquad \begin{cases} y'' + \kappa\cdot y = 0, \\ y(0) = 0, \\ y'(0) = 1. \end{cases} \qquad \begin{cases} z'' + \kappa\cdot z = 1, \\ z(0) = 0, \\ z'(0) = 0. \end{cases}$$

Namely we set $\mathrm{cs}^\kappa(t) = x(t)$, $\mathrm{sn}^\kappa(t) = y(t)$ and

$$\mathrm{md}^\kappa(t) = \begin{cases} z(t) & \text{if } 0 \leqslant t \leqslant \varpi^\kappa, \\ \frac{2}{\kappa} & \text{if } t > \varpi^\kappa. \end{cases}$$

Here are the tables which relate our trigonometric functions to the standard ones, where we take $\kappa > 0$:

$$\begin{aligned} &\mathrm{sn}^{\pm\kappa} = \tfrac{1}{\sqrt{\kappa}}\cdot\mathrm{sn}^{\pm1}(x\cdot\sqrt{\kappa}); && \mathrm{cs}^{\pm\kappa} = \mathrm{cs}^{\pm1}(x\cdot\sqrt{\kappa}); \\ &\mathrm{sn}^{-1}x = \sinh x; && \mathrm{cs}^{-1}x = \cosh x; \\ &\mathrm{sn}^{0}x = x; && \mathrm{cs}^{0}x = 1; \\ &\mathrm{sn}^{1}x = \sin x; && \mathrm{cs}^{1}x = \cos x. \end{aligned}$$

$$\begin{aligned} \mathrm{md}^{\pm\kappa} &= \tfrac{1}{\kappa}\cdot\mathrm{md}^{\pm1}(x\cdot\sqrt{\kappa}); \\ \mathrm{md}^{-1}x &= \cosh x - 1; \\ \mathrm{md}^{0}x &= \tfrac{1}{2}\cdot x^2; \\ \mathrm{md}^{1}x &= \begin{bmatrix} 1 - \cos x & \text{for} & x \leqslant \pi, \\ 2 & \text{for} & x > \pi. \end{bmatrix} \end{aligned}$$

Note that

$$\mathrm{md}^\kappa(x) = \int_0^x \mathrm{sn}^\kappa(x)\cdot\boldsymbol{d}x \quad \text{for} \quad x \leqslant \varpi^\kappa$$



Let φ be the angle of $\tilde\triangle^\kappa\{a,b,c\}$ opposite to a. In this case we will write

$$a = \tilde\gamma^\kappa\{\varphi; b, c\} \quad \text{or} \quad \varphi = \measuredangle^\kappa\{a; b, c\}.$$

The functions $\tilde{\Upsilon}^{\kappa}$ and $\tilde{\angle}^{\kappa}$ will be called correspondingly the *model side* and the *model angle*. Set

$$\tilde{\Upsilon}^{\kappa}\{\varphi; b, -c\} = \tilde{\Upsilon}^{\kappa}\{\varphi; -b, c\} := \tilde{\Upsilon}^{\kappa}\{\pi - \varphi; b, c\},$$

this way we define $\tilde{\Upsilon}^{\kappa}\{\varphi; b, c\}$ for some negative values of b and c.

1.1.1. Properties of standard functions.

a) For fixed a and φ, the function

$$y(t) = \mathrm{md}^{\kappa}\left(\tilde{\Upsilon}^{\kappa}\{\varphi; a, t\}\right)$$

satisfies the following differential equation:

$$y'' + \kappa \cdot y = 1.$$

b) Let $\alpha\colon [a, b] \to \mathbb{M}^2(\kappa)$ be a unit-speed geodesic, and A be the image of a complete geodesic. If $f(t)$ is the distance from $\alpha(t)$ to A, the function

$$y(t) = \mathrm{sn}^{\kappa}(f(t))$$

satisfies the following differential equation:

$$y'' + \kappa \cdot y = 0$$

for $y \neq 0$.

c) For fixed κ, b and c, the function

$$a \mapsto \tilde{\angle}^{\kappa}\{a; b, c\}$$

is increasing and defined on a real interval. Equivalently, the function

$$\varphi \mapsto \tilde{\Upsilon}^{\kappa}\{\varphi; b, c\}$$

is increasing and defined if $b, c < \varpi^{\kappa}$ and $\varphi \in [0, \pi]$[1]*.*

d) For fixed φ, a, b, c, the functions

$$\kappa \mapsto \tilde{\angle}^{\kappa}\{a; b, c\} \quad \text{and} \quad \kappa \mapsto \tilde{\Upsilon}^{\kappa}\{\varphi; b, c\}$$

are respectively non-decreasing (in fact, decreasing, if $a \notin \{|b - c|, b + c\}$) and non-increasing (in fact, increasing, if $\varphi \notin \{0, \pi\}$).

e) (Alexandrov's lemma) Assume that for real numbers a, b, a', b', x and κ, the following two expressions are defined

[1] Formally speaking, if $\kappa > 0$ and $b + c \geqslant \varpi^{\kappa}$, it is defined only for $\varphi \in [0, \pi)$, but $\tilde{\Upsilon}^{\kappa}\{\varphi; b, c\}$ can be extended to $[0, \pi]$ as a continuos function.

1. $\measuredangle^\kappa\{a;b,x\} + \measuredangle^\kappa\{a';b',x\} - \pi$,

2. $\measuredangle^\kappa\{a';b+b',a\} - \measuredangle^\kappa\{x;a,b\}$,

Then they have the same sign.

All the properties except Alexandrov's lemma (e) can be shown by direct calculation. Alexandrov's lemma is reformulated in 5.2.1 and is proved there.

Cosine law.

The above formulas easily imply the cosine law in $\mathbb{M}^2(\kappa)$, which can be expressed as follows

$$\cos\varphi = \begin{cases} \dfrac{b^2+c^2-a^2}{2\cdot b\cdot c} & \text{if } \kappa = 0 \\ \dfrac{\operatorname{cs}^\kappa a - \operatorname{cs}^\kappa b\cdot\operatorname{cs}^\kappa c}{\kappa\cdot\operatorname{sn}^\kappa b\cdot\operatorname{sn}^\kappa c} & \text{if } \kappa \neq 0 \end{cases} \qquad \frac{\operatorname{cs}^\kappa a - \operatorname{cs}^\kappa b\cdot\operatorname{cs}^\kappa c}{\kappa\cdot\operatorname{sn}^\kappa b\cdot\operatorname{sn}^\kappa c} \text{if} \quad \kappa \neq 0$$

However, rather than using these explicit formulas, we mainly will use the properties of $\measuredangle^\kappa$ and $\tilde{\Upsilon}^\kappa$ listed in 1.1.1.

1.2 Hemisphere lemma

1.2.1. Hemisphere lemma. *For $\kappa > 0$, any closed path of length $< 2\cdot\varpi^\kappa$ (respectively, $\leqslant 2\cdot\varpi^\kappa$) in $\mathbb{M}^2(\kappa)$ lies in an open (respectively, closed) hemisphere.*

Proof. By rescaling, we may assume that $\kappa = 1$ and thus $\varpi^\kappa = \pi$ and $\mathbb{M}^2(\kappa) = \mathbb{S}^2$. Let α be a closed curve in $\mathbb{S}^2$ of length $2\cdot\ell$.

Assume $\ell < \pi$. Let $\check{\alpha}$ be a subarc of α of length ℓ, with endpoints p and q. Since $|p-q| \leqslant \ell < \pi$, there is a unique geodesic $[pq]$ in $\mathbb{S}^2$. Let z be the midpoint of $[pq]$. We claim that α lies in the open hemisphere centered at z. If not, α intersects the boundary great circle in a point, say r. Without loss of generality we may assume that $r \in \check{\alpha}$. The arc $\check{\alpha}$ together with its reflection in z form a closed curve of length $2\cdot\ell$ that contains r and its antipodal point r'. Thus

$$\ell = \operatorname{length}\check{\alpha} \geqslant |r-r'| = \pi,$$

a contradiction.

If $\ell = \pi$, then either α is a local geodesic, and hence a great circle, or α may be strictly shortened by substituting a geodesic arc for a subarc of α whose endpoints p^1, p^2 are arbitrarily close to some point p on α. In the latter case, α lies in a closed hemisphere obtained as a limit of closures of open hemispheres containing the shortened curves as p^1, p^2 approach p. $\square$

1.2.2. Exercise. *Build a proof of Hemisphere lemma 1.2.1 based on Crofton's formula.*

Chapter 2

Metric spaces

In this chapter we fix some conventions, assuming that the reader is familiar with basic notions in metric geometry.

2.1 Metrics and their relatives

Definitions. Let $\mathbb{I}$ be a subinterval of $[0,\infty]$. A function ρ defined on $\mathcal{X}\times\mathcal{X}$ is called an $\mathbb{I}$-valued metric if the following conditions hold;

- ◇ $\rho(x,x)=0$ for any x;
- ◇ $\rho(x,y)\in\mathbb{I}$ for any pair $x\neq y$;
- ◇ $\rho(x,y)+\rho(x,z)\geqslant\rho(y,z)$ for any triple of points x,y,z.

The value $\rho(x,y)$ is also called the *distance* between x and y.

The above definition will be used for four choices of the interval $\mathbb{I}$: $(0,\infty)$, $(0,\infty]$, $[0,\infty]$ and $[0,\infty]$. Any $\mathbb{I}$-valued metric can be refereed briefly as a metric; the interval should be apparent from the context, but by default a metric is $(0,\infty)$-valued. If we need to be more specific, we may also use the following names:

- ◇ a $(0,\infty)$-valued metric may be called a *genuine metric*.
- ◇ a $(0,\infty]$-valued metric may be called an ∞-*metric*.
- ◇ a $[0,\infty)$-valued metric may be called a *genuine pseudometric*.
- ◇ A $[0,\infty]$-valued metric may be called a *pseudometric* or ∞-*pseudometric*.

A metric space is a set equipped with a metric; the distance between points x and y in a metric space $\mathcal{X}$ will usually be denoted by

$$|x-y| \qquad \text{or} \quad |x-y|_{\mathcal{X}};$$

the latter will be used if we need to emphasize that we are working in the space $\mathcal{X}$.

The function $\operatorname{dist}_x : \mathcal{X} \to \mathbb{R}$ defined as

$$\operatorname{dist}_x : y \mapsto |x - y|$$

will be called the *distance function* from x.

Any subset A in a metric space $\mathcal{X}$ will be also considered as a *subspace*; that is, a metric space with the metric defined by restricting the metric of $\mathcal{X}$ to $A \times A \subset \mathcal{X} \times \mathcal{X}$.

Zero and infinity. Genuine metric spaces are the main objects of study in this book. However, the generalizations above are useful in various definitions and constructions. For example, the construction of length-metric (see Section 2.3) uses infinite distances. The following definition gives another example.

2.1.1. Definition. *Assume $\{\mathcal{X}_\alpha\}_{\alpha\in\mathcal{A}}$ is a collection of ∞-metric spaces. The disjoint union*

$$\boldsymbol{X} = \bigsqcup_{\alpha\in\mathcal{A}} \mathcal{X}_\alpha$$

has a natural ∞-metric on it defined as follows: given two points $x \in \mathcal{X}_\alpha$ and $y \in \mathcal{X}_\beta$ set

$$\begin{aligned} |x-y|_{\boldsymbol{X}} &= \infty && \text{if } \alpha \neq \beta, \\ |x-y|_{\boldsymbol{X}} &= |x-y|_{\mathcal{X}_\alpha} && \text{if } \alpha = \beta. \end{aligned}$$

The resulting ∞-metric space $\boldsymbol{X}$ will be called the disjoint union *of $\{\mathcal{X}_\alpha\}_{\alpha\in\mathcal{A}}$, denoted by*

$$\bigsqcup_{\alpha\in\mathcal{A}} \mathcal{X}_\alpha.$$

Now let us give examples when vanishing distance between distinct points is useful:

Suppose a set $\mathcal{X}$ comes with a set of metrics $|*-*|_\alpha$ for $\alpha \in \mathcal{A}$. Then

$$|x-y| = \inf\{\, |x-y|_\alpha : \alpha \in \mathcal{A} \,\}.$$

is only a pseudometric; that is, if the metrics $|*-*|_\alpha$ are genuine, then $|*-*|$ has to be $[0,\infty)$-valued.

Let $\mathcal{X}$ be a set, $\mathcal{Y}$ be a metric space and $\Phi\colon \mathcal{X} \to \mathcal{Y}$ be a map. If Φ is not injective, then the *pullback*

$$|x-y|_{\mathcal{X}} = |\Phi(x) - \Phi(y)|_{\mathcal{Y}}$$

defines only a pseudometric on $\mathcal{X}$.

Corresponding metric space and metric component. The following two observations show that nearly any question about metric spaces can be reduced to a question about genuine metric spaces.

Assume $\mathcal{X}$ is a pseudometric space. Set $x \sim y$ if $|x-y|=0$. Note that if $x \sim x'$, then $|y-x| = |y-x'|$ for any $y \in \mathcal{X}$. Thus, $|*-*|$ defines a metric on the quotient set $\mathcal{X}/{\sim}$. In this way we obtain a metric space $\mathcal{X}'$. The space $\mathcal{X}'$ is called the *corresponding metric space* for the pseudometric space $\mathcal{X}$. Often we do not distinguish between $\mathcal{X}'$ and $\mathcal{X}$.

Set $x \approx y$ if and only if $|x-y| < \infty$; this is an other equivalence relation on $\mathcal{X}$. The equivalence class of a point $x \in \mathcal{X}$ will be called the *metric component* of x; it will be denoted as $\mathcal{X}_x$. One could think of $\mathcal{X}_x$ as $\mathrm{B}(x,\infty)_{\mathcal{X}}$, the open ball centered at x and radius ∞ in $\mathcal{X}$; see definition below.

It follows that any ∞-metric space is a *disjoint union* of genuine metric spaces, the metric components of the original ∞-metric space; see Definition 2.1.1

To summarize this discussion: Given a $[0,\infty]$-valued metric space $\mathcal{X}$, we may pass to the corresponding $(0,\infty]$-valued metric space $\mathcal{X}'$ and break the latter into a disjoint union of metric components, each of which is a genuine metric space.

2.2 Notations

Balls. Given $R \in [0,\infty]$ and a point x in a metric space $\mathcal{X}$, the sets

$$\begin{aligned}\mathrm{B}(x,R) &= \{\, y \in \mathcal{X} : |x-y| < R \,\},\\ \overline{\mathrm{B}}[x,R] &= \{\, y \in \mathcal{X} : |x-y| \leqslant R \,\}\end{aligned}$$

are called respectively the *open* and the *closed balls* of radius R with center x.

If we need to emphasize that these balls are taken in the space $\mathcal{X}$, we write

$$\mathrm{B}(x,R)_{\mathcal{X}} \quad \text{and} \quad \overline{\mathrm{B}}[x,R]_{\mathcal{X}}$$

correspondingly.

Since in a model space $\mathbb{M}^m(\kappa)$ all balls of the same radiuses are isometric; often will not need to specify the center of the ball, so we may write

$$\mathrm{B}(R)_{\mathbb{M}^m(\kappa)} \quad \text{and} \quad \overline{\mathrm{B}}[R]_{\mathbb{M}^m(\kappa)}$$

correspondingly.

A set $A \subset \mathcal{X}$ is called *bounded* if $A \subset \mathrm{B}(x, R)$ for some $x \in \mathcal{X}$ and $R < \infty$.

Distances to sets. For subset $A \subset \mathcal{X}$, let us denote the distance from A to a point x in $\mathcal{X}$ as $\operatorname{dist}_A x$; that is,

$$\operatorname{dist}_A x := \inf \{ |a - x| : a \in A \},$$

and we define the distance between sets A and B as

$$|A - B| := \inf \{ |a - b| : a \in A,\ b \in B \}.$$

For any subset $A \subset \mathcal{X}$, the sets

$$\mathrm{B}(A, R) = \{ y \in \mathcal{X} : \operatorname{dist}_A y < R \},$$
$$\overline{\mathrm{B}}[A, R] = \{ y \in \mathcal{X} : \operatorname{dist}_A y \leqslant R \}$$

are called respectively the *open* and *closed R-neighborhoods* of A.

Diameter, radius and packing. Let $\mathcal{X}$ be a metric space. Then the *diameter* of $\mathcal{X}$ is defined as

$$\operatorname{diam} \mathcal{X} = \sup \{ |x - y| : x, y \in \mathcal{X} \}.$$

The *radius* of $\mathcal{X}$ is defined as

$$\operatorname{rad} \mathcal{X} = \inf \{ R > 0 : \mathrm{B}(x, R) = \mathcal{X} \text{ for some } x \in \mathcal{X} \}.$$

The packing number *ε-pack* of $\mathcal{X}$ is the maximal number (possibly infinite) of points in $\mathcal{X}$ at distance $> \varepsilon$ from each other; it is denoted by $\operatorname{pack}_\varepsilon \mathcal{X}$. If $m = \operatorname{pack}_\varepsilon \mathcal{X} < \infty$, then a set $\{x^1, x^2, \dots, x^m\}$ in $\mathcal{X}$ such that $|x^i - x^j| > \varepsilon$ is called a *maximal ε-packing* in $\mathcal{X}$.

G-delta sets. Recall that arbitrary union of open balls in a metric space is called *open set*. A subset of a metric space is called G-delta set if it can be presented as an intersection of countable number of open subset. Often we will use the following classical result:

2.2.1. Baire's theorem. *Let $\mathcal{X}$ be a complete metric space and $\{\Omega_n\}$, $n \in \mathbb{N}$ be a collection of open dense subsets of $\mathcal{X}$. Then $\bigcap_{n=1}^{\infty} \Omega_n$ is dense in $\mathcal{X}$.*

Proper spaces. A metric space $\mathcal{X}$ is called *proper* if all closed bounded sets in $\mathcal{X}$ are compact. This condition is equivalent to each of the following statements:

1. For some (and therefore any) point $p \in \mathcal{X}$ and any $R < \infty$, the closed ball $\overline{\mathrm{B}}[p, R] \subset \mathcal{X}$ is compact.
2. The function $\operatorname{dist}_p : \mathcal{X} \to \mathbb{R}$ is proper for some (and therefore any) point $p \in \mathcal{X}$.

We will often use the following two classical statements:

2.2.2. Proposition. *Proper metric spaces are separable and second countable.*

2.2.3. Proposition. *Let $\mathcal{X}$ be a metric space. Then the following are equivalent*

i) *$\mathcal{X}$ is compact;*

ii) *$\mathcal{X}$ is* sequentially compact*; that is, any sequence of points in $\mathcal{X}$ contains a convergent subsequence;*

iii) *$\mathcal{X}$ is complete and for any $\varepsilon > 0$ there is a finite ε-net in $\mathcal{X}$; that is, there is a finite collection of points $p_1, \dots, p_N$ such that $\bigcup_i \mathrm{B}(p_i, \varepsilon) = \mathcal{X}$.*

2.3 Length spaces

2.3.1. Definition. *Let $\mathcal{X}$ be a metric space. Given a curve $\alpha\colon \mathbb{I} \to \mathcal{X}$ we define its* length *as*

$$\operatorname{length}\alpha := \sup_{a\leqslant t_0\leqslant\dots\leqslant t_n\leqslant b}\left\{\sum_{i\geqslant 1}|\alpha(t_i)-\alpha_{i-1}|\right\}$$

The following lemma is an easy exercise.

2.3.2. Lower semicontinuity of length. *Assume $\alpha_n\colon \mathbb{I}\to\mathcal{X}$ is a sequence of curves that converges pointwise to a curve $\alpha_\infty\colon \mathbb{I}\to\mathcal{X}$. Then*

$$\operatorname{length}\alpha_\infty \leqslant \varliminf_{n\to\infty}\operatorname{length}\alpha_n.$$

Given two points x and y in a metric space $\mathcal{X}$ consider the value

$$\|x-y\| = \inf_\alpha\{\operatorname{length}\alpha\},$$

where infimum is taken for all paths α from x to y.

It is easy to see that $\|*-*\|$ defines a $(0,\infty]$-valued metric on $\mathcal{X}$; it will be called the *length-metric* on $\mathcal{X}$. Clearly

$$\|x-y\| \geqslant |x-y|$$

for any $x, y \in \mathcal{X}$.

2.3.3. Definition. *If $\|x-y\| = |x-y|$ for any pair of points x, y in a metric space $\mathcal{X}$, then $\mathcal{X}$ is called a* length space.

In other words, a metric space $\mathcal{X}$ is a *length space* if for any $\varepsilon > 0$ and any two points $x, y \in \mathcal{X}$ with $|x-y| < \infty$ there is a path $\alpha\colon [0,1] \to \mathcal{X}$ connecting[1] x to y such that

$$\operatorname{length} \alpha < |x-y| + \varepsilon.$$

In this book, most of the time we consider length spaces. If $\mathcal{X}$ is length space, and $A \subset \mathcal{X}$. The set A comes with the inherited metric from $\mathcal{X}$ which might be not a length-metric. The corresponding length-metric on A will be denoted as $|*-*|_A$.

Variations of the definition. We will need the following variations of Definition 2.3.3:

- ⋄ Assume $R > 0$. If $\|x-y\| = |x-y|$ for any pair $|x-y| < R$, then $\mathcal{X}$ is called an *R-length space*.
- ⋄ If any point in $\mathcal{X}$ admits a neighborhood Ω such that $\|x-y\| = |x-y|$ for any pair of points $x, y \in \Omega$ then $\mathcal{X}$ is called a *locally length space*.
- ⋄ A metric space $\mathcal{X}$ is called *geodesic* if for any two points $x, y \in \mathcal{X}$ with $|x-y| < \infty$ there is a geodesic $[xy]$ in $\mathcal{X}$.
- ⋄ Assume $R > 0$. A metric space $\mathcal{X}$ is called *R-geodesic* if for any two points $x, y \in \mathcal{X}$ such that $|x-y| < R$ there is a geodesic $[xy]$ in $\mathcal{X}$.

Note that the notions of ∞-length spaces and length spaces are the same. Clearly, any geodesic space is a length space and any R-geodesic space is R-length.

2.3.4. Example. *Let $\mathcal{X}$ be obtained by gluing a countable collection of disjoint intervals $\mathbb{I}_n$ of length $1 + \frac{1}{n}$ where for each $\mathbb{I}_n$ one end is glued to p and the other to q. Then $\mathcal{X}$ carries a natural complete length metric such that $|p-q| = 1$, but there is no geodesic connecting p to q.*

2.3.5. Exercise. *Let $\mathcal{X}$ be a metric space and $\|*-*\|$ is the length metric on it. Show the following*

a) If $\mathcal{X}$ is complete, then $(\mathcal{X}, \|*-*\|)$ is complete.

b) If $\mathcal{X}$ is compact, then $(\mathcal{X}, \|*-*\|)$ is geodesic.

2.3.6. Exercise. *Give an example of a complete length space such that no pair of distinct points can be joined by a geodesic.*

2.3.7. Definition. *Let $\mathcal{X}$ be a metric space and $x, y \in \mathcal{X}$.*

(i) A point $z \in \mathcal{X}$ is called a midpoint *of x and y if*

$$|x-z| = |y-z| = \tfrac{1}{2}\cdot|x-y|.$$

[1] That is, such that $\alpha(0) = x$ and $\alpha(1) = y$.

(ii) Assume $\varepsilon \geqslant 0$. A point $z \in \mathcal{X}$ is called ε-midpoint *of x and y if*

$$|x-z|, \quad |y-z| \leqslant \tfrac{1}{2}\cdot|x-y| + \varepsilon.$$

Note that a 0-midpoint is the same as a midpoint.

2.3.8. Lemma. *Let $\mathcal{X}$ be a complete metric space.*

a) Assume that for any pair of points $x, y \in \mathcal{X}$ and any $\varepsilon > 0$ there is a ε-midpoint z. Then the space $\mathcal{X}$ is a length space.

b) Assume that for any pair of points $x, y \in \mathcal{X}$ there is a midpoint z. Then the space $\mathcal{X}$ is a geodesic space.

c) If for some $R > 0$, the assumptions (a) or (b) hold only for pairs of points $x, y \in \mathcal{X}$ such that $|x-y| < R$, then the space $\mathcal{X}$ is a R-length or R-geodesic correspondingly.

Proof. Fix a pair of points $x, y \in \mathcal{X}$. Set $\varepsilon_n = \frac{\varepsilon}{2^{2\cdot n}}$, $\alpha(0) = x$ and $\alpha(1) = y$.

Set $\alpha(\frac{1}{2})$ to be an ε_1-midpoint of $\alpha(0)$ and $\alpha(1)$. Further, set $\alpha(\frac{1}{4})$ and $\alpha(\frac{3}{4})$ to be ε_2-midpoints for the pairs $(\alpha(0), \alpha(\frac{1}{2})$ and $(\alpha(\frac{1}{2}), \alpha(1)$ respectively. Applying the above procedure recursively, on the n-th step we define $\alpha(\frac{k}{2^n})$ for every odd integer k such that $0 < \frac{k}{2^n} < 1$, as an ε_n-midpoint of the already defined $\alpha(\frac{k-1}{2^n})$ and $\alpha(\frac{k+1}{2^n})$.

In this way we define $\alpha(t)$ for all dyadic rationals t in $[0,1]$. If $t \in$ $\in [0,1]$ is not a dyadic rational, consider a sequence of dyadic rationals $t_n \to t$ as $n \to \infty$. Note that the sequence $\alpha(t_n)$ converges; define $\alpha(t)$ as its limit. It is easy to see that $\alpha(t)$ does not depend on the choice of the sequence t_n and $\alpha\colon [0,1] \to \mathcal{X}$ is a path from x to y. Moreover,

$$\begin{aligned} \text{length}\,\alpha &\leqslant |x-y| + \sum_{n=1}^{\infty} 2^{n-1}\cdot\varepsilon_n \leqslant \\ &\leqslant |x-y| + \tfrac{\varepsilon}{2}. \end{aligned} \tag{1}$$

Since $\varepsilon > 0$ is arbitrary, we get that (a).

To prove (b), one should repeat the same argument taking midpoints instead of ε_n-midpoints. In this case ❶ holds for $\varepsilon_n = \varepsilon = 0$.

The proof of (2.3.8c) obtained by a straightforward modifications of the proofs above. □

2.3.9. Corollary. *A proper length space is geodesic.*

It follows from Lemma 2.3.8, since in a compact set a sequence of $\frac{1}{n}$-midpoints z_n contains a convergent subsequence.

2.3.10. Hopf–Rinow theorem. *Any complete, locally compact length space is proper.*

Proof. Let $\mathcal{X}$ be a locally compact length space. Given $x \in \mathcal{X}$, denote by $\rho(x)$ the supremum of all $R > 0$ such that the closed ball $\overline{\mathrm{B}}[x, R]$ is compact. Since $\mathcal{X}$ is locally compact

❷ $$\rho(x) > 0 \quad \text{for any} \quad x \in \mathcal{X}.$$

It is sufficient to show that $\rho(x) = \infty$ for some (and therefore any) point $x \in \mathcal{X}$.

Assume the contrary; that is, $\rho(x) < \infty$.

❸ $B = \overline{\mathrm{B}}[x, \rho(x)]$ *is compact for any* x.

Indeed, $\mathcal{X}$ is a length space; therefore for any $\varepsilon > 0$, the set $\overline{\mathrm{B}}[x, \rho(x) - \varepsilon]$ is a compact ε-net in B. Since B is closed and hence complete, it has to be compact by Proposition 2.2.3. △

❹ $|\rho(x) - \rho(y)| \leqslant |x - y|_{\mathcal{X}}$, *in particular* $\rho\colon \mathcal{X} \to \mathbb{R}$ *is a continuous function.*

Indeed, assume the contrary; that is, $\rho(x) + |x - y| < \rho(y)$ for some $x, y \in \mathcal{X}$. Then $\overline{\mathrm{B}}[x, \rho(x) + \varepsilon]$ is a closed subset of $\overline{\mathrm{B}}[y, \rho(y)]$ for some $\varepsilon > 0$. Then compactness of $\overline{\mathrm{B}}[y, \rho(y)]$ implies compactness of $\overline{\mathrm{B}}[x, \rho(x) + \varepsilon]$, a contradiction. △

Set $\varepsilon = \min_{y \in B}\{\rho(y)\}$; the minimum is defined since B is compact. From ❷, we have $\varepsilon > 0$.

Choose a finite $\frac{\varepsilon}{10}$-net $\{a_1, a_2, \dots, a_n\}$ in B. The union W of the closed balls $\overline{\mathrm{B}}[a_i, \varepsilon]$ is compact. Clearly $\overline{\mathrm{B}}[x, \rho(x) + \frac{\varepsilon}{10}] \subset W$. Therefore $\overline{\mathrm{B}}[x, \rho(x) + \frac{\varepsilon}{10}]$ is compact; a contradiction. □

2.3.11. Exercise. *Construct a geodesic space that is locally compact, but whose completion is neither geodesic nor locally compact.*

2.4 Convex sets

2.4.1. Definition. *Let $\mathcal{X}$ be a geodesic space and $A \subset \mathcal{X}$.*

We say that A convex *if for every two points $p, q \in A$ any geodesic $[pq]$ lies in A.*

We say that A weakly convex *if for every two points $p, q \in A$ there is a geodesic $[pq]$ that lies in A.*

We say that A is totally convex *if for every two points $p, q \in A$, every local geodesic from p to q lies in A.*

If for some $R \in (0, \infty]$ these definitions are applied only for pairs of points such that $|p - q| < R$ and only for the geodesics of length $< R$ then A is called correspondingly R-convex, weakly R-convex *and* totally R-convex *.*

A set $A \subset \mathcal{X}$ is called locally convex *if every point $a \in A$ admits an open neighborhood $\Omega \ni a$ such that for every two points $p, q \in A \cap \Omega$ every geodesic $[pq] \subset \Omega$ lies in A. Similarly one defines locally weakly convex and locally totally convex sets.*

Remarks. Let us state few observations from the definition.

- ◇ The notion of *(weakly) convex set* is the same as *(weakly) ∞-convex set.*
- ◇ The inherited metric on a weakly convex set coincides with its length-metric.
- ◇ Any open set is locally convex by definition.

The following proposition states that weak convexity survive under ulralimit. An analogus statement about convexity does not hold; for example there is a sequence of convex discs in $\mathbb{S}^2$ that converges to the half sphere $\mathbb{S}^2_+$, which is not convex.

2.4.2. Proposition. *Let $\mathcal{X}_n$, be a sequence of geodesic spaces. Assume $A_n \subset \mathcal{X}_n$ be a sequence of weakly convex sets and $\mathcal{X}_n \to \mathcal{X}_\omega$ and $A_n \to A_\omega \subset \mathcal{X}_\omega$ as $n \to \omega$. Then A_ω is a weakly convex set of $\mathcal{X}_\omega$.*

Proof. Fix $x_\omega, y_\omega \in A_\omega$. Consider the sequences $x_n, y_n \in A_n$ such that $x_n \to x_\omega$ and $y_n \to y_\omega$ as $n \to \omega$.

Denote by α_n a geodesic path from x_n to y_n that lies in A_n. Set

$$\alpha_\omega(t) = \lim_{n \to \omega} \alpha_n(t).$$

Note that α_ω is a geodesic path that lies in A_ω. Hence the result follows. □

2.5 Quotient spaces

Quotient spaces. Assume $\mathcal{X}$ is a metric space with an equivalence relation $\sim$. Note that given two pseudometrics ρ_1 and ρ_2 on $\mathcal{X}/\sim$ their maximum

$$\rho(x, y) = \max\{\rho_1(x, y), \rho_2(x, y)\}$$

is also a pseudometric; also if for these two pseudometric ρ_1 and ρ_2 the projections $\mathcal{X} \to (\mathcal{X}/\sim, \rho_i)$ are short, then so it true for $\rho(x, y)$.

It follows that the quotient space $\mathcal{X}/{\sim}$ admits natural pseudometric; this is the maximal pseudometric on $\mathcal{X}/{\sim}$ that makes the quotient map $\mathcal{X} \to \mathcal{X}/{\sim}$ to be short. The corresponding metric space will be also denoted as $\mathcal{X}/{\sim}$ and will be called the *quotient space* of $\mathcal{X}$ by the equivalence relation $\sim$.

In general case the points of the metric space $\mathcal{X}/{\sim}$ are formed by equivalence classes in $\mathcal{X}$ for a wider equivalence relation. However, in the most of the cases we will consider, the set of equivalence classes will coincide with the set of points in the metric space $\mathcal{X}/{\sim}$.

2.5.1. Proposition. *Let $\mathcal{X}$ be a length space and $\sim$ be an equivalence relation on $\mathcal{X}$, then $\mathcal{X}/{\sim}$ is a length space.*

Proof. Let $\mathcal{Y}$ be arbitrary metric space. Since $\mathcal{X}$ is a length space, the map $f\colon \mathcal{X} \to \mathcal{Y}$ is short if and only if

$$\operatorname{length}(f \circ \alpha) \leqslant \operatorname{length} \alpha$$

for any curve $\alpha\colon \mathbb{I} \to \mathcal{X}$.

Denote by $\|{*}-{*}\|$ the length-metric on $\mathcal{Y}$. It follows that if $f\colon \mathcal{X} \to \mathcal{Y}$ is short then so is $f\colon \mathcal{X} \to (\mathcal{Y}, \|{*}-{*}\|)$.

Consider the quotient map $f\colon \mathcal{X} \to \mathcal{X}/{\sim}$. Recall that the space $\mathcal{X}/{\sim}$ is defined by the maximal pseudometric that makes f to be short.

Denote by $\|{*}-{*}\|$ the length-metric on $\mathcal{X}/{\sim}$. It follows that

$$f\colon \mathcal{X} \to (\mathcal{X}/{\sim}, \|{*}-{*}\|)$$

is also short.

Note that

$$\|x-y\| \geqslant |x-y|_{\mathcal{X}/\sim}$$

for any $x, y \in \mathcal{X}/{\sim}$. From maximality of $|{*}-{*}|_{\mathcal{X}/\sim}$, we get

$$\|x-y\| = |x-y|_{\mathcal{X}/\sim}$$

for any $x, y \in \mathcal{X}/{\sim}$; that is, $\mathcal{X}/{\sim}$ is a length space. $\square$

Group actions. Assume a group G acts on a metric space $\mathcal{X}$. Consider relation $\sim$ on $\mathcal{X}$ defined as $x \sim y$ if there is $g \in G$ such that $x = g{\cdot}y$. Note that $\sim$ is an equivalence relation.

In this case the quotient space $\mathcal{X}/{\sim}$ will be also denoted as $\mathcal{X}/G$; it could be also regarded as space of G-orbits in $\mathcal{X}$.

Assume that the group G acts on $\mathcal{X}$ by isometries. In this case the distance between orbits $G{\cdot}x$ and $G{\cdot}y$ in $\mathcal{X}/G$ can be defined directly

$$|G{\cdot}x - G{\cdot}y|_{\mathcal{X}/G} = \inf\left\{\, |x - g{\cdot}y|_{\mathcal{X}} = |g^{-1}{\cdot}x - y|_{\mathcal{X}} \,:\, g \in G \,\right\}.$$

If the G-orbits are closed, then $|G{\cdot}x - G{\cdot}y|_{\mathcal{X}/G} = 0$ if and only if $G{\cdot}x = G{\cdot}y$. In this case the quotient space $\mathcal{X}/G$ is a genuine metric space.

The following proposition follows from the definition of quotient space:

2.5.2. Proposition. *Assume $\mathcal{X}$ is metric space and group G acts on $\mathcal{X}$ by isometries. Then the projection $\mathcal{X} \to \mathcal{X}/G$ is a submetry; that is $\mathcal{X} \to \mathcal{X}/G$ is short and 1-co-Lipschitz.*

2.6 Gluing and doubling

Gluing. Recall that disjoint union of metric spaces can be also considered as a metric space; see Definition 2.1.1. Therefore the quotient-space construction works as well for an equivalence relation on a disjoint union of metric spaces.

Consider two metric spaces $\mathcal{X}_1$ and $\mathcal{X}_2$ with subset $A_1 \subset \mathcal{X}_1$ and $A_2 \subset \mathcal{X}_2$ and a bijection $\varphi\colon A_1 \to A_2$. Consider the minimal equivalence relation on $\mathcal{X}_1 \sqcup \mathcal{X}_2$ such that $a \sim \varphi(a)$ for any $a \in A_1$. In this case, the corresponding quotient space $(\mathcal{X}_1 \sqcup \mathcal{X}_2)/{\sim}$ will be called *gluing* of $\mathcal{X}$ and $\mathcal{Y}$ along φ and denoted as

$$\mathcal{X}_1 \sqcup_\varphi \mathcal{X}_2.$$

Note that if the map $\varphi\colon A_1 \to A_2$ is distance preserving, then the projections $\iota_i\colon \mathcal{X}_i \to \mathcal{X}_1 \sqcup_\varphi \mathcal{X}_2$ are also distance preserving and

$$|\iota_1(x_1) - \iota_2(x_2)|_{\mathcal{X}_1 \sqcup_\varphi \mathcal{X}_2} = \inf_{a_2 = \varphi(a_1)} \{\, |x_1 - a_1|_{\mathcal{X}_1} + |x_2 - a_2|_{\mathcal{X}_2} \,\}$$

for any $x_1 \in \mathcal{X}_1$ and $x_2 \in \mathcal{X}_2$.

Doubling. Let $\mathcal{V}$ be a metric space and $A \subset \mathcal{V}$ be a closed subset. A metric space $\mathcal{W}$ glued from two copies of $\mathcal{V}$ along A is called *doubling of $\mathcal{V}$ in A*.

The space $\mathcal{W}$ is completely described by the following properties:

- The space $\mathcal{W}$ contains $\mathcal{V}$ as a subspace; in particular the set A can be treated as a subset of $\mathcal{W}$.
- There is an isometric involution of $\mathcal{W}$ which is called *reflection in A*; further it will be denoted as $x \mapsto x'$.
- For any $x \in \mathcal{W}$ we have $x \in \mathcal{V}$ or $x' \in \mathcal{V}$ and

  $$|x' - y|_{\mathcal{W}} = |x - y'|_{\mathcal{W}} = \inf_{a \in A} \{|x - a|_{\mathcal{V}} + |a - y|_{\mathcal{V}}\}.$$

  for any $x, y \in \mathcal{V}$.

The image of $\mathcal{V}$ under the reflection in A will be denoted by $\mathcal{V}'$. The subspace $\mathcal{V}'$ is an isometric copy of $\mathcal{V}$. Clearly $\mathcal{V} \cup \mathcal{V}' = \mathcal{W}$ and $\mathcal{V} \cap \mathcal{V}' = A$. Moreover $a = a' \iff a \in A$.

The following proposition follows directly from the definitions.

2.6.1. Proposition. *Assume $\mathcal{W}$ is the doubling of metric space $\mathcal{V}$ in its closed subset A. Then*

a) If $\mathcal{V}$ is a complete length space, then so is $\mathcal{W}$.

b) If $\mathcal{V}$ is proper, then so is $\mathcal{W}$. In this case, for any $x, y \in \mathcal{V}$ there is $a \in A$ such that

$$|x - a|_{\mathcal{V}} + |a - y|_{\mathcal{V}} = |x - y'|_{\mathcal{W}}.$$

c) Given $x \in \mathcal{W}$ set $\bar{x} = x$ if $x \in \mathcal{V}$ and $\bar{x} = x'$ otherwise. The map $\mathcal{W} \to \mathcal{V}$ defined by $x \mapsto \bar{x}$ is short and length-preserving. In particular if γ is a geodesic in $\mathcal{W}$ with ends in $\mathcal{V}$, then $\bar{\gamma}$ is a geodesic in $\mathcal{V}$ with the same ends.

2.7 Kuratowsky embedding

Given a metric space $\mathcal{X}$, let us denote by $\operatorname{Bnd}(\mathcal{X}, \mathbb{R})$ the space of all bounded functions on $\mathcal{X}$ equipped with the sup-norm,

$$\|f\| = \sup_{x \in \mathcal{X}} \{|f(x)|\}.$$

Kuratowski embedding. Given a point $p \in \mathcal{X}$, consider map $\operatorname{kur}_p \colon \mathcal{X} \to \operatorname{Bnd}(\mathcal{X}, \mathbb{R})$, defined as $\operatorname{kur}_p x = \operatorname{dist}_x - \operatorname{dist}_p$. The map kur_p will be called *Kuratowski map at p*.

From the triangle inequality, we have

$$\|\operatorname{kur}_p x - \operatorname{kur}_p y\| = \sup_{z \in \mathcal{X}} \{||x - z| - |y - z||\} = |x - y|.$$

Therefore, for any $p \in \mathcal{X}$, the Kuratowski map gives a distance preserving map $\operatorname{kur}_p \colon \mathcal{X} \hookrightarrow \operatorname{Bnd}(\mathcal{X}, \mathbb{R})$. Thus, we can (and often will) consider space $\mathcal{X}$ as a subset of $\operatorname{Bnd}(\mathcal{X}, \mathbb{R})$.

2.7.1. Exercise. *Show that any compact space is isometric to a subspace in a compact length space.*

Chapter 3

Ultralimits

Here we introduce ultralimits of sequences of points, metric spaces and functions. The ultralimits of metric spaces can be considered as a variation of Gromov–Hausdorff convergence. Our presentation is based on [1].

In this book, the use of ultralimits is very limited; we use them only as a canonical way to pass to a convergent subsequence. (In principle, we could avoid selling our souls to the set-theoretical devil, but in this case we must say "pass to convergent subsequence" too many times.)

3.1 Ultrafilters

We will need the existence of a selective ultrafilter ω, which we fix once and for all. Existence follows from the axiom of choice and continuum hypothesis.

Recall that $\mathbb{N}$ denotes the set of natural numbers, $\mathbb{N} = \{1, 2, \dots\}$

3.1.1. Definition. *A finitely additive measure ω on $\mathbb{N}$ is called an* ultrafilter *if it satisfies*

a) $\omega(S) = 0$ *or* 1 *for any subset* $S \subset \mathbb{N}$.

An ultrafilter ω is called nonprinciple *if in addition*

b) $\omega(F) = 0$ *for any finite subset* $F \subset \mathbb{N}$.

A nonprinciple ultrafilter ω is called selective *if in addition*

c) for any partition of $\mathbb{N}$ into sets $\{C_\alpha\}_{\alpha\in\mathcal{A}}$ such that $\omega(C_\alpha) = 0$ for each α, there is a set $S \subset \mathbb{N}$ such that $\omega(S) = 1$ and $S \cap C_\alpha$ is a one-point set for each $\alpha \in \mathcal{A}$.

If $\omega(S) = 0$ for some subset $S \subset \mathbb{N}$, we say that S is ω-*small*. If $\omega(S) = 1$, we say that S contains ω-*almost all* elements of $\mathbb{N}$.

Classical definition. More commonly, a nonprinciple ultrafilter is defined as a collection, say $\mathfrak{F}$, of sets in $\mathbb{N}$ such that

1. if $P \in \mathfrak{F}$ and $Q \supset P$, then $Q \in \mathfrak{F}$,
2. if $P, Q \in \mathfrak{F}$, then $P \cap Q \in \mathfrak{F}$,
3. for any subset $P \subset \mathbb{N}$, either P or its complement is an element of $\mathfrak{F}$.
4. if $F \subset \mathbb{N}$ is finite, then $F \notin \mathfrak{F}$.

Setting $P \in \mathfrak{F} \Leftrightarrow \omega(P) = 1$ makes these two definitions equivalent.

A nonempty collection of sets $\mathfrak{F}$ that does not include the empty set and satisfies only conditions 1 and 2 is called a *filter*; if in addition $\mathfrak{F}$ satisfies Condition 3 it is called an *ultrafilter*. From Zorn's lemma, it follows that every filter contains an ultrafilter. Thus there is an ultrafilter $\mathfrak{F}$ contained in the filter of all complements of finite sets; clearly this $\mathfrak{F}$ is nonprinciple.

The existence of a selective ultrafilter follows from the continuum hypothesis; it was proved by Walter Rudin in [2].

Stone–Čech compactification. Given a set $S \subset \mathbb{N}$, consider subset Ω_S of all ultrafilters ω such that $\omega(S) = 1$. It is straightforward to check that the sets Ω_S for all $S \subset \mathbb{N}$ form a topology on the set of ultrafilters on $\mathbb{N}$. The obtained space is called *Stone–Čech compactification* of $\mathbb{N}$; it is usually denoted as $\beta\mathbb{N}$.

There is a natural embedding $\mathbb{N} \hookrightarrow \beta\mathbb{N}$ defined as $n \mapsto \omega_n$, where ω_n is the principle ultrafilter such that $\omega_n(S) = 1$ if and only if $n \in S$. Using the described embedding, we can (and will) consider $\mathbb{N}$ as a subset of $\beta\mathbb{N}$.

The space $\beta\mathbb{N}$ is the maximal compact Hausdorff space that contains $\mathbb{N}$ as an everywhere dense subset. More precisely, for any compact Hausdorff space $\mathcal{X}$ and a map $f\colon \mathbb{N} \to \mathcal{X}$ there is unique continuous map $\bar{f}\colon \beta\mathbb{N} \to X$ such that the restriction $\bar{f}|_{\mathbb{N}}$ coincides with f.

3.2 Ultralimits of points

Fix an ultrafilter ω. Assume (x_n) is a sequence of points in a metric space $\mathcal{X}$. Let us define the ω-*limit* of (x_n) as the point x_ω such that for any $\varepsilon > 0$, ω-almost all elements of (x_n) lie in $\mathrm{B}(x_\omega, \varepsilon)$; that is,

$$\omega\{\, n \in \mathbb{N} : |x_\omega - x_n| < \varepsilon \,\} = 1.$$

In this case, we will write

$$x_\omega = \lim_{n\to\omega} x_n \quad \text{or} \quad x_n \to x_\omega \text{ as } n \to \omega.$$

For example if ω is the principle ultrafilter such that $\omega(\{n\}) = 1$ for some $n \in \mathbb{N}$, then $x_\omega = x_n$.

Note that ω-limits of a sequence and its subsequence may differ. For example, in general

$$\lim_{n\to\omega} x_n \neq \lim_{n\to\omega} x_{2\cdot n}.$$

3.2.1. Proposition. *Let ω be a nonprinciple ultrafilter. Assume (x_n) is a sequence of points in a metric space $\mathcal{X}$ and $x_n \to x_\omega$ as $n \to \omega$. Then x_ω is a partial limit of the sequence (x_n); that is, there is a subsequence $(x_n)_{n\in S}$ that converges to x_ω in the usual sense.*

Moreover, if ω is selective, then the subsequence $(x_n)_{n\in S}$ can be chosen so that $\omega(S) = 1$.

Proof. Given $\varepsilon > 0$, set $S_\varepsilon = \{\, n \in \mathbb{N} : |x_n - x_\omega| < \varepsilon \,\}$.

Note that $\omega(S_\varepsilon) = 1$ for any $\varepsilon > 0$. Since ω is nonprinciple, the set S_ε is infinite. Therefore we can choose an increasing sequence (n_k) such that $n_k \in S_{\frac{1}{k}}$ for each $k \in \mathbb{N}$. Clearly $x_{n_k} \to x_\omega$ as $k \to \infty$.

Now assume that ω is selective. Consider the sets

$$C_k = \left\{\, n \in \mathbb{N} : \tfrac{1}{n} < |x_n - x_\omega| \leqslant \tfrac{1}{n-1} \,\right\},$$

where we assume $\frac{1}{0} = \infty$ and the set

$$C_\infty = \{\, n \in \mathbb{N} : x_n = x_\omega \,\},$$

Note that $\omega(C_k) = 0$ for any $k \in \mathbb{N}$.

If $\omega(C_\infty) = 1$, we can take the sequence $(x_n)_{n\in C_\infty}$.

Otherwise discarding all empty sets among C_k and C_∞ gives a partition of $\mathbb{N}$ into countable collection of ω-small sets. Since ω is selective, we can choose a set $S \subset \mathbb{N}$ such that S meets each set of the partition at one point and $\omega(S) = 1$. Clearly the sequence $(x_n)_{n\in S}$ converges to x_ω in the usual sense. $\square$

The following proposition is analogous to the statement that any sequence in a compact metric space has a convergent subsequence; it can be proved the same way.

3.2.2. Proposition. *Let $\mathcal{X}$ be a compact metric space. Then any sequence of points (x_n) in $\mathcal{X}$ has unique ω-limit x_ω.*

In particular, a bounded sequence of real numbers has a unique ω-limit.

The following lemma is an ultralimit analog of Cauchy convergence test.

3.2.3. Lemma. *Let (x_n) be a sequence of points in a complete space $\mathcal{X}$. Assume for each subsequence (y_n) of (x_n), the ω-limit*

$$y_\omega = \lim_{n\to\omega} y_n \in \mathcal{X}$$

is defined and does not depend on the choice of subsequence, then the sequence (x_n) converges in the usual sense.

Proof. Assume that (x_n) does not converge in itself. Then for some $\varepsilon > 0$, there is a subsequence (y_n) of (x_n) such that $|x_n - y_n| \geqslant \varepsilon$ for all n.

It follows that $|x_\omega - y_\omega| \geqslant \varepsilon$, a contradiction. □

3.3 Ultralimits of spaces

From now on, ω denotes a selective ultrafilter on the set of natural numbers.

Let $\mathcal{X}_n$ be a sequence of metric spaces $\mathcal{X}_n$. Consider all sequences $x_n \in \mathcal{X}_n$. On the set of all such sequences, define a pseudometric on these subsequences by

❶ $$|(x_n) - (y_n)| = \lim_{n\to\omega} |x_n - y_n|.$$

Note that the ω-limit on the right hand side is always defined and takes a value in $[0, \infty]$.

Set $\mathcal{X}_\omega$ to be the corresponding metric space; that is, the underlying set of $\mathcal{X}_\omega$ is formed by classes of equivalence of sequences of points $x_n \in \in \mathcal{X}_n$ defined by

$$(x_n) \sim (y_n) \;\Leftrightarrow\; \lim_{n\to\omega} |x_n - y_n| = 0$$

and the distance is defined as in ❶.

The space $\mathcal{X}_\omega$ is called ω-*limit* of $\mathcal{X}_n$. Typically $\mathcal{X}_\omega$ will denote the ω-limit of sequence $\mathcal{X}_n$; we may also write

$$\mathcal{X}_n \to \mathcal{X}_\omega \;\text{ as }\; n \to \omega \;\text{ or }\; \mathcal{X}_\omega = \lim_{n\to\omega} \mathcal{X}_n.$$

Given a sequence $x_n \in \mathcal{X}_n$, we will denote by x_ω its equivalence class which is a point in $\mathcal{X}_\omega$; equivalently we will write

$$x_n \to x_\omega \;\text{ as }\; n \to \omega \;\text{ or }\; x_\omega = \lim_{n\to\omega} x_n.$$

3.3.1. Observation. *The ω-limit of any sequence of metric spaces is complete.*

Proof. Let $\mathcal{X}_n$ be a sequence of metric spaces and $\mathcal{X}_n \to \mathcal{X}_\omega$ as $n \to \omega$.

Fix a sequence $x_m \in \mathcal{X}_\omega$ converging in itself. Passing to a subsequence we can assume that $|x_m - x_{m-1}|_{\mathcal{X}_\omega} < \frac{1}{2^m}$ for any m.

Let us choose double sequence $x_{n,m} \in \mathcal{X}_n$ such that for any fixed m we have $x_{n,m} \to x_m$ as $n \to \omega$.

Set $S_1 = \mathbb{N}$ and

$$S_m = \left\{ n \in \mathbb{N} : |x_{n,k} - x_{n,k-1}| < \tfrac{1}{2^k} \quad \text{for} \quad k \leqslant m \right\}.$$

For each n consider the largest value $m(n)$ such that $m(n) \leqslant n$ and $m(n) \in S_m$. Consider the sequence $y_n = x_{n,m(n)}$; denote by $y \in \mathcal{X}^\omega$ its ω-limit. By construction $x_n \to y$ as $n \to \infty$. Hence the statement follows. □

3.3.2. Observation. *The ω-limit of any sequence of length spaces is geodesic.*

Proof. If $\mathcal{X}_n$ is a sequence length spaces, then for any sequence of pairs $x_n, y_n \in X_n$ there is a sequence of $\frac{1}{n}$-midpoints z_n.

Let $x_n \to x_\omega$, $y_n \to y_\omega$ and $z_n \to z_\omega$ as $n \to \omega$. Note that z_ω is a midpoint of x_ω and y_ω in $\mathcal{X}^\omega$.

By Observation 3.3.1, $\mathcal{X}^\omega$ is complete. Applying Lemma 2.3.8 we get the statement. □

Ultrapower. If all the metric spaces in the sequence are identical $\mathcal{X}_n = \mathcal{X}$, its ω-limit $\lim_{n\to\omega} \mathcal{X}_n$ is denoted by $\mathcal{X}^\omega$ and called ω-power of $\mathcal{X}$.

According to Theorem 3.5.1, there is a distance preserving map $\iota : \mathcal{X} \hookrightarrow \mathcal{X}^\omega$, here $\iota(y)$ is the equivalence class of constant sequence $y_n = y$.

The image $\iota(\mathcal{X})$ might be proper subset of $\mathcal{X}^\omega$. For example $\mathbb{R}^\omega$ has pairs of points on distance ∞ from each other; at the same time, each metric component of $\mathbb{R}^\omega$ is isometric to $\mathbb{R}$.

According to Theorem 3.5.1, if $\mathcal{X}$ is compact, then $\iota(\mathcal{X}) = \mathcal{X}^\omega$; in particular, $\mathcal{X}^\omega$ is isometric to $\mathcal{X}$. If $\mathcal{X}$ is proper, then $\iota(\mathcal{X})$ forms a metric component of $\mathcal{X}^\omega$.

The embedding ι gives us right to treat $\mathcal{X}$ as a subset of its ultrapower $\mathcal{X}^\omega$.

3.3.3. Observation. *Let $\mathcal{X}$ be a complete metric space. Then $\mathcal{X}^\omega$ is geodesic space if and only if $\mathcal{X}$ is a length space.*

Proof. Assume $\mathcal{X}^\omega$ is geodesic space. Then any pair of points $x, y \in \mathcal{X}$ has a midpoint $z_\omega \in \mathcal{X}^\omega$. Fix a sequence of points $z_n \in \mathcal{X}$ such that $z_n \to z_\omega$ as $n \to \omega$.

Note that $|x - z_n|_{\mathcal{X}} \to \frac{1}{2}\cdot|x - y|_{\mathcal{X}}$ and $|y - z_n|_{\mathcal{X}} \to \frac{1}{2}\cdot|x - y|_{\mathcal{X}}$ as $n \to \omega$. In particular, for any $\varepsilon > 0$, the point z_n is an ε-midpoint of x and y for ω-almost all n. It remains to apply Lemma 2.3.8.

The "if"-part follows from Observation 3.3.2. □

3.3.4. Corollary. *Assume $\mathcal{X}$ is a complete length space and $p, q \in$ $\in \mathcal{X}$ cannot be joined by a geodesic in $\mathcal{X}$. Then there are at least two distinct geodesics between p and q in the ultrapower $\mathcal{X}^\omega$.*

3.4 Ultratangent space

Recall that we assume that ω is a once for all fixed choice of a selective ultrafilter.

For a metric space $\mathcal{X}$ and a positive real number $\mathfrak{z}$, we will denote by $\mathfrak{z}\cdot\mathcal{X}$ its $\mathfrak{z}$-*blowup*, which is a metric space with the same underlying set as $\mathcal{X}$ and the metric multiplied by $\mathfrak{z}$. The tautological bijection $\mathcal{X} \to \mathfrak{z}\cdot\mathcal{X}$ will be denoted as $x \mapsto x^{\mathfrak{z}}$, so

$$|x^{\mathfrak{z}} - y^{\mathfrak{z}}| = \mathfrak{z}\cdot|x - y|$$

for any $x, y \in \mathcal{X}$.

The ω-blowup $\omega\cdot\mathcal{X}$ of $\mathcal{X}$ is defined as the ω-limit of the n-blowups $n\cdot\mathcal{X}$; that is,

$$\omega\cdot\mathcal{X} := \lim_{n\to\omega} n\cdot\mathcal{X}.$$

Given a point $x \in \mathcal{X}$ we can consider the sequence $x^n \in n\cdot\mathcal{X}$; it corresponds to a point $x^\omega \in \omega\cdot\mathcal{X}$. Note that if $x \neq y$, then

$$|x^\omega - y^\omega|_{\omega\cdot\mathcal{X}} = \infty;$$

that is, x^ω and y^ω belong to different metric components of $\omega\cdot\mathcal{X}$.

The metric component of x^ω in $\omega\cdot\mathcal{X}$ is called ultratangent space of $\mathcal{X}$ at x and it is denoted as $\mathrm{T}_x^\omega\mathcal{X}$.

Equivalently, ultratangent space $\mathrm{T}_x^\omega\mathcal{X}$ can be defined the following way. Consider all the sequences of points $x_n \in \mathcal{X}$ such that the sequence $\ell_n = n\cdot|x - x_n|_{\mathcal{X}}$ is bounded. Define the pseudodistance between two such sequences as

$$|(x_n) - (y_n)| = \lim_{n\to\omega} n\cdot|x_n - y_n|_{\mathcal{X}}.$$

Then $\mathrm{T}_x^\omega\mathcal{X}$ is the corresponding metric space.

Tangent space as well as ultratangent space, generalize the notion of tangent space of Riemannian manifold. In the simplest cases these two notions define the same space. In general, they are different and both useful — often lack of a property in one is compensated by the other.

It is clear from the definition that tangent space has cone structure. On the other hand, in general, ultratangent space does not have a cone structure; the Hilbert's cube $\prod_{n=1}^{\infty}[0,2^{-n}]$ is an example — it is CBB(0) as well as CAT(0).

The next theorem shows that the tangent space T_p can be (and often will be) considered as a subset of T_p^ω.

3.4.1. Theorem. *Let $\mathcal{X}$ be a metric space with defined angles. Then for any $p\in\mathcal{L}$, there is an distance preserving map*

$$\iota:\mathrm{T}_p\hookrightarrow\mathrm{T}_p^\omega$$

such that for any geodesic γ starting at p we have

$$\gamma^+(0)\mapsto\lim_{n\to\omega}[\gamma(\tfrac{1}{n})]^n.$$

Proof. Given $v\in\mathrm{T}_p'$ choose a geodesic γ that starts at p such that $\gamma^+(0)=v$. Set $v^n=[\gamma(\frac{1}{n})]^n\in n\cdot\mathcal{X}$ and

$$v^\omega=\lim_{n\to\omega}v^n.$$

Note that the value $v^\omega\in\mathrm{T}_p^\omega$ does not depend on choice of γ; that is, if γ_1 is an other geodesic starting at p such that $\gamma_1^+(0)=v$, then

$$\lim_{n\to\omega}v^n=\lim_{n\to\omega}v_1^n,$$

where $v_1^n=[\gamma_1(\frac{1}{n})]^n\in n\cdot\mathcal{X}$. The latter follows since

$$|\gamma(t)-\gamma_1(t)|_{\mathcal{X}}=o(t)$$

and therefore $|v^n-v_1^n|_{n\cdot\mathcal{X}}\to 0$ s $n\to\infty$.

Set $\iota(v)=v^\omega$. Since angles between geodesics in $\mathcal{X}$ are defined, for any $v,w\in\mathrm{T}_p'$ we have $n\cdot|v_n-w_n|\to|v-w|$. Thus $|v_\omega-w_\omega|=|v-w|$; that is, ι is a global isometry of T_p'.

Since T_p' is dense in T_p, we can extend ι to a global isometry $\mathrm{T}_p\to\mathrm{T}_p^\omega$. □

3.5 Gromov–Hausdorff convergence and ultralimits

3.5.1. Theorem. *Assume $\mathcal{X}_n$ is a sequence of complete spaces. Let $\mathcal{X}_n\to\mathcal{X}_\omega$ as $n\to\omega$, and $\mathcal{Y}_n\subset\mathcal{X}_n$ be a sequence of subsets such that $\mathcal{Y}_n\xrightarrow[\mathrm{GH}]{}\mathcal{Y}_\infty$. Then there is a distance preserving map $\iota:\mathcal{Y}_\infty\to\mathcal{X}_\omega$.*

Moreover:

a) If $\mathcal{X}_n \xrightarrow[\text{GH}]{} \mathcal{X}_\infty$ and $\mathcal{X}_\infty$ is compact, then $\mathcal{X}_\infty$ is isometric to $\mathcal{X}_\omega$.

b) If $\mathcal{X}_n \xrightarrow[\text{GH}]{} \mathcal{X}_\infty$ and $\mathcal{X}_\infty$ is proper, then $\mathcal{X}_\infty$ is isometric to a metric component of $\mathcal{X}_\omega$.

Proof. For each point $y_\infty \in \mathcal{Y}_\infty$ choose a lifting $y_n \in \mathcal{Y}_n$. Pass to the ω-limit $y_\omega \in \mathcal{X}_\omega$ of (y_n). Clearly for any $y_\infty, z_\infty \in \mathcal{Y}_\infty$, we have

$$|y_\infty - z_\infty|_{\mathcal{Y}_\infty} = |y_\omega - z_\omega|_{\mathcal{X}_\omega};$$

that is, the map $y_\infty \mapsto y_\omega$ gives a distance preserving map $\iota : \mathcal{Y}_\infty \to \to \mathcal{X}_\omega$.

(a)+(b). Fix $x_\omega \in \mathcal{X}_\omega$. Choose a sequence $x_n \in \mathcal{X}_n$ such that $x_n \to x_\omega$ as $n \to \omega$.

Denote by $\boldsymbol{X} = \mathcal{X}_\infty \sqcup \mathcal{X}_1 \sqcup \mathcal{X}_2 \sqcup \dots$ the common space for the convergence $\mathcal{X}_n \xrightarrow[\text{GH}]{} \mathcal{X}_\infty$; as in the definition of Gromov–Hausdorff convergence. Consider the sequence (x_n) as a sequence of points in $\boldsymbol{X}$.

If the ω-limit x_∞ of (x_n) exists, it must lie in $\mathcal{X}_\infty$.

The point x_∞, if defined, does not depend on the choice of (x_n). Indeed, if $y_n \in \mathcal{X}_n$ is an other sequence such that $y_n \to x_\omega$ as $n \to \omega$, then

$$|y_\infty - x_\infty| = \lim_{n\to\omega} |y_n - x_n| = 0;$$

that is, $x_\infty = y_\infty$.

In this way we obtain a map $\nu\colon x_\omega \to x_\infty$; it is defined on a subset of $\operatorname{Dom}\nu \subset \mathcal{X}_\omega$. By construction of ι, we get $\iota \circ \nu(x_\omega) = x_\omega$ for any $x_\omega \in \operatorname{Dom}\nu$.

Finally note that if $\mathcal{X}_\infty$ is compact, then ν is defined on all of $\mathcal{X}_\omega$; this proves (a).

If $\mathcal{X}_\infty$ is proper, choose any point $z_\infty \in \mathcal{X}_\infty$ and set $z_\omega = \iota(z_\infty)$. For any point $x_\omega \in \mathcal{X}_\omega$ at finite distance from z_ω, for the sequence x_n as above we have that $|z_n - x_n|$ is bounded for ω-almost all n. Since $\mathcal{X}_\infty$ is proper, $\nu(x_\omega)$ is defined; in other words ν is defined on the metric component of z_ω. Hence (b) follows. □

3.5.2. Corollary. *The ω-limit of a sequence of complete length spaces is geodesic.*

Proof. Given two points $x_\omega, y_\omega \in \mathcal{X}_\omega$, find two bounded sequences of points $x_n, y_n \in \mathcal{X}_n$, $x_n \to x_\omega$, $y_n \to y_\omega$ as $n \to \omega$. Consider a sequence of paths $\gamma_n\colon [0,1] \to \mathcal{X}_n$ from $_n$ to y_n such that

$$\operatorname{length}\gamma_n \leqslant |x_n - y_n| + \tfrac{1}{n}.$$

Apply Theorem 3.5.1 for the images $\mathcal{Y}_n = \gamma_n([0,1]) \subset \mathcal{X}_n$. □

3.6 Ultralimits of sets

Let $\mathcal{X}_n$ be a sequence of metric spaces and $\mathcal{X}_n \to \mathcal{X}_\omega$ as $n \to \omega$.

For a sequence of sets $\Omega_n \subset \mathcal{X}_n$, consider the maximal set $\Omega_\omega \subset \subset \mathcal{X}_\omega$ such that for any $x_\omega \in \Omega_\omega$ and any sequence $x_n \in \mathcal{X}_n$ such that $x_n \to x_\omega$ as $n \to \omega$, we have $x_n \in \Omega_n$ for ω-almost all n.

The set Ω_ω is called the *open ω-limit* of Ω_n; we could also write $\Omega_n \to \Omega_\omega$ as $n \to \omega$ or $\Omega_\omega = \lim_{n\to\omega} \Omega_n$.

Applying Observation 3.3.1 to the sequence of complements $\mathcal{X}_n \backslash \Omega_n$, we see that Ω_ω is open for any sequence Ω_n. The definition can be applied for arbitrary sequences of sets, but open ω-convergence will be applied here only for sequences of open sets.

3.7 Ultralimits of functions

Recall that a family of submaps between metric spaces $\{f_\alpha \colon \mathcal{X} \looparrowright \mathcal{Y}\}_{\alpha\in\mathcal{A}}$ is called *equicontinuous* if for any $\varepsilon > 0$ there is $\delta > 0$ such that for any $p, q \in \mathcal{X}$ with $|p - q| < \delta$ and any $\alpha \in \mathcal{A}$ it holds that $|f(p) - f(q)| < \varepsilon$.

Let $f_n \colon \mathcal{X}_n \looparrowright \mathbb{R}$ be a sequence of subfunctions.

Set $\Omega_n = \operatorname{Dom} f_n$. Consider the open ω-limit set $\Omega_\omega \subset \mathcal{X}_\omega$ of Ω_n.

Assume there is a subfunction $f_\omega \colon \mathcal{X}_\omega \looparrowright \mathbb{R}$ that satisfies the following conditions: (1) $\operatorname{Dom} f_\omega = \Omega_\omega$, (2) if $x_n \to x_\omega \in \Omega_\omega$ for a sequence of points $x_n \in \mathcal{X}_n$, then $f_n(x_n) \to f_\omega(x_\omega)$ as $n \to \omega$. In this case the subfunction $f_\omega \colon \mathcal{X}_\omega \to \mathbb{R}$ is said to be the ω-limit of $f_n \colon \mathcal{X}_n \to \mathbb{R}$.

The following lemma gives a mild condition on a sequence of functions f_n guaranteeing the existence of its ω-limit.

3.7.1. Lemma. *Let $\mathcal{X}_n$ be a sequence of metric spaces and $f_n \colon \mathcal{X}_n \looparrowright \mathbb{R}$ be a sequence of subfunctions.*

Assume for any positive integer k, there is an open set $\Omega_n(k) \subset \subset \operatorname{Dom} f_n$ such that the restrictions $f_n|_{\Omega_n(k)}$ are uniformly bounded and continuous and the open ω-limit of $\Omega_n(n)$ coincides with the open ω-limit of $\operatorname{Dom} f_n$. Then the ω-limit of f_n is defined.

In particular, if the f_n are uniformly bounded and continuous, then the ω-limit is defined.

The proof is straightforward.

3.7.2. Exercise. *Construct a sequence of compact length spaces $\mathcal{X}_n$ with a converging sequence of $\pounds$-Lipschitz concave functions $f_n \colon \mathcal{X}_n \to \mathbb{R}$ such that the ω-limit $\mathcal{X}_\omega$ of $\mathcal{X}_n$ is compact and the ω-limit $f_\omega \colon \mathcal{X}_\omega \to \mathbb{R}$ of f_n is not concave.*

If $f\colon \mathcal{X} \multimap \mathbb{R}$ is a subfunction, the ω-limit of the constant sequence $f_n = f$ is called the ω-power of f and denoted by f^ω. So

$$f^\omega\colon \mathcal{X} \multimap \mathbb{R}, \quad f^\omega(x_\omega) = \lim_{n\to\omega} f(x_n).$$

Recall that we treat $\mathcal{X}$ as a subset of its ω-power $\mathcal{X}^\omega$. Note that $\operatorname{Dom} f = \mathcal{X} \cap \operatorname{Dom} f^\omega$. Moreover, if $\mathrm{B}(x,\varepsilon)_{\mathcal{X}} \subset \operatorname{Dom} f$ then $\mathrm{B}(x,\varepsilon)_{\mathcal{X}^\omega} \subset \operatorname{Dom} f^\omega$.

Ultradifferential. Given a function $f\colon \mathcal{L} \to \mathbb{R}$, consider sequence of functions $f_n\colon n{\cdot}\mathcal{L} \to \mathbb{R}$, defined by

$$f_n(x^n) = n{\cdot}(f(x) - f(p)),$$

here $x^n \in n{\cdot}\mathcal{L}$ is the point corresopnding to $x \in \mathcal{L}$. While $n{\cdot}(\mathcal{L},p) \to$ $\to (\mathrm{T}^\omega, 0)$ as $n \to \omega$, functions f_n converge to ω-differential of f at p. It will be denoted by $\boldsymbol{d}_p^\omega f$;

$$\boldsymbol{d}_p^\omega f\colon \mathrm{T}_p^\omega \to \mathbb{R}, \quad \boldsymbol{d}_p^\omega f = \lim_{n\to\omega} f_n.$$

Clearly, the ω-differential $\boldsymbol{d}_p^\omega f$ of a locally Lipschitz subfunction f is defined at each point $p \in \operatorname{Dom} f$.

Chapter 4

Maps and functions

Here we introduce some classes of maps between metric spaces and develop a language to describe different levels of convexity/concavity of real-valued functions on general metric space.

Concave functions are used mainly in the CBB spaces, while convex ones are used mainly in CAT spaces.

4.1 Submaps

We will often need maps and functions defined on subsets of a metric space. We call them *submaps* and *subfunction*. Thus given a metric space $\mathcal{X}$ and $\mathcal{Y}$, a submap $\Phi\colon \mathcal{X} \multimap \mathcal{Y}$ is a map defined on a subset $\operatorname{Dom}\Phi \subset \mathcal{X}$.

A submap is defined to be *continuous* if the inverse image of any open set is open. Note that for a continuous submap Φ, the domain $\operatorname{Dom}\Phi$ is automatically open. Indeed, a submap $\Phi\colon \mathcal{X} \multimap \mathcal{Y}$ is continuous if the inverse image of any open set is open. Since $\mathcal{Y}$ is open then so is $\operatorname{Dom}\Phi = \Phi^{-1}(\mathcal{Y})$. The same is true for upper and lower semicontinuous functions $f\colon \mathcal{X} \multimap \mathbb{R}$ since it is a continuous function for special topology on $\mathbb{R}$.

(The continuous partially defined maps could be defined via closed sets; namely, one may request that inverse images of closed sets are closed. While this condition is equivalent to the continuity for the functions defined on whole space, it is different for partially defined functions. In particular with this definition domain of the continuous submap would have to be closed.)

4.2 Lipschitz conditions

4.2.1. Lipschitz maps. *Let $\mathcal{X}$ and $\mathcal{Y}$ be metric spaces $\Phi\colon \mathcal{X} \looparrowright \mathcal{Y}$ be a continuous submap and $\pounds \in \mathbb{R}$.*

*a) The submap Φ is called $\pounds$-*Lipschitz *if*

$$|\Phi(x) - \Phi(y)|_{\mathcal{Y}} \leqslant \pounds \cdot |x - y|_{\mathcal{X}}$$

for any two points $x, y \in \operatorname{Dom}\Phi$.

- *The 1-Lipschitz maps will be also called* short.

b) We say that Φ is Lipschitz *if it is $\pounds$-Lipschitz for some constant $\pounds$. The minimal such constant is denoted by* $\operatorname{lip}\Phi$.

c) We say that Φ is locally Lipschitz *if any point $x \in \operatorname{Dom}\Phi$ admits a neighborhood $\Omega \subset \operatorname{Dom}\Phi$ such that the restriction $\Phi|_\Omega$ is Lipschitz.*

d) Given $p \in \operatorname{Dom}\Phi$, we denote by $\operatorname{lip}_p\Phi$ the infimum of the real values $\pounds$ such that p admits a neighborhood $\Omega \subset \operatorname{Dom}\Phi$ such that the restriction $\Phi|_\Omega$ is $\pounds$-Lipschitz.

Note that $\Phi\colon \mathcal{X} \to \mathcal{Y}$ is $\pounds$-Lipschitz if

$$\Phi(\mathrm{B}(x, R)_{\mathcal{X}}) \subset \mathrm{B}(\Phi(x), \pounds \cdot R)_{\mathcal{Y}}$$

for any $R \geqslant 0$ and $x \in \mathcal{X}$. The following definition gives a dual version of it.

4.2.2. Definitions. *Let $\mathcal{X}$ and $\mathcal{Y}$ be metric spaces, $\Phi\colon \mathcal{X} \to \mathcal{Y}$ be a map and $\pounds \in \mathbb{R}$.*

*a) The map Φ is called $\pounds$-*co-Lipshitz *if*

$$\Phi(\mathrm{B}(x, \pounds \cdot R)_{\mathcal{X}}) \supset \mathrm{B}(\Phi(x), R)_{\mathcal{Y}}$$

for any $x \in \mathcal{X}$ and $R > 0$.

b) The map Φ is called co-Lipshitz *if it is $\pounds$-co-Lipshitz for some constant $\pounds$. The minimal such constant is denoted by* $\operatorname{colip}\Phi$.

From the definition of co-Lipschitz map we get the following.

4.2.3. Proposition. *Any co-Lipschitz map is open and surjective.*

In other words, the $\pounds$-co-Lipschitz maps can be considered as a quantitative version of open maps. By that reason they also called $\pounds$-open [3]. Also, be aware that some authors, refer to our $\pounds$-co-Lipschitz maps as to $\frac{1}{\pounds}$-co-Lipschitz.

4.2.4. Proposition. *Let $\mathcal{X}$ and $\mathcal{Y}$ be metric spaces and $\Phi\colon \mathcal{X} \to \mathcal{Y}$ a continuous co-Lipschitz map. Then $\mathcal{Y}$ is complete.*

Proof. Choose a converging in itself sequence $y_1, y_2, \dots$ in $\mathcal{Y}$. Passing to a subsequence if nececury, we may assume that $|y_n - y_{n+1}|_{\mathcal{Y}} < \frac{1}{2^n}$ for each n.

Denote by $\pounds$ a co-Lipschitz constant of Φ. Note that there is a sequence $x_1, x_2, \dots$ in $\mathcal{X}$ such that

$$\Phi(x_n) = y_n \quad \text{and} \quad |x_n - x_{n+1}|_{\mathcal{X}} < \tfrac{\pounds}{2^n}. \tag{❶}$$

for each n. Indeed, such a sequence (x_n) can be constructed recursively. Assuming that the points $x_1, \dots, x_{n-1}$ are already constructed, the existence of x_n satisfying ❶ follows since Φ is $\pounds$-co-Lipschitz.

Notice that (x_n) converges in itself; since $\mathcal{X}$ is complete, it converges in $\mathcal{X}$. Denote by x_∞ its limit and set $y_\infty = \Phi(x_\infty)$. Since Φ is continuous, $y_n \to y_\infty$ as $n \to \infty$. Hence the result. □

4.2.5. Lemma. *Let $\mathcal{X}$ be a metric space and $f\colon \mathcal{X} \to \mathbb{R}$ be a continuous function. Then for any $\varepsilon > 0$ there is a locally Lipschitz function $f_\varepsilon\colon \mathcal{X} \to \mathbb{R}$ such that $|f(x) - f_\varepsilon(x)| < \varepsilon$ for any $x \in \mathcal{X}$.*

Proof. Assume that $f \geqslant 1$. Construct a continous positive function $\rho\colon \mathcal{X} \to \mathbb{R}_{>0}$ such that

$$|x - y| < \rho(x) \quad \Rightarrow \quad |f(x) - f(y)| < \varepsilon.$$

Consider function

$$f_\varepsilon(x) = \sup\left\{ f(x)\cdot\left(1 - \tfrac{|x-y|}{\rho(x)}\right) : x \in \mathcal{X} \right\}.$$

It is straightforward to check that each f_ε is locally Lipschtz and $0 \leqslant$ $\leqslant f_\varepsilon - f < \varepsilon$.

Note that any continuous function can be presented as difference of two continuous functions bounded below by 1. Hence the result. □

4.3 Isometries and sumbetries

4.3.1. Isometry. *Let $\mathcal{X}$ and $\mathcal{Y}$ be metric spaces and $\Phi\colon \mathcal{X} \to \mathcal{Y}$ be a map*

a) The map Φ is distance preserving *if*

$$|\Phi(x) - \Phi(x')|_{\mathcal{Y}} = |x - x'|_{\mathcal{X}}$$

for any $x, x' \in X$.

b) A distance preserving bijection Φ is called an isometry*.*

c) The spaces X and Y are called isometric *(briefly $X \overset{iso}{=\!=} Y$) if there is an isometry $\Phi\colon X \to Y$.*

4.3.2. Submetry. *A map $\sigma\colon \mathcal{L} \to \mathcal{M}$ between the metric spaces $\mathcal{L}$ and $\mathcal{M}$ is called* submetry *if*

$$\sigma(\mathrm{B}(p,r)_{\mathcal{L}}) = \mathrm{B}(\sigma(p),r)_{\mathcal{M}}$$

for any $p \in \mathcal{L}$ and $r \geqslant 0$.

Note $\sigma\colon \mathcal{L} \to \mathcal{M}$ is a sumbetry if it is 1-Lipshitz and 1-co-Lipschitz at the same time.

Note that any submetry is an onto map.

The main souse of examples of submetries comes from isometric group actions.

Nemely, assume $\mathcal{L}$ is a metric space and G be a subgroup of isometries of $\mathcal{L}$. Denote by $\mathcal{L}/G$ the set of G-orbits; let us equip it with the Hausdorff metric.

In general, $\mathcal{L}/G$ is a pseudometric space, but if all the G-orbits form closed sets in $\mathcal{L}$ then $\mathcal{L}/G$ is a genuine metric space.

4.3.3. Proposition. *Let $\mathcal{L}$ be a metric space. Assume that a group G acts on $\mathcal{L}$ by isometries and in such a way that every G-orbit is closed. Then the projection map $\mathcal{L} \to \mathcal{L}/G$ is a sumbetry.*

Proof. Denote by $\hat{x}$ the projection of $x \in \mathcal{L}$ in $\mathcal{L}/G$. We need to show that the map $x \mapsto \hat{x}$ is 1-Lipschitz and 1-co-Lipschitz. The co-Lipschitz part follows directly from the definition of Hausdorff distance and co-Lipschitz map.

Assume $|x-y|_{\mathcal{L}} \leqslant r$; equivalently $\overline{\mathrm{B}}[x,r]_{\mathcal{L}} \ni y$. Since the action $G \curvearrowright \mathcal{L}$ is isometric, $\overline{\mathrm{B}}[g{\cdot}x,r]_{\mathcal{L}} \ni g{\cdot}y$ for any $g \in G$.

In particular the orbit $G{\cdot}y$ lies in the closed r-neighborhood of the orbit $G{\cdot}x$. The same way we can prove that the orbit $G{\cdot}x$ lies in the closed r-neighborhood of the orbit $G{\cdot}y$. That is, the Hausdorff distance between the orbits $G{\cdot}x$ and $G{\cdot}y$ is at most r or, equivalently, $|\hat{x}-\hat{y}|_{\mathcal{L}/G} \leqslant r$. Since x and y are arbitrary, the map $x \mapsto \hat{x}$ is 1-Lipschitz. $\square$

4.3.4. Proposition. *Let $\mathcal{L}$ be a length space and $\sigma\colon \mathcal{L} \to \mathcal{M}$ is a submetry. Then $\mathcal{M}$ is a length space.*

Proof. Fix $\varepsilon > 0$ and a pair of points $x, y \in \mathcal{M}$.

Since σ is 1-co-Lipschitz, there are points $\hat{x}, \hat{y} \in \mathcal{L}$ such that $\sigma(\hat{x}) = x$, $\sigma(\hat{y}) = y$ and $|\hat{x}-\hat{y}|_{\mathcal{L}} < |x-y|_{\mathcal{M}} + \varepsilon$.

Since $\mathcal{L}$ is a length space, there is a curve γ joining $\hat{x}$ to $\hat{y}$ in $\mathcal{L}$ such that

$$\operatorname{length}\gamma \leqslant |x-y|_{\mathcal{M}} + \varepsilon.$$

Since σ is 1-Lipschitz, there

$$\operatorname{length}\sigma\circ\gamma \leqslant \operatorname{length}\gamma.$$

Note that the curve $\sigma\circ\gamma$ joins x to y and from above

$$\operatorname{length}\sigma\circ\gamma < |x-y|_{\mathcal{M}} + \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, we have that $\mathcal{M}$ is a length space. □

4.4 Speed of curves

Let $\mathcal{X}$ be a metric space. A *curve* in $\mathcal{X}$ is a continuous map $\alpha\colon \mathbb{I}\to\mathcal{X}$, where $\mathbb{I}$ is a *real interval* (that is, an arbitrary convex subset of $\mathbb{R}$); the curve it called *Lipschitz* or *locally Lipschitz* if the map $\alpha\colon \mathbb{I}\to\mathcal{X}$ is.

Recall that length of curves is defined in 2.3.1. The following theorem follows from [4, p. 2.7.4].

4.4.1. Theorem. *Let $\mathcal{X}$ be a metric space and $\alpha\colon \mathbb{I}\to\mathcal{X}$ be a locally Lipschitz curve. Then the speed function*

$$\operatorname{speed}_{t_0}\alpha = \lim_{t\to t_0}\frac{|\alpha(t)-\alpha(t_0)|}{|t-t_0|}$$

is defined for almost all $t_0\in\mathbb{I}$ and

$$\operatorname{length}\alpha = \int\limits_{\mathbb{I}}\operatorname{speed}_t\alpha\cdot\boldsymbol{d}t,$$

where $\int$ denotes the Lebesgue integral.

A curve $\alpha\colon \mathbb{I}\to\mathcal{X}$ is called *unit-speed curve*, if for any subinterval $[a,b]\subset\mathbb{I}$, we have

$$b-a = \operatorname{length}(\alpha|_{[a,b]}).$$

According to the above theorem, this is equivalent to the condition that α is Lipschitz and $\operatorname{speed}\alpha \overset{a.e.}{=\!=} 1$.

The following theorem of Bernd Kirchheim [5] generalizes the one above.

4.4.2. Theorem. *Let $\mathcal{X}$ be a metric space and $f\colon \mathbb{R}^n \looparrowright \mathcal{X}$ be 1-Lipschitz. Then for almost all $x\in\operatorname{Dom}f$ there is a pseudonorm $\|*\|_x$ on $\mathbb{R}^n$ such that we have*

$$|f(y)-f(z)|_{\mathcal{X}} = \|z-y\|_x + o(|y-x|+|z-x|).$$

The (pseudo)norm $\|*\|_x$ in the above theorem will be called *differential of induced metric* at x.

4.5 Convex real-to-real functions

We will be interested in the generalized solutions of the following differential inequalities

❶ $$y'' + \kappa\cdot y \geqslant \lambda \quad \text{and respectively} \quad y'' + \kappa\cdot y \leqslant \lambda$$

for fixed $\kappa, \lambda \in \mathbb{R}$. The solution $y\colon \mathbb{R} \circ\!\!\to \mathbb{R}$ are only assumed to be upper (respectively lower) semicontinuous subfunctions.

The inequalities ❶ are understood in the sense of distributions. That is, for any smooth function φ with compact support $\operatorname{Supp}\varphi \subset \subset \operatorname{Dom} y$ the following inequality should be satisfied:

❷ $$\int\limits_{\operatorname{Dom} y} [y(t)\cdot\varphi''(t) + \kappa\cdot y(t)\cdot\varphi(t) - \lambda]\cdot \boldsymbol{d}t \geqslant 0$$

$$\text{respectively} \quad \leqslant 0.$$

The integral is understood in the sense of Lebesgue, in particular the inequality ❷ makes sense for any Borel-measurable subfunction y. The proofs of the following propositions are straightforward.

4.5.1. Proposition. *Let $\mathbb{I} \subset \mathbb{R}$ be an open interval and $y_n\colon \mathbb{I} \to \to \mathbb{R}$ be a sequence of solutions of one of the inequality in ❶. Assume $y_n(t) \to y_\infty(t)$ as $n \to \infty$ for any $t \in \mathbb{I}$. Then y_∞ is a solution of the same inequality in ❶.*

Assume y is a solution of one of the inequality in ❶. For $t_0 \in \operatorname{Dom} y$, let us define right (left) derivative $y^+(t_0)$ ($y^-(t_0)$) at t_0 by

$$y^\pm(t_0) = \lim_{t\to t_0\pm} \frac{y(t) - y(t_0)}{|t - t_0|}.$$

Note that our sign convention for y^- is not standard — for $y(t) = t$ we have $y^+(t) = 1$ and $y^-(t) = -1$.

4.5.2. Proposition. *Let $\mathbb{I} \subset \mathbb{R}$ be an open interval and $y\colon \mathbb{I} \to \mathbb{R}$ be a solution of the inequality in ❶. Then y is locally Lipschitz; its right and left derivatives $y^+(t_0)$ and $y^-(t_0)$ are defined for any $t_0 \in \mathbb{I}$. Moreover*

$$y^+(t_0) + y^-(t_0) \geqslant 0 \quad \textit{or respectively} \quad y^+(t_0) + y^-(t_0) \leqslant 0$$

The next theorem gives a number of equivalent ways to define such a generalized solution.

4.5.3. Theorem. *Let $\mathbb{I}$ be an open real interval and $y\colon \mathbb{I} \to \mathbb{R}$ be a locally Lipschitz function then the following conditions are equivalent:*

a) $y'' \geqslant \lambda - \kappa\cdot y$ *(respectively* $y'' \leqslant \lambda - \kappa\cdot y$*)*

b) (barrier inequality) For any $t_0 \in \mathbb{I}$*, there is a solution* $\bar{y}$ *of ordinary differential equation* $\bar{y}'' = \lambda - \kappa\cdot\bar{y}$ *with* $\bar{y}(t_0) = y(t_0)$ *such that* $\bar{y} \geqslant y$ *(respectively* $\bar{y} \leqslant y$*) for all* $t \in [t_0 - \varpi^\kappa, t_0 + \varpi^\kappa] \cap \mathbb{I}$*.*

The function $\bar{y}$ *is called* lower *(respectively* upper*)* barrier *of* y *at* t_0*.*

c) Suppose $y'' \leqslant \lambda - \kappa\cdot y$ *. Let* $t_0 \in \mathbb{I}$ *and let* $\bar{y}$ *be a solution of the ordinary differential equation* $\bar{y}'' = \lambda - \kappa\cdot\bar{y}$ *such that* $\bar{y}(t_0) = y(t_0)$ *and* $y^+(t_0) \leqslant y(t_0) \leqslant -y^-(t_0)$*. (note that such* $\bar{y}$ *is unique if* y *is differentiable at* t_0*).*

Then $\bar{y} \geqslant y$ *for all* $t \in [t_0 - \varpi^\kappa, t_0 + \varpi^\kappa] \cap \mathbb{I}$*; that is* $\bar{y}$ *is a barrier of* y *at* t_0*.*

d) (Jensen's inequality) For any pair of values $t_1 < t_2$ *in* $\mathbb{I}$*, such that* $|t_2 - t_1| < \varpi^\kappa$ *the unique solutions* $z(t)$ *of*

$$z'' = \lambda - \kappa\cdot z$$

such that

$$z(t_1) = y(t_1), \quad z(t_2) = y(t_2)$$

satisfies $y(t) \leqslant z(t)$ *(respectively* $y(t) \geqslant z(t)$*) for all* $t \in$ $\in [t_1, t_2]$*.*

The proof is left to the reader.

We will often need the following fact about convergence of derivative of convex functions:

4.5.4. Lemma on equilibrium. *Let* $\mathbb{I}$ *be an open interval and* $f_n\colon \mathbb{I} \to \mathbb{R}$ *be a sequence of concave functions. Assume the functions* f_n *pointwise converge to a function* $f\colon \mathbb{I} \to \mathbb{R}$*. Then for any* $t_0 \in \mathbb{I}$*,*

$$f^\pm(t_0) \leqslant \varliminf_{n\to\infty} f_n^\pm(t_0).$$

Proof. Since f_n are convex, we have $f_n^+(t_0) + f_n^-(t_0) \geqslant 0$, and for any t,

$$f_n(t) \geqslant f_n(t_0) \pm f^\pm(t_0)\cdot(t - t_0).$$

Passing to the limit, we get

$$f(t) \geqslant f(t_0) + \left[\varlimsup_{n\to\infty} f_n^+(t_0)\right]\cdot(t - t_0)$$

for $t \geqslant t_0$, and

$$f(t) \geqslant f(t_0) - \left[\varlimsup_{n\to\infty} f_n^-(t_0)\right]\cdot(t - t_0)$$

for $t \leqslant t_0$. Hence the result. $\square$

4.5.5. Corollary. *Let $\mathbb{I}$ be an open interval and $f_n\colon \mathbb{I} \to \mathbb{R}$ be a sequence of functions such that $f_n'' \leqslant \lambda$ and converge pointwise to a function $f\colon \mathbb{I} \to \mathbb{R}$. Then:*

a) If f is differentiable at $t_0 \in \mathbb{I}$, then

$$f'(t_0) = \pm \lim_{n\to\infty} f_n^{\pm}(t_0).$$

b) If all f_n and f are differentiable at $t_0 \in \mathbb{I}$, then

$$f'(t_0) = \lim_{n\to\infty} f_n'(t_0).$$

Proof. Set $\hat f_n(t) = f_n(t) - \frac{\lambda}{2}\cdot t^2$ and $\hat f(t) = f(t) - \frac{\lambda}{2}\cdot t^2$. Note that $\hat f_n$ are concave and $\hat f_n \to \hat f$ pointwise. Thus, theorem follows from the lemma on equilibrium (4.5.4). $\square$

4.6 Convex functions on a metric space

The following notion will be used for geodesic spaces most of the time.

4.6.1. Definition. *Let $\mathcal{X}$ be a metric space. We say that an upper semicontinuous subfunction $f\colon \mathcal{X} \looparrowright (-\infty, \infty]$ satisfies the inequality*

$$f'' + \kappa\cdot f \geqslant \lambda$$

if for any unit-speed geodesic γ in $\operatorname{Dom} f$ the real-to-real function $y(t) = f\circ\gamma(t)$ satisfies

$$y'' + \kappa\cdot y \geqslant \lambda$$

in the domain $\{\, t : y(t) < \infty \,\}$; see the definition in Section 4.5.

We say that a lower semicontinuous subfunction $f\colon \mathcal{X} \looparrowright [-\infty, \infty)$ satisfies the inequality

$$f'' + \kappa\cdot f \leqslant \lambda$$

if the subfunction $h = -f$ satisfies

$$h'' - \kappa\cdot h \geqslant -\lambda.$$

Functions satisfying the inequalities

$$f'' \geqslant \lambda \quad \text{and} \quad f'' \leqslant \lambda$$

are called λ-convex and λ-concave correspondingly.

0-*convex and* 0-*concave subfunctions will be also called* convex *and* concave *correspondingly.*

If f is λ-convex for some $\lambda > 0$ it will be called strongly convex*; correspondingly if f is λ-concave for some $\lambda < 0$ it will be called* strongly concave*.*

If for any point $p \in \operatorname{Dom} f$ there is a neighborhood $\Omega \ni p$ and a real number λ such that the restriction $f|_\Omega$ is λ-convex (or λ-concave), then f is called semiconvex (correspondingly semiconcave).

Further, we give a meaning of certain type of second order ordinary differential inequalities in context of metric spaces.

Various authors define the class of λ-convex (λ-concave) function differently. It may correspond to $\pm\lambda$-convex ($\pm\lambda$-concave) or $\pm\frac{\lambda}{2}$-convex ($\pm\frac{\lambda}{2}$-concave) function in our definitions.

4.6.2. Proposition. *Let $\mathcal{X}$ be a metric space. Assume that $f\colon \mathcal{X} \multimap \multimap \mathbb{R}$ is a semiconvex subfunction and $\varphi\colon \mathbb{R} \to \mathbb{R}$ is a nondecreasing seimconvex function. Then the composition $\varphi \circ f$ is a semiconvex subfunction.*

The proof is straightforward.

Chapter 5

The ghost of Euclid

5.1 Geodesics, triangles and hinges

Geodesics and their relatives. Let $\mathcal{X}$ be a metric space and $\mathbb{I}$ be a real interval. A globally isometric map $\gamma\colon \mathbb{I} \to \mathcal{X}$ is called a *unit-speed geodesic*[1]; in other words, $\gamma\colon \mathbb{I} \to \mathcal{X}$ is a unit-speed geodesic if the equality

$$|\gamma(s) - \gamma(t)|_{\mathcal{X}} = |s - t|$$

holds for any pair $s, t \in \mathbb{I}$.

A unit-speed geodesic $\gamma\colon \mathbb{R}_{\geqslant 0} \to \mathcal{X}$ is called a *ray*.

A unit-speed geodesic $\gamma\colon \mathbb{R} \to \mathcal{X}$ is called a *line*.

A unit-speed geodesic between p and q in $\mathcal{X}$ will be denoted by $\mathrm{geod}_{[pq]}$. We will always assume $\mathrm{geod}_{[pq]}$ is parametrized starting at p; that is, $\mathrm{geod}_{[pq]}(0) = p$ and $\mathrm{geod}_{[pq]}(|p-q|) = q$. The image of $\mathrm{geod}_{[pq]}$ will be denoted by $[pq]$ and called a *geodesic*. The term *geodesic* will also be used for a linear reparametrization of a unit-speed geodesic; when a confusion is possible we call the latter a *constant-speed geodesic*. With slight abuse of notation, we will use $[pq]$ also for the class of all linear reparametrizations of $\mathrm{geod}_{[pq]}$.

We may write $[pq]_{\mathcal{X}}$ to emphasize that the geodesic $[pq]$ is in the space $\mathcal{X}$. Also we use the following short-cut notation:

$$]pq[= [pq]\backslash\{p, q\}, \qquad]pq] = [pq]\backslash\{p\}, \qquad [pq[= [pq]\backslash\{q\}.$$

In general, a geodesic between p and q need not exist and if it exists, it need not to be unique. However, once we write $\mathrm{geod}_{[pq]}$ or $[pq]$ we mean that we fixed a choice of geodesic.

[1] Various authors call it differently: *shortest path*, *minimizing geodesic*.

A constant-speed geodesic $\gamma\colon [0,1] \to \mathcal{X}$ is called a *geodesic path*. Given a geodesic $[pq]$, we denote by $\mathrm{path}_{[pq]}$ the corresponding geodesic path; that is,

$$\mathrm{path}_{[pq]}(t) \equiv \mathrm{geod}_{[pq]}(t\cdot|p-q|).$$

A curve $\gamma\colon \mathbb{I} \to \mathcal{X}$ is called a *local geodesic*, if for any $t \in \mathbb{I}$ there is a neighborhood $U \ni t$ in $\mathbb{I}$ such that the restriction $\gamma|_U$ is a constant-speed geodesic. If $\mathbb{I} = [0,1]$, then γ is called a *local geodesic path*.

5.1.1. Proposition. *Suppose $\mathcal{X}$ is a metric space and $\gamma\colon [0,\infty) \to \mathcal{X}$ is a ray. Then the* Busemann function $\mathrm{bus}_\gamma\colon \mathcal{X} \to \mathbb{R}$

❶ $$\mathrm{bus}_\gamma(x) = \lim_{t\to\infty} |\gamma(t) - x| - t$$

is defined and 1*-Lipschitz.*

Proof. As follows from the triangle inequality, the function

$$t \mapsto |\gamma(t) - x| - t$$

is nonincreasing in t. Clearly $|\gamma(t) - x| - t \geqslant -|\gamma(0) - x|$. Thus the limit in ❶ is defined. □

Triangles. For a triple of points $p, q, r \in \mathcal{X}$, a choice of triple of geodesics $([qr], [rp], [pq])$ will be called a *triangle*, and we will use the short notation $[pqr] = ([qr], [rp], [pq])$. Again, given a triple $p, q, r \in$ $\in \mathcal{X}$, there may be no triangle $[pqr]$, simply because one of the pairs of these points cannot be joined by a geodesic. Or there may be many different triangles, any of which can be denoted by $[pqr]$. Once we write $[pqr]$, it means we have chosen such a triangle; that is, made a choice of each $[qr], [rp]$ and $[pq]$.

The value

$$|p-q| + |q-r| + |r-p|$$

will be called the *perimeter of triangle* $[pqr]$; it obviously coincides with perimeter of triple p, q, r as defined below.

Hinges. Let $p, x, y \in \mathcal{X}$ be a triple of points such that p is distinct from x and y. A pair of geodesics $([px], [py])$ will be called a *hinge*, and will be denoted by $[p\,{}^{x}_{y}] = ([px], [py])$.

5.2 Model angles and triangles

Let $\mathcal{X}$ be a metric space, $p, q, r \in \mathcal{X}$ and $\kappa \in \mathbb{R}$. Let us define the *model triangle* $[\tilde{p}\tilde{q}\tilde{r}]$ (briefly, $[\tilde{p}\tilde{q}\tilde{r}] = \tilde{\triangle}^\kappa(pqr)$) to be a triangle in the model plane $\mathbb{M}^2(\kappa)$ such that

$$|\tilde{p} - \tilde{q}| = |p-q|, \quad |\tilde{q} - \tilde{r}| = |q-r|, \quad |\tilde{r} - \tilde{p}| = |r-p|.$$

In the notation of Section 1.1, $\tilde\triangle^\kappa(pqr) = \tilde\triangle^\kappa\{|q-r|, |r-p|, |p-q|\}$.

If $\kappa \leqslant 0$, the model triangle is always defined, that is, exists and is unique up to isometry of $\mathbb{M}^2(\kappa)$. If $\kappa > 0$, the model triangle is said to be defined if in addition

$$|p-q| + |q-r| + |r-p| < 2\cdot\varpi^\kappa.$$

In this case it also exists and is unique up to isometry of $\mathbb{M}^2(\kappa)$. The value $|p-q| + |q-r| + |r-p|$ will be called the *perimeter of triple* p, q, r.

If for $p, q, r \in \mathcal{X}$, $[\tilde p\tilde q\tilde r] = \tilde\triangle^\kappa(pqr)$ is defined and $|p-q|, |p-r| > 0$, the angle measure of $[\tilde p\tilde q\tilde r]$ at $\tilde p$ will be called the *model angle* of triple p, q, r, and will be denoted by $\measuredangle^\kappa(p\,{}^q_r)$.

In the notation of Section 1.1,

$$\measuredangle^\kappa(p\,{}^q_r) = \measuredangle^\kappa\{|q-r|; |p-q|, |p-r|\}.$$

5.2.1. Alexandrov's lemma. *Let p, q, r, z be distinct points in a metric space such that $z \in\,]pr[$ and*

$$|p-q| + |q-r| + |r-p| < 2\cdot\varpi^\kappa.$$



Then the following expressions have the same sign:

a) $\measuredangle^\kappa(p\,{}^q_z) - \measuredangle^\kappa(p\,{}^q_r)$,

b) $\measuredangle^\kappa(z\,{}^q_p) + \measuredangle^\kappa(z\,{}^q_r) - \pi$.

Moreover,

$$\measuredangle^\kappa(q\,{}^p_r) \geqslant \measuredangle^\kappa(q\,{}^p_z) + \measuredangle^\kappa(q\,{}^z_r),$$

with equality if and only if the expressions in (a) and (b) vanish.

Proof. By the triangle inequality,

$$|p-q| + |q-z| + |z-p| \leqslant |p-q| + |q-r| + |r-p| < 2\cdot\varpi^\kappa.$$

Therefore the model triangle $[\tilde p\tilde q\tilde z] = \tilde\triangle^\kappa pqz$ is defined. Take a point $\tilde r$ on the extension of $[\tilde p\tilde z]$ beyond $\tilde z$ so that $|\tilde p - \tilde r| = |p-r|$ (and therefore $|\tilde p - \tilde z| = |p-z|$).

From monotonicity of function $a \mapsto \measuredangle^\kappa\{a; b, c\}$ (1.1.1c), the following expressions have the same sign:

(i) $\measuredangle[\tilde p\,{}^{\tilde q}_{\tilde r}] - \measuredangle^\kappa(p\,{}^q_r)$;

(ii) $|\tilde p - \tilde r| - |p-r|$;

(iii) $\measuredangle[\tilde z\,{}^{\tilde q}_{\tilde r}] - \measuredangle^\kappa(z\,{}^q_r)$.

Since

$$\measuredangle[\tilde p\,{}^{\tilde q}_{\tilde r}] = \measuredangle[\tilde p\,{}^{\tilde q}_{\tilde z}] = \mathcal{Z}^{\kappa}(p\,{}^{q}_{z})$$

and

$$\measuredangle[\tilde z\,{}^{\tilde q}_{\tilde r}] = \pi - \measuredangle[\tilde z\,{}^{\tilde p}_{\tilde q}] = \pi - \mathcal{Z}^{\kappa}(z\,{}^{p}_{q}),$$

the first statement follows.

For the second statement, construct $[\tilde q\tilde z r'] = \tilde\triangle^{\kappa} qzr$ on the opposite side of $[\tilde q\tilde z]$ from $[\tilde p\tilde q\tilde z]$. Since

$$|\tilde p - r'| \leqslant |\tilde p - \tilde z| + |\tilde z - r'| = |p - z| + |z - r| = |p - r|,$$

then

$$\begin{aligned}\mathcal{Z}^{\kappa}(q\,{}^{p}_{z}) + \mathcal{Z}^{\kappa}(q\,{}^{z}_{r}) &= \measuredangle[\tilde q\,{}^{\tilde p}_{\tilde z}] + \measuredangle[\tilde q\,{}^{\tilde z}_{r'}] = \\ &= \measuredangle[\tilde q\,{}^{\tilde p}_{r'}] \leqslant \\ &\leqslant \mathcal{Z}^{\kappa}(q\,{}^{p}_{r}).\end{aligned}$$

Equality holds if and only if $|\tilde p - r'| = |p - r|$, as required. □

5.3 Angles and the first variation

Given a hinge $[p\,{}^{x}_{y}]$, we define its *angle* to be

❶ $$\measuredangle[p\,{}^{x}_{y}] := \lim_{\bar x,\bar y\to p} \mathcal{Z}^{\kappa}(p\,{}^{\bar x}_{\bar y}),$$

for $\bar x \in\,]px]$ and $\bar y \in\,]py]$, if this limit exists.

Similarly to $\mathcal{Z}^{\kappa}(p\,{}^{x}_{y})$, we will use the short notation

$$\tilde\Upsilon^{\kappa}[p\,{}^{x}_{y}] = \tilde\Upsilon^{\kappa}\left\{\measuredangle[p\,{}^{x}_{y}]; |p - x|, |p - y|\right\},$$

where the right hand side is defined on page 4. The value $\tilde\Upsilon^{\kappa}[p\,{}^{x}_{y}]$ will be called the *model side* of hinge $[p\,{}^{x}_{y}]$.

5.3.1. Lemma. *For any $\kappa, \mathrm{K} \in \mathbb{R}$, there exists $c \in \mathbb{R}$ such that*

❷ $$|\mathcal{Z}^{\mathrm{K}}(p\,{}^{x}_{y}) - \mathcal{Z}^{\kappa}(p\,{}^{x}_{y})| \leqslant c\cdot|p - x|\cdot|p - y|,$$

whenever the lefthand side is defined.

Lemma 5.3.1 implies that the definition of angle is independent of κ. In particular, one can take $\kappa = 0$ in ❶; thus the angle can be calculated from the cosine law:

$$\cos\mathcal{Z}^{0}(p\,{}^{x}_{y}) = \frac{|p - x|^2 + |p - y|^2 - |x - y|^2}{2\cdot|p - x|\cdot|p - y|}.$$

Proof. The function $\kappa \mapsto \measuredangle^{\kappa}(p\,{}^{x}_{y})$ is nondecreasing (1.1.1d). Thus, for $\mathrm{K} > \kappa$, we have

$$\begin{aligned}0 \leqslant \measuredangle^{\mathrm{K}}(p\,{}^{x}_{y}) - \measuredangle^{\kappa}(p\,{}^{x}_{y}) \leqslant & \measuredangle^{\mathrm{K}}(p\,{}^{x}_{y}) + \measuredangle^{\mathrm{K}}(x\,{}^{p}_{y}) + \measuredangle^{\mathrm{K}}(y\,{}^{p}_{x}) - \\ & - \measuredangle^{\kappa}(p\,{}^{x}_{y}) - \measuredangle^{\kappa}(x\,{}^{p}_{y}) - \measuredangle^{\kappa}(y\,{}^{p}_{x}) = \\ = & \mathrm{K}\cdot \operatorname{area} \tilde{\triangle}^{\mathrm{K}}(pxy) - \kappa\cdot \operatorname{area} \tilde{\triangle}^{\kappa}(pxy).\end{aligned}$$

Thus, ❷ follows since

$$0 \leqslant \operatorname{area} \tilde{\triangle}^{\kappa}(pxy) \leqslant \operatorname{area} \tilde{\triangle}^{\mathrm{K}}(pxy),$$

where the latter area is at most $|p-x|\cdot|p-y|$. □

5.3.2. Triangle inequality for angles. *Let $[px^1]$, $[px^2]$ and $[px^3]$ be three geodesics in a metric space. If all of the angles $\alpha^{ij} = \measuredangle[p\,{}^{x^i}_{x^j}]$ are defined then they satisfy the triangle inequality:*

$$\alpha^{13} \leqslant \alpha^{12} + \alpha^{23}.$$

Proof. Since $\alpha^{13} \leqslant \pi$, we can assume that $\alpha^{12} + \alpha^{23} < \pi$. Set $\gamma^i = \operatorname{geod}_{[px^i]}$. Given any $\varepsilon > 0$, for all sufficiently small $t, \tau, s \in \mathbb{R}_+$ we have

$$\begin{aligned}|\gamma^1(t) - \gamma^3(\tau)| \leqslant & |\gamma^1(t) - \gamma^2(s)| + |\gamma^2(s) - \gamma^3(\tau)| < \\ < & \sqrt{t^2 + s^2 - 2\cdot t\cdot s\cdot \cos(\alpha^{12} + \varepsilon)} + \\ & + \sqrt{s^2 + \tau^2 - 2\cdot s\cdot \tau\cdot \cos(\alpha^{23} + \varepsilon)} \leqslant\end{aligned}$$

Below we define $s(t,\tau)$ so that for $s = s(t,\tau)$, this chain of inequalities can be continued as follows:

$$\leqslant \sqrt{t^2 + \tau^2 - 2\cdot t\cdot \tau\cdot \cos(\alpha^{12} + \alpha^{23} + 2\cdot\varepsilon)}.$$

Thus for any $\varepsilon > 0$,

$$\alpha^{13} \leqslant \alpha^{12} + \alpha^{23} + 2\cdot\varepsilon.$$



Hence the result follows.

To define $s(t,\tau)$, consider three rays $\tilde\gamma^1$, $\tilde\gamma^2$, $\tilde\gamma^3$ on a Euclidean plane starting at one point, such that $\measuredangle(\tilde\gamma^1, \tilde\gamma^2) = \alpha^{12} + \varepsilon$, $\measuredangle(\tilde\gamma^2, \tilde\gamma^3) = \alpha^{23} + \varepsilon$ and $\measuredangle(\tilde\gamma^1, \tilde\gamma^3) = \alpha^{12} + \alpha^{23} + 2\cdot\varepsilon$. We parametrize each ray by the distance from the starting point. Given two positive numbers $t, \tau \in \mathbb{R}_+$, let $s = s(t,\tau)$ be

the number such that $\tilde\gamma^2(s) \in [\tilde\gamma^1(t)\,\tilde\gamma^3(\tau)]$. Clearly $s \leqslant \max\{t,\tau\}$, so t,τ,s may be taken sufficiently small. $\square$

5.3.3. Exercise. *Prove that the sum of adjacent angles is at least* π.

More precisely: let $\mathcal{X}$ *be a complete length space and* $p,x,y,z \in \mathcal{X}$. *If* $p \in\,]xy[$, *then*

$$\measuredangle[p\,{}^{x}_{z}] + \measuredangle[p\,{}^{y}_{z}] \geqslant \pi$$

whenever each angle on the left-hand side is defined.

5.3.4. First variation inequality. *Assume for hinge* $[q\,{}^{p}_{x}]$, *the angle* $\alpha = \measuredangle[q\,{}^{p}_{x}]$ *is defined. Then*

$$|p - \operatorname{geod}_{[qx]}(t)| \leqslant |q-p| - t\cdot\cos\alpha + o(t).$$

Proof. Take sufficiently small $\varepsilon > 0$. For all sufficiently small $t > 0$, we have

$$\begin{aligned}|\operatorname{geod}_{[qp]}(t/\varepsilon) - \operatorname{geod}_{[qx]}(t)| &\leqslant \tfrac{t}{\varepsilon}\cdot\sqrt{1+\varepsilon^2-2\cdot\varepsilon\cdot\cos\alpha} + o(t) \leqslant \\ &\leqslant \tfrac{t}{\varepsilon} - t\cdot\cos\alpha + t\cdot\varepsilon.\end{aligned}$$

Applying the triangle inequality, we get

$$\begin{aligned}|p - \operatorname{geod}_{[qx]}(t)| &\leqslant |p - \operatorname{geod}_{[qp]}(t/\varepsilon)| + |\operatorname{geod}_{[qp]}(t/\varepsilon) - \operatorname{geod}_{[qx]}(t)| \leqslant \\ &\leqslant |p-q| - t\cdot\cos\alpha + t\cdot\varepsilon\end{aligned}$$

for any $\varepsilon > 0$ and all sufficiently small t. Hence the result. $\square$

5.4 Constructions

Cone. The *Euclidean cone* $\mathcal{Y} = \operatorname{Cone}\mathcal{X}$ over a metric space $\mathcal{X}$ is defined as the metric space whose underlying set consists of equivalence classes in $[0,\infty)\times\mathcal{X}$ with the equivalence relation "$\sim$" given by $(0,p) \sim (0,q)$ for any points $p,q \in \mathcal{X}$, and whose metric is given by the cosine rule

$$|(s,p)-(t,q)|_{\mathcal{Y}} = \sqrt{s^2+t^2-2\cdot s\cdot t\cdot\cos\alpha},$$

where $\alpha = \min\{\pi, |p-q|_{\mathcal{X}}\}$.

The point in $\operatorname{Cone}\mathcal{X}$ formed by the equivalence class of $\{0\}\times\mathcal{X}$ is called the *tip of the cone* and is denoted by 0 or $0_{\mathcal{Y}}$. The distance $|0-v|_{\mathcal{Y}}$ is called the norm of v and is denoted by $|v|$ or $|v|_{\mathcal{Y}}$.

5.5 Space of directions and tangent space

Let $\mathcal{X}$ be a metric space. If the angle $\measuredangle[p\,{}^{x}_{y}]$ is defined for any hinge $[p\,{}^{x}_{y}]$ in $\mathcal{X}$, then we will say that the space $\mathcal{X}$ has *defined angles*.

Let $\mathcal{X}$ be a space with defined angles. For $p \in \mathcal{X}$, consider the set $\mathfrak{S}_p$ of all nontrivial unit-speed geodesics starting at p. By 5.3.2, the triangle inequality holds for $\measuredangle$ on $\mathfrak{S}_p$, that is, $(\mathfrak{S}_p, \measuredangle)$ forms a pseudometric space.

The metric space corresponding to $(\mathfrak{S}_p, \measuredangle)$ is called the *space of geodesic directions* at p, denoted by Σ'_p or $\Sigma'_p\mathcal{X}$. The elements of Σ'_p are called *geodesic directions* at p. Each geodesic direction is formed by an equivalence class of geodesics starting from p for the equivalence relation

$$[px] \sim [py] \quad \Longleftrightarrow \quad \measuredangle[p\,{}^{x}_{y}] = 0.$$

The completion of Σ'_p is called the *space of directions* at p and is denoted by Σ_p or $\Sigma_p\mathcal{X}$. The elements of Σ_p are called *directions* at p.

The Euclidean cone $\operatorname{Cone}\Sigma_p$ over the space of directions Σ_p is called the *tangent space* at p and denoted by T_p or $\mathrm{T}_p\mathcal{X}$.

The tangent space T_p could be also defined directly, without introducing the space of directions. To do so, consider the set $\mathfrak{T}_p$ of all geodesics starting at p, with arbitrary speed. Given $\alpha, \beta \in \mathfrak{T}_p$, set

$$|\alpha - \beta|_{\mathfrak{T}_p} = \lim_{\varepsilon \to 0} \frac{|\alpha(\varepsilon) - \beta(\varepsilon)|_{\mathcal{X}}}{\varepsilon}. \tag{❶}$$

If the angles in $\mathcal{X}$ are defined, then so is the limit in ❶. In this case it defines a pseudometric on $\mathfrak{T}_p$.

The corresponding metric space admits a natuaral isometric identification with the cone $\mathrm{T}'_p = \operatorname{Cone}\Sigma'_p$. The elements of T'_p are formed by the equivalence classes for the relation

$$\alpha \sim \beta \quad \Longleftrightarrow \quad |\alpha(t) - \beta(t)|_{\mathcal{X}} = o(t).$$

The completion of T'_p is therefore naturally isometric to T_p.

The elements of T_p will be called *tangent vectors* at p, despite that T_p is only a cone — not a vector space. The elements of T'_p will be called *geodesic tangent vectors* at p.

5.6 Velocity of curves

5.6.1. Definition. *Let $\mathcal{X}$ be a metric space. Assume $\alpha\colon [0, a) \to \mathcal{X}$ for some $a > 0$ is a function, not necessarily continuous, such that $\alpha(0) = p$. We say that $v \in \mathrm{T}_p$ is the right derivative of α at 0, briefly*

$\alpha^+(0) = v$, if for some (and therefore any) sequence of vectors $v_n \in \in \mathrm{T}'_p$, with corresponding geodesics γ_n, such that $v_n \to v$ as $n \to \infty$, we have

$$\varlimsup_{\varepsilon\to 0+} \frac{|\alpha(\varepsilon) - \gamma_n(\varepsilon)|_{\mathcal{X}}}{\varepsilon} \to 0 \quad \text{as} \quad n \to \infty.$$

We define right and left derivatives $\alpha^+(t_0)$ and $\alpha^-(t_0)$ of α at $t_0 \in \mathbb{I}$ as

$$\alpha^{\pm}(t_0) = \check{\alpha}^+(0),$$

where $\check{\alpha}(t) = \alpha(t_0 \pm t)$.

The sign convention is not quite standard; if α is a smooth curve in a Riemannian manifold then we have

$$\alpha^+(t) = -\alpha^-(t).$$

Note that if γ is a geodesic starting at p and the tangent vector $v \in \mathrm{T}'_p$ corresponds to γ, then $\gamma^+(0) = v$.

5.6.2. Exercise. *Assume $\mathcal{X}$ is a metric space with defined angles, and let $\alpha, \beta\colon [0, a) \to \mathcal{X}$ be two maps such that the right derivatives $\alpha^+(0)$, $\beta^+(0)$ are defined and $\alpha^+(0) = \beta^+(0)$. Show that*

$$|\alpha(t) - \beta(t)|_{\mathcal{X}} = o(t).$$

5.6.3. Proposition. *Let $\mathcal{X}$ be a metric space with defined angles and $p \in \mathcal{X}$. Then for any tangent vector $v \in \mathrm{T}_p\mathcal{X}$ there is a map $\alpha\colon [0, \varepsilon) \to \mathcal{X}$ such that $\alpha^+(0) = v$.*

Proof. If $v \in \mathrm{T}'_p$, then for the corresponding geodesic α we have $\alpha^+(0) = v$.

Given $v \in \mathrm{T}_p$, construct a sequence $v_n \in \mathrm{T}'_p$ such that $v_n \to v$, and let γ_n be the sequence of corresponding geodesic.

The needed map α can be found among the maps such that $\alpha(0) = = p$ and

$$\alpha(t) = \gamma_n(t) \quad \text{if} \quad \varepsilon_{n+1} \leqslant t < \varepsilon_n,$$

where (ε_n) is a decreasing sequence converging to 0 as $n \to \infty$. In order to satify the condition one has to choose the sequence ε_n converging to 0 very fast. □

5.6.4. Definition. *Let $\mathcal{X}$ be a metric space and $\alpha\colon \mathbb{I} \to \mathcal{X}$ be a curve.*

For $t_0 \in \mathbb{I}$, if $\alpha^+(t_0)$ or $\alpha^-(t_0)$ or both are defined, we say correspondingly that α is right *or* left *or* both-sided differentiable *at t_0. In the exceptional cases where t_0 is the left (respectively right) end of $\mathbb{I}$, α is by definition left (respectively right) differentiable at t_0.*

5.6.5. Exercise. *Assume $\mathcal{X}$ is a metric space with defined angles. Show that any geodesic $\gamma\colon \mathbb{I} \to \mathcal{X}$ is both-sided differentiable everywhere.*

5.7 Differential

5.7.1. Definition. *Let $\mathcal{X}$ be a metric space with defined angles and $f\colon \mathcal{X} \to\!\!\!\!\circ\, \mathbb{R}$ be a subfunction, $p \in \operatorname{Dom} f$ and $\mathbb{I}$ be a real interval. A function $\varphi\colon \mathrm{T}_p \to \mathbb{R}$ is called differential of f at p (briefly $\varphi = \boldsymbol{d}_p f$) if for any map $\alpha\colon \mathbb{I} \to \mathcal{X}$ such that $\alpha(0) = p$ and $\alpha^+(0)$ is defined, we have*

$$(f \circ \alpha)^+(0) = \varphi(\alpha^+(0)).$$

5.7.2. Proposition. *Let $f\colon \mathcal{X} \to\!\!\!\!\circ\, \mathbb{R}$ be a locally Lipschitz semiconcave subfunction. Then differential $\boldsymbol{d}_p f$ is uniquely defined for any $p \in \operatorname{Dom} f$. Moreover,*

a) The differential $\boldsymbol{d}_p f\colon \mathrm{T}_p \to \mathbb{R}$ is Lipschitz and the Lipschitz constant of $\boldsymbol{d}_p f\colon \mathrm{T}_p \to \mathbb{R}$ does not exceed the Lipschitz constant of f in a neighborhood of p.

b) $\boldsymbol{d}_p f\colon \mathrm{T}_p \to \mathbb{R}$ is a positive homogenius function; that is, for any $\mathfrak{z} \geqslant 0$ and $v \in \mathrm{T}_p$ we have

$$\mathfrak{z}\cdot \boldsymbol{d}_p f(v) = \boldsymbol{d}_p f(\mathfrak{z}\cdot v).$$

Proof. Passing to a subdomain of f if nesessary, we can assume that f is $\pounds$-Lipschitz and λ-concave for some $\pounds, \lambda \in \mathbb{R}$.

Take a geodessic γ starting at p that lies in $\operatorname{Dom} f$. Since $f \circ \gamma$ is semiconcave, the rigth derivative $(f \circ \gamma)^+(0)$ is defined. Since f is $\pounds$-Lipschitz, we have

❶ $$|(f \circ \gamma)^+(0) - (f \circ \gamma_1)^+(0)| \leqslant \pounds\cdot|\gamma^+(0) - \gamma_1^+(0)|$$

for any other geodesic γ_1 starting at p.

Define $\varphi\colon \mathrm{T}'_p \to \mathbb{R}\colon \gamma^+(0) \mapsto (f \circ \gamma)^+(0)$. From ❶, φ is a $\pounds$-Lipschtz function defined on T'_p. Thus, we can extend φ to a whole T_p as a $\pounds$-Lipschitz function.

It remains to show that φ is differential of f at p. Assume $\alpha\colon [0, a) \to \mathcal{X}$ is a map such that $\alpha(0) = p$ and $\alpha^+(0) = v \in \mathrm{T}_p$. Let $\gamma_n \in \Gamma_p$ be a sequence of geodesics as in the definition 5.6.1; that is, if

$$v_n = \gamma_n^+(0) \quad \text{and} \quad a_n = \varlimsup_{t \to 0+} |\alpha(t) - \gamma_n(t)|/t$$

then $a_n \to 0$ and $v_n \to v$ as $n \to \infty$. Then

$$\varphi(v) = \lim_{n\to\infty} \varphi(v_n),$$

$$f \circ \gamma_n(t) = f(p) + \varphi(v_n)\cdot t + o(t),$$
$$|f \circ \alpha(t) - f \circ \gamma_n(t)| \leqslant \pounds\cdot|\alpha(t) - \gamma_n(t)|.$$

Hence

$$f \circ \alpha(t) = f(p) + \varphi(v)\cdot t + o(t)$$

□

5.8 Remarks

In Alexandrov geometry, angles defined as in Section 5.3 always exist (see Theorem 7.3.1*c* and Corollary 8.3.2*b*).

For general metric spaces, this angle may not exist, and it is more natural to consider the *upper angle* defined as

$$\measuredangle^{\mathrm{up}}[p\,{}^{x}_{y}] := \varlimsup_{\bar x,\bar y\to p} \tilde\measuredangle^{\kappa}(p\,{}^{\bar x}_{\bar y}),$$

where $\bar x \in\,]px]$ and $\bar y \in\,]py]$. The triangle inequality (5.3.2) holds for upper angles as well.

Chapter 6

Dimension theory

In Section 6.1, we give definitions of different type of dimension-like invariants of metric spaces and state general relations between them.

6.1 Definitions

The proof of most of the statements in this section can be found in the book of Witold Hurewicz and Henry Wallman [6], the rest follows directly from the definitions.

6.1.1. Hausdorff dimension. *Let $\mathcal{X}$ be a metric space. Its Hausdorff dimension is defined as*

$$\operatorname{HausDim}\mathcal{X} = \sup\{\,\alpha\in\mathbb{R} : \operatorname{HausMes}_\alpha(\mathcal{X}) > 0\,\},$$

where $\operatorname{HausMes}_\alpha$ *denotes* α*-dimensional Hausdorff measure.*

Let $\mathcal{X}$ be a metric space and $\{V_\beta\}_{\beta\in\mathcal{B}}$ be its open cover. Let us recall two notions in general topology:

- ◇ The *order* of $\{V_\beta\}$ is supremum of all integers n such that there is a collection of $n+1$ elements of $\{V_\beta\}$ with nonempy intersection.
- ◇ An open cover $\{W_\alpha\}_{\alpha\in\mathcal{A}}$ of $\mathcal{X}$ is called *refinement* of $\{V_\beta\}_{\beta\in\mathcal{B}}$ if for any $\alpha\in\mathcal{A}$ there is $\beta\in\mathcal{B}$ such that $W_\alpha\subset V_\beta$.

6.1.2. Topological dimension. *Let $\mathcal{X}$ be a metric space. The topological dimension of $\mathcal{X}$ is defined as the minimum of nonnegative integers n, such that for any finite open cover of $\mathcal{X}$ there is a finite open refinement with order n.*

If no such n exists, the topological dimension of $\mathcal{X}$ is infinite.

The topological dimension of $\mathcal{X}$ will be denoted by $\operatorname{TopDim}\mathcal{X}$.

The invariants satisfying the following two statements 6.1.3 and 6.1.4 are commonly called "dimension"; by that reason we call them axioms.

6.1.3. Normalization axiom. *For any $m \in \mathbb{Z}_{\geqslant 0}$,*

$$\operatorname{TopDim} \mathbb{E}^m = \operatorname{HausDim} \mathbb{E}^m = m.$$

6.1.4. Cover axiom. *If $\{A_n\}_{n=1}^{\infty}$ is a countable closed cover of $\mathcal{X}$, then*

$$\operatorname{TopDim} \mathcal{X} = \sup_n \{\operatorname{TopDim} A_n\},$$
$$\operatorname{HausDim} \mathcal{X} = \sup_n \{\operatorname{HausDim} A_n\}.$$

On product spaces. Let us mention that the following two inequalities

$$\operatorname{TopDim}(\mathcal{X} \times \mathcal{Y}) \leqslant \operatorname{TopDim} \mathcal{X} + \operatorname{TopDim} \mathcal{Y},$$

and

$$\operatorname{HausDim}(\mathcal{X} \times \mathcal{Y}) \geqslant \operatorname{HausDim} \mathcal{X} + \operatorname{HausDim} \mathcal{Y},$$

hold for any pair of metric spaces $\mathcal{X}$ and $\mathcal{Y}$.

These inequalities in might be strict. For the topological dimension it holds for apair of Pontryagin surfaces constructed in [7]. For Hausdorff dimension, an example was constructed by Abram Besicovitch and Pat Moran in [8].

The following theorem follows from [6, theorems V 8 and VII 2].

6.1.5. Szpilrajn's theorem. *Let $\mathcal{X}$ be a separable metric space. Assume* $\operatorname{TopDim} \mathcal{X} \geqslant m$. *Then* $\operatorname{HausMes}_m \mathcal{X} > 0$.

In particular, $\operatorname{TopDim} \mathcal{X} \leqslant \operatorname{HausDim} \mathcal{X}$.

In fact it is true that for any separable metric space $\mathcal{X}$ we have

$$\operatorname{TopDim} \mathcal{X} = \inf\{\operatorname{HausDim} \mathcal{Y}\},$$

where infimum is taken over all metric spaces $\mathcal{Y}$ homeomorphic to $\mathcal{X}$.

6.1.6. Definition. *Let $\mathcal{X}$ be a metric space and $F\colon \mathcal{X} \to \mathbb{R}^m$ be a continuous map. A point $\boldsymbol{z} \in \operatorname{Im} F$ is called a* stable value *of F if there is $\varepsilon > 0$ such that $\boldsymbol{z} \in \operatorname{Im} F'$ for any ε-close continuous map $F'\colon \mathcal{X} \to \mathbb{R}^m$; that is, $|F'(x) - F(x)| < \varepsilon$ for all $x \in \mathcal{X}$.*

The following theorem follows from [6, theorems VI 1&2]. (This theorem also holds for non-separable metric spaces [9] or [10, 3.2.10]).

6.1.7. Stable value theorem. *Let $\mathcal{X}$ be a separable metric space. Then* $\operatorname{TopDim}\mathcal{X} \geqslant m$ *if and only if there is a map $F\colon \mathcal{X} \to \mathbb{R}^m$ with a stable value.*

6.1.8. Proposition. *Suppose $\mathcal{X}$ and $\mathcal{Y}$ are metric spaces and $\Phi\colon \mathcal{X} \to \mathcal{Y}$ satisfies*

$$|\Phi(x) - \Phi(x')| \geqslant \varepsilon\cdot|x - x'|$$

for some fixed $\varepsilon > 0$ and any pair $x, x' \in \mathcal{X}$. Then

$$\operatorname{HausDim}\mathcal{X} \leqslant \operatorname{HausDim}\mathcal{Y}.$$

In particular, if there is a Lipschitz onto map $\mathcal{Y} \to \mathcal{X}$, then

$$\operatorname{HausDim}\mathcal{X} \leqslant \operatorname{HausDim}\mathcal{Y}.$$

6.2 Linear dimension

In addition to HausDim and TopDim, we will use yet another one, which we call *linear dimension.*

Recall that *cone map* is a map between cones respecting the cone multiplication.

6.2.1. Definition of linear dimension. *Let $\mathcal{X}$ be a metric space. The* linear dimension *of $\mathcal{X}$ (denoted by* $\operatorname{LinDim}\mathcal{X}$*) is defined exact upper bound on $m \in \mathbb{Z}_{\geqslant 0}$ such that there is an isometric cone embedding $\mathbb{E}^m \hookrightarrow \mathrm{T}_p\mathcal{X}$ for some $p \in \mathcal{X}$; here $\mathbb{E}^m$ denotes m-dimensional Euclidean space and $\mathrm{T}_p\mathcal{X}$ denotes tangent space of $\mathcal{X}$ at p (defined in Section 5.5).*

Note that LinDim takes values in $\mathbb{Z}_{\geqslant 0} \cup \{\infty\}$.

For general metric spaces, LinDim has no relations to HausDim and TopDim. Also, LinDim does not satisfy the cover axiom (6.1.4). For LinDim, an inverse of the product inequality holds; that is,

$$\operatorname{LinDim}(\mathcal{X} \times \mathcal{Y}) \geqslant \operatorname{LinDim}\mathcal{X} + \operatorname{LinDim}\mathcal{Y} \qquad ❶$$

for any two metric spaces $\mathcal{X}$ and $\mathcal{Y}$; that is easy to check. According to the following exercise, the inequality ❶ might be strict for some spaces [11].

6.2.2. Exercise. *Construct two norms on $\mathbb{R}^{10}$ such that non of the corresponding metric spaces $\mathcal{X}$ and $\mathcal{Y}$ have a isometric copy of $\mathbb{E}^2$ but $\mathcal{X} \times \mathcal{Y}$ has isometric copy of $\mathbb{E}^{10}$.*

The linear dimension will be applied only to Alexandrov spaces and to their open subsets (in both cases of curvature bounded below and above). As we shall see, in all these cases, LinDim behaves nicely and is easy to work with.

Remarks. Linear dimension was first introduced by Conrad Plaut in [12] under the name *local dimension*. *Geometric dimension* introduced in by Bruce Kleiner in [13] is closely related; it coincides with the linear dimension for CBB and CAT spaces.

One could modify the definition of linear dimension by taking an arbitrary n-dimensional Banach space instead of the Euclidean n-space. Such a definition makes more sense for general metric spaces. This dimension is additive with respect to the direct product (that is, we always have equality in ❶).

For Alexandrov spaces (either CBB or CAT) this modification is equivalent to the definition we use in this book.

Part II

Definitions and constructions

Chapter 7

Definitions of curvature bounded below

7.1 Four-point comparison

Recall (Section 5.2) that the model angle $\measuredangle^{\kappa}(p\,{}^{x}_{y})$ is defined if

$$|p-x|+|p-y|+|x-y|<\varpi^{\kappa}.$$

7.1.1. Four-point comparison. *A quadruple of points p,x^1,x^2,x^3 in a metric space satisfies* CBB(κ) *comparison if*

$$\measuredangle^{\kappa}(p\,{}^{x^1}_{x^2})+\measuredangle^{\kappa}(p\,{}^{x^2}_{x^3})+\measuredangle^{\kappa}(p\,{}^{x^3}_{x^1})\leqslant 2\cdot\pi. \tag{1}$$

or at least one of the model angles $\measuredangle^{\kappa}(p\,{}^{x^i}_{x^j})$ is not defined.

7.1.2. Definition. *Let $\mathcal{L}$ be a metric space.*

a) $\mathcal{L}$ is CBB(κ) *if any quadruple $p,x^1,x^2,x^3\in\mathcal{L}$ satisfies* CBB(κ) *comparison.*

b) $\mathcal{L}$ is locally CBB(κ) *if any point $q\in\mathcal{L}$ admits a neighborhood $\Omega\ni q$ such that any quadruple $p,x^1,x^2,x^3\in\Omega$ satisfies* CBB(κ) *comparison.*

c) $\mathcal{L}$ is a CBB *space if $\mathcal{L}$ is* CBB(κ) *for some $\kappa\in\mathbb{R}$.*

Remarks

- CBB(κ) length spaces are often called *spaces with curvature $\geqslant\kappa$ in the sence of Alexandrov.* These spaces will usually be denoted by $\mathcal{L}$, for $\mathcal{L}$ower curvature bound.

- ◇ In the definition of CBB(κ), when $\kappa > 0$ most authors assume in addition that the diameter is at most ϖ^κ. For complete length spaces, the latter means that it is not isometric to one of the exceptional spaces, see 7.8.1. We do not make this assumption. In particular, we consider the real line to have curvature $\geqslant 1$.
- ◇ If $\kappa < \mathrm{K}$, then any complete length CBB(K) space is CBB(κ). Moreover directly from the definition it follows that if $\mathrm{K} \leqslant 0$, then any CBB(K) is CBB(κ). However, in the case $\mathrm{K} > 0$ the latter statement does not hold and the former statement is not trivial; it will be proved in 7.6.4.

7.1.3. Exercise. *Let $\mathcal{L}$ be a metric space. Show that $\mathcal{L}$ is* CBB(κ) *if for any quadruple of points $p, x^1, x^2, x^3 \in \mathcal{L}$ such that*

$$|p - x^1|, |p - x^2|, |p - x^3| < \tfrac{\varpi^\kappa}{2},$$

there is a quadruple of points $q, y^1, y^2, y^3 \in \mathbb{M}^2(\kappa)$ such that

$$|p - x^i| = |q - y^i| \quad \text{and} \quad |x^i - x^j| \leqslant |y^i - y^j|$$

for all i and j.

The exercise above is a partial case of $(1{+}n)$-point comparison (9.2.1).

7.1.4. Exercise. *Let $\mathcal{L}$ be a metric space. Show that $\mathcal{L}$ is* CBB(0) *if and only if*

$$\operatorname{area} \tilde{\triangle}^0(xyz) \leqslant \operatorname{area} \tilde{\triangle}^0(pxy) + \operatorname{area} \tilde{\triangle}^0(pyz) + \operatorname{area} \tilde{\triangle}^0(pzx)$$

for any 4 distinct points $p, x, y, z \in \mathcal{L}$.

The following exercise decribes the first known definition of spaces with curvature bounded below; it was given by Abraham Wald in [14].

7.1.5. Exercise. *Let $\mathcal{L}$ be a metric space and $\kappa \leqslant 0$. Prove that $\mathcal{L}$ is* CBB(κ) *if and only if any quadruple of points $p, q, r, s \in \mathcal{L}$ can be isometrically embedded into some $\mathbb{M}^2(\mathrm{K})$ for some $\mathrm{K} \geqslant \kappa$.*

Is the same true for $\kappa > 0$, what is the difference?

7.1.6. Exercise. *Let $\mathcal{L}$ be $\mathbb{R}^m$ with the metric defined by a norm. Show that $\mathcal{L}$ is a complete length* CBB *space if and only if $\mathcal{L} \overset{iso}{=\!=} \mathbb{E}^m$.*

Recall that ω denotes a selective ultrafilter on $\mathbb{N}$, which is fixed once for all. The following proposition follows directly from the definition of CBB(κ) comparison and the definitions of ω-limit and ω-power given in Section 3.2.

7.1.7. Proposition. *Let $\mathcal{L}_n$ be a* CBB(κ_n) *space for each n. Assume $\mathcal{L}_n \to \mathcal{L}_\omega$ and $\kappa_n \to \kappa_\omega$ as $n \to \omega$. Then $\mathcal{L}_\omega$ is* CBB(κ_ω).

Moreover, a metric space $\mathcal{L}$ is CBB(κ) *if and only if so is its ultrapower $\mathcal{L}^\omega$.*

7.1.8. Theorem. *Let $\mathcal{L}$ be a* CBB(κ) *space, $\mathcal{M}$ be a metric space and $\sigma\colon \mathcal{L} \to \mathcal{M}$ be a submetry. Assume p, x^1, x^2, x^3 is a quadruple of points in $\mathcal{M}$ such that $|p - x^i| < \frac{\varpi^\kappa}{2}$ for any i. Then the quadruple satisfies* CBB(κ) *comparison.*

In particular,

a) The space $\mathcal{M}$ is locally CBB(κ)*. Moreover, any open ball of radius $\frac{\varpi^\kappa}{4}$ in $\mathcal{M}$ is* CBB(κ).

b) If $\kappa \leqslant 0$, then $\mathcal{M}$ is CBB(κ).

Corollary 7.6.5 gives a stronger statement; it states that if $\mathcal{L}$ is a complete length space, then $\mathcal{M}$ is always CBB(κ). The theorem above together with Proposition 4.3.3 imply the following:

7.1.9. Corollary. *Assume that the group G acts isometrically on a* CBB(κ) *space $\mathcal{L}$ and has closed orbits. Then the quotient space $\mathcal{L}/G$ is locally* CBB(κ).

Proof of 7.1.8. Fix a quadruple of points $p, x^1, x^2, x^3 \in \mathcal{M}$ such that $|p - x^i|_< \frac{\varpi^\kappa}{2}$ for any i. Choose arbitrary $\hat{p} \in \mathcal{L}$ such that $\sigma(\hat{p}) = p$.

Since σ is submetry, we can choose the points $\hat{x}^1, \hat{x}^2, \hat{x}^3 \in \mathcal{L}$ such that $\sigma(\hat{x}_i) = x_i$ and

$$|p - x^i|_{\mathcal{M}} \lesseqgtr |\hat{p} - \hat{x}^i|_{\mathcal{L}} \pm \delta$$

for all i and any fixed $\delta > 0$.

Note that

$$|x^i - x^j|_{\mathcal{M}} \leqslant |\hat{x}^i - \hat{x}^j|_{\mathcal{L}} \leqslant |p - x^i|_{\mathcal{M}} + |p - x^j|_{\mathcal{M}} + 2{\cdot}\delta$$

for all i and j.

Since $|p - x^i| < \frac{\varpi^\kappa}{2}$, we can choose $\delta > 0$ above so that the angles $\measuredangle^\kappa(\hat{p}\,{}^{\hat{x}^i}_{\hat{x}^j})$ are defined. Moreover, given $\varepsilon > 0$, the value δ can be chosen in such a way that the inequality

❷ $$\measuredangle^\kappa(p\,{}^{x^i}_{x^j}) < \measuredangle^\kappa(\hat{p}\,{}^{\hat{x}^i}_{\hat{x}^j}) + \varepsilon$$

holds for all i and j.

By CBB(κ) comparison in $\mathcal{L}$, we have

$$\measuredangle^\kappa(\hat{p}\,{}^{\hat{x}^1}_{\hat{x}^2}) + \measuredangle^\kappa(\hat{p}\,{}^{\hat{x}^2}_{\hat{x}^3}) + \measuredangle^\kappa(\hat{p}\,{}^{\hat{x}^3}_{\hat{x}^1}) \leqslant 2{\cdot}\pi.$$

Applying ❷, we get

$$\measuredangle^{\kappa}(p\,{}^{x^1}_{x^2}) + \measuredangle^{\kappa}(p\,{}^{x^2}_{x^3}) + \measuredangle^{\kappa}(p\,{}^{x^3}_{x^1}) < 2\cdot\pi + 3\cdot\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary we have

$$\measuredangle^{\kappa}(p\,{}^{x^1}_{x^2}) + \measuredangle^{\kappa}(p\,{}^{x^2}_{x^3}) + \measuredangle^{\kappa}(p\,{}^{x^3}_{x^1}) \leqslant 2\cdot\pi;$$

that is, the CBB(κ) comparison holds for the quadruple. □

7.2 Geodesics

We are going to show that all complete length CBB spaces have plenty of geodesics in the following sense.

7.2.1. Definition. *A metric space $\mathcal{X}$ is called* G-delta geodesic *if for any point $p \in \mathcal{X}$ there is a dense G-delta set $W_p \subset \mathcal{X}$ such that for any $q \in W_p$ there is a geodesic $[pq]$.*

A metric space $\mathcal{X}$ is called locally G-delta geodesic *if for any point $p \in \mathcal{X}$ there is a G-delta set $W_p \subset \mathcal{X}$ such that W_p is dense in a neighborhood of p and for any $q \in W_p$ there is a geodesic $[pq]$.*

Recall that general complete length spaces might have no geodesics; see Exercise 2.3.6.

7.2.2. Exercise. *Construct a complete length* CBB(0) *space that is not geodesic.*

7.2.3. Definition. *Let $\mathcal{X}$ be a metric space and $p \in \mathcal{X}$. A point $q \in \mathcal{X}$ is called p-*straight *(briefly, $q \in \mathrm{Str}(p)$) if*

$$\varlimsup_{r\to q} \frac{|p-r| - |p-q|}{|q-r|} = 1.$$

For an array of points $x^1, x^2, \dots, x^k$, we use the notation

$$\mathrm{Str}(x^1, x^2, \dots, x^k) = \bigcap_{i=1}^{k} \mathrm{Str}(x^i).$$

7.2.4. Theorem. *Let $\mathcal{L}$ be a complete length* CBB *space and $p \in \mathcal{L}$. Then the set $\mathrm{Str}(p)$ is a dense G-delta set, and for any $q \in \mathrm{Str}(p)$ there is a unique geodesic $[pq]$.*

In particular, $\mathcal{L}$ is G-delta geodesic.

The proof below is very close to the original proof given by Conrad Plaut [12, Th. 27].

Proof. Given a positive integer n, consider the set Ω_n of all points $q \in \mathcal{L}$ such that

$$(1-\tfrac{1}{n})\cdot|q-r| < |p-r| - |p-q| < \tfrac{1}{n}$$

for some $r \in \mathcal{L}$. Clearly Ω_n is open; let us show that Ω_n is dense in $\mathcal{L}$.

Assuming the contrary, there is a point $x \in \mathcal{L}$ such that

$$\mathrm{B}(x,\varepsilon) \cap \Omega_n = \varnothing$$

for some $\varepsilon > 0$. Since $\mathcal{L}$ is a length space, for any $\delta > 0$, there exists a point $y \in \mathcal{L}$ such that

$$|x-y| < \tfrac{\varepsilon}{2} + \delta \quad \text{and} \quad |p-y| < |p-x| - \tfrac{\varepsilon}{2} + \delta.$$

If ε and δ are sufficiently small, then

$$(1-\tfrac{1}{n})\cdot|y-x| < |p-x| - |p-y| < \tfrac{1}{n};$$

that is, $y \in \Omega_n$, a contradiction.

Note that $\mathrm{Str}(p) = \bigcap_{n\in\mathbb{N}} \Omega_n$. It follows that $\mathrm{Str}(p)$ is a dense G-delta set.

Assume $q \in \mathrm{Str}(p)$; let us show that there is a unique geodesic connecting p and q. Note that it is sufficient to show that for all sufficiently small $t > 0$ there is a unique point z such that

❶ $$t = |q-z| = |p-q| - |p-z|.$$

First let us show uniqueness. Assume z and z' both satisfy ❶. Take a sequence $r_n \to q$ such that

$$\frac{|p-r_n| - |p-q|}{|q-r_n|} \to 1.$$

By the triangle inequality,

$$|z-r| - |z-q|, \quad |z'-r| - |z'-q| \geqslant |p-r| - |p-q|;$$

thus, as $n \to \infty$,

$$\frac{|z-r_n| - |z-q|}{|q-r_n|}, \quad \frac{|z'-r_n| - |z'-q|}{|q-r_n|} \to 1.$$

Therefore $\measuredangle^{\kappa}(q\,{}^{z}_{r_n}),\ \measuredangle^{\kappa}(q\,{}^{z'}_{r_n}) \to \pi$. (Here we use that t is small, otherwise if $\kappa > 0$ the angles might be undefined.)

From CBB(κ) comparison (7.1.2), $\measuredangle^\kappa(q\,{}^{z}_{z'}) = 0$ and thus $z = z'$.

The proof of existence is similar. Choose a sequence r_n as above. Since $\mathcal{L}$ is a complete length space, there is a sequence $z_{\bar{k}} \in \mathcal{L}$ such that $|q - z_{\bar{k}}|,\ |p - q| - |p - z_{\bar{k}}| \to t$ as $\bar{k} \to \infty$. Then

$$\lim_{n\to\infty} \lim_{\bar{k}\to\infty} \measuredangle^\kappa(q\,{}^{z_{\bar{k}}}_{r_n}) = \pi.$$

Thus, for any $\varepsilon > 0$ and sufficiently large $n, \bar{k}$, we have $\measuredangle^\kappa(q\,{}^{z_{\bar{k}}}_{r_n}) >$ $> \pi - \varepsilon$. From CBB(κ) comparison (7.1.2), for all large $\bar{k}$ and j, we have $\measuredangle^\kappa(q\,{}^{z_{\bar{k}}}_{z_j}) < 2\cdot\varepsilon$ and thus

$$|z_{\bar{k}} - z_j| < \varepsilon\cdot c(\kappa, t);$$

that is, $\{z_n\}$ converges in itself, and $z = \lim_n z_n$ satisfies ❶. □

7.2.5. Exercise. *Let $\mathcal{L}$ be a complete length* CBB *space and $A \subset \mathcal{L}$ be a closed subset. Show that there is a dense G-delta set $W \subset \mathcal{L}$ such that for any $q \in W$, there is a unique geodesic $[pq]$ with $p \in A$ that realize the distance from q to A; that is, $|p - q| = \operatorname{dist}_A q$.*

7.2.6. Exercise. *Give an alternative proof of Theorem 7.2.4 built on the fact that geodesics do not split (7.7.1) and Corollary 3.3.4.*

Recall that directions of geodesics are defined in Section 5.5.

7.2.7. Exercise. *Assume $\mathcal{L}$ is a complete length* CBB *space, and $[px]$, $[py]$ be two geodesics in the same geodesic direction $\xi \in \Sigma'_p$. Show that*

$$[px] \subset [py] \quad \text{or} \quad [px] \supset [py].$$

7.2.8. Exercise. *Construct a complete length* CBB *space $\mathcal{L}$ with an everywhere dense G-delta set A such that $A \cap]xy[= \varnothing$ for any geodesic $[xy]$ in $\mathcal{L}$.*

7.3 More comparisons

The following theorem makes it easier to use Euclidean intuition in the Alexandrov universe.

7.3.1. Theorem. *If $\mathcal{L}$ is a* CBB(κ) *space, then the following conditions hold for all $p, x, y \in \mathcal{L}$, once the model triangle $\tilde{\triangle}^\kappa(pxy)$ is defined.*

a) (adjacent angle comparison) for any geodesic $[xy]$ and $z \in$ $\in\,]xy[$, $z \neq p$ we have

$$\measuredangle^\kappa(z\,{}^{p}_{x}) + \measuredangle^\kappa(z\,{}^{p}_{y}) \leqslant \pi.$$

b) (point-on-side comparison) for any geodesic $[xy]$ *and* $z \in$ $\in\,]xy[$*, we have*

$$\measuredangle^{\kappa}(x\,{}^{p}_{y}) \leqslant \measuredangle^{\kappa}(x\,{}^{p}_{z});$$

or, equivalently,

$$|\tilde{p} - \tilde{z}| \leqslant |p - z|,$$

where $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^{\kappa}(pxy)$*,* $\tilde{z} \in\,]\tilde{x}\tilde{y}[$*,* $|\tilde{x} - \tilde{z}| = |x - z|$*.*

c) (hinge comparison) for any hinge $[x\,{}^{p}_{y}]$*, the angle* $\measuredangle[x\,{}^{p}_{y}]$ *is defined and*

$$\measuredangle[x\,{}^{p}_{y}] \geqslant \measuredangle^{\kappa}(x\,{}^{p}_{y}),$$

or equivalently

$$\tilde{\Upsilon}^{\kappa}[x\,{}^{p}_{y}] \geqslant |p - y|.$$

Moreover, if $z \in\,]xy[$ *and* $z \neq p$*, then*

$$\measuredangle[z\,{}^{p}_{y}] + \measuredangle[z\,{}^{p}_{x}] \leqslant \pi$$

for any two hinges $[z\,{}^{p}_{y}]$ *and* $[z\,{}^{p}_{x}]$ *with common side* $[zp]$*.*

Moreover, in each case the converse holds if $\mathcal{L}$ *is G-delta geodesic. That is if one of the conditions (a), (b) or (c) holds in a G-delta geodesic space* $\mathcal{L}$*, then* $\mathcal{L}$ *is* CBB(κ)*.*

Remarks. Monotonicity of the model angle with respect to adjacent sidelengths (7.3.2) was named the *convexity property* by Alexandrov.

A bit weaker form of (c) is given in 7.5.4. See also Problem 7.9.1.

Proof; (a). Since $z \in\,]xy[$, we have $\measuredangle^{\kappa}(z\,{}^{x}_{y}) = \pi$. Thus, CBB($\kappa$) comparison

$$\measuredangle^{\kappa}(z\,{}^{x}_{y}) + \measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) \leqslant 2\cdot\pi$$

implies

$$\measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) \leqslant \pi.$$

(a) $\Leftrightarrow$ *(b).* Follows from Alexandrov's lemma (5.2.1).

(a) + (b) $\Rightarrow$ *(c).* From (b) we get that for $\bar{p} \in\,]xp]$ and $\bar{y} \in\,]xy]$, the function $(|x-\bar{p}|, |x-\bar{y}|) \mapsto \measuredangle^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}})$ is nonincreasing in each argument. In particular, $\measuredangle[x\,{}^{p}_{y}] = \sup\{\measuredangle^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}})\}$. Hence $\measuredangle[x\,{}^{p}_{y}]$ is defined and is at least $\measuredangle^{\kappa}(x\,{}^{p}_{y})$.

From above and (a), it follows that

$$\measuredangle[z\,{}^{p}_{y}] + \measuredangle[z\,{}^{p}_{x}] \leqslant \pi.$$



Converse. Assume first that $\mathcal{L}$ is geodesic. Consider a point $w \in\,]pz[$ close to p. From (c), it follows that

$$\measuredangle[w\,{}^{x}_{z}] + \measuredangle[w\,{}^{x}_{p}] \leqslant \pi \quad\text{and}\quad \measuredangle[w\,{}^{y}_{z}] + \measuredangle[w\,{}^{y}_{p}] \leqslant \pi.$$

Since $\measuredangle[w\,{}^{x}_{y}] \leqslant \measuredangle[w\,{}^{x}_{p}] + \measuredangle[w\,{}^{y}_{p}]$ (see 5.3.2), we get

$$\measuredangle[w\,{}^{x}_{z}] + \measuredangle[w\,{}^{y}_{z}] + \measuredangle[w\,{}^{x}_{y}] \leqslant 2\cdot\pi.$$

Applying the first inequality in (c),

$$\tilde{\measuredangle}^{\kappa}(w\,{}^{x}_{z}) + \tilde{\measuredangle}^{\kappa}(w\,{}^{y}_{z}) + \tilde{\measuredangle}^{\kappa}(w\,{}^{x}_{y}) \leqslant 2\cdot\pi.$$

Passing to the limits $w \to p$, we have

$$\tilde{\measuredangle}^{\kappa}(p\,{}^{x}_{z}) + \tilde{\measuredangle}^{\kappa}(p\,{}^{y}_{z}) + \tilde{\measuredangle}^{\kappa}(p\,{}^{x}_{y}) \leqslant 2\cdot\pi.$$

If $\mathcal{L}$ is only G-delta geodesic, we can apply the above arguments to sequences of points $p_n, w_n \to p$, $x_n \to x$, $y_n \to y$ such that $[p_n z]$, $w_n \in\,]zp_n[$ and $[x_n w_n]$, $[y_n w_n]$ exist, and then pass to the limit as $n \to \infty$. $\square$

7.3.2. Angle-sidelength monotonicity. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $p, x, y \in \mathcal{L}$, $\tilde{\triangle}^{\kappa}(pxy)$ is defined and there is a geodesic $[xy]$. Then for $\bar{y} \in\,]xy]$ the function*

$$|x - \bar{y}| \mapsto \tilde{\measuredangle}^{\kappa}(x\,{}^{p}_{\bar{y}})$$

is nonincreasing.

In particular, if geodesic $[xp]$ exists and $\bar{p} \in\,]xp]$, then

a) the function

$$(|x - \bar{y}|, |x - \bar{p}|) \mapsto \tilde{\measuredangle}^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}})$$

is nonincresing in each argument

b) The angle $\measuredangle[x\,{}^{p}_{y}]$ is defined and

$$\measuredangle[x\,{}^{p}_{y}] = \sup\left\{\, \tilde{\measuredangle}^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}}) : \bar{p} \in\,]xp],\ \bar{y} \in\,]xy] \,\right\}.$$

The proof is contained in the first part of (a)+(b)⇒(c) of Theorem 7.3.1.

7.3.3. Exercise. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $p, x, y \in$ $\in \mathcal{L}$ and $v, w \in\,]xy[$. Prove that*

$$\tilde{\measuredangle}^{\kappa}(x\,{}^{y}_{p}) = \tilde{\measuredangle}^{\kappa}(x\,{}^{v}_{p}) \quad\Leftrightarrow\quad \tilde{\measuredangle}^{\kappa}(x\,{}^{y}_{p}) = \tilde{\measuredangle}^{\kappa}(x\,{}^{w}_{p}).$$

7.3.4. Advanced exercise. *Construct a geodesic space $\mathcal{X}$ that is not* CBB(0)*, but meets the following condition: for any 3 points $p, x, y \in$ $\in \mathcal{X}$ there is a geodesic $[xy]$ such that for any $z \in\,]xy[$*

$$\tilde{\measuredangle}^{0}(z\,{}^{p}_{x}) + \tilde{\measuredangle}^{0}(z\,{}^{p}_{y}) \leqslant \pi.$$

7.3.5. Advanced exercise. *Let $\mathcal{L}$ be a complete length space such that for any quadrouple $p, x, y, z \in \mathcal{L}$ the following inequality holds*

❶ $|p-x|^2 + |p-y|^2 + |p-z|^2 \geqslant \frac{1}{3}\cdot\left[|x-y|^2 + |y-z|^2 + |z-x|^2\right].$

Prove that $\mathcal{L}$ is CBB(0).

Construct a 4-point metric space $\mathcal{X}$ that satisfies inequality ❶ *for any relabeling of its points by p, x, y, z, such that $\mathcal{X}$ is not* CBB(0).

Assume that for a given triangle $[x^1x^2x^3]$ in a metric space its κ-model triangle $[\tilde{x}^1\tilde{x}^2\tilde{x}^3] = \tilde{\triangle}^\kappa(x^1x^2x^3)$ is defined. We say the triangle $[x^1x^2x^3]$ is κ-*fat* if the natural map (see definition 8.4.1) $[\tilde{x}^1\tilde{x}^2\tilde{x}^3] \to$ $\to [x^1x^2x^3]$ is distance non contracting.

7.3.6. Exercise. *Prove that any triangle with perimeter $< \varpi^\kappa$ in a* CBB(κ) *space is κ-fat.*

The following exercise is inspired by Busemann's definition given in [15].

7.3.7. Exercise.

a) Show that any CBB(0) *space $\mathcal{L}$ satisfies the following condition: for any three points $p, q, r \in \mathcal{L}$, if $\bar{q}$ and $\bar{r}$ are midpoints of geodesics $[pq]$ and $[pr]$ correspondingly, then $2\cdot|\bar{q}-\bar{r}| \geqslant |q-r|$.*

b) Show that there is a metric on $\mathbb{R}^2$ defined by a norm that satisfies the above condition, but not CBB(0).

7.4 Function comparison

In this section we will translate the angle comparison definitions (Theorem 7.3.1) to a concavity-like property of the distance functions as defined in Section 4.6. This a conceptual step — we reformulate a global geometric condition into an infinitesimal condition on distance functions.

7.4.1. Theorem. *Let $\mathcal{L}$ be a complete length space. Then the following statements are equivalent:*

a) $\mathcal{L}$ is CBB(κ).

b) (function comparison) $\mathcal{L}$ is G-delta geodesic and for any $p \in \mathcal{L}$, the function $f = \mathrm{md}^\kappa \circ \mathrm{dist}_p$ satisfies the differential inequality

$$f'' \leqslant 1 - \kappa\cdot f.$$

in $\mathrm{B}(p, \varpi^\kappa)$.

In particular, a complete G-delta geodesic space $\mathcal{L}$ is CBB(0) if and only if for any $p \in \mathcal{L}$, the function $\operatorname{dist}_p^2 : \mathcal{L} \to \mathbb{R}$ is 2-concave as defined in Section 4.6.

Proof. Let $[xy]$ be a geodesic in $\mathrm{B}(p, \varpi^\kappa)$ and $\ell = |x - y|$. Consider the model triangle $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^\kappa(pxy)$. Set

$$\tilde{r}(t) = |\tilde{p} - \operatorname{geod}_{[\tilde{x}\tilde{y}]}(t)|, \qquad r(t) = |p - \operatorname{geod}_{[xy]}(t)|.$$

Clearly $\tilde{r}(0) = r(0)$ and $\tilde{r}(\ell) = r(\ell)$. Set $\tilde{f} = \operatorname{md}^\kappa \circ \tilde{r}$ and $f = \operatorname{md}^\kappa \circ r$. From 1.1.1*a* we get that $\tilde{f}'' = 1 - \kappa \cdot \tilde{f}$.

Note that the point-on-side comparison (7.3.1*b*) for point p and geodesic $[xy]$ is equivalent to $\tilde{r} \leqslant r$. Since md^κ is increasing on $[0, \varpi^\kappa)$, $\tilde{r} \leqslant r$ is equivalent to $\tilde{f} \leqslant f$. The latter is the Jensen's inequality (4.5.3*d*) for the function $t \mapsto \operatorname{md}^\kappa |p - \operatorname{geod}_{[xy]}(t)|$ on interval $[0, \ell]$. Hence the result. □

Recall that Busemann functions are defined in Proposition 5.1.1. Compare the following exercise to Exercise 8.5.2.

7.4.2. Exercise. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and* $\operatorname{bus}_\gamma \colon \mathcal{L} \to \mathbb{R}$ *is the Busemann function for a ray* $\gamma \colon [0, \infty) \to \mathcal{L}$.

a) If $\kappa = 0$, then the Busemann function $\operatorname{bus}_\gamma$ *is concave.*

b) If $\kappa = -1$, then the function

$$f = \exp \circ \operatorname{bus}_\gamma$$

satisfies

$$f'' - f \leqslant 0$$

7.5 Development

In this section we reformulate the function comparison using a more geometric language based on the definition of development given below.

This definition appears in [16] and an earlier form of it can be found in [17]. The definition is somewhat lengthy, but it defines a useful comparison object for a curve. Often it is easier to write proofs in terms of function comparison, but think in terms of developments.

7.5.1. Lemma-definition. *Let $\kappa \in \mathbb{R}$, $\mathcal{X}$ be a metric space, $\gamma \colon \mathbb{I} \to \mathcal{X}$ be a 1-Lipschitz curve, $p \in \mathcal{X}$, $\tilde{p} \in \mathbb{M}^2(\kappa)$. Assume $0 < |p - \gamma(t)| < \varpi^\kappa$ for all $t \in \mathbb{I}$. Then there exists a unique up to rotation curve $\tilde{\gamma} \colon \mathbb{I} \to \mathbb{M}^2(\kappa)$, parameterized by the arc-length such that $|\tilde{p} - \tilde{\gamma}(t)| = |p - \gamma(t)|$ for all t and the direction of $[\tilde{p}\tilde{\gamma}(t)]$ monotonically turns around $\tilde{p}$ conterclockwise, as t increases.*

If $p, \tilde{p}, \gamma, \tilde{\gamma}$ be as in above, then $\tilde{\gamma}$ is called the κ-development of γ with respect to p; *the point $\tilde{p}$ is called* the basepoint of development. *When we say that the κ-development of γ with respect to p is* defined *we allways assume that $0 < |p - \gamma(t)| < \varpi^\kappa$ for all $t \in \mathbb{I}$.*

Proof. Consider functions $\rho, \vartheta \colon \mathbb{I} \to \mathbb{R}$ defined as

$$\rho(t) = |p - \gamma(t)|, \qquad \vartheta(t) = \int\limits_{t_0}^{t} \frac{\sqrt{1 - (\rho'(\mathfrak{t}))^2}}{\operatorname{sn}^\kappa \circ \rho(\mathfrak{t})} \cdot \boldsymbol{d}\mathfrak{t},$$

where $t_0 \in \mathbb{I}$ is a fixed number and $\int$ stays for Lebesgue integral. Since γ is 1-Lipshitz, so is $\rho(t)$ and thus, the function ϑ is defined and non decreasing.

It is straightforward to check that (ρ, ϑ) describe $\tilde{\gamma}$ in polar coordinates on $\mathbb{M}^2(\kappa)$ with center at $\tilde{p}$. $\square$

We need the following analogues of sub- and super-graphs and convex/concave functions, adapted to polar coordinates in $\mathbb{M}^2(\kappa)$.

7.5.2. Definition. *Let $\tilde{\gamma} \colon \mathbb{I} \to \mathbb{M}^2(\kappa)$ be a curve and $\tilde{p} \in \mathbb{M}^2(\kappa)$ such that there is unique geodesic $[\tilde{p}\tilde{\gamma}(t)]$ for any $t \in \mathbb{I}$ and the direction of $[\tilde{p}\tilde{\gamma}(t)]$ turns monotonically as t grows.*



The set formed by all geodesics from $\tilde{p}$ to the points on $\tilde{\gamma}$ is called the subgraph *of $\tilde{\gamma}$ with respect to $\tilde{p}$.*

The set of all points $\tilde{x} \in \mathbb{M}^2(\kappa)$ such that a geodesic $[\tilde{p}\tilde{x}]$ intersect $\tilde{\gamma}$ is called the supergraph *of $\tilde{\gamma}$ with respect to $\tilde{p}$.*

The curve $\tilde{\gamma}$ is called convex (concave) with respect to $\tilde{p}$ *if the subgraph (supergraph) of $\tilde{\gamma}$ with respect to $\tilde{p}$ is convex.*

The curve $\tilde{\gamma}$ is called locally convex (concave) with respect to $\tilde{p}$ *if for any interior value t_0 in $\mathbb{I}$ there is a subsegment $(a, b) \subset \mathbb{I}$, $(a, b) \ni$ $\ni t_0$, such that the restriction $\tilde{\gamma}|_{(a,b)}$ is convex (concave) with respect to $\tilde{p}$.*

Note that if $\kappa > 0$, then the supergraph of a curve is a subgraph with respect to the opposite point.

For developments, all the notions above will be considered with respect to their basepoints. In particular, if $\tilde{\gamma}$ is a development, we will say it is *(locally) convex* if it is (locally) convex with respect to its basepoint.

7.5.3. Development comparison. *A complete G-delta geodesic space $\mathcal{L}$ is* $\mathrm{CBB}(\kappa)$ *if and only if for any point $p \in \mathcal{L}$ and any geodesic γ in $\mathrm{B}(p, \varpi^\kappa) \backslash \{p\}$, its κ-development with respect to p is convex.*



A simpler proof of "if"-part can be build on the adjacent angle comparison (7.3.1a). We use a longer proof since it also implies the short hinge lemma (7.5.4).

Proof; "if"-part. Let $\gamma\colon [0,T] \to \mathrm{B}(p,\varpi^\kappa)\backslash\{p\}$ be a unit-speed geodesic in $\mathcal{L}$.

Consider fine partition

$$0 = t_0 < t_1 < \ldots < t_n = T.$$

Set $x_i = \gamma(t_i)$ and choose a point

$$p' \in \mathrm{Str}(x_0, x_1, \ldots, x_n)$$

sufficiently close to p.

Let us construct a chain of model triangles $[\tilde{p}'\tilde{x}_{i-1}\tilde{x}_i] = \tilde{\triangle}^\kappa(p'x_{i-1}x_i)$ in such a way that direction $[\tilde{p}'\tilde{x}_i]$ turns counterclockwise as i grows. By hinge comparison (7.3.1c), we have

❶
$$\begin{aligned}\measuredangle[\tilde{x}_i\,{}^{\tilde{x}_{i-1}}_{\tilde{p}'}] + \measuredangle[\tilde{x}_i\,{}^{\tilde{x}_{i+1}}_{\tilde{p}'}] &= \tilde{\measuredangle}^\kappa(x_i\,{}^{x_{i-1}}_{p'}) + \tilde{\measuredangle}^\kappa(x_i\,{}^{x_{i+1}}_{p'}) \leqslant \\ &\leqslant \measuredangle[x_i\,{}^{x_{i-1}}_{p'}] + \measuredangle[x_i\,{}^{x_{i+1}}_{p'}] \leqslant \\ &\leqslant \pi.\end{aligned}$$

Further, since γ is a unit-speed geodesic, we have that

❷
$$\sum_{i=1}^n |x_{i-1} - x_i| \leqslant |p' - x_0| + |p' - x_n|.$$

If $\kappa \leqslant 0$, then ❷ implies that

❸
$$\sum_{i=1}^n \measuredangle[\tilde{p}'\,{}^{\tilde{x}_i}_{\tilde{x}_{i-1}}] \leqslant \pi.$$

If $\kappa > 0$, then ❸ require more work. Applying rescaling we can assume that $\kappa = 1$. Assume the contrary; that is, $\sum_{i=1}^n \measuredangle[\tilde{p}'\,{}^{\tilde{x}_i}_{\tilde{x}_{i-1}}] > \pi$, then for some $\bar{k}$ there is a point $\tilde{y} \in [\tilde{x}_{\bar{k}}\tilde{x}_{\bar{k}+1}]$ such that

$$\measuredangle[\tilde{p}'\,{}^{\tilde{x}_0}_{\tilde{x}_1}] + \ldots + \measuredangle[\tilde{p}'\,{}^{\tilde{x}_{\bar{k}-1}}_{\tilde{x}_{\bar{k}}}] + \measuredangle[\tilde{p}'\,{}^{\tilde{x}_{\bar{k}}}_{\tilde{y}}] = \pi.$$

By ❷, the broken line $\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_{\bar{k}}\tilde{y}$ is convex; in particular,

$$|\tilde{p}' - \tilde{x}_0|_{\mathbb{S}^2} + |\tilde{p}' - \tilde{y}|_{\mathbb{S}^2} \leqslant \pi$$

By triangle inequality

$$|\tilde{x}_0-\tilde{x}_1|_{\mathbb{S}^2}+\ldots+|\tilde{x}_{k-1}-\tilde{x}_k|_{\mathbb{S}^2}+|\tilde{x}_k-\tilde{y}|_{\mathbb{S}^2}\geqslant|\tilde{p}'-\tilde{x}_0|_{\mathbb{S}^2}+|\tilde{p}'-\tilde{y}|_{\mathbb{S}^2}$$

Applying the triangle inequality again, we get

$$|\tilde{y}-\tilde{x}_{k+1}|_{\mathbb{S}^2}+|\tilde{x}_{k+1}-\tilde{x}_{k+2}|_{\mathbb{S}^2}+\ldots+|\tilde{x}_{n-1}-\tilde{x}_n|_{\mathbb{S}^2}>|\tilde{p}'-\tilde{x}_n|_{\mathbb{S}^2}-|\tilde{p}'-\tilde{y}|_{\mathbb{S}^2}$$

Hence

$$T=\sum_{i=1}^{n}|\tilde{x}_i-\tilde{x}_{i-1}|_{\mathbb{S}^2}>|\tilde{p}'-\tilde{x}_0|_{\mathbb{S}^2}+|\tilde{p}'-\tilde{x}_n|_{\mathbb{S}^2}.$$

The later contradicts that γ is length minimizing.

Inequalities ❶ and ❸ imply that the polygon $[\tilde{p}'\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_n]$ is convex.

Let us take finer and finer partitions and pass to the limit of the polygon $\tilde{p}'\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_n$ as $p'\to p$. We obtain a convex curvelinear triangle formed by a curve $\tilde{\gamma}\colon[0,T]\to\mathbb{M}^2(\kappa)$ — the limit of broken line $\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_n$ and two geodesics $[\tilde{p}'\,\tilde{\gamma}(0)]$, $[\tilde{p}'\,\tilde{\gamma}(T)]$. Since $[\tilde{p}'\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_n]$ is convex, the natural parametrization of $\tilde{x}_0\tilde{x}_1\ldots\tilde{x}_n$ converges to a natural parametrization of $\tilde{\gamma}$. Thus $\tilde{\gamma}$ is the κ-development of γ with respect to p.

($\Leftarrow$). Assuming convexity of the development, we will prove the point-on-side comparison (7.3.1*b*). We can assume that $p\notin[xy]$; otherwise the statement is trivial.

Set $T=|x-y|$ and $\gamma(t)=\operatorname{geod}_{[xy]}(t)$; note that γ is a geodesic in $\mathrm{B}(p,\varpi^\kappa)\backslash\{p\}$. Let $\tilde{\gamma}\colon[0,T]\to\mathbb{M}^2(\kappa)$ be κ-development with base $\tilde{p}$ of γ with respect to p. Take a partition $0=t_0<t_1<\ldots<t_n=T$ and set

$$\tilde{y}_i=\tilde{\gamma}(t_i)\quad\text{and}\quad\tau_i=|\tilde{y}_0-\tilde{y}_1|+|\tilde{y}_1-\tilde{y}_2|+\ldots+|\tilde{y}_{i-1}-\tilde{y}_i|.$$

Since $\tilde{\gamma}$ is convex, for a fine partition we have that broken line $\tilde{y}_0\tilde{y}_1\ldots$ $\ldots\,\tilde{y}_n$ is also convex. Applying Alexandrov's lemma (5.2.1) inductively to pairs of model triangles

$$\tilde{\triangle}^\kappa\{\tau_{i-1},|\tilde{p}-\tilde{y}_0|,|\tilde{p}-\tilde{y}_{i-1}|\}$$

and

$$\tilde{\triangle}^\kappa\{|\tilde{y}_{i-1}-\tilde{y}_i|,|\tilde{p}-\tilde{y}_{i-1}|,|\tilde{p}-\tilde{y}_i|\}$$

we obtain that sequence $\tilde{\measuredangle}^\kappa\{|\tilde{p}-\tilde{y}_i|;|\tilde{p}-\tilde{y}_0|,\tau_i\}$ is non increasing.

For finer and finer partitions we have

$$\max_i\{|\tau_i-t_i|\}\to0.$$

Thus, point-on-side comparison (7.3.1b) follows. □

Note that in the proof of "if"-part we could use a slightly weaker version of the hinge comparison (7.3.1c). Namely we proved the following lemma which will be needed later in the proof of the globalization theorem (7.6.2).

7.5.4. Short hinge lemma. *Let $\mathcal{L}$ be a complete G-delta geodesic space such that for any hinge $[x\,{}^p_y]$ in $\mathcal{L}$ the angle $\measuredangle[x\,{}^p_y]$ is defined and moreover, if $x \in\,]yz[$, then*

$$\measuredangle[x\,{}^p_y] + \measuredangle[x\,{}^p_z] \leqslant \pi.$$

Assume that for any hinge $[x\,{}^p_y]$ in $\mathcal{L}$ we have

$$|p-x| + |x-y| < \varpi^\kappa \quad \Rightarrow \quad \measuredangle[x\,{}^p_y] \geqslant \tilde\measuredangle^\kappa(x\,{}^p_y),$$

then $\mathcal{L}$ is CBB(κ).

7.6 Local definitions and globalization

In this section we discuss locally CBB(κ) spaces; in particular, we prove the *globalization theorem* — equivalence of local and global definitions for complete length spaces.

The following theorem summarizes equivalent definitions of locally CBB(κ) spaces

7.6.1. Theorem. *Let $\mathcal{X}$ be a complete length space and $p \in \mathcal{X}$. Then the following conditions are equivalent:*

1) *(local* CBB(κ) *comparison) there is $R_1 > 0$ such that comparison*

$$\tilde\measuredangle^\kappa(q\,{}^{x^1}_{x^2}) + \tilde\measuredangle^\kappa(q\,{}^{x^2}_{x^3}) + \tilde\measuredangle^\kappa(q\,{}^{x^3}_{x^1}) \leqslant 2\cdot\pi$$

holds for any $q, x^1, x^2, x^3 \in \mathrm{B}(p, R_1)$.

2) *(local Kirszbraun property) there is $R_2 > 0$, such that for any 3-point subset F_3 and any 4-point subset $F_4 \supset F_3$ in $\mathrm{B}(p, R_2)$, any short map $f\colon F_3 \to \mathbb{M}^2(\kappa)$ can be extended as a short map $\bar f\colon F_4 \to \mathbb{M}^2(\kappa)$ (so $f = \bar f|_{F_3}$).*

3) *(local function comparison) there is $R_3 > 0$, such that $\mathrm{B}(p, R_3)$ is G-delta geodesic and for any $q \in \mathrm{B}(p, R_3)$, the function $f = \mathrm{md}^\kappa \circ \mathrm{dist}_q$ satisfies $f'' \leqslant 1 - \kappa\cdot f$ in $\mathrm{B}(p, R_3)$.*

4) *(local adjacent angle comparison) there is $R_4 > 0$ such that $\mathrm{B}(p, R_4)$ is G-delta geodesic and if q and geodesic $[xy]$ lie in $\mathrm{B}(p, R_4)$ and $z \in\,]xy[$, then*

$$\tilde\measuredangle^\kappa(z\,{}^q_x) + \tilde\measuredangle^\kappa(z\,{}^q_y) \leqslant \pi.$$

5) *(local point-on-side comparison) there is* $R_5 > 0$, *such that* $\mathrm{B}(p, R_5)$ *is G-delta geodesic and if* q *and geodesic* $[xy]$ *lie in* $\mathrm{B}(p, R_5)$ *and* $z \in\,]xy[$, *we have*

$$\tilde{\measuredangle}^{\kappa}(x\,{}^{q}_{y}) \leqslant \tilde{\measuredangle}^{\kappa}(x\,{}^{q}_{z});$$

or, equivalently,

$$|\tilde{p} - \tilde{z}| \leqslant |p - z|,$$

where $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^{\kappa}(pxy)$, $\tilde{z} \in\,]\tilde{x}\tilde{y}[$, $|\tilde{x} - \tilde{z}| = |x - z|$.

6) *(local hinge comparison) there is* $R_6 > 0$, *such that* $\mathrm{B}(p, R_6)$ *is G-delta geodesic and if* $x \in \mathrm{B}(p, R_6)$, *then (1) for any hinge* $[x\,{}^{q}_{y}]$, *the angle* $\measuredangle[x\,{}^{q}_{y}]$ *is defined and (2) if* $x \in\,]yz[$, *then*[1]

$$\measuredangle[x\,{}^{q}_{y}] + \measuredangle[x\,{}^{q}_{z}] \leqslant \pi.$$

Moreover, if hinge $[x\,{}^{q}_{y}]$ *lies in* $\mathrm{B}(p, R_6)$, *then*

$$\measuredangle[x\,{}^{q}_{y}] \geqslant \tilde{\measuredangle}^{\kappa}(x\,{}^{q}_{y}).$$

or equivalently

$$\tilde{\gamma}^{\kappa}[x\,{}^{q}_{y}] \geqslant |q - y|.$$

7) *(local development comparison) there is* $R_7 > 0$, *such that* $\mathrm{B}(p, R_7)$ *is G-delta geodesic and if a geodesic* γ *lies in* $\mathrm{B}(p, R_7)$ *and* $q \in \mathrm{B}(p, R_7)\backslash\gamma$, *then the* κ*-development* $\tilde{\gamma}$ *with respect to* q *is convex.*

Moreover, for each pair $i, j \in \{1, 2, \dots, 7\}$ *we can assume that*

$$R_i > \tfrac{1}{9} \cdot R_j.$$

The proof of each equivalence repeats the proof of corresponding global equivalence in localized form; see proofs of theorems 7.3.1, 7.4.1, 7.5.1, 9.1.1.

7.6.2. Globalization theorem. *Any complete length locally* $\mathrm{CBB}(\kappa)$ *space is* $\mathrm{CBB}(\kappa)$.

In the two-dimensional case this theorem was proved by Paolo Pizzetti [18]; latter it was reproved independently by Alexandr Alexandrov [16].Victor Toponogov [19] proved it for Riemannian manifolds of all dimensions. In the above generality, the theorem first appears in the paper of Mikhael Gromov, Yuri Burago and Grigory Perelman [3]; simplifications and modifications were given by Conrad Plaut [20],

[1] Let us remind that $[x\,{}^{q}_{y}]$ and $[x\,{}^{q}_{z}]$ are short notations for pairs $([xq], [xy])$ and $([xq], [xz])$, thus these two hinges automatically have common side $[xq]$.

Katsuhiro Shiohama [21] and in the book of Dmitry Burago, Yuri Burago and Sergei Ivanov [4]. A generalization for non-complete but geodesic spaces was obtained by the third author [22]; namely it solves the following exercise:

7.6.3. Advanced exercise. *Assume $\mathcal{X}$ is a geodesic locally* $\mathrm{CBB}(\kappa)$ *space. Prove that the completion of $\mathcal{X}$ is* $\mathrm{CBB}(\kappa)$.

Our proof, is based on presentations in [20] and [4]; this proof was rediscovered independently by Urs Lang and Viktor Schroeder [23].

The following corollary says that the expression "space with curvature $\geqslant \kappa$" makes sense.

7.6.4. Corollary. *Let $\mathcal{L}$ be a complete length space. Then $\mathcal{L}$ is* $\mathrm{CBB}(\mathrm{K})$ *if and only if $\mathcal{L}$ is* $\mathrm{CBB}(\kappa)$ *for any $\kappa < \mathrm{K}$.*

Proof. Note that if $\mathrm{K} \leqslant 0$, this statement follows directly from definition of Alexandrov space (7.1.2) and monotonicity of the function $\kappa \mapsto \measuredangle^{\kappa}(x\,{}^{y}_{z})$ (1.1.1d).

The "if"-part also follows directly from the definition.

For $\mathrm{K} > 0$, the angle $\measuredangle^{\mathrm{K}}(x\,{}^{y}_{z})$ might be undefined while $\measuredangle^{\kappa}(x\,{}^{y}_{z})$ is defined. However, $\measuredangle^{\mathrm{K}}(x\,{}^{y}_{z})$ is defined if x, y and z are sufficiently close to each other. Thus, if $\mathrm{K} > \kappa$, then any $\mathrm{CBB}(\mathrm{K})$ space is locally $\mathrm{CBB}(\kappa)$. It remains to apply the globalization theorem (7.6.2). □

7.6.5. Corollary. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space. Assume $\mathcal{M}$ is a target space of a sumetry from $\mathcal{L}$. Then $\mathcal{M}$ is a complete length space* $\mathrm{CBB}(\kappa)$ *space.*

In particular, if $G \curvearrowright \mathcal{L}$ an isometric group action with closed orbits, then the quotient space $\mathcal{L}/G$ is a complete length space $\mathrm{CBB}(\kappa)$ *space.*

Proof. Follows from the globalization theorem and Theorem 7.1.8. □

In the proof of the globalization theorem, we will use three lemmas. First is the short hinge lemma (7.5.4) — it gives a characterization of $\mathrm{CBB}(\kappa)$ spaces which is the same as the hinge comparison 7.3.1c if $\kappa \leqslant 0$ and slightly weaker in case $\kappa > 0$.

The following lemma says that if comparison holds for all small hinges, then it holds for slightly bigger hinges near the given point.

7.6.6. Key lemma. *Let $\kappa \in \mathbb{R}$, $0 < \ell \leqslant \varpi^{\kappa}$, $\mathcal{X}$ be a complete geodesic space and $p \in \mathcal{X}$ be a point such that* $\mathrm{B}(p, 2{\cdot}\ell)$ *is locally* $\mathrm{CBB}(\kappa)$.

Assume that for any point $q \in \mathrm{B}(p, \ell)$, comparison

$$\measuredangle[x\,{}^{y}_{q}] \geqslant \measuredangle^{\kappa}(x\,{}^{y}_{q})$$

holds for any hinge $[x\,{}^{y}_{q}]$ *with* $|x-y|+|x-q|<\frac{2}{3}\cdot\ell$. *Then comparison*

$$\measuredangle[x\,{}^{p}_{q}]\geqslant \mathcal{Z}^{\kappa}(x\,{}^{p}_{q})$$

holds for any hinge $[x\,{}^{p}_{q}]$ *with* $|x-p|+|x-q|<\ell$.

Proof. We will prove an equivalent inequality:

❶ $$\tilde{\Upsilon}^{\kappa}[x\,{}^{p}_{q}]\geqslant |p-q|.$$

for any hinge $[x\,{}^{p}_{q}]$ with $|x-p|+|x-q|<\ell$.

Fix q. Consider a hinge $[x\,{}^{p}_{q}]$ such that

$$\tfrac{2}{3}\cdot\ell\leqslant |p-x|+|x-q|<\ell.$$

Let us construct a new smaller hinge $[x'\,{}^{p}_{q}]$; that is,



❷ $$|p-x|+|x-q|\geqslant |p-x'|+|x'-q|$$

and such that

❸ $$\tilde{\Upsilon}^{\kappa}[x\,{}^{p}_{q}]\geqslant \tilde{\Upsilon}^{\kappa}[x'\,{}^{p}_{q}].$$

Assume $|x-q|\geqslant|x-p|$, otherwise switch the roles of p and q in the following construction. Take $x'\in[xq]$ such that

❹ $$|p-x|+3\cdot|x-x'|=\tfrac{2}{3}\cdot\ell$$

Choose a geodesic $[x'p]$ and consider the hinge $[x'\,{}^{p}_{q}]$ fromed by $[x'p]$ and $[x'q]\subset[xq]$. (In fact the same argument as in 7.7.1 shows that condition $[x'q]\subset[xq]$ always holds.) Then ❷ follows from the triangle inequality.

Further, note that we have $x,x'\in \mathrm{B}(p,\ell)\cap\mathrm{B}(q,\ell)$ and moreover $|p-x|+|x-x'|,|p-x'|+|x'-x|<\frac{2}{3}\cdot\ell$. In particular,

❺ $$\measuredangle[x\,{}^{p}_{x'}]\geqslant \mathcal{Z}^{\kappa}(x\,{}^{p}_{x'})\quad\text{and}\quad \measuredangle[x'\,{}^{p}_{x}]\geqslant \mathcal{Z}^{\kappa}(x'\,{}^{p}_{x}).$$

Now, let $[\tilde{x}\tilde{x}'\tilde{p}]=\tilde{\triangle}^{\kappa}(xx'p)$. Take $\tilde{q}$ on the extension of $[\tilde{x}\tilde{x}']$ beyond x' such that $|\tilde{x}-\tilde{q}|=|x-q|$ (and therefore $|\tilde{x}'-\tilde{q}|=|x'-q|$). From ❺,

$$\measuredangle[x\,{}^{p}_{q}]=\measuredangle[x\,{}^{p}_{x'}]\geqslant \mathcal{Z}^{\kappa}(x\,{}^{p}_{x'})\quad\Rightarrow\quad \tilde{\Upsilon}^{\kappa}[x\,{}^{q}_{p}]\geqslant|\tilde{p}-\tilde{q}|.$$

From the assumptions of the lemma, we have $\measuredangle[x'\,{}^{p}_{x}]+\measuredangle[x'\,{}^{p}_{q}]\leqslant\pi$; thus ❺ implies

$$\pi-\mathcal{Z}^{\kappa}(x'\,{}^{p}_{x})\geqslant\pi-\measuredangle[x'\,{}^{p}_{x}]\geqslant\measuredangle[x'\,{}^{p}_{q}].$$

Therefore $|\tilde{p}-\tilde{q}|\geqslant\tilde{\Upsilon}^{\kappa}[x'\,{}^{q}_{p}]$ and ❸ follows.



Let us continue the proof. Set $x_0 = x$. Let us apply inductively the above construction to get a sequence of hinges $[x_n\,{}^{p}_{q}]$ with $x_{n+1} = x'_n$.

The sequence might terminate at some n only if $|p-x_n|+|x_n-q| < \frac{2}{3}\cdot\ell$. In this case, by the assumptions of the lemma, $\tilde\Upsilon^{\kappa}[x_n\,{}^{p}_{q}] \geqslant |p-q|$. From ❸, we get that the sequence $s_n = \tilde\Upsilon^{\kappa}[x_n\,{}^{p}_{q}]$ is decreasing. Hence inequality ❶ follows.

Otherwise, the sequence $r_n = |p-x_n| + |x_n - q|$ is non-increasing and $r_n \geqslant \frac{2}{3}\cdot\ell$ for all n. Hence $r_n \to r$ as $n\to\infty$. Pass to a subsequence $y_k = x_{n_k}$ such that $y'_k = x_{n_k+1} \in [y_k q]$. (In particular, $|y_k - q| \geqslant |y_k - p|$.) Clearly,

$$❻ \qquad |p-y_k| + |y_k - y'_k| - |p - y'_k| = r_k - r_{k+1} \underset{k\to\infty}{\to} 0.$$

From ❹, it follows that $|y_k - y'_k| > \frac{\ell}{100}$. From ❹ and ❻, it follows that $|p-y_k| > \frac{\ell}{100}$ for all large k. Therefore, $\tilde\measuredangle^{\kappa}(y_k\,{}^{p}_{y'_k}) \to \pi$. Since $\measuredangle[y_k\,{}^{p}_{y'_k}] \geqslant \tilde\measuredangle^{\kappa}(y_k\,{}^{p}_{y'_k})$, we have $\measuredangle[y_k\,{}^{p}_{q}] = \measuredangle[y_k\,{}^{p}_{y'_k}] \to \pi$. Therefore,

$$|p-y_k| + |y_k - q| - \tilde\Upsilon^{\kappa}[y_k\,{}^{p}_{q}] \to 0.$$

(Here we used that $\ell \leqslant \varpi^{\kappa}$.) Together with the triangle inequality

$$|p-y_k| + |y_k - q| \geqslant |p-q|$$

this yields

$$\lim_{n\to\infty} \tilde\Upsilon^{\kappa}[y_k\,{}^{p}_{q}] \geqslant |p-q|.$$

Applying monotonicity of sequence $s_n = \tilde\Upsilon^{\kappa}[y_k\,{}^{p}_{q}]$ we obtain ❶. □

The following lemma works in complete spaces; it will used the same way as existence of minimum point in a compact.

7.6.7. Lemma on almost minimum. *Let $\mathcal{X}$ be a complete metric space, $p\in\mathcal{X}$ and $r\colon \mathcal{X}\to\mathbb{R}$ be a function and $\varepsilon>0$. Assume that the function r is strictly positive in $\overline{\mathrm{B}}[p,\frac{1}{\varepsilon^2}\cdot r(p)]$ and $\varliminf_n r(x_n) > 0$ for any convergent sequence $x_n\to x\in\overline{\mathrm{B}}[p,\frac{1}{\varepsilon^2}\cdot r(p)]$.*

Then, there is a point $p^*\in\overline{\mathrm{B}}[p,\frac{1}{\varepsilon^2}\cdot r(p)]$ such that

a) $r(p^*)\leqslant r(p)$ and

b) $r(x) > (1-\varepsilon)\cdot r(p^*)$ for any $x\in\overline{\mathrm{B}}[p^*,\frac{1}{\varepsilon}\cdot r(p^*)]$.

Proof. Assume the statement is wrong. Then for any $x \in \mathrm{B}(p, \frac{1}{\varepsilon^2}\cdot r(p))$ with $r(x) \leqslant r(p)$, there is a point $x' \in \mathcal{X}$ such that

$$|x - x'| < \tfrac{1}{\varepsilon}\cdot r(x) \quad \text{and} \quad r(x') \leqslant (1-\varepsilon)\cdot r(x).$$

Take $x_0 = p$ and consider a sequence of points (x_n) such that $x_{n+1} = x'_n$. Clearly

$$|x_{n+1} - x_n| \leqslant \tfrac{r(p)}{\varepsilon}\cdot(1-\varepsilon)^n \quad \text{and} \quad r(x_n) \leqslant r(p)\cdot(1-\varepsilon)^n.$$

In particular, $|p - x_n| < \frac{1}{\varepsilon^2}\cdot r(p)$. Therefore (x_n) converges in itself, $x_n \to x \in \overline{\mathrm{B}}[p, \frac{1}{\varepsilon^2}\cdot r(p)]$ and $\lim_n r(x_n) = 0$, a contradiction. □

Proof of the globalization theorem (7.6.2). Exactly the same argument as in the proof of Theorem 7.2.4 shows that $\mathcal{L}$ is G-delta geodesic. By Theorem 7.6.1-6, for any hinge $[x\,{}^p_y]$ in $\mathcal{L}$ the angle $\measuredangle[x\,{}^p_y]$ is defined and moreover, if $x \in\,]yz[$, then

$$\measuredangle[x\,{}^p_y] + \measuredangle[x\,{}^p_z] \leqslant \pi.$$

Let us denote by $\mathrm{ComRad}(p, \mathcal{L})$ (which stands for *comparison radius* of $\mathcal{L}$ at p) the maximal value (possibly ∞) such that the comparison

$$\measuredangle[x\,{}^p_y] \geqslant \measuredangle^{\kappa}(x\,{}^p_y)$$

holds for any hinge $[x\,{}^p_y]$ with $|p - x| + |x - y| < \mathrm{ComRad}(p, \mathcal{L})$.

As follows from 7.6.1-3, $\mathrm{ComRad}(p, \mathcal{L}) > 0$ for any $p \in \mathcal{L}$ and

$$\varliminf_{n\to\infty} \mathrm{ComRad}(p_n, \mathcal{L}) > 0$$

for any converging sequence of points $p_n \to p$. That makes it possible to apply the lemma on almost minimum (7.6.7) to the function $p \mapsto \mathrm{ComRad}(p, \mathcal{L})$.

According to the short hinge lemma (7.5.4), it is sufficient to show that

❼ $$s_0 = \inf_{p\in\mathcal{L}} \mathrm{ComRad}(p, \mathcal{L}) \geqslant \varpi^{\kappa} \quad \text{for any} \quad p \in \mathcal{L}.$$

We argue by contradiction, assuming that ❼ does not hold.

The rest of the proof is easier for geodesic spaces and yet easier for compact spaces. Thus we give three different arguments for each of these cases.

Compact space. Assume $\mathcal{L}$ is compact.

By Theorem 7.6.1-3, $s_0 > 0$. Take a point $p^* \in \mathcal{L}$ such that $r^* = \mathrm{ComRad}(p^*, \mathcal{L})$ is sufficiently close to s_0 (p^* such that $s_0 \leqslant r^* <$

$\min\{\varpi^\kappa, \frac{3}{2}\cdot s_0\}$ will do). Then the key lemma (7.6.6) applied for p^* and ℓ slightly bigger than r^* (say, such that $r^* < \ell < \min\{\varpi^\kappa, \frac{3}{2}\cdot s_0\}$) implies that

$$\measuredangle[x\,{}^{p^*}_{q}] \geqslant \tilde{\measuredangle}^\kappa(x\,{}^{p^*}_{q})$$

for any hinge $[x\,{}^{p^*}_{q}]$ such that $|p^* - x| + |x - q| < \ell$. Thus $r^* \geqslant \ell$, a contradiction.

Geodesic case. Assume $\mathcal{L}$ is geodesic.

Fix small $\varepsilon > 0$ ($\varepsilon = 0.0001$ will do). Apply the lemma on almost minimum (7.6.7) to find a point $p^* \in \mathcal{L}$ such that

$$r^* = \mathrm{ComRad}(p^*, \mathcal{L}) < \varpi^\kappa$$

and

❽ $$\mathrm{ComRad}(q, \mathcal{L}) > (1 - \varepsilon)\cdot r^*$$

for any $q \in \overline{\mathrm{B}}[p^*, \frac{1}{\varepsilon}\cdot r^*]$.

Applying the key lemma (7.6.6) for p^* and ℓ slightly bigger than r^* leads to a contradiction.

General case. Let us construct $p^* \in \mathcal{L}$ as in the previous case. Since $\mathcal{L}$ is not geodesic, we can not apply the key lemma directly. Instead, let us pass to the ultrapower $\mathcal{L}^\omega$ which is a geodesic space (see 3.5.2).

According to Theorem 7.6.1, inequality ❽ implies that the ondition in 7.6.1-1 holds for some fixed $R_1 = \frac{r^*}{100} > 0$ at any point $q \in$ $\in \overline{\mathrm{B}}[p^*, \frac{1}{2\cdot\varepsilon}\cdot r^*] \subset \mathcal{L}$. Therefore a similar statement is true in the ultrapower $\mathcal{L}^\omega$; that is, for any point $q_\omega \in \overline{\mathrm{B}}[p^*, \frac{1}{2\cdot\varepsilon}\cdot r^*] \subset \mathcal{L}^\omega$, Condition 7.6.1-1 holds for say $R_1 = \frac{r^*}{101}$.

Note that $r^* \geqslant \mathrm{ComRad}(p^*, \mathcal{L}^\omega)$. Therefore we can apply the lemma on almost minimum at point p^* for space $\mathcal{L}^\omega$ with function $x \mapsto \mathrm{ComRad}(x, \mathcal{L}^\omega)$ and $\varepsilon' = \sqrt{\varepsilon} = 0.01$.

For the obtained point $p^{**} \in \mathcal{L}^\omega$, we have $r^{**} = \mathrm{ComRad}(p^{**}, \mathcal{L}) <$ $< \varpi^\kappa$ and $\mathrm{ComRad}(q_\omega, \mathcal{L}^\omega) > (1 - \varepsilon')\cdot r^{**}$ for any $q_\omega \in \overline{\mathrm{B}}[p^{**}, \frac{1}{\varepsilon'}\cdot r^{**}]$. Thus applying the key lemma (7.6.6) for p^{**} and ℓ slightly bigger than r^{**} leads to a contradiction. □

7.7 Properties of geodesics and angles

Remark. All proofs in this section can be easely modified to use only the local definition of CBB spaces without use of the globalization theorem (7.6.2).

7.7.1. Geodesics do not split. *In* CBB *space, geodesics do not bifurcate.*

More precisely: let $\mathcal{L}$ be a CBB *space and $[px]$, $[py]$ be two geodesics, then*

a) If there is an $\varepsilon > 0$, such that $\operatorname{geod}_{[px]}(t) = \operatorname{geod}_{[py]}(t)$ for all $t \in [0, \varepsilon)$, then $[px] \subset [py]$ or $[py] \subset [px]$.

b) If $\measuredangle[p\,{}^{x}_{y}] = 0$, then $[px] \subset [py]$ or $[py] \subset [px]$.

7.7.2. Corollary. *Let $\mathcal{L}$ be a* CBB *space. Then restriction of any geodesic in $\mathcal{L}$ to a proper segment is the unique minimal geodesic joining its endpoints.*

In case $\kappa \leqslant 0$, the proof is easier, since the model triangles are always defined. To deal with $\kappa > 0$ we have to argue locally.

Proof of 7.7.1; a. Let $t_{\max}$ be the maximal value such that $\operatorname{geod}_{[px]}(t) = \operatorname{geod}_{[py]}(t)$ for all $t \in [0, t_{\max})$. Since geodesics are continuous $\operatorname{geod}_{[px]}(t_{\max}) = \operatorname{geod}_{[py]}(t_{\max})$; set



$$q = \operatorname{geod}_{[px]}(t_{\max}) = \operatorname{geod}_{[py]}(t_{\max}).$$

We have to show that $t_{\max} = \min\{|p - x|, |p - y|\}$.

If that is not true, choose sufficiently small $\varepsilon > 0$ such that points

$$x_\varepsilon = \operatorname{geod}_{[px]}(t_{\max} + \varepsilon) \quad \text{and} \quad y_\varepsilon = \operatorname{geod}_{[py]}(t_{\max} + \varepsilon)$$

are distinct. Set

$$z_\varepsilon = \operatorname{geod}_{[px]}(t_{\max} - \varepsilon) = \operatorname{geod}_{[py]}(t_{\max} - \varepsilon).$$

Clearly, $\tilde\angle^\kappa(q\,{}^{z_\varepsilon}_{x_\varepsilon}) = \tilde\angle^\kappa(q\,{}^{z_\varepsilon}_{y_\varepsilon}) = \pi$. Thus from CBB($\kappa$) comparison (7.1.2), $\tilde\angle^\kappa(q\,{}^{x_\varepsilon}_{y_\varepsilon}) = 0$ and thus $x_\varepsilon = y_\varepsilon$, a contradiction.

(b). From hinge comparison 7.3.1*c*

$$\measuredangle[p\,{}^{x}_{y}] = 0 \quad \Rightarrow \quad \tilde\angle^\kappa\left(p\,{}^{\operatorname{geod}_{[px]}(t)}_{\operatorname{geod}_{[py]}(t)}\right) = 0$$

and thus $\operatorname{geod}_{[px]}(t) = \operatorname{geod}_{[py]}(t)$ for all small t. Therefore we can apply (*a*). □

7.7.3. Adjacent angle lemma. *Let $\mathcal{L}$ be a* CBB *space. Assume that two hinges $[z\,{}^{x}_{p}]$ and $[z\,{}^{y}_{p}]$ in $\mathcal{L}$ are* adjacent*; that is, they share a common side $[zp]$ and $z \in\,]xy[$. Then*

$$\measuredangle[z\,{}^{p}_{y}] + \measuredangle[z\,{}^{p}_{x}] = \pi.$$

Proof. From hinge comparison (7.3.1*c*) we have that both angles $\measuredangle[z\,{}^{p}_{y}]$ and $\measuredangle[z\,{}^{p}_{x}]$ are defined and

$$\measuredangle[z\,{}^{p}_{y}] + \measuredangle[z\,{}^{p}_{x}] \leqslant \pi.$$

Clearly $\measuredangle[z\,{}^{x}_{y}] = \pi$. Thus, the result follows from the triangle inequality for angles (5.3.2). $\square$

7.7.4. Angle semicontinuity. *Let $\mathcal{L}_n$ be a* CBB(κ) *space for each $n \in \mathbb{N}$ and $\mathcal{L}_n \to \mathcal{L}_\omega$ as $n \to \omega$. Assume that the sequence of hinges $[p_n\,{}^{x_n}_{y_n}]$ in $\mathcal{L}_n$ converges to a hinge $[p_\omega\,{}^{x_\omega}_{y_\omega}]$ in $\mathcal{L}_\omega$. Then*

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] \leqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}].$$

Proof. From 7.3.2,

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] = \sup\left\{\, \tilde{\measuredangle}^\kappa(p_\omega\,{}^{\bar{x}_\omega}_{\bar{y}_\omega}) : \bar{x}_\omega \in\,]p_\omega x_\omega],\ \bar{y}_\omega \in\,]p_\omega x_\omega] \,\right\}.$$

For fixed $\bar{x}_\omega \in\,]p_\omega x_\omega]$ and $\bar{y}_\omega \in\,]p_\omega x_\omega]$, choose $\bar{x}_n \in\,]p x_n]$ and $\bar{y}_n \in\,]p y_n]$ so that $\bar{x}_n \to \bar{x}_\omega$ and $\bar{y}_n \to \bar{y}_\omega$ as $n \to \omega$. Clearly

$$\tilde{\measuredangle}^\kappa(p_n\,{}^{\bar{x}_n}_{\bar{y}_n}) \to \tilde{\measuredangle}^\kappa(p_\omega\,{}^{\bar{x}_\omega}_{\bar{y}_\omega})$$

as $n \to \omega$.



From the hinge comparison (7.3.1*c*), $\measuredangle[p_n\,{}^{x_n}_{y_n}] \geqslant$ $\geqslant \tilde{\measuredangle}^\kappa(p_n\,{}^{\bar{x}_n}_{\bar{y}_n})$. Hence the result. $\square$

7.7.5. Angle continuity. *Let $\mathcal{L}_n$ be a complete length* CBB(κ) *space and $\mathcal{L}_n \to \mathcal{L}_\omega$ as $n \to \omega$. Assume that sequences p_n, x_n, y_n in $\mathcal{L}_n$ converges correspondingly to the points $p_\omega, x_\omega, y_\omega$ in $\mathcal{L}_\omega$ and the following two conditions hold*

a) $p_\omega \in \mathrm{Str}(x_\omega)$,

b) $p_\omega \in \mathrm{Str}(y_\omega)$ *or* $y_\omega \in \mathrm{Str}(p_\omega)$.

Then

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] = \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}].$$

Proof. By Corollary 7.6.4, we may assume that $\kappa \leqslant 0$.

By Plaut's theorem (7.2.4), the hinge $[p_\omega\,{}^{x_\omega}_{y_\omega}]$ is uniquely defined. Therefore the hinges $[p_n\,{}^{x_n}_{y_n}]$ converge to $[p_\omega\,{}^{x_\omega}_{y_\omega}]$ as $n \to \omega$. Hence by angle semicontinuity (7.7.4), we get

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] \leqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}].$$

It remains to show that

❶ $$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] \geqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}].$$

Fix $\varepsilon > 0$. Since $p_\omega \in \mathrm{Str}(x_\omega)$, there is a point $q_\omega \in \mathcal{L}_\omega$ such that

$$\tilde{\measuredangle}^\kappa(p_\omega\,{}^{x_\omega}_{q_\omega}) > \pi - \varepsilon.$$

The hinge comparison (7.3.1c) implies that

❷ $$\measuredangle[p_\omega\,{}^{x_\omega}_{q_\omega}] > \pi - \varepsilon.$$

The triangle inequality for angles (5.3.2)

❸ $$\begin{aligned}\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] \geqslant \measuredangle[p_\omega\,{}^{x_\omega}_{q_\omega}] - \measuredangle[p_\omega\,{}^{y_\omega}_{q_\omega}] >\\ > \pi - \varepsilon - \measuredangle[p_\omega\,{}^{y_\omega}_{q_\omega}].\end{aligned}$$

Note that we can assume in addition that $q_\omega \in \mathrm{Str}(p_\omega)$. Choose $q_n \in \mathcal{L}_n$ such that $q_n \to q_\omega$ as $n \to \omega$. Note that the angle semicontinuity, again we get

❹ $$\begin{aligned}\measuredangle[p_\omega\,{}^{x_\omega}_{q_\omega}] \leqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{q_n}],\\ \measuredangle[p_\omega\,{}^{y_\omega}_{q_\omega}] \leqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{y_n}_{q_n}].\end{aligned}$$

By CBB(κ) comparison (7.1.2) and 7.3.2b, we get

$$\measuredangle[p_n\,{}^{x_n}_{y_n}] + \measuredangle[p_n\,{}^{y_n}_{q_n}] + \measuredangle[p_n\,{}^{x_n}_{q_n}] \leqslant 2\cdot\pi.$$

for all n. Together with ❹, ❷ and ❸, it implies

$$\begin{aligned}\lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}] \leqslant 2\cdot\pi - \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{q_n}] - \lim_{n\to\omega} \measuredangle[p_n\,{}^{y_n}_{q_n}] \leqslant\\ \leqslant 2\cdot\pi - \measuredangle[p_\omega\,{}^{x_\omega}_{q_\omega}] - \measuredangle[p_\omega\,{}^{y_\omega}_{q_\omega}] <\\ < \measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] + 2\cdot\varepsilon.\end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, ❶ follows. □

7.7.6. First variation formula. *Let $\mathcal{L}$ be a complete length* CBB *space. For any point q and geodesic $[px]$ in $\mathcal{L}$, we have*

❺ $$|q - \mathrm{geod}_{[px]}(t)| = |q - p| - t\cdot\cos\varphi + o(t),$$

where φ is the infimum of angles between $[px]$ and all geodesics from p to q in the ultrapower $\mathcal{L}^\omega$.

Remark. If $\mathcal{L}$ is proper space, then $\mathcal{L}^\omega = \mathcal{L}$, see Section 3.2. Therefore the infimum φ is achieved on some particular geodesic from p to q.

As a corollary we obtain the following classical result:

7.7.7. Strong angle lemma. *Let $\mathcal{L}$ be a complete length* CBB *space and $p, q \in \mathcal{L}$ be such that there is unique geodesic from p to q in the ultrapower $\mathcal{L}^\omega$. Then for any hinge $[p\,{}^q_x]$ we have*

❻ $$\measuredangle[p\,{}^q_x] = \lim_{\substack{\bar{x}\to p\\ \bar{x}\in\,]px]}} \tilde{\measuredangle}^\kappa(p\,{}^q_{\bar{x}}).$$

for any $\kappa \in \mathbb{R}$ such that $|p-q| < \varpi^\kappa$.

In partiqular ❻ *holds if $p \in \mathrm{Str}(q)$ as well as if $q \in \mathrm{Str}(p)$*

Remark.

- ◇ The above lemma is essentially due to Alexandrov. The right hand side in ❻ is called *strong angle* of hinge $[p\,{}^q_x]$. Note that in general metric space angle and strong angle of the same hinge might differ.
- ◇ As it follows from Corollary 3.3.4, if there is a unique geodesic $[pq]$ in the ultrapower $\mathcal{L}^\omega$, then $[pq]$ lies in $\mathcal{L}$.

Proof of 7.7.7. The first statement follows directly from the first variation formula (7.7.6) and definition of model angle (see Section 5.3). The second statement follows from the Plaut's theorem (7.2.4) applyied to $\mathcal{L}^\omega$. (Note that according to Proposition 7.1.7 $\mathcal{L}^\omega$ is a complete length CBB space.) □

Proof of 7.7.6. By Corollary 7.6.4, we can assume that $\kappa \leqslant 0$. The inequality

$$|q - \mathrm{geod}_{[px]}(t)| \leqslant |q-p| - t\cdot\cos\varphi + o(t)$$

follows from the first variation inequality (5.3.4). Thus, it is sufficient to show that

$$|q - \mathrm{geod}_{[px]}(t)| \geqslant |q-p| - t\cdot\cos\varphi + o(t).$$

Assume the contrary, then there is $\varepsilon > 0$, such that $\varphi + \varepsilon < \pi$ and for a sequence $t_n \to 0+$ we have

❼ $$|q - \mathrm{geod}_{[px]}(t_n)| < |q-p| - t_n\cdot\cos(\varphi-\varepsilon).$$

Set $x_n = \mathrm{geod}_{[px]}(t_n)$. Clearly

$$\tilde{\measuredangle}^\kappa(x_n\,{}^p_q) > \pi - \varphi + \tfrac{\varepsilon}{2}$$

for all large n.

Assume $\mathcal{L}$ is geodesic. Choose a sequence of geodesics $[x_nq]$. Let $[x_nq] \to [pq]_{\mathcal{L}^\omega}$ as $n \to \omega$ (in general $[pq]$ might lie in $\mathcal{L}^\omega$). Applying both parts of hinge comparison (7.3.1c), we get that $\measuredangle[x_n\,{}^q_x] < \varphi - \frac{\varepsilon}{2}$

for all large n. According to 7.7.4, the angle between $[pq]$ and $[px]$ is at most $\varphi - \frac{\varepsilon}{2}$, a contradiction.

Finally, if $\mathcal{L}$ is not geodesic choose a sequence $q_n \in \operatorname{Str}(x_n)$, such that $q_n \to q$ and the inequality

$$\measuredangle^{\kappa}(x_n\,{}^{p}_{q_n}) > \pi - \varphi + \tfrac{\varepsilon}{2}$$

still holds. Then the same argument as above shows that $[x_n q_n]$ ω-converges to a geodesic $[pq]_{\mathcal{L}^\omega}$ from p to q having angle at most $\varphi - \frac{\varepsilon}{2}$ with $[px]$. $\square$

7.8 On positive lower bound

In this section we consider CBB(κ) spaces for $\kappa > 0$. Applying rescaling we can assume that $\kappa = 1$.

The following theorem states that if one ignores a few exceptional spaces, then the diameter of a space with positive lower curvature bound is bounded. Note that many authors (but not us) exclude these spaces in the definition of Alexandrov space with positive lower curvature bound.

7.8.1. On diameter of a space. *Let $\mathcal{L}$ be a complete length* CBB(1) *space. Then either*

a) $\operatorname{diam}\mathcal{L} \leqslant \pi$*;*

b) or $\mathcal{L}$ is isometric to one of the following exceptional spaces:

1. *real line $\mathbb{R}$,*
2. *a ray $\mathbb{R}_{\geqslant 0}$,*
3. *a closed interval $[0, a] \in \mathbb{R}$, $a > \pi$,*
4. *a circle $\mathbb{S}^1_a$ of length $a > 2\cdot\pi$.*

Proof. Assume that $\mathcal{L}$ is a geodesic space and $\operatorname{diam}\mathcal{L} > \pi$. Choose $x, y \in \mathcal{L}$ so that $|x - y| = \pi + \varepsilon$, $0 < \varepsilon < \frac{\pi}{4}$. By moving y slightly, we can also assume that geodesic $[xy]$ is unique; to prove it, use either Plaut's theorem (7.2.4) or that geodesics do not split (7.7.1). Let z be the midpoint of a geodesic $[xy]$.

Consider the function $f = \operatorname{dist}_x + \operatorname{dist}_y$. As it follows from Lemma 7.8.2, f is concave in $\mathrm{B}(z, \frac{\varepsilon}{4})$. Let $p \in \mathrm{B}(z, \frac{\varepsilon}{4})$. Choose a geodesic $[zp]$. Set $h(t) = f \circ \operatorname{geod}_{[zp]}(t)$ and $\ell = |z - p|$. Clearly h is concave. From Adjacent angle lemma (7.7.3), we have that $h^+(0) = 0$. Therefore h is nonincreasing which means that

$$|x - p| + |y - p| = h(\ell) \leqslant h(0) = |x - y|.$$

Since the geodesic $[xy]$ is unique this means that $p \in [xy]$ and hence $\mathrm{B}(z, \frac{\varepsilon}{4})$ only contains points of $[xy]$. Since in CBB spaces, geodesics do not bifurcate (7.7.1a), it follows that all of $\mathcal{L}$ coincides with the maximal extension of $[xy]$ as a local geodesic γ (which might be not minimizing). In other words, $\mathcal{L}$ is isometric to a 1-dimensional Riemannian manifold with possibly non empty boundary. From this, it is easy to see that $\mathcal{L}$ falls into one of the exceptional spaces described in the theorem.

Lastly, if $\mathcal{L}$ is not geodesic and $\operatorname{diam}\mathcal{L} > \pi$, then the above argument applied to $\mathcal{L}^\omega$ yields that $\mathcal{L}^\omega$ is isometric to one of the exceptional spaces. As all of those spaces are proper it means that $\mathcal{L} = \mathcal{L}^\omega$. □

7.8.2. Lemma. *Let $\mathcal{L}$ be a complete length* CBB(1) *space and $p \in \mathcal{L}$. Then $\operatorname{dist}_p : \mathcal{L} \to \mathbb{R}$ is concave in $\mathrm{B}(p,\pi)\backslash\mathrm{B}(p, \frac{\pi}{2})$.*

In particular, if $\operatorname{diam}\mathcal{L} \leqslant \pi$ then the complements $\mathcal{L}\backslash\mathrm{B}(p,r)$ and $\mathcal{L}\backslash\overline{\mathrm{B}}[p,r]$ are convex for any $r > \frac{\pi}{2}$.

Proof. This is a consequence of 7.4.1b. □

7.8.3. Exercise. *Let $\mathcal{L}$ be an m-dimensional complete length* CBB(1) *space and $\operatorname{diam}\mathcal{L} \leqslant \pi$. Assume that the group G acts on $\mathcal{L}$ by isometries and has closed orbits and*

$$\operatorname{diam}(\mathcal{L}/G) > \tfrac{\pi}{2}.$$

Show that the action of G has a fixed point in $\mathcal{L}$.

7.8.4. Advanced exercise. *Let $\mathcal{L}$ be a complete length* CBB(1) *space. Show that $\mathcal{L}$ contains at most 3 points with space of directions $\leqslant \frac{1}{2}\cdot\mathbb{S}^n$.*

7.8.5. On perimeter of a triple. *Let $\mathcal{L}$ be a complete length* CBB(1) *space and $\operatorname{diam}\mathcal{L} \leqslant \pi$. Then perimeter of any triple of points $p, q, r \in \mathcal{L}$ is at most $2\cdot\pi$.*

Proof. Arguing by contradiction, suppose

❶ $$|p-q| + |q-r| + |r-p| > 2\cdot\pi$$

for some $p, q, r \in \mathcal{L}$. Rescaling the space slightly, we can assume that $\operatorname{diam}\mathcal{L} < \pi$, but the inequality ❶ still holds.

By Corollary 7.6.4, after rescaling $\mathcal{L}$ is still CBB(1).

Since $\mathcal{L}$ is G-delta geodesic (7.2.4), it is sufficient to consider case when there is a geodesic $[qr]$.



First note that since $\operatorname{diam}\mathcal{L} < \pi$, by 7.4.1*b* we have that the function

$$y(t) = \mathrm{md}^1|p - \mathrm{geod}_{[qr]}(t)|$$

satisfies the differential inequality $y'' \leqslant 1 - y$.

Take $z_0 \in [qr]$ so that restriction $\mathrm{dist}_p|_{[qr]}$ attains its maximum at z_0, set $t_0 = |q - z_0|$ so $z_0 = \mathrm{geod}_{[qr]}(t_0)$. Consider the following model configuration: two geodesics $[\tilde p\tilde z_0]$, $[\tilde q\tilde r]$ in $\mathbb{S}^2$ such that

$$|\tilde p - \tilde z_0| = |p - z_0|, \qquad |\tilde q - \tilde r| = |q - r|,$$
$$|\tilde z_0 - \tilde q| = |z_0 - q|, \qquad |\tilde z_0 - \tilde r| = |z_0 - q|$$

and

$$\measuredangle[\tilde z_0\,{}^{\tilde q}_{\tilde p}] = \measuredangle[\tilde z_0\,{}^{\tilde r}_{\tilde p}] = \tfrac{\pi}{2}.$$

Clearly, $\bar y(t) = \mathrm{md}^1|\tilde p - \mathrm{geod}_{[\tilde q\tilde r]}(t)|$ satisfies $\bar y'' = 1 - \bar y$ and $\bar y'(t_0) = 0$, $\bar y(t_0) = y(t_0)$. Since z_0 is a maximum point, $y(t) \leqslant y(t_0) + o(t - t_0)$; thus, $\bar y(t)$ is a barrier for $y(t) = \mathrm{md}^1|p - \mathrm{geod}_{[qr]}(t)|$ at t_0 by 4.5.3*c*. From barrier inequality (4.5.3*c*), we get

$$|\tilde p - \mathrm{geod}_{[\tilde q\tilde r]}(t)| \geqslant |p - \mathrm{geod}_{[qr]}(t)|,$$

thus, $|\tilde p - \tilde q| \geqslant |p - q|$ and $|\tilde p - \tilde r| \geqslant |p - r|$.

Therefore, $|p - q| + |q - r| + |r - p|$ can not exceed the perimeter of the spherical triangle $[\tilde p\tilde q\tilde r]$; that is,

$$|p - q| + |q - r| + |r - p| \leqslant 2\cdot\pi,$$

a contradiction. □

Let $\kappa > 0$. Consider the following extension $\measuredangle^{\kappa+}(*\,{}^{*}_{*})$ of the model angle function $\measuredangle^{\kappa}(*\,{}^{*}_{*})$. Some authors define comparison angle to be $\measuredangle^{\kappa+}(*\,{}^{*}_{*})$ but that leads to two different definition of model angles — one is good for CAT spaces and the other for CBB spaces.

7.8.6. Definition of extended angle. *Let p, q, r be points in a metric space such that $p \neq q$, $p \neq r$. Then set*

$$\measuredangle^{\kappa+}(p\,{}^{q}_{r}) = \sup\left\{\,\measuredangle^{\mathrm{K}}(p\,{}^{q}_{r}) : \mathrm{K} \leqslant \kappa\,\right\}.$$

The value $\measuredangle^{\kappa+}(p\,{}^{q}_{r})$ is called extended model angle *of triple p, q, r.*

7.8.7. Extended angle comparison. *Let $\kappa > 0$ and $\mathcal{L}$ be a complete length* CBB *space. Then for any hinge $[p\,{}^{q}_{r}]$ we have $\measuredangle[p\,{}^{q}_{r}] \geqslant$ $\geqslant \measuredangle^{\kappa+}(p\,{}^{q}_{r})$.*

Moreover, the extended model angle can be calculated $\measuredangle^{\kappa+}(p\,{}^{q}_{r})$ using the following rules:

a) $\measuredangle^{\kappa+}(p\,{}^{q}_{r}) = \measuredangle^{\kappa}(p\,{}^{q}_{r})$ *once* $\measuredangle^{\kappa}(p\,{}^{q}_{r})$ *is defined;*
b) $\measuredangle^{\kappa+}(p\,{}^{q}_{r}) = \measuredangle^{\kappa+}(p\,{}^{r}_{q}) = 0$ *if* $|p-q|+|q-r| = |p-r|$*;*
c) $\measuredangle^{\kappa+}(p\,{}^{q}_{r}) = \pi$ *if none of above is applicable.*

Proof. From Corollary 7.6.4, $\mathrm{K} < \kappa$ implies that any complete length CBB(K) space is CBB(κ); thus the extended angle comparison follows from the definition.

The rules for calculating extended angle is an easy consequence of its definition. □

7.9 Remarks and open problems

The question whether the first part of 7.3.1*c* suffices to conclude that $\mathcal{L}$ is CBB(κ) is a long standing open problem (possibly dating back to Alexandrov), but as far as we know, it was first formally stated in print in [4, footnote in 4.1.5].

7.9.1. Open question. *Let* $\mathcal{L}$ *be a complete geodesic space (you can also assume that* $\mathcal{L}$ *is homeomorphic to* $\mathbb{S}^2$ *or* $\mathbb{R}^2$*) such that for any hinge* $[x\,{}^{p}_{y}]$ *in* $\mathcal{L}$*, the angle* $\measuredangle[x\,{}^{p}_{y}]$ *is defined and*

$$\measuredangle[x\,{}^{p}_{y}] \geqslant \measuredangle^{0}(x\,{}^{p}_{y}).$$

Is it true that $\mathcal{L}$ *is* CBB(0)*?*

Chapter 8

Definitions of curvature bounded above

8.1 Four-point comparison.

8.1.1. Four-point comparison. *A quadruple of points* p^1, p^2, x^1, x^2 *in a metric space satisfies* CAT(κ) comparison *if*

a) $\measuredangle^\kappa(p^1\,{}^{x^1}_{x^2}) \leqslant \measuredangle^\kappa(p^1\,{}^{p^2}_{x^1}) + \measuredangle^\kappa(p^1\,{}^{p^2}_{x^2})$, *or*

b) $\measuredangle^\kappa(p^2\,{}^{x^1}_{x^2}) \leqslant \measuredangle^\kappa(p^2\,{}^{p^1}_{x^1}) + \measuredangle^\kappa(p^2\,{}^{p^1}_{x^2})$, *or*

c) one of the six model angles

$$\measuredangle^\kappa(p^1\,{}^{x^1}_{x^2}),\ \measuredangle^\kappa(p^1\,{}^{p^2}_{x^1}),\ \measuredangle^\kappa(p^1\,{}^{p^2}_{x^2}),\ \measuredangle^\kappa(p^2\,{}^{x^1}_{x^2}),\ \measuredangle^\kappa(p^2\,{}^{p^1}_{x^1}),\ \measuredangle^\kappa(p^2\,{}^{p^1}_{x^2})$$

is undefined.

Here is more intuitive reformulation.



8.1.2. Reformulation. *Let* $\mathcal{X}$ *be a metric space. A quadruple* $p^1, p^2, x^1, x^2 \in \mathcal{X}$ *satisfies* CAT(κ) *point comparison if one of the following holds:*

a) One of the triples (p^1, p^2, x^1) *or* (p^1, p^2, x^2) *has perimeter* $> 2\cdot\varpi^\kappa$.

b) If $[\tilde p^1\tilde p^2\tilde x^1] = \tilde\triangle^\kappa(p^1p^2x^1)$ *and* $[\tilde p\tilde p^2\tilde x^2] = \tilde\triangle^\kappa p^1p^2x^2$, *then*

$$|\tilde x^1 - \tilde z| + |\tilde z - \tilde x^2| \geqslant |x^1 - x^2|,$$

for any $\tilde z \in [\tilde p^1\tilde p^2]$.

8.1.3. Definition. *A metric space $\mathcal{U}$ is called* CAT(κ) *if every quadruple p^1, p^2, x^1, x^2 satisfies the* CAT(κ) *comparison (8.1.1).*

We say that $\mathcal{U}$ is CAT *if it is* CAT(κ) *for some $\kappa \in \mathbb{R}$.*

The condition $\mathcal{U}$ is CAT(κ) should be understood as "$\mathcal{U}$ has global curvature $\leqslant \kappa$". In Proposition 8.3.4, it will be shown that this formulation makes sense; in particular, if $\kappa \leqslant \mathrm{K}$, then any CAT($\kappa$) space is CAT(K).

This terminology was introduced by Gromov; CAT stands for Élie Cartan, Alexandr Alexandrov and Victor Toponogov. Originally these spaces were called $\mathfrak{R}_\kappa$ *domains*; this is Alexandrov's terminology and is still in use.

8.1.4. Exercise. *Let $\mathcal{U}$ be a metric space. Show that $\mathcal{U}$ is* CAT(κ) *if and only if every quadruple of points in $\mathcal{U}$ admits one labeling by (p, x^1, x^2, x^3) so that the three angles $\measuredangle^\kappa(p\,{}^{x^1}_{x^2})$, $\measuredangle^\kappa(p\,{}^{x^2}_{x^3})$ and $\measuredangle^\kappa(p\,{}^{x^1}_{x^3})$ satisfy all three triangle inequalities or one of these angels is undefined.*

8.1.5. Exercise. *Show that $\mathcal{U}$ is* CAT(κ) *if and only if for any quadruple of points p^1, p^2, x^1, x^2 in $\mathcal{U}$ such that $|p^1 - p^2|, |x^1 - x^2| \leqslant \varpi^\kappa$ there is a quadruple q^1, q^2, y^1, y^2 in $\mathbb{M}^m(\kappa)$ such that*

$$|q^1 - q^2| = |p^1 - p^2|, \quad |y^1 - y^2| = |x^1 - x^2|, \quad |q^i - y^j| \leqslant |p^i - x^j|$$

for any i and j.

The next proposition follows directly from Definition 8.1.3 and the definitions of ultralimit and ultrapower; see Section 3.2 for the related definitions. Recall that ω denotes a selective ultrafilter on $\mathbb{N}$ which is fixed once for all.

8.1.6. Proposition. *Let $\mathcal{U}_n$ be a* CAT(κ_n) *space for each $n \in \mathbb{N}$. Assume $\mathcal{U}_n \to \mathcal{U}_\omega$ and $\kappa_n \to \kappa_\omega$ as $n \to \omega$. Then $\mathcal{U}_\omega$ is* CAT(κ_ω).

Moreover, a metric space $\mathcal{U}$ is CAT(κ) *if and only if so is its ultrapower $\mathcal{U}^\omega$.*

8.2 Geodesics

8.2.1. Uniqueness of geodesics. *In a complete length* CAT(κ) *space, pairs of points at distance $< \varpi^\kappa$ are joined by unique geodesics, and these geodesics depend continuously on their endpoint pairs.*

Proof. Fix a complete length CAT(κ) space $\mathcal{U}$. Fix two points $p^1, p^2 \in \mathcal{U}$ such that

$$|p^1 - p^2|_{\mathcal{U}} < \varpi^\kappa.$$

Choose a sequence of approximate midpoints z_n for p^1 and p^2; that is,

❶ $$|p^1 - z_n|, |p^2 - z_n| \to \tfrac{1}{2}\cdot|p^1 - p^2| \text{ as } n \to \infty.$$

By the law of cosines, $\measuredangle^\kappa(p^1\,{}^{z_n}_{p^2})$ and $\measuredangle^\kappa(p^2\,{}^{z_n}_{p^1})$ are arbitrarily small when n is sufficiently large.

Let us apply $\mathrm{CAT}(\kappa)$ comparison (8.1.1) to the quadruple p^1, p^2, z_n, $z_{\bar{k}}$ with large n and $\bar{k}$. We conclude that $\measuredangle^\kappa(p\,{}^{z_n}_{z_{\bar{k}}})$ is arbitrarily small when $n, \bar{k}$ are sufficiently large and p is either p^1 or p^2. By ❶ and the law of cosines, (z_n) converges.

Since $\mathcal{U}$ is complete, the sequence (z_n) converges to a midpoint of p^1 and p^2. By Lemma 2.3.8 we obtain the existence of a geodesic $[p^1p^2]$.

Now suppose $p^1_n \to p^1$, $p^2_n \to p^2$ as $n \to \infty$. Let z_n be the midpoint of a geodesic $[p^1_np^2_n]$ and z be the midpoint of a geodesic $[p^1p^2]$.

It suffices to show that

❷ $$|z_n - z| \to 0 \quad \text{as} \quad n \to \infty.$$

By the triangle inequality, the z_n are approximate midpoints of p^1 and p^2. Apply $\mathrm{CAT}(\kappa)$ comparison (8.1.1) to the quadruple p^1, p^2, z_n, z. For $p = p^1$ or $p = p^2$, we see that $\measuredangle^\kappa(p\,{}^{z_n}_{z})$ is arbitrarily small when n is sufficiently large. By the law of cosines, ❷ follows. □

8.2.2. Exercise. *Let $\mathcal{U}$ be a complete length* CAT *space. Assume $\mathcal{U}$ is a topological manifold. Show that any geodesic in $\mathcal{U}$ can be extended as a two-side infinite local geodesic.*

Moreover the same holds for any locally geodesic locally CAT *space $\mathcal{U}$ with nontrivial local homology groups at any point; the latter holds in particular, if $\mathcal{U}$ is a homological manifold.*

8.2.3. Exercise. *Assume $\mathcal{U}$ is a locally compact geodesic* CAT *space with extandable geodesics; that is, any geodesic in $\mathcal{U}$ can be extended to a both-side infinite local geodesic.*

Show that the space of geodesic directions Σ'_p is complete for any $p \in \mathcal{U}$.

By uniqueness of geodesics (8.2.1), we have the following.

8.2.4. Corollary. *Any complete length* $\mathrm{CAT}(\kappa)$ *space is ϖ^κ-geodesic.*

8.2.5. Proposition. *The completion of any geodesic* $\mathrm{CAT}(\kappa)$ *space is a complete length* $\mathrm{CAT}(\kappa)$ *space.*

Moreover, $\mathcal{U}$ is a geodesic CAT(κ) *space if and only if there is a complete length* CAT(κ) *space $\bar{\mathcal{U}}$ that contains a ϖ^κ-convex dense set isometric to $\mathcal{U}$.*

Proof. By Theorem 8.2.1, in order to show that the completion $\bar{\mathcal{U}}$ of any geodesic CAT(κ) space $\mathcal{U}$ is CAT(κ), it is sufficient to verify that the completion of a length space is a length space; this is straightforward.

For the second part of the proposition, note that the completion $\bar{\mathcal{U}}$ contains the original space $\mathcal{U}$ as a dense ϖ^κ-convex subset, and the metric on $\mathcal{U}$ coincides with the induced length metric from $\bar{\mathcal{U}}$. □

Here is a corollary from Proposition 8.2.5 and Theorem 8.2.1.

8.2.6. Corollary. *Let $\mathcal{U}$ be a ϖ^κ-geodesic* CAT(κ) *space. Then pairs of points in $\mathcal{U}$ at distance less than ϖ^κ are joined by unique geodesics, and these geodesics depend continuously on their endpoint pairs.*

Moreover for any pair of points $p, q \in \mathcal{U}$ and any value

$$\pounds > \sup\left\{ \frac{\operatorname{sn}^\kappa r}{\operatorname{sn}^\kappa |p-q|} : 0 \leqslant r \leqslant |p-q| \right\}$$

there are neighborhoods $\Omega_p \ni p$ and $\Omega_q \ni q$ such that the map

$$(x, y, t) \mapsto \operatorname{path}_{[xy]}(t)$$

is $\pounds$-Lipschitz in $\Omega_p \times \Omega_q \times [0,1]$.

Proof. By Proposition 8.2.5 any geodesic CAT(κ) space is isometric to a convex dense subset of a complete length CAT(κ) space. It remains to apply Theorem 8.2.1. □

8.3 More comparisons

Here we give a few reformulations of Definition 8.1.3.

8.3.1. Theorem. *If $\mathcal{U}$ is a* CAT(κ) *space, then the following conditions hold for all triples $p, x, y \in \mathcal{U}$ of perimeter $< 2\cdot\varpi^\kappa$:*

a) (adjacent-angles comparison) for any geodesic $[xy]$ and $z \in\,]xy[$, we have

$$\measuredangle^\kappa(z\,{}^p_x) + \measuredangle^\kappa(z\,{}^p_y) \geqslant \pi.$$



b) (point-on-side comparison) for any geodesic $[xy]$ and $z \in$ $\in\,]xy[$, we have

$$\measuredangle^{\kappa}(x\,{}^{p}_{y}) \geqslant \measuredangle^{\kappa}(x\,{}^{p}_{z}),$$

or equivalently,

$$|\tilde{p} - \tilde{z}| \geqslant |p - z|,$$

where $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^{\kappa}(pxy)$, $\tilde{z} \in\,]\tilde{x}\tilde{y}[$, $|\tilde{x} - \tilde{z}| = |x - z|$.

c) (angle comparison) for any hinge $[x\,{}^{p}_{y}]$, the angle $\measuredangle[x\,{}^{p}_{y}]$ exists and

$$\measuredangle[x\,{}^{p}_{y}] \leqslant \measuredangle^{\kappa}(x\,{}^{p}_{y}),$$

or equivalently,

$$\tilde{\gamma}^{\kappa}[x\,{}^{p}_{y}] \leqslant |p - y|.$$

Moreover, if $\mathcal{U}$ is ϖ^{κ}-geodesic, then the converse holds in each case.

Remark. In the following proof, the part $(c)\Rightarrow(a)$ only requires that the CAT(κ) comparison (8.1.1) hold for any quadruple, and does not require the existence of geodesics at distance $< \varpi^{\kappa}$. The same is true of the parts $(a)\Leftrightarrow(b)$ and $(b)\Rightarrow(c)$. Thus the conditions (a), (b) and (c) are valid for any metric space (not necessarily length space) that satisfies CAT(κ) comparison (8.1.1). The converse does not hold; for example, all these conditions are vacuously true in a totally disconnected space, while CAT(κ) comparison is not.

Proof. (a). Since the perimeter of p, x, y is $< 2\cdot\varpi^{\kappa}$, so is the perimeter of any subtriple of p, z, x, y by the triangle inequality. By Alexandrov's lemma (5.2.1),

$$\measuredangle^{\kappa}(p\,{}^{z}_{x}) + \measuredangle^{\kappa}(p\,{}^{z}_{y}) < \measuredangle^{\kappa}(p\,{}^{x}_{y}) \text{ or } \measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) = \pi.$$

In the former case, CAT(κ) comparison (8.1.1) applied to the quadruple p, z, x, y implies

$$\measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) \geqslant \measuredangle^{\kappa}(z\,{}^{x}_{y}) = \pi.$$

(a) $\Leftrightarrow$ (b). Follows from Alexandrov's lemma (5.2.1).

(b) $\Rightarrow$ (c). By (b), for $\bar{p} \in\,]xp]$ and $\bar{y} \in\,]xy]$ the function $(|x - \bar{p}|, |x - \bar{y}|) \mapsto \measuredangle^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}})$ is nondecreasing in each argument. In particular, $\measuredangle[x\,{}^{p}_{y}] = \inf \measuredangle^{\kappa}(x\,{}^{\bar{p}}_{\bar{y}})$. Thus $\measuredangle[x\,{}^{p}_{y}]$ exists and is at most $\measuredangle^{\kappa}(x\,{}^{p}_{y})$.

Converse. Assume $\mathcal{U}$ is ϖ^{κ}-geodesic. Let us first show that in this case $(c) \Rightarrow (a)$.

Indeed, by (c) and the triangle inequality for angles (5.3.2),

$$\measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) \geqslant \measuredangle[z\,{}^{p}_{x}] + \measuredangle[z\,{}^{p}_{y}] \geqslant \pi.$$



Therefore, it is sufficient to prove the converse for *(b)*.

Given a quadruple p^1, p^2, x^1, x^2 whose subtriples have perimeter $< 2\cdot\varpi^\kappa$, we must verify CAT(κ) comparison (8.1.1). In $\mathbb{M}^2(\kappa)$, construct the model triangles $[\tilde p^1\tilde p^2\tilde x^1] = \tilde\triangle^\kappa(p^1p^2x^1)$ and $[\tilde p^1\tilde p^2\tilde x^2] = \tilde\triangle^\kappa(p^1p^2x^2)$, lying on either side of a common segment $[\tilde p^1\tilde p^2]$. We may suppose

$$\measuredangle^\kappa(p^1\,{}^{p^2}_{x^1}) + \measuredangle^\kappa(p^1\,{}^{p^2}_{x^2}) \leqslant \pi \quad\text{and}\quad \measuredangle^\kappa(p^2\,{}^{p^1}_{x^1}) + \measuredangle^\kappa(p^2\,{}^{p^1}_{x^2}) \leqslant \pi,$$

since otherwise CAT(κ) comparison holds trivially. Then $[\tilde p^1\tilde p^2]$ and $[\tilde x^1\tilde x^2]$ intersect, say at $\tilde q$.

By assumption, there is a geodesic $[p^1p^2]$. Choose $q \in [p^1p^2]$ corresponding to $\tilde q$; that is, $|p^1 - q| = |\tilde p^1 - \tilde q|$. Then

$$|x^1 - x^2| \leqslant |x^1 - q| + |q - x^2| \leqslant |\tilde x^1 - \tilde q| + |\tilde q - \tilde x^2| = |\tilde x^1 - \tilde x^2|,$$

where the second inequality follows from (b). Therefore by monotonicity of the function $a \mapsto \measuredangle^\kappa\{a; b, c\}$ (1.1.1c),

$$\measuredangle^\kappa(p^1\,{}^{x^1}_{x^2}) \leqslant \measuredangle[\tilde p^1\,{}^{\tilde x^1}_{\tilde x^2}] = \measuredangle^\kappa(p^1\,{}^{p^2}_{x^1}) + \measuredangle^\kappa(p^1\,{}^{p^2}_{x^2}).$$

□

Let us display a corollary from the proof of 8.3.1, namely, monotonicity of the model angle with respect to adjacent sidelengths.

8.3.2. Angle-sidelength monotonicity. *Suppose $\mathcal{U}$ is a ϖ^κ-geodesic* CAT(κ) *space, and $p, x, y \in \mathcal{U}$ have perimeter $< 2\cdot\varpi^\kappa$. Then for $\bar y \in$ $\in\,]xy]$, the function*

$$|x - \bar y| \mapsto \measuredangle^\kappa(x\,{}^{p}_{\bar y})$$

is nondecreasing.

In particular, if $\bar p \in\,]xp]$, then

a) the function

$$(|x - \bar y|, |x - \bar p|) \mapsto \measuredangle^\kappa(x\,{}^{\bar p}_{\bar y})$$

is nondecreasing in each argument,

b) the angle $\measuredangle[x\,{}^{p}_{y}]$ exists and

$$\measuredangle[x\,{}^{p}_{y}] = \inf\left\{\, \measuredangle^\kappa(x\,{}^{\bar p}_{\bar y}) : \bar p \in\,]xp],\ \bar y \in\,]xy]\,\right\}.$$

8.3.3. Exercise. *Assume $\mathcal{U}$ is a geodesic* CAT(0) *space. Show that for any two geodesic paths $\gamma, \sigma\colon [0,1] \to \mathcal{U}$ the function*

$$t \mapsto |\gamma(t) - \sigma(t)|$$

is convex.

8.3.4. Proposition. *Assume* $\kappa < \mathrm{K}$. *Then any complete length* CAT(κ) *space is* CAT(K).

Moreover a space $\mathcal{U}$ *is* CAT(κ) *if* $\mathcal{U}$ *is* CAT(K) *for all* $\mathrm{K} > \kappa$.

Proof. The first statement follows from Corollary 8.2.4, the adjacent-angles comparison (8.3.1a) and the monotonicity of the function $\kappa \mapsto$ $\mapsto \measuredangle^\kappa(x\,{}^y_z)$ (1.1.1d).

The second statement follows since the function $\kappa \mapsto \measuredangle^\kappa(x\,{}^y_z)$ is continuous. □

8.4 Thin triangles

In this section we define thin triangles and use them to characterize CAT spaces. Inheritance for thin triangles with respect to decomposition is the main result of this section. It will lead to two fundamental constructions: Alexandrov's patchwork globalization (8.8.1) and Reshetnyak gluing (8.9.1).

8.4.1. Definition of κ-thin triangles. *Let* $[x^1x^2x^3]$ *be a triangle of perimeter* $< 2{\cdot}\varpi^\kappa$ *in a metric space. Consider its model triangle* $[\tilde{x}^1\tilde{x}^2\tilde{x}^3] = \tilde{\triangle}^\kappa(x^1x^2x^3)$ *and the* natural map $[\tilde{x}^1\tilde{x}^2\tilde{x}^3] \to [x^1x^2x^3]$ *that sends a point* $\tilde{z} \in [\tilde{x}^i\tilde{x}^j]$ *to the corresponding point* $z \in [x^ix^j]$ *(that is, such that* $|\tilde{x}^i - \tilde{z}| = |x^i - z|$ *and therefore* $|\tilde{x}^j - \tilde{z}| = |x^j - z|$*).*

We say the triangle $[x^1x^2x^3]$ *is* κ-thin *if the natural map* $[\tilde{x}^1\tilde{x}^2\tilde{x}^3] \to$ $\to [x^1x^2x^3]$ *is short.*

8.4.2. Exercise. *Let* $\mathcal{U}$ *be a* ϖ^κ*-geodesic* CAT(κ) *space,* $[xyz]$ *be a triangle in* $\mathcal{U}$ *and* $[\tilde{x}\tilde{y}\tilde{z}]$ *its model triangle in* $\mathbb{M}^2(\kappa)$. *Prove that the natural map* $f\colon [\tilde{x}\tilde{y}\tilde{z}] \to [xyz]$ *is distance-preserving if and ony if one of the following conditions hold:*

a) $\measuredangle[x\,{}^y_z] = \measuredangle^\kappa(x\,{}^y_z)$,

b) $|x - w| = |\tilde{x} - \tilde{w}|$ *for some* $\tilde{w} \in]\tilde{y}\tilde{z}[$ *and* $w = f(\tilde{w})$,

c) $|v - w| = |\tilde{v} - \tilde{w}|$ *for some* $\tilde{v} \in]\tilde{x}\tilde{y}[$, $\tilde{w} \in]\tilde{x}\tilde{z}[$ *and* $v = f(\tilde{v})$, $w = f(\tilde{w})$.

8.4.3. Proposition. *Let* $\mathcal{U}$ *be a* ϖ^κ*-geodesic space. Then* $\mathcal{U}$ *is* CAT(κ) *if and only if every triangle of perimeter* $< 2{\cdot}\varpi^\kappa$ *in* $\mathcal{U}$ *is* κ*-thin.*

Proof. "If" follows from point-on-side comparison 8.3.1b. "Only if" follows from the angle-sidelength monotonicity 8.3.2a. □

8.4.4. Corollary. *Suppose $\mathcal{U}$ is a ϖ^κ-geodesic* CAT(κ) *space. Then any local geodesic in $\mathcal{U}$ of length $< \varpi^\kappa$ is length-minimizing.*

Proof. Suppose $\gamma\colon [0,\ell] \to \mathcal{U}$ is a local geodesic that is not minimizing, with $\ell < \varpi^\kappa$. Choose a to be the maximal value such that γ is minimizing on $[0,a]$. Further choose $b > a$ so that γ is minimizing on $[a,b]$.

Since triangle $[\gamma(0)\gamma(a)\gamma(b)]$ is κ-thin, we have

$$|\gamma(a-\varepsilon) - \gamma(a+\varepsilon)| < 2\cdot\varepsilon$$

for all small $\varepsilon > 0$, a contradiction. □

Now let us formulate the main result of this section. The inheritance lemma states that in any metric space, a triangle is κ-thin if it decomposes into κ-thin triangles. In contrast, CBB(κ) comparisons are not inherited in this way.

8.4.5. Inheritance lemma. *In a metric space, consider a triangle $[pxy]$ that* decomposes *into two triangles $[pxz]$ and $[pyz]$; that is, $[pxz]$ and $[pyz]$ have common side $[pz]$, and the sides $[xz]$ and $[zy]$ together form the side $[xy]$ of $[pxy]$.*

If the triangle $[pxy]$ has perimeter $< 2\cdot\varpi^\kappa$ and both triangles $[pxz]$ and $[pyz]$ are κ-thin, then triangle $[pxy]$ is κ-thin.



The following model-space lemma, is an indeterminate statement in the proof of [24, Lemma 2].

8.4.6. Lemma. *Let $[\tilde p\tilde x\tilde y]$ be a triangle in $\mathbb{M}^2(\kappa)$ and $\tilde z \in [\tilde x\tilde y]$. Consider the solid triangle $\tilde D = \operatorname{Conv}[\tilde p\tilde x\tilde y]$. Construct points $\dot p, \dot x, \dot z, \dot y \in$ $\in \mathbb{M}^2(\kappa)$ such that*

$$|\dot p - \dot x| = |\tilde p - \tilde x|, \qquad |\dot p - \dot y| = |\tilde p - \tilde y|, \qquad |\dot p - \dot z| \leqslant |\tilde p - \tilde z|,$$
$$|\dot x - \dot z| = |\tilde x - \tilde z|, \qquad |\dot y - \dot z| = |\tilde y - \tilde z|,$$

where points $\dot x$ and $\dot y$ lie on either side of $[\dot p\dot z]$. Set

$$\dot D = \operatorname{Conv}[\dot p\dot x\dot z] \cup \operatorname{Conv}[\dot p\dot y\dot z].$$

Then there is a short map $F\colon \tilde D \to \dot D$ that maps $\tilde p$, $\tilde x$, $\tilde y$ and $\tilde z$ to $\dot p$, $\dot x$, $\dot y$ and $\dot z$ respectively.



Proof. By Alexandrov's lemma (5.2.1), there are nonoverlapping triangles $[\tilde{p}\tilde{x}\tilde{z}_y] \overset{iso}{=\!=} [\dot{p}\dot{x}\dot{z}]$ and $[\tilde{p}\tilde{y}\tilde{z}_x] \overset{iso}{=\!=} [\dot{p}\dot{y}\dot{z}]$ inside triangle $[\tilde{p}\tilde{x}\tilde{y}]$.

Connect points in each pair $(\tilde{z}, \tilde{z}_x)$, $(\tilde{z}_x, \tilde{z}_y)$ and $(\tilde{z}_y, \tilde{z})$ with arcs of circles centered at $\tilde{y}$, $\tilde{p}$, and $\tilde{x}$ respectively. Define F as follows.

- ◇ Map $\operatorname{Conv}[\tilde{p}\tilde{x}\tilde{z}_y]$ isometrically onto $\operatorname{Conv}[\dot{p}\dot{x}\dot{y}]$; similarly map $\operatorname{Conv}[\tilde{p}\tilde{y}\tilde{z}_x]$ onto $\operatorname{Conv}[\dot{p}\dot{y}\dot{z}]$.
- ◇ If w is in one of the three circular sectors, say at distance r from center of the circle, let $F(w)$ be the point on the corresponding segment $[\dot{p}\dot{z}]$, $[\dot{x}\dot{z}]$ or $[\dot{y}\dot{z}]$ whose distance from the lefthand endpoint of the segment is r.
- ◇ Finally, if w lies in the remaining curvilinear triangle $\tilde{z}\tilde{z}_x\tilde{z}_y$, set $F(w) = \dot{z}$.

By construction, F satisfies the conditions of the lemma. □

Proof of 8.4.5. Construct model triangles $[\dot{p}\dot{x}\dot{z}] = \tilde{\triangle}^{\kappa}(pxz)$ and $[\dot{p}\dot{y}\dot{z}] = \tilde{\triangle}^{\kappa}(pyz)$ so that $\dot{x}$ and $\dot{y}$ lie on opposite sides of $[\dot{p}\dot{z}]$.

Suppose

$$\measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) < \pi.$$



Then for some point $\dot{w} \in [\dot{p}\dot{z}]$, we have

$$|\dot{x} - \dot{w}| + |\dot{w} - \dot{y}| < |\dot{x} - \dot{z}| + |\dot{z} - \dot{y}| = |x - y|.$$

Let $w \in [pz]$ correspond to $\dot{w}$; that is, $|z - w| = |\dot{z} - \dot{w}|$. Since $[pxz]$ and $[pyz]$ are κ-thin, we have

$$|x - w| + |w - y| < |x - y|,$$

contradicting the triangle inequality.

Thus

$$\measuredangle^{\kappa}(z\,{}^{p}_{x}) + \measuredangle^{\kappa}(z\,{}^{p}_{y}) \geqslant \pi.$$

By Alexandrov's lemma (5.2.1), this is equivalent to

$$\measuredangle^{\kappa}(x\,{}^{p}_{z}) \leqslant \measuredangle^{\kappa}(x\,{}^{p}_{y}). \qquad ❶$$

Let $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^{\kappa}(pxy)$ and $\tilde{z} \in [\tilde{x}\tilde{y}]$ correspond to z; that is, $|x - z| = |\tilde{x} - \tilde{z}|$. Inequality ❶ is equivalent to $|p - z| \leqslant |\tilde{p} - \tilde{z}|$. Hence Lemma 8.4.6 applies. Therefore there is a short map F that sends $[\tilde{p}\tilde{x}\tilde{y}]$ to $\dot{D} = \operatorname{Conv}[\dot{p}\dot{x}\dot{z}] \cup \operatorname{Conv}[\dot{p}\dot{y}\dot{z}]$ in such a way that $\tilde{p} \mapsto \dot{p}$, $\tilde{x} \mapsto \dot{x}$, $\tilde{z} \mapsto \dot{z}$ and $\tilde{y} \mapsto \dot{y}$.

By assumption, the natural maps $[\dot{p}\dot{x}\dot{z}] \to [pxz]$ and $[\dot{p}\dot{y}\dot{z}] \to [pyz]$ are short. By composition, the natural map from $[\tilde{p}\tilde{x}\tilde{y}]$ to $[pyz]$ is short, as claimed. □

8.5 Function comparison

In this section, we give analytic and geometric ways of viewing the point-on-side comparison (8.3.1*b*) as a convexity condition.

First we obtain a corresponding differential inequality for the distance function in $\mathcal{U}$. In particular, a geodesic space $\mathcal{U}$ is CAT(0) if and only if for any $p \in \mathcal{U}$, the function $\operatorname{dist}_p^2 : \mathcal{U} \to \mathbb{R}$ is 2-convex; see Section 4.6 for the definition.

8.5.1. Theorem. *Suppose $\mathcal{U}$ is a ϖ^κ-geodesic space. Then the following are equivalent:*

a) $\mathcal{U}$ is CAT(κ),

b) for any $p \in \mathcal{U}$, the function $f = \operatorname{md}^\kappa \circ \operatorname{dist}_p$ satisfies

$$f'' + \kappa \cdot f \geqslant 1$$

in $\mathrm{B}(p, \varpi^\kappa)$.

Proof. Fix a sufficiently short geodesic $[xy]$ in $\mathrm{B}(p, \varpi^\kappa)$. We can assume that the model triangle $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\triangle}^\kappa(pxy)$ is defined. Set

$$\tilde{r}(t) = |\tilde{p} - \operatorname{geod}_{[\tilde{x}\tilde{y}]}(t)|, \qquad r(t) = |p - \operatorname{geod}_{[xy]}(t)|.$$

Set $\tilde{f} = \operatorname{md}^\kappa \circ \tilde{r}$ and $f = \operatorname{md}^\kappa \circ r$. By Property 1.1.1*a*, we have $\tilde{f}'' = 1 - \kappa \cdot \tilde{f}$. Clearly $\tilde{f}(t)$ and $f(t)$ agree at $t = 0$ and $t = |x - y|$. The point-on-side comparison (8.3.1*b*) is the condition $r(t) \leqslant \tilde{r}(t)$ for all $t \in [0, |x - y|]$. Since md^κ is increasing on $[0, \varpi^\kappa)$, $r \leqslant \tilde{r}$ and $f \leqslant \tilde{f}$ are equivalent. Thus the claim follows by Jensen's inequality (4.5.3*d*). □

Recall that Busemann functions defined in Proposition 5.1.1. The following exercise is analogous to Exercise 7.4.2.

8.5.2. Exercise. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space and* $\operatorname{bus}_\gamma : \mathcal{U} \to \mathbb{R}$ *be the Busemann function for a ray $\gamma\colon [0, \infty) \to \mathcal{L}$.*

a) If $\kappa = 0$, then the Busemann function $\operatorname{bus}_\gamma$ *is convex.*

b) If $\kappa = -1$, then the function

$$f = \exp \circ \operatorname{bus}_\gamma$$

satisfies

$$f'' - f \geqslant 0.$$

8.6 Convex sets

The following is a corollary of Theorem 8.5.1.

8.6.1. Corollary. *Suppose $\mathcal{U}$ is a ϖ^κ-geodesic* CAT(κ) *space. Then any ball (close or open) of radius $R < \frac{\varpi^\kappa}{2}$ in $\mathcal{U}$ is convex.*

Moreover, any open ball of radius $\frac{\varpi^\kappa}{2}$ is convex and any closed ball of radius $\frac{\varpi^\kappa}{2}$ is ϖ^κ-convex.

Proof. Suppose $p \in \mathcal{U}$, $R \leqslant \varpi^\kappa/2$, and two points x and y lie in $\overline{\mathrm{B}}[p,R]$ or $\mathrm{B}(p,R)$. By the triangle inequality, if $|x-y| < \varpi^\kappa$, then any geodesic $[xy]$ lies in $\mathrm{B}(p,\varpi^\kappa)$.

By function comparison (8.5.1), the geodesic $[xy]$ lies in $\overline{\mathrm{B}}[p,R]$ or $\mathrm{B}(p,R)$ respectively.

Thus any ball (close or open) of radius $R < \frac{\varpi^\kappa}{2}$ is ϖ^κ-convex. This implies convexity unless there is a pair of points in the ball at distance at least ϖ^κ. By the triangle inequality, the latter is possible only for the closed ball of radius $\frac{\varpi^\kappa}{2}$. □

8.6.2. Proposition. *Any weakly ϖ^κ-convex set in a complete length* CAT(κ) *space is ϖ^κ-convex.*

Proof. Follows from the uniqueness of geodesics in CAT(κ) spaces (8.2.1). □

8.6.3. Closest-point projection lemma. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space and $K \subset \mathcal{U}$ be a closed ϖ^κ-convex set. Assume that $\operatorname{dist}_K p < \frac{\varpi^\kappa}{2}$ for some point $p \in \mathcal{U}$. Then there is unique point $p^* \in K$ that minimizes the distance to p; that is, $|p^* - p| = \operatorname{dist}_K p$.*

Proof. Fix r properly between $\operatorname{dist}_K p$ and $\frac{\varpi^\kappa}{2}$. By function comparison (8.5.1) the function $f = \mathrm{md}^\kappa \circ \operatorname{dist}_p$ is strongly convex in $\overline{\mathrm{B}}[p,r]$.

The lemma follows from Lemma 12.1.4 applied to the subspace $K' = K \cap \overline{\mathrm{B}}[p,r]$ and the restriction $f|_{K'}$. □

8.6.4. Exercise. *Let $\mathcal{U}$ be a complete length* CAT(0) *space and $K \subset \mathcal{U}$ be a closed convex set. Show that the closest-point projection $\mathcal{U} \to K$ is short.*

8.6.5. Advanced exercise. *Let $\mathcal{U}$ be a complete length* CAT(1) *space and $K \subset \mathcal{U}$ be a closed π-convex set. Assume $K \subset \overline{\mathrm{B}}[p,\frac{\pi}{2}]$ for some point $p \in \mathcal{U}$. Show that there is a short retraction of $\mathcal{U}$ to K.*

8.6.6. Lemma. *Let $\mathcal{U}$ be a ϖ^κ-geodesic* CAT(κ) *space and $K \subset \mathcal{U}$ be a closed ϖ^κ-convex set. Set*

$$f = \mathrm{sn}^\kappa \circ \mathrm{dist}_K\,.$$

Then

$$f'' + \kappa\cdot f \geqslant 0$$

holds in $\mathrm{B}(K, \frac{\varpi^\kappa}{2})$.

Proof. It is sufficient to show that Jensen's inequality (4.5.3d) holds on a sufficiently short geodsic $[pq]$ in $\mathrm{B}(K, \frac{\varpi^\kappa}{2})$. Since

$$\mathrm{dist}_K\, p, \mathrm{dist}_K\, q < \tfrac{\varpi^\kappa}{2},$$

we may assume that

❶ $$|p-q| + \mathrm{dist}_K\, p + \mathrm{dist}_K\, q < \varpi^\kappa.$$

Namely, for each $x \in [pq]$, we need to find a value $h(x) \in \mathbb{R}$ such that for any x we have

$$h(x) \leqslant f(x), \qquad h(p) = f(p), \qquad h(q) = f(q),$$

and

❷ $$h'' + \kappa\cdot h \geqslant 0$$

along $[pq]$.

Denote by p^* and q^* the closest-point projections of p and q on K; they are provided by lemma 8.6.3. From ❶ and triangle inequality, we get

$$|p^* - q^*| < \varpi^\kappa.$$

Since K is ϖ^κ-convex, $K \supset [pq]$; in particular

$$\mathrm{dist}_K\, x \leqslant \mathrm{dist}_{[p^*q^*]}\, x$$

for any $x \in \mathcal{U}$.

Consider a majorization F for quadraliteral $[pp^*q^*q]$. By Majoriztion theorem 8.12.4 and Proposition 8.12.2, the figure F is a solid convex model quadraliteral $[\tilde p\tilde p^*\tilde q^*\tilde q]$ in $\mathbb{M}^2(\kappa)$ such that

$$|\tilde p - \tilde p^*|_{\mathbb{M}^2(\kappa)} = |p - p^*|_{\mathcal{U}} \qquad |\tilde p - \tilde q|_{\mathbb{M}^2(\kappa)} = |p - q|_{\mathcal{U}}$$
$$|\tilde q - \tilde q^*|_{\mathbb{M}^2(\kappa)} = |q - q^*|_{\mathcal{U}} \qquad |\tilde p^* - \tilde q^*|_{\mathbb{M}^2(\kappa)} = |p^* - q^*|_{\mathcal{U}}$$

Given $x \in [pq]$, denote by $\tilde{x}$ the corresponding point on $[\tilde{p}\tilde{q}]$. According to Majorization theorem,

$$|[pq] - x|_{\mathcal{U}} \leqslant |[\tilde{p}\tilde{q}] - \tilde{x}|_{\mathbb{M}^2(\kappa)}$$

for any x. Set

$$h(x) = \operatorname{sn}^{\kappa}|[\tilde{p}\tilde{q}] - \tilde{x}|_{\mathbb{M}^2(\kappa)}.$$

By straghtforward calculations ❷ hods and hence the staement follows. □

8.6.7. Corollary. *Let $\mathcal{U}$ be a complete length* $\mathrm{CAT}(\kappa)$ *space and $K \subset \subset \mathcal{U}$ be a closed locally convex set. Then there is an open set $\Omega \supset K$ such that the function*

$$f = \operatorname{sn}^{\kappa} \circ \operatorname{dist}_K$$

satisfies

$$f'' + \kappa \cdot f \geqslant 0$$

in Ω.

Proof. Fix $p \in K$. By Corollary 8.6.1, all $\overline{\mathrm{B}}[p, r]$ is convex for all small $r > 0$.

Since K is locally convex there is $r_p > 0$ such that the intesection $K' = K \cap \mathrm{B}(p, r_p)$ is convex.

Note that

$$\operatorname{dist}_K x = \operatorname{dist}_{K'} x$$

for any $x \in \mathrm{B}(p, \frac{r_p}{2})$. Therefore the statement holds for

$$\Omega = \bigcup_{p \in K} \mathrm{B}(p, \tfrac{r_p}{2}).$$

□

8.6.8. Theorem. *Assume $\mathcal{U}$ is a complete length* $\mathrm{CAT}(\kappa)$ *space and $K \subset \mathcal{U}$ is a closed connected locally convex set. Assume $|x - y| < \varpi^{\kappa}$ for any $x, y \in K$. Then K is convex.*

In particular, if $\kappa \leqslant 0$, then any closed connected locally convex set in a $\mathcal{U}$ is convex.

The following proof is inspired by the answer of Sergei Ivanov to the question of Nathan Reading [25].

Proof. Since K is locally convex, it is locally path connected. Since K is connected and locally path connected it is path connected.

Fix two points $x, y \in K$. Let us connect x to y by a path $\alpha\colon [0,1] \to \to K$. Since $|x - \alpha(s)| < \varpi^{\kappa}$ for any s, Theorem 8.2.1 implies that the geodesic $[x\alpha(s)]$ uniquely defined and depends continuously on s.

Let $\Omega \supset K$ be the open set proveded by Corollary 8.6.7. If $[xy] = [x\alpha(1)]$ does not completely lie in K, then there is a value $s \in [0,1]$ such that $[x\alpha(s)]$ lies in Ω, but does not completely lie in K. By Corollary 8.6.7, the function $f = \mathrm{sn}^\kappa \circ \mathrm{dist}_K \mathcal{U}$ satisfies differential inequality

❸ $$f'' + \kappa \cdot f \geqslant 0$$

along $[x\alpha(s)]$.

Since

$$|x - \alpha(s)| < \varpi^\kappa, \qquad f(x) = f(\alpha(s)) = 0,$$

the barrier inequality (4.5.3*b*) implies that $f(z) \leqslant 0$ for $z \in [x\alpha(s)]$; that is $[x\alpha(s)] \subset K$, a contradiction. □

8.7 Development

Geometrically, the development construction (7.5.1) translates distance comparison into a local convexity statement for subsets of $\mathbb{M}^2(\kappa)$. Recall that a curve in $\mathbb{M}^2(\kappa)$ is *(locally) concave with respect to* p if (locally) its supergraph with respect to p is a convex subset of $\mathbb{M}^2(\kappa)$; see Definition 7.5.2.

8.7.1. Development criterion. *For a ϖ^κ-geodesic space $\mathcal{U}$, the following statements hold:*

a) For any $p \in \mathcal{U}$ and any geodesic $\gamma\colon [0,T] \to \mathrm{B}(p, \varpi^\kappa)$, suppose the κ-development $\tilde\gamma$ in $\mathbb{M}^2(\kappa)$ of γ with respect to p is locally concave. Then $\mathcal{U}$ is CAT(κ).

b) If $\mathcal{U}$ is CAT(κ)*, then for any $p \in \mathcal{U}$ and any geodesic $\gamma\colon [0,T] \to \mathcal{U}$ such that the triangle $[p\gamma(0)\gamma(T)]$ has perimeter $< 2\cdot\varpi^\kappa$, the κ-development $\tilde\gamma$ in $\mathbb{M}^2(\kappa)$ of γ with respect to p is concave.*

Proof. (a). Set $\gamma = \mathrm{geod}_{[xy]}$ and $T = |x - y|$. Let $\tilde\gamma\colon [0,T] \to \mathbb{M}^2(\kappa)$ be the concave κ-development based at $\tilde p$ of γ with respect to p. Let us show that the function

❶ $$t \mapsto \measuredangle^\kappa(x\,{}^p_{\gamma(t)})$$

is nondecreasing.

For a partition $0 = t^0 < t^1 < \ldots < t^n = T$, set

$$\tilde y^i = \tilde\gamma(t^i) \;\text{ and }\; \tau^i = |\tilde y^0 - \tilde y^1| + |\tilde y^1 - \tilde y^2| + \ldots + |\tilde y^{i-1} - \tilde y^i|.$$

Since $\tilde\gamma$ is locally concave, for a sufficiently fine partition the broken geodesic $\tilde y^0\tilde y^1\dots\tilde y^n$ is locally convex with respect to $\tilde p$. Alexandrov's lemma (5.2.1), applied inductively to pairs of triangles $\tilde\triangle^\kappa\{\tau^{i-1},|p-\tilde y^0|,|p-\tilde y^{i-1}|\}$ and $\tilde\triangle^\kappa\{|\tilde y^{i-1}-\tilde y^i|,|p-\tilde y^{i-1}|,|p-\tilde y^i|\}$, shows that the sequence $\tilde\measuredangle^\kappa\{|p-\tilde y^i|;|p-\tilde y^0|,\tau^i\}$ is nondecreasing.

Taking finer partitions and passing to the limit,

$$\max_i\{|\tau^i-t^i|\}\to 0,$$

we get ❶ and the point-on-side comparison (8.3.1*b*) follows.

(b). Consider a partition $0=t^0<t^1<\dots<t^n=T$, and set $x^i=\gamma(t^i)$. Construct a chain of model triangles $[\tilde p\tilde x^{i-1}\tilde x^i]=\tilde\triangle^\kappa(px^{i-1}x^i)$ with the direction of $[\tilde p\tilde x^i]$ turning counterclockwise as i grows. By angle comparison (8.3.1*c*),

$$❷\qquad \measuredangle[\tilde x^i\,{}^{\tilde x^{i-1}}_{\tilde p}]+\measuredangle[\tilde x^i\,{}^{\tilde x^{i+1}}_{\tilde p}]\geqslant\pi.$$

Since γ is a geodesic,

$$❸\qquad \operatorname{length}\gamma=\sum_{i=1}^n|x^{i-1}-x^i|\leqslant|p-x^0|+|p-x^n|.$$

By repeated application of Alexandrov's Lemma (5.2.1), and inequality ❸,

$$\sum_{i=1}^n\measuredangle[\tilde p\,{}^{\tilde x^{i-1}}_{\tilde x^i}]\leqslant\tilde\measuredangle^\kappa(p\,{}^{x^0}_{x^n})\leqslant\pi.$$

Then by ❷, the broken geodesic $\tilde p\tilde x^0\tilde x^1\dots\tilde x^n$ is concave with respect to $\tilde p$.

Note that the broken geodesics $\tilde x^0\tilde x^1\dots\tilde x^n$ approach the development of γ with respect to p. Since the broken geodesics is convex, its length converges to the length of γ. Hence the result. □

8.8 Patchwork globalization

If $\mathcal U$ is CAT(κ) space, then it is locally CAT(κ). The converse, does not hold even for complete length space. For example, $\mathbb S^1$ is locally isometric to $\mathbb R$, and so is locally CAT(0), but it is easy to find a quadruple of points in $\mathbb S^1$ that violates CAT(0) comparison.

The following theorem was essentially proved by Alexandrov [16, Satz 9]; it gives a global condition on geodesics that is necessary and sufficient for a locally CAT(κ) space to be globally CAT(κ). The proof use a thin-triangle decompositions, and the inheritance lemma (8.4.5).

8.8.1. Patchwork globalization theorem. *For a complete length space $\mathcal U$, the following two statements are equivalent:*

a) $\mathcal{U}$ *is* $\mathrm{CAT}(\kappa)$.
b) $\mathcal{U}$ *is locally* $\mathrm{CAT}(\kappa)$*; moreover, pairs of points in* $\mathcal{U}$ *at distance* $< \varpi^\kappa$ *are joined by unique geodesics, and these geodesics depend continuously on their endpoint pairs.*

Note that the implication $(a)\Rightarrow(b)$ follows from Theorem 8.2.1.

8.8.2. Corollary. *Let* $\mathcal{U}$ *be a complete length space and* $\Omega \subset \mathcal{U}$ *be an open locally* $\mathrm{CAT}(\kappa)$ *subset. Then for any point* $p \in \Omega$ *there is* $R > 0$ *such that* $\overline{\mathrm{B}}[p, R]$ *is a convex subset of* $\mathcal{U}$ *and* $\overline{\mathrm{B}}[p, R]$ *is* $\mathrm{CAT}(\kappa)$.

Proof. Fix $R > 0$ such that $\mathrm{CAT}(\kappa)$ comparison holds in $\mathrm{B}(p, R)$.

We may assume that $\mathrm{B}(p, R) \subset \Omega$ and $R < \varpi^{\mathrm{K}}$. The same argument as in the proof of theorem on uniqueness of geodesics (8.2.1) shows that any two points in $\overline{\mathrm{B}}[p, \frac{R}{2}]$ can be joint by a unique geodesic that depends continuously on the end points.

The same argument as in the proof of Corollary 8.6.1 shows that $\overline{\mathrm{B}}[p, \frac{R}{2}]$ is a convex set. Then $(b)\Rightarrow(a)$ of Patchwork globalization implies that $\overline{\mathrm{B}}[p, \frac{R}{2}]$ is $\mathrm{CAT}(\kappa)$. □

The proof of Patchwork globalization uses the following construction:

8.8.3. Definition (Line-of-sight map). *Let* p *be a point and* α *be a curve of finite length in a length space* $\mathcal{X}$*. Let* $\bar\alpha : [0,1] \to \mathcal{U}$ *be the constant-speed parameterization of* α*. If* $\gamma_t \colon [0,1] \to \mathcal{U}$ *is a geodesic from* p *to* $\bar\alpha(t)$*, we say*

$$[0,1] \times [0,1] \to \mathcal{U} \colon (t,s) \mapsto \gamma_t(s)$$

is a line-of-sight map from p to α.

Proof of 8.8.1. As was already noted, it only remains to prove $(b)\Rightarrow(a)$.

Let $[pxy]$ be a triangle of perimeter $< 2\cdot\varpi^\kappa$ in $\mathcal{U}$. According to propositions 8.4.3 and 8.3.4, it is sufficient to show the triangle $[pxy]$ is κ-thin.

Since pairs of points at distance $< \varpi^\kappa$ are joined by unique geodesics and these geodesics depend continuously on their endpoint pairs, there is a unique and continuous line-of-sight map (8.8.3) for $[xy]$ from p.



For a partition

$$0 = t^0 < t^1 < \ldots < t^N = 1,$$

set $x^{i,j} = \gamma_{t^i}(t^j)$. Since the line-of-sight map is continuous, we may assume each triangle $[x^{i,j}x^{i,j+1}x^{i+1,j+1}]$ and $[x^{i,j}x^{i+1,j}x^{i+1,j+1}]$ is κ-thin (see Proposition 8.4.3).

Now we show that the κ-thin property propagates to $[pxy]$ by repeated application of the inheritance lemma (8.4.5):

- First, for fixed i, sequentially applying the lemma shows that the triangles $[xx^{i,1}x^{i+1,2}]$, $[xx^{i,2}x^{i+1,2}]$, $[xx^{i,2}x^{i+1,3}]$, and so on are κ-thin.

In particular, for each i, the long triangle $[xx^{i,N}x^{i+1,N}]$ is κ-thin.

- Applying the lemma again shows that the triangles $[xx^{0,N}x^{2,N}]$, $[xx^{0,N}x^{3,N}]$, and so on are κ-thin.

In particular, $[pxy] = [px^{0,N}x^{N,N}]$ is κ-thin. □

8.9 Reshetnyak's gluing theorem

The following theorem was proved by Yurii Reshetnyak [24], assuming $\mathcal{U}_1$, $\mathcal{U}_2$ are proper and complete. In the following form the theorem appears in the book of Martin Bridson and André Haefliger.[26].

8.9.1. Reshetnyak's gluing theorem. *Suppose $\mathcal{U}_1$ and $\mathcal{U}_2$ are ϖ^κ-geodesic spaces with isometric complete ϖ^κ-convex sets $A_i \subset \mathcal{U}_i$. Let $\iota\colon A_1 \to A_2$ be an isometry. Let $\mathcal{W} = \mathcal{U}_1 \sqcup_\iota \mathcal{U}_2$; that is, $\mathcal{W}$ is the gluing of $\mathcal{U}_1$ and $\mathcal{U}_2$ along ι (see Section 2.5).*

Then:

a) Both canonical mappings $\jmath_i\colon \mathcal{U}_i \to \mathcal{W}$ are isometric and the images $\jmath_i(\mathcal{U}_i)$ are ϖ^κ-convex subsets in $\mathcal{W}$.

b) If $\mathcal{U}_1, \mathcal{U}_2$ are CAT(κ) *spaces, then $\mathcal{W}$ is a* CAT(κ) *space.*

Proof. Part (a) follows directly from ϖ^κ-convexity of A_i.

(b). According to (a), we can identify $\mathcal{U}_i$ with its image $\jmath_i(\mathcal{U}_i)$ in $\mathcal{W}$; in this way, the subsets $A_i \subset \mathcal{U}_i$ will be identified and denoted further by A. Thus $A = \mathcal{U}_1 \cap \mathcal{U}_2 \subset \mathcal{W}$, and A is ϖ^κ-convex in $\mathcal{W}$.

Part (b) can be reformulated as follows:

8.9.2. Reformulation of 8.9.1*b*. *Let $\mathcal{W}$ be a length space having two ϖ^κ-convex subsets $\mathcal{U}_1, \mathcal{U}_2 \subset \mathcal{W}$ such that $\mathcal{W} = \mathcal{U}_1 \cup \mathcal{U}_2$. Assume the subset $A = \mathcal{U}_1 \cap \mathcal{U}_2$ is complete and ϖ^κ-convex in $\mathcal{W}$, and $\mathcal{U}_1$, $\mathcal{U}_2$ are* CAT(κ) *spaces. Then $\mathcal{W}$ is a* CAT(κ) *space.*

❶ *If $\mathcal{W}$ is ϖ^κ-geodesic, then $\mathcal{W}$ is* CAT(κ).



Indeed, according to 8.4.3, it is sufficient to show that any triangle $[x^0x^1x^2]$ of perimeter $< 2\cdot\varpi^\kappa$ in $\mathcal{W}$ is κ-thin. This is obviously true if all three points x^0, x^1, x^2 lie in a single $\mathcal{U}_i$. Thus, without loss of generality, we may assume that $x^0 \in \mathcal{U}_1$ and $x^1, x^2 \in \mathcal{U}_2$.

Choose points $z^1, z^2 \in A = \mathcal{U}_1 \cap \mathcal{U}_2$ lying respectively on the sides $[x^0x^1], [x^0x^2]$. Note that all distances between any pair of points from x^0, x^1, x^2, z^1, z^2 are less than ϖ^κ. Therefore

- ⋄ triangle $[x^0z^1z^2]$ lies in $\mathcal{U}_1$,
- ⋄ both triangles $[x^1z^1z^2]$ and $[x^1z^2x^2]$ lie in $\mathcal{U}_2$.

In particular each triangle $[x^0z^1z^2]$, $[x^1z^1z^2]$, $[x^1z^2x^2]$ is κ-thin.

Applying the inheritance lemma for thin triangles (8.4.5) twice, we get that $[x^0x^1z^2]$ and consequently $[x^0x^1x^2]$ is κ-thin. △

❷ *$\mathcal{W}$ is geodesic if $\kappa \leqslant 0$.*

For $p_1 \in \mathcal{U}_1$, $p_2 \in \mathcal{U}_2$, we may choose a sequence $z^i \in A$ such that $|p_1 - z^i| + |p_2 - z^i|$ converges to $|p_1 - p_2|$, and $|p_1 - z^i|$ and $|p_2 - z^i|$ converge. Since A is complete, it suffices to show z^i is a Cauchy sequence. In that case, the limit point z of z^i satisfies $|p_1 - z| + |p_2 - z| = |p_1 - p_2|$, so the geodesics $[p_1z]$ in $\mathcal{U}_1$ and $[p_2z]$ in $\mathcal{U}_2$ together give a geodesic $[p_1p_2]$ in $\mathcal{U}$.

Suppose z^i is not a Cauchy sequence. Then there are subsequences x^i and y^i of z^i satisfying $\lim |x^i - y^i| > 0$. Let m^i be the midpoint of $[x^iy^i]$. Since $|p_1 - m^i| + |p_2 - m^i| \geqslant |p_1 - p_2|$, and $|p_1 - x^i| + |p_2 - x^i|$ and $|p_1 - y^i| + |p_2 - y^i|$ converge to $|p_1 - p_2|$, then for any $\varepsilon > 0$, we may assume (taking subsequences and possibly relabeling p_1 and p_2)

$$|p_1 - m^i| \geqslant |p_1 - x^i| - \varepsilon, |p_1 - m^i| \geqslant |p_1 - y^i| - \varepsilon.$$

Since triangle $[p^1x^iy^i]$ is thin, the analogous inequalities hold for the Euclidean model triangle $[\tilde{p}^1\tilde{x}^i\tilde{y}^i]$. Then there is a nondegenerate limit triangle $[pxy]$ in the Euclidean plane satisfying $|p - x| = |p - y| \leqslant |p - m|$ where m is the midpoint of $[xy]$. This contradiction proves the claim.

△

Finally suppose $\kappa > 0$; by scaling, take $\kappa = 1$. Consider the Euclidean cones $\operatorname{Cone}\mathcal{U}_i$ (see Section 5.4). By Theorem 10.4.2*a*, $\operatorname{Cone}\mathcal{U}_i$ is a CAT(0) space for $i = 1, 2$. Geodesics of $\operatorname{Cone}\mathcal{U}_i$ that do not pass through the vertex of the cone correspond, under projection from the vertex, to geodesics of $\mathcal{U}_i$ of length $< \pi$. It follows that $\operatorname{Cone} A$ is convex in $\operatorname{Cone}\mathcal{U}_1$ and $\operatorname{Cone}\mathcal{U}_2$. By the cone distance formula , $\operatorname{Cone} A$ is complete since A is complete. Gluing on $\operatorname{Cone} A$ and applying ❶ and

❷ for $\kappa = 0$, we find that $\operatorname{Cone}\mathcal{W}$ is a CAT(0) space. By Theorem 10.4.2a, $\mathcal{W}$ is a CAT(1) space. □

8.9.3. Exercise. *Let Q be the nonconvex subset of the plane bounded by two rays γ_1 and γ_2 with common starting point and angle α between them. Assume $\mathcal{U}$ is a complete length* CAT(0) *space and γ_1', γ_2' are two rays in $\mathcal{U}$ with common starting point and angle α between them. Show that the space glued from Q and $\mathcal{U}$ along the corresponding rays is a* CAT(0) *space.*

8.9.4. Exercise. *Assume $\mathcal{U}$ is complete length* CAT(0) *space and $A \subset \mathcal{U}$ is a closed subset. Assume that the doubling of $\mathcal{U}$ in A is* CAT(0)*. Show that A is a convex set of $\mathcal{U}$.*

8.9.5. Exercise. *Let $\mathcal{U}$ be a complete length* CAT(0) *space. Assume $\tilde{\mathcal{U}} \to \mathcal{U}$ is a metric covering branching along a geodesic. Show that $\tilde{\mathcal{U}}$ is* CAT(0)*.*

More generally, assume $A \subset \mathcal{U}$ is a closed convex subset and $f\colon \mathcal{X} \to \mathcal{U}\backslash A$ is a metric cover. Denote by $\bar{\mathcal{X}}$ the completion of $\mathcal{X}$, and $\bar{f}\colon \bar{\mathcal{X}} \to \mathcal{U}$ the continuous extension of f. Let $\tilde{\mathcal{U}}$ be the space glued from $\bar{\mathcal{X}}$ and A by identifying x and $\bar{f}(x)$ if $\bar{f}(x) \in A$. Show that $\tilde{\mathcal{U}}$ is CAT(0)*.*

8.10 Space of geodesics

In this section we prove a "no-conjugate-point" theorem for spaces with upper curvature bounds, and derive from it a number of statements about local geodesics. These statements will be used in the proof of the Hadamard–Cartan theorem (8.13.1) and Lifting globalization theorem (8.11.1), in much the same way as the exponential map is used in Riemannian geometry.

8.10.1. Proposition. *Let $\mathcal{U}$ be a locally* CAT(κ) *space. Let $\gamma_n\colon [0,1] \to \mathcal{U}$ be a sequence of local geodesic paths converging to a path $\gamma_\infty\colon [0,1] \to \mathcal{U}$. Then γ_∞ is a local geodesic path. Moreover*

$$\operatorname{length}\gamma_n \to \operatorname{length}\gamma_\infty$$

as $n \to \infty$.

Proof. Fix $t \in [0,1]$. By Corollary 8.8.2, we may choose R satisfying $0 < R < \varpi^{\mathrm{K}}$, and such that the ball $\mathcal{B} = \mathrm{B}(\gamma_\infty(t), R)$ is a convex subset of $\mathcal{U}$ and forms a CAT(κ) space.

A local geodesic segment with length less than $R/2$ that intersects $\mathrm{B}(\gamma_\infty(t), R/2)$, cannot leave $\mathcal{B}$ and hence is minimizing by Corollary 8.4.4. In particular, for all sufficiently large n, any subsegment of γ_n through $\gamma_n(t)$ with length $< R/2$ is a geodesic.

Since $\mathcal{B}$ is $\mathrm{CAT}(\kappa)$, geodesic segments in $\mathcal{B}$ depend uniquely and continuously on their endpoint pairs by Theorem 8.2.1. Thus there is a subinterval $\mathbb{I}$ of $[0,1]$, that contains a neighborhood of t in $[0,1]$ and such that $\gamma_n|_{\mathbb{I}}$ is minimizing for all large n. It follows that the restriction $\gamma_\infty|_{\mathbb{I}}$ is a geodesic, and therefore γ_∞ is a local geodesic. $\square$

The following theorem was proved by the first author and Richard Bishop [27]. In analogy with Riemannian geometry, the main statement of the following theorem could be restated as: *In a space of curvature $\leqslant \kappa$, two points cannot be conjugate along a local geodesic of length $< \varpi^\kappa$.*

8.10.2. No-conjugate-point theorem. *Suppose $\mathcal{U}$ is a locally complete, length, locally* $\mathrm{CAT}(\kappa)$ *space. Let $\gamma\colon [0,1] \to \mathcal{U}$ be a local geodesic path with length $< \varpi^\kappa$. Then for some neighborhoods $\Omega^0 \ni \gamma(0)$ and $\Omega^1 \ni \gamma(1)$ there is a unique continuous map $\Omega^0 \times \Omega^1 \times \times [0,1] \to \mathcal{U}$*

$$(x,y,t) \mapsto \gamma_{xy}(t)$$

such that $\gamma_{xy}\colon [0,1] \to \mathcal{U}$ is a local geodesic path with $\gamma_{xy}(0) = x$ and $\gamma_{xy}(1) = y$ for each $(x,y) \in \Omega^0 \times \Omega^1$, and the family γ_{xy} contains γ. Moreover, we can assume that the map

$$(x,y,t) \mapsto \gamma_{xy}(t)\colon \Omega^0 \times \Omega^1 \times [0,1] \to \mathcal{U}$$

is $\pounds$-Lipschitz for any $\pounds > \max\left\{ \frac{\mathrm{sn}^\kappa r}{\mathrm{sn}^\kappa \ell} : 0 \leqslant r \leqslant \ell \right\}$.

The following lemma was suggested to us by Alexander Lytchak. The proof proceeds by piecing together $\mathrm{CAT}(\kappa)$ neighborhoods of points on a curve to construct a new $\mathrm{CAT}(\kappa)$ space. Exercise 8.13.7 is inspired by the original idea of the proof of No-conjugate-point theorem (8.10.2) given in [27].

8.10.3. Patchwork along a curve. *Let $\mathcal{U}$ be a locally complete, length, locally* $\mathrm{CAT}(\kappa)$ *space, and $\alpha\colon [a,b] \to \mathcal{U}$ be a curve.*

Then there is a complete length $\mathrm{CAT}(\kappa)$ *space $\mathcal{N}$ with an open set $\hat\Omega \subset \mathcal{N}$, a curve $\hat\alpha\colon [0,1] \to \hat\Omega$, and an open locally isometric immersion $\Phi\colon \hat\Omega \looparrowright \mathcal{U}$ such that $\Phi \circ \hat\alpha = \alpha$.*

Moreover if α is simple, then one can assume in addition that Φ is an open embedding; thus $\hat\Omega$ is locally isometric to a neighborhood of $\Omega = \Phi(\hat\Omega)$ of α.

Proof. According to Corollary 8.8.2, for any $t \in [a,b]$ there is $r(t) > 0$ such that the closed ball $\overline{\mathrm{B}}[\alpha(t), r(t)]$ is a convex set that forms a complete length $\mathrm{CAT}(\kappa)$ space.

Choose balls $\mathcal{B}^i = \overline{\mathrm{B}}[\alpha(t^i), r(t^i)]$ for some partition $a = t^0 < t^1 < \ldots < t^n = b$ in such a way that $\operatorname{Int}\mathcal{B}^i \supset \alpha([t^{i-1}, t^i])$ for all $i > 0$.

Consider the disjoint union $\bigsqcup_i \mathcal{B}^i = \left\{ (i,x) : x \in \mathcal{B}^i \right\}$ with the minimal equivalence relation $\sim$ such that $(i,x) \sim (i-1,x)$ for all $i > 0$. Let $\mathcal{N}$ be the space obtained by gluing the $\mathcal{B}^i$ along $\sim$. Note that $A^i = \mathcal{B}^i \cap \mathcal{B}^{i-1}$ is convex in $\mathcal{B}^i$ and in $\mathcal{B}^{i-1}$. Applying the Reshetnyak gluing theorem (8.9.1) n times, we conclude that $\mathcal{N}$ is a complete length $\mathrm{CAT}(\kappa)$ space.



For $t \in [t^{i-1}, t^i]$, let $\hat\alpha(t)$ be the equivalence class of $(i, \alpha(t))$ in $\mathcal{N}$. Let $\hat\Omega$ be the ε-neighborhood of $\hat\alpha$ in $\mathcal{N}$, where $\varepsilon > 0$ is chosen so that $\mathrm{B}(\alpha(t), \varepsilon) \subset \mathcal{B}^i$ for all $t \in [t^{i-1}, t^i]$.

Define $\Phi\colon \hat\Omega \to \mathcal{U}$ by sending the equivalence class of (i,x) to x. It is straightforward to check that $\Phi\colon \mathcal{N} \to \mathcal{U}$, $\hat\alpha\colon [0,1] \to \mathcal{N}$ and $\hat\Omega \subset \mathcal{N}$ satisfy the conclusion of the main part of the lemma.

To prove the final statement in the lemma, we only have to choose $\varepsilon > 0$ so that in addition $|\alpha(\tau) - \alpha(\tau')| > 2\cdot\varepsilon$ if $\tau \leqslant t^{i-1}$ and $t^i \leqslant \tau'$ for some i. □

Proof of 8.10.2. Apply patchwork along γ (8.10.3). □

The No-conjugate-point theorem (8.10.2) allows us to move a local geodesic path so that its endpoints follow given trajectories. The following corollary describe how this process might terminate.

8.10.4. Corollary. *Let $\mathcal{U}$ be a locally complete, length, locally* $\mathrm{CAT}(\kappa)$ *space. Suppose $\gamma\colon [0,1] \to \mathcal{U}$ is a local geodesic path with length $< \varpi^\kappa$. Let $\alpha^i\colon [0,1] \to \mathcal{U}$, for $i = 0, 1$, be paths starting at $\gamma(0)$ and $\gamma(1)$ respectively.*

Then there is a uniquely determined pair consisting of an interval $\mathbb{I}$ satisfying $0 \in \mathbb{I} \subset [0,1]$, and a continuous family of local geodesic paths $\gamma_t\colon [0,1] \to \mathcal{U}$ for $t \in \mathbb{I}$, such that

a) $\gamma_0 = \gamma$, $\gamma_t(0) = \alpha^0(t)$, $\gamma_t(1) = \alpha^1(t)$, and γ_t has length $< \varpi^\kappa$,

b) if $\mathbb{I} \neq [0,1]$, then $\mathbb{I} = [0,a)$, where either γ_t converges uniformly to a local geodesic γ_a of length ϖ^κ, or for some fixed $s \in [0,1]$ the curve $\gamma_t(s) : [0,a) \to \mathcal{U}$ is a Lipschitz curve with no limit as $t \to a-$.

Proof. Uniqueness follows from Theorem 8.10.2.

Let $\mathbb{I}$ be the maximal interval for which there is a family γ_t satisfying condition (a). By Theorem 8.10.2, such an interval exists and is open in $[0,1]$. Suppose $\mathbb{I} \neq [0,1]$. Then $\mathbb{I} = [0,a)$ for some $0 < a \leqslant 1$. It suffices to show that $\mathbb{I}$ satisfies condition (b).

For each fixed $s \in [0,1]$, define the curve $\alpha_s : [0,a) \to \mathcal{U}$ by $\alpha_s(t) = \gamma_t(s)$. By Theorem 8.10.2, α_s is $\pounds$-Lipschitz for some $\pounds$.

If α_s for some value of s does not converge as $t \to a-$, then condiion (b) holds. If each α_s converges as $t \to a-$, then γ_t converges as $t \to$ $\to a-$, say to γ_a. By Proposition 8.10.1, γ_a is a local geodesic path and

$$\operatorname{length}\gamma_t \to \operatorname{length}\gamma_a \leqslant \varpi^\kappa.$$

By maximality of $\mathbb{I}$, $\operatorname{length}\gamma_a = \varpi^\kappa$ and so condition (b) again holds. □

8.10.5. Corollary. *Let $\mathcal{U}$ be a complete locally* CAT(κ) *length space, $\alpha\colon [0,1] \to \mathcal{U}$ be a path that starts at p and ends at q. Then*

a) There is a unique homotopy of local geodesic paths $\gamma_t\colon [0,1] \to$ $\to \mathcal{U}$ such that $\gamma_0(t) = \gamma_t(0) = p$ and $\gamma_t(1) = \alpha(t)$ for any t;

b) We have that

$$\operatorname{length}\gamma_t \leqslant \operatorname{length}(\alpha|_{[0,t]}),$$

for any $t \in [0,1]$, and equality holds for given t if and only if the restriction $\alpha|_{[0,t]}$ is a reparametrization of γ_t.

Moreover instead of completeness of $\mathcal{U}$ one can assume that the set

$$W = \{\, x \in \mathcal{U} : |x-p| + |x-q| \leqslant \ell \,\}$$

is complete.

Proof. Apply Corollary 8.10.4, setting $\alpha^0(t) = \alpha(0)$ for all $t \in [0,1]$, and $\alpha^1 = \alpha$. By (a), there is an interval $\mathbb{I}$ such that statement holds for all $t \in \mathbb{I}$, and either $\mathbb{I} = [0,1]$ or $\mathbb{I} = [0,a)$ for some $a \leqslant 1$.

By the patchwork along a curve (8.10.3), the values of t for which condition (b) holds form an open subset of $\mathbb{I}$ containing 0; clearly this subset is also closed in $\mathbb{I}$. Therefore (b) holds on all of $\mathbb{I}$.

Since $\ell < \varpi^\kappa$, Corollary 8.10.4 implies that $\mathbb{I} = [0,1]$. Indeed if $\mathbb{I} = [0,a)$, then either $\operatorname{length}\gamma_t \to \varpi^\kappa$ as $t \to a-$ or for some fixed $s \in$ $\in [0,1]$ the Lipschitz curve $\gamma_t(s) : [0,a) \to \mathcal{U}$ with no limit as $t \to a-$. Since $\ell < \varpi^\kappa$, the condition (b) implies that both of these scenarios are not possible. □

The no-conjugate-point theorem (8.10.2) gives the following extension of the first variation formula (8.14.3) from geodesics in CAT(κ) spaces to local geodesics in locally CAT(κ) spaces.

8.10.6. Corollary. *Suppose $\mathcal{U}$ is a locally complete locally* CAT(κ) *space. For any geodesic $[px]$ in $\mathcal{U}$ and local geodesic $\gamma\colon [0,1] \to \mathcal{U}$ from p to q of length $< \varpi^\kappa$, let $\gamma_t\colon [0,1] \to \mathcal{U}$ be a continuous family of local geodesic paths with $\gamma_0 = \gamma$, $\gamma_t(0) = \operatorname{geod}_{[px]}(t)$ and $\gamma_t(1) = q$. Then*

$$\operatorname{length}(\gamma_t) = \operatorname{length}(\gamma) - t\cdot \cos\measuredangle(\gamma,[px]) + o(t).$$

8.11 Lifting globalization

The Hadamard–Cartan theorem (8.13.1) states that the universal metric cover of a complete space locally CAT(0) space is CAT(0). The lifting globalization theorem gives an appropriate generalization of the above statement to arbitrary curvature bounds; it could be also described as a global version of Gauss's lemma.

8.11.1. Lifting globalization theorem. *Suppose $\mathcal{U}$ is a complete length locally* CAT(κ) *space and $p \in \mathcal{U}$. Then there is a complete length* CAT(κ) *space $\mathcal{B}$, with a point $\hat p$ such that $|\hat p - \hat x|_{\mathcal{B}} \leqslant \frac{\varpi^\kappa}{2}$ for any $\hat x \in \mathcal{B}$ and there is a locally isometric map $\Phi\colon \mathcal{B} \to \mathcal{U}$ such that $\Phi(\hat p) = p$ and the following lifting property holds: for any path $\alpha\colon [0,1] \to \mathcal{U}$ with $\alpha(0) = p$ and $\operatorname{length}\alpha < \varpi^\kappa/2$, there is a unique path $\hat\alpha\colon [0,1] \to \mathcal{B}$ such that $\hat\alpha(0) = \hat p$ and $\Phi\circ\hat\alpha \equiv \alpha$.*

Before proving the theorem we state and prove its corollary.

8.11.2. Corollary. *Suppose $\mathcal{U}$ is a complete length locally* CAT(κ) *space. Then for any $p \in \mathcal{U}$ there is $\rho_p > 0$ such that $\overline{\mathrm{B}}[p,\rho_p]$ is a complete length* CAT(κ) *space.*

Moreover, we can assume that $\rho_p < \frac{\varpi^\kappa}{2}$ for any p and the function $p \mapsto \rho_p$ is 1-Lipschitz.

Proof. Assume $\Phi\colon \mathcal{B} \to \mathcal{U}$ and $\hat p \in \mathcal{B}$ are provided by Lifting globalization theorem (8.11.1).

Since Φ is local isometry, we can choose $r > 0$ so that the restriction of Φ to $\overline{\mathrm{B}}[\hat p, r]$ is distance preserving. By Lifting globalization, the

image $\Phi(\overline{\mathrm{B}}[\hat{p},r])$ coincides with the ball $\overline{\mathrm{B}}[p,r]$. This proves the first part of theorem.

To prove the second part, lat us choose ρ_p to be maximal value $\leqslant \frac{\varpi^\kappa}{2}$ such that $\overline{\mathrm{B}}[p,\rho_p]$ is a complete length $\mathrm{CAT}(\kappa)$ space. By Corollary 8.6.1, the ball

$$\overline{\mathrm{B}}[q,\rho_p-|p-q|]$$

is convex in $\overline{\mathrm{B}}[p,\rho_p]$. Therefore

$$\overline{\mathrm{B}}[q,\rho_p-|p-q|]$$

is a complete length $\mathrm{CAT}(\kappa)$ space for any $q\in\mathrm{B}(p,\rho_p)$. In particular $\rho_q\geqslant\rho_p-|p-q|$ for any $p,q\in\mathcal{U}$. Hence the second statement follows. □

The proof of the lifting globalization theorem relies heavily on the properties of the space of local geodesic paths discussed in Section 8.10. The following lemma proved by the first author and Richard Bishop [28] is a key step in the proof.

8.11.3. Radial lemma. *Let $\mathcal{U}$ be a length locally* $\mathrm{CAT}(\kappa)$ *space, and suppose $p\in\mathcal{U}$, $R\leqslant\varpi^\kappa$. Assume the ball $\overline{\mathrm{B}}[p,\bar{R}]$ is complete for any $\bar{R}<R$, and there is a unique geodesic path,* $\mathrm{path}_{[px]}$*, from p to any point $x\in\mathrm{B}(p,R)$, that depends continuously on x. Then $\mathrm{B}(p,\frac{R}{2})$ is a ϖ^κ-geodesic* $\mathrm{CAT}(\kappa)$ *space.*

Proof. Without loss of generality, we may assume $\mathcal{U}=\mathrm{B}(p,R)$.

Set $f=\mathrm{md}^\kappa\circ\mathrm{dist}_p$. Let us show that

❶ $$f''+\kappa\cdot f\geqslant 1.$$

Fix $z\in\mathcal{U}$. We will apply Theorem 8.10.2 for the unique geodesic path γ from p to z. The notations Ω^0, Ω^1, γ_{xy}, $\mathcal{N}$, $\hat{x}$, $\hat{y}$ will be as in the theorem; in particular, $z\in\Omega^1$.

By assumption, $\gamma_{py}=\mathrm{path}_{[py]}$ for any $y\in\Omega^1$. Consequently, $f(y)=\mathrm{md}^\kappa|\hat{p}-\hat{y}|_{\mathcal{N}}$. Applying the function comparison (8.5.1) in $\mathcal{N}$, we have that $f''+\kappa\cdot f\geqslant 1$ in Ω^1; whence ❶ follows.

Fix $r<\frac{R}{2}$; the proof of the following claim takes most of the remaining proof.

❷ $\overline{\mathrm{B}}[p,r]$ *is a convex set in $\mathcal{U}$.*

Choose arbitrary $x,z\in\overline{\mathrm{B}}[p,r]$. First note that ❶ implies the following claim.

❸ *If $\gamma\colon[0,1]\to\mathcal{U}$ is a local geodesic path from x to z and* $\mathrm{length}\,\gamma<$ $<\varpi^\kappa$*, then* $\mathrm{length}\,\gamma\leqslant 2\cdot r$ *and γ lies completely in* $\overline{\mathrm{B}}[p,r]$.

Note that $|x - z| < \varpi^\kappa$. Thus, to prove Claim ❷, it is sufficient to show that there is a geodesic path from x to z. Note that by assumption $\overline{\mathrm{B}}[p, 2{\cdot}r]$ is complete; therefore Corollary 8.10.5 implies the following.

❹ *Given a path* $\alpha\colon [0,1] \to \mathcal{U}$ *from* x *to* z *with* $\operatorname{length}\alpha < 2{\cdot}r$, *there is a local geodesic path* γ *from* x *to* z *such that*

$$\operatorname{length}\gamma \leqslant \operatorname{length}\alpha.$$

Further, let us prove the following.

❺ *There is unique local geodesic path* γ_{xz} *in* $\overline{\mathrm{B}}[p,r]$ *from* x *to* z.

Denote by Δ_{xz} the set of all local geodesic paths in $\overline{\mathrm{B}}[p,r]$ from x to z. By Corollary 8.10.4, there is a bijection $\Delta_{xz} \to \Delta_{pp}$. According to ❶, Δ_{pp} contains only the constant path. Claim ❺ follows.

Note that claims ❸, ❹ and ❺ imply that γ_{xz} is minimizing; hence Claim ❷.

Further, Claim ❸ and the no-conjugate-point theorem (8.10.2) together imply that the map $(x,z) \mapsto \gamma_{xz}$ is continuous.

Therefore by the patchwork globalization theorem (8.8.1), $\overline{\mathrm{B}}[p,r]$ is a ϖ^κ-geodesic $\mathrm{CAT}(\kappa)$ space.

Since

$$\mathrm{B}(p,R) = \bigcup_{r<R} \overline{\mathrm{B}}[p,r],$$

then $\mathrm{B}(p,R)$ is convex in $\mathcal{U}$ and $\mathrm{CAT}(\kappa)$ comparison holds for any quadruple in $\mathrm{B}(p,R)$. Therefore $\mathrm{B}(p,\varpi^\kappa/2)$ is $\mathrm{CAT}(\kappa)$. □

In the following proof, we construct a space $\mathfrak{G}_p$ of local geodesic paths that start at p. The space $\mathfrak{G}_p$ comes with a marked point $\hat{p}$ and the endpoint map $\Phi\colon \mathfrak{G}_p \to \mathcal{U}$. One can think of the map Φ as an analog of $\exp_p$ in the Riemannian case; in this case the space $\mathfrak{G}_p$ is the ball of radius ϖ^κ in the tangent space at p, equipped with the metric pulled back by $\exp_p$.

We are going to set $\mathcal{B} = \mathrm{B}(\hat{p},\varpi^\kappa/2) \subset \mathfrak{G}_p$, and use the radial lemma (8.11.3) to prove that $\mathcal{B}$ is a ϖ^κ-geodesic $\mathrm{CAT}(\kappa)$ space.

Proof of 8.11.1. Suppose $\hat{\gamma}$ is a homotopy of local geodesic paths that start at p. Thus the map

$$\hat{\gamma}\colon (t,\tau) \mapsto \hat{\gamma}_t(\tau)\colon [0,1]\times[0,1] \to \mathcal{U}$$

is continuous, and the following holds for each t:

- ⋄ $\hat{\gamma}_t(0) = p$,
- ⋄ $\hat{\gamma}_t\colon [0,1] \to \mathcal{U}$ is a local geodesic path in $\mathcal{U}$.

Let $\vartheta(\hat\gamma)$ be the length tranced by the ends of $\hat\gamma_t$; that is, $\vartheta(\hat\gamma)$ is the length of the path $t\mapsto\hat\gamma_t(1)$.

Let $\mathfrak{G}_p$ be the set of all local geodesic paths with length $<\varpi^\kappa$ in $\mathcal{U}$ that start at p. Denote by $\hat p\in\mathfrak{G}_p$ the constant path $\hat p(t)\equiv p$. Given $\alpha,\beta\in\mathfrak{G}_p$ define

$$|\alpha-\beta|_{\mathfrak{G}_p}=\inf_{\hat\gamma}\{\vartheta(\hat\gamma)\},$$

with infimum taken along all homotopies $\hat\gamma\colon[0,1]\times[0,1]\to\mathcal{U}$ such that $\hat\gamma_0=\alpha$, $\hat\gamma_1=\beta$ and $\hat\gamma_t\in\mathfrak{G}_p$ for all $t\in[0,1]$.

From Theorem 8.10.2, we have $|\alpha-\beta|_{\mathfrak{G}_p}>0$ for distinct α and β; that is,

❻ $|*-*|_{\mathfrak{G}_p}$ *is a metric on* $\mathfrak{G}_p$.

Further, again from Theorem 8.10.2, we have

❼ *The map*

$$\Phi\colon\xi\mapsto\xi(1)\colon\mathfrak{G}_p\to\mathcal{U}$$

is a local isometry. In particular, $\mathfrak{G}_p$ *is locally* $\mathrm{CAT}(\kappa)$.

Let $\alpha\colon[0,1]\to\mathcal{U}$ be a path with $\operatorname{length}\alpha<\varpi^\kappa$ and $\alpha(0)=p$. The homotopy constructed in Corollary 8.10.5 can be regarded as a path in $\mathfrak{G}_p$, say $\hat\alpha\colon[0,1]\to\mathfrak{G}_p$, such that $\hat\alpha(0)=\hat p$ and $\Phi\circ\hat\alpha=\alpha$; in particular $\hat\alpha_t(1)\equiv\alpha(t)$. By ❼,

$$\operatorname{length}(\hat\alpha)_{\mathfrak{G}_p}=\operatorname{length}(\alpha)_{\mathcal{U}}.$$

Moreover, it follows that α is a local geodesic path of $\mathcal{U}$ if and only if $\hat\alpha$ is a local geodesic path of $\mathfrak{G}_p$.

Further, from Corollary 8.10.5, for any $\xi\in\mathfrak{G}_p$ and path $\hat\alpha\colon[0,1]\to\to\mathfrak{G}_p$ from $\hat p$ to ξ, we have

$$\begin{aligned}\operatorname{length}\hat\alpha=\operatorname{length}\Phi\circ\hat\alpha&\geqslant\\ \geqslant\operatorname{length}\xi&=\operatorname{length}\hat\xi\end{aligned}$$

where equality holds only if $\hat\alpha$ is a reparametrization of $\hat\xi$. In particular,

❽ $$|\hat p-\xi|_{\mathfrak{G}_p}=\operatorname{length}\xi$$

and $\hat\xi\colon[0,1]\to\mathfrak{G}_p$ is the unique geodesic path from $\hat p$ to ξ. Clearly, the map $\xi\mapsto\hat\xi$ is continuous.

From ❽ and Proposition 8.10.1, it follows

❾ *For any* $\bar R<\varpi^\kappa$, *the closed ball* $\overline{\mathrm{B}}[\hat p,\bar R]$ *in* $\mathfrak{G}_p$ *is complete.*

Take $\mathcal{B}=\mathrm{B}(\hat p,\varpi^\kappa/2)$ and Φ constructed above. According to the Radial lemma (8.11.3), $\mathcal{B}$ is a ϖ^κ-geodesic $\mathrm{CAT}(\kappa)$ space. The remaining statements are already proved. □

8.12 Reshetnyak majorization

8.12.1. Definition. *Let $\mathcal{X}$ be a metric space, $\tilde\alpha$ be a simple closed curve of finite length in $\mathbb{M}^2(\kappa)$, and $D \subset \mathbb{M}^2(\kappa)$ be a closed region bounded by $\tilde\alpha$. A length-nonincreasing map $F\colon D \to \mathcal{X}$ is called* majorizing *if it is length-preserving on $\tilde\alpha$.*

In this case, we say that D majorizes *the curve $\alpha = F \circ \tilde\alpha$ under the map F.*

The following proposition is a consequence of the definition.

8.12.2. Proposition. *Let α be a closed curve in a metric space $\mathcal{X}$. Suppose $D \subset \mathbb{M}^2(\kappa)$ majorizes α under $F\colon D \to \mathcal{X}$. Then any geodesic subarc of α is the image under F of a subarc of $\partial_{\mathbb{M}^2(\kappa)} D$ that is geodesic in the length-metric of D.*

In particular, if D is convex, then the corresponding subarc is a geodesic in $\mathbb{M}^2(\kappa)$.

Proof. For a geodesic subarc $\gamma\colon [a,b] \to \mathcal{X}$ of $\alpha = F \circ \tilde\alpha$, set

$$\begin{aligned} \tilde r &= |\tilde\gamma(a) - \tilde\gamma(b)|_D, & \tilde\gamma &= (F|_{\partial D})^{-1} \circ \gamma, \\ s &= \operatorname{length}\gamma, & \tilde s &= \operatorname{length}\tilde\gamma. \end{aligned}$$

Then

$$\tilde r \geqslant r = s = \tilde s \geqslant \tilde r.$$

Therefore $\tilde s = \tilde r$. □

8.12.3. Corollary. *Let $[pxy]$ be a triangle of perimeter $< 2\cdot\varpi^\kappa$ in a metric space $\mathcal{X}$. Assume a convex region $D \subset \mathbb{M}^2(\kappa)$ majorizes $[pxy]$. Then $D = \operatorname{Conv}[\tilde p\tilde x\tilde y]$ for a model triangle $[\tilde p\tilde x\tilde y] = \tilde\triangle^\kappa(pxy)$, and the majorizing map sends $\tilde p$, $\tilde x$ and $\tilde y$ respectively to p, x and y.*

Now we come to the main theorem of this section.

8.12.4. Majorization theorem. *Any closed curve α with length smaller than $2\cdot\varpi^\kappa$ in a ϖ^κ-geodesic* CAT(κ) *space is majorized by a convex region in $\mathbb{M}^2(\kappa)$.*

This theorem is proved by Yurii Reshetnyak in [24]; our proof uses a trick which we learned from the lectures of Werner Ballmann [29]. An other proof can be build on Kirszbraun's theorem (9.4.1), but it works only for complete spaces.

The case when α is a triangle, say $[pxy]$, is leading and nontrivial. In this case, by Proposition 8.12.3, the majorizing convex region has to be isometric to $\operatorname{Conv}[\tilde p\tilde x\tilde y]$, where $[\tilde p\tilde x\tilde y] = \tilde\triangle^\kappa(pxy)$. There is a

majorizing map for $[pxy]$ which image W is the image of the line-of-sight map (definition 8.8.3) for $[xy]$ from p, but as one can see from the following example, the line-of-sight map is not majorizing in general.

Example. Let $\mathcal{Q}$ be a bounded region in $\mathbb{E}^2$ formed by two congruent triangles, where the bounding quadrangle $[pxzy]$ is non-convex at z (as in the picture). Equip $\mathcal{Q}$ with the length-metric. Then $\mathcal{Q}$ is CAT(0) by Reshetnyak gluing (8.9.1). For triangle $[pxy]_{\mathcal{Q}}$ in $\mathcal{Q}$ and its model triangle $[\tilde{p}\tilde{x}\tilde{y}]$ in $\mathbb{E}^2$, we have



$$|\tilde{x}-\tilde{y}| = |x-y|_{\mathcal{Q}} = |x-z| + |z-y|.$$

Then the map F defined by matching line-of-sight parameters satisfies $F(\tilde{x}) = x$ and $|x - F(\tilde{w})| > |\tilde{x}-\tilde{w}|$ if $\tilde{w}$ is near the midpoint $\tilde{z}$ of $[\tilde{x}\tilde{y}]$ and lies on $[\tilde{p}\tilde{z}]$. Indeed, by the first variation formula (7.7.6), for $\varepsilon = 1 - s$ we have

$$|\tilde{x}-\tilde{w}| = |\tilde{x}-\tilde{\gamma}_{\frac{1}{2}}(s)| = |x-z| + o(\varepsilon)$$

and

$$|x - F(\tilde{w})| = |x - \gamma_{\frac{1}{2}}(s)| = |x-z| - \varepsilon\cdot\cos\measuredangle[z\,{}^{p}_{x}] + o(\varepsilon).$$

Thus F is not majorizing.

In the following proofs, $x^1\dots x^n$ ($n \geqslant 3$) denotes a broken geodesic with vertices $x^1,\dots,x^n$, and $[x^1\dots x^n]$ denotes the corresponding (closed) polygon. For a subset R of the ambient metric space, we denote by $[x^1\dots x^n]_R$ a polygon in the length metric of R.

Our first lemma gives a model space construction based on repeated application of Lemma 8.4.6. Recall that convex and concave curves with respect to a point are defined in 7.5.2.

8.12.5. Lemma. *In $\mathbb{M}^2(\kappa)$, let β be a curve from x to y that is concave with respect to p. Let D be the subgraph of β with respect to p. Assume*

$$\operatorname{length}\beta + |p-x| + |p-y| < 2\cdot\varpi^{\kappa}.$$

a) Then β forms a geodesic $[xy]_D$ in D and therefore β, $[px]$ and $[py]$ form a triangle $[pxy]_D$ in the length-metric of D.

b) Let $[\tilde{p}\tilde{x}\tilde{y}]$ be the model triangle for triangle $[pxy]_D$. Then there is a short map $G\colon \operatorname{Conv}[\tilde{p}\tilde{x}\tilde{y}] \to D$ such that $\tilde{p}\mapsto p$, $\tilde{x}\mapsto x$, $\tilde{y}\mapsto y$, and G is length-preserving on each side of $[\tilde{p}\tilde{x}\tilde{y}]$. In particular, $\operatorname{Conv}[\tilde{p}\tilde{x}\tilde{y}]$ majorizes triangle $[pxy]_D$ in D under G.

Proof. We prove the lemma for a broken geodesic β; the general case then follows by approximation. Namely, since β is concave it can be approximated by broken geodesics that are concave with respect to p, with their lengths converging to $\operatorname{length}\beta$. Passing to a partial limit we will obtain the needed map G.

Suppose $\beta = x^0x^1\dots x^n$ is a broken geodesic with $x^0 = x$ and $x^n = y$. Consider a sequence of broken geodesics $\beta_i = x^0x^1\dots x^{i-1}y_i$ such that $|p - y_i| = |p - y|$ and β_i has same length as β; that is,

$$|x^{i-1} - y_i| = |x^{i-1} - x^i| + |x^i - x^{i+1}| + \dots + |x^{n-1} - x^n|.$$



Clearly $\beta_n = \beta$. Sequentially applying Alexandrov's lemma (5.2.1) shows that each of the broken geodesics $\beta_{n-1}, \beta_{n-2}, \dots, \beta_1$ is concave with respect to p. Let D_i to be subgraph of β_i with respect to p. Applying Lemma 8.4.6 gives a short map $G_i\colon D_i \to D_{i+1}$ that maps $y_i \mapsto y_{i+1}$ and does not move p and x (in fact, G_i is the identity everywhere except on $\operatorname{Conv}[px^{i-1}y_i]$). Thus the composition

$$G_{n-1} \circ \dots \circ G_1\colon D_1 \to D_n$$

is short. The result follows since $D_1 \overset{iso}{=\!=} \operatorname{Conv}[\tilde p\tilde x\tilde y]$. $\square$

8.12.6. Lemma. *Let $[pxy]$ be a triangle of perimeter $< 2\cdot\varpi^\kappa$ in a ϖ^κ-geodesic $\mathrm{CAT}(\kappa)$ space $\mathcal{U}$. In $\mathbb{M}^2(\kappa)$, let $\tilde\gamma$ be the κ-development of $[xy]$ with respect to p, where $\tilde\gamma$ has basepoint $\tilde p$ and subgraph D. Consider the map $H\colon D \to \mathcal{U}$ that sends the point with parameter (t,s) under the line-of-sight map for $\tilde\gamma$ from $\tilde p$, to the point with the same parameter under the line-of-sight map f for $[xy]$ from p. Then H is length-nonincreasing. In particular, D majorizes triangle $[pxy]$.*

Proof. Let $\gamma = \operatorname{geod}_{[xy]}$ and $T = |x - y|$. As in the proof of the development criterion (8.7.1), take a partition

$$0 = t^0 < t^1 < \dots < t^n = T,$$

and set $x^i = \gamma(t^i)$. Construct a chain of model triangles $[\tilde{p}\tilde{x}^{i-1}\tilde{x}^i] = \tilde{\triangle}^\kappa(px^{i-1}x^i)$, with $\tilde{x}^0 = \tilde{x}$ and the direction of $[\tilde{p}\tilde{x}^i]$ turning counterclockwise as i grows. Let D_n be the subgraph with respect to $\tilde{p}$ of the broken geodesic $\tilde{x}^0 \dots \tilde{x}^n$.

Let δ_n be the maximum radius of a circle inscribed in any of the triangles $[\tilde{p}\tilde{x}^{i-1}\tilde{x}^i]$.

Now we construct a map $H_n\colon D_n \to \mathcal{U}$ that increases distances by at most $2{\cdot}\delta_n$.

Suppose $x \in D_n$. Then x lies on or inside some triangle $[\tilde{p}\tilde{x}^{i-1}\tilde{x}^i]$. Define $H_n(x)$ by first mapping x to a nearest point on $[\tilde{p}\tilde{x}^{i-1}\tilde{x}^i]$ (choosing one if there are several), followed by the natural map to the triangle $[px^{i-1}x^i]$.

Since triangles in $\mathcal{U}$ are κ-thin (8.4.3), the restriction of H_n to each triangle $[\tilde{p}\tilde{x}^{i-1}\tilde{x}^i]$ is short. Then the triangle inequality implies that the restriction of H_n to

$$U_n = \bigcup_{1\leqslant i\leqslant n} [\tilde{p}\tilde{x}^{i-1}\tilde{x}^i]$$

is short with respect to the length-metric on D_n. Since nearest-point projection from D_n to U_n increases the D_n-distance between two points by at most $2{\cdot}\delta_n$, the map H_n also increases the D_n-distance by at most $2{\cdot}\delta_n$.

Consider $y_n \in D_n$ with $y_n \to y \in D$ and $z_n \in D_n$ with $z_n \to \to z \in D$. Since $\delta_n \to 0$ under increasingly finer partitions, and geodesics in $\mathcal{U}$ vary continuously with their endpoints (8.8.1), we have $H_n(x_n) \to H(x)$ and $H_n(y_n) \to H(y)$. Since

$$|H_n(x_n) - H_n(y_n)| \leqslant |x_n - y_n|_{D_n} + 2{\cdot}\delta_n,$$

where the left hand side converges to $|H(x)-H(y)|$ and the right hand side converges to $|x-y|_D$, it follows that H is short. □

Proof of 8.12.4. We begin by proving the theorem in case α is polygonal.

First suppose α is a triangle, say $[pxy]$. By assumption, the perimeter of $[pxy]$ is less than $2{\cdot}\varpi^\kappa$. This is the base case for the induction.

Let $\tilde{\gamma}$ be the κ-development of $[xy]$ with respect to p, where $\tilde{\gamma}$ has basepoint $\tilde{p}$ and subgraph D. By the development criterion (8.7.1), $\tilde{\gamma}$ is concave. By Lemma 8.12.5, there is a short map $G\colon \operatorname{Conv}\tilde{\triangle}^\kappa(pxy) \to \to D$. Further, by Lemma 8.12.6, D majorizes $[pxy]$ under a majorizing map $H\colon D \to \mathcal{U}$. Clearly $H \circ G$ is a majorizing map for $[pxy]$.



Now we claim that any closed n-gon $[x^1x^2\dots x^n]$ of perimeter less than $2\cdot\varpi^\kappa$ in a CAT(κ) space is majorized by a convex polygonal region

$$R_n = \operatorname{Conv}[\tilde{x}^1\tilde{x}^2\dots\tilde{x}^n]$$

under a map F_n such that $F_n\colon \tilde{x}^i \mapsto x^i$ for each i.

Assume the statement is true for $(n-1)$-gons, $n \geqslant 4$. Then $[x^1x^2\dots x^{n-1}]$ is majorized by a convex polygonal region

$$R_{n-1} = \operatorname{Conv}(\tilde{x}^1, \tilde{x}^2, \dots, \tilde{x}^{n-1}),$$

in $\mathbb{M}^2(\kappa)$ under a map F_{n-1} satisfying $F_{n-1}(\tilde{x}^i) = x^i$ for all i. Take $\dot{x}^n \in \mathbb{M}^2(\kappa)$ such that $[\tilde{x}^1\tilde{x}^{n-1}\dot{x}^n] = \tilde{\triangle}^\kappa(x^1x^{n-1}x^n)$ and this triangle lies on the other side of $[\tilde{x}^1\tilde{x}^{n-1}]$ from R_{n-1}. Let $\dot{R} = \operatorname{Conv}[\tilde{x}^1\tilde{x}^{n-1}\dot{x}^n]$, and $\dot{F}\colon \dot{R} \to \mathcal{U}$ be a majorizing map for $[x^1x^{n-1}x^n]$ as provided above.

Set $R = R_{n-1} \cup \dot{R}$, where R carries its length-metric. Since F_n and F agree on $[\tilde{x}^1\tilde{x}^{n-1}]$, we may define $F\colon R \to \mathcal{U}$ by

$$F(x) = \begin{cases} F_{n-1}(x), & x \in R_{n-1}, \\ \dot{F}(x), & x \in \dot{R}. \end{cases}$$

Then F is length-nonincreasing, and is a majorizing map for $[x^1x^2\dots$ $\dots x^n]$ (as in Definition 8.12.1).

If R is a convex subset of $\mathbb{M}^2(\kappa)$, we are done.

If R is not convex, the total internal angle of R at $\tilde{x}^1$ or $\tilde{x}^{n-1}$ or both is $> \pi$. By relabeling we may suppose that it holds for $\tilde{x}^{n-1}$.

The region R is obtained by glueing R_{n-1} to $\dot{R}$ by $[x^1x^{n-1}]$. Thus, by Reshetnyak gluing (8.9.1), R in its length-metric is a CAT(κ)-space. Moreover $[\tilde{x}^{n-2}\tilde{x}^{n-1}] \cup [\tilde{x}^{n-1}\dot{x}^n]$ is a geodesic of R. Thus $[\tilde{x}^1, \tilde{x}^2, \dots$ $\dots, \tilde{x}^{n-1}\dot{x}^n]_R$ is a closed $(n-1)$-gon in R, to which the induction hypothesis applies. The resulting short map from a convex region in $\mathbb{M}^2(\kappa)$ to R, followed by F, is the desired majorizing map.

Note that in fact we proved the following:

❶ *Let F_{n-1} be a majorizing map for the polygon $[x^1x^2\dots x^{n-1}]$, and $\dot{F}$ be a majorizing map for the triangle $[x^1x^{n-1}x^n]$. Then there is a majorizing map F_n for the polygon $[x^1x^2\dots x^n]$ such that*

$$\operatorname{Im} F_{n+1} = \operatorname{Im} F_n \cup \operatorname{Im} \dot{F}.$$

We now use this claim to prove the theorem for general curves.

Assume $\alpha\colon [0,\ell] \to \mathcal{U}$ is an arbitrary closed curve with natural parameter. Choose a sequence of partitions $0 = t_n^0 < t_n^2 < \ldots < t_n^n = \ell$ so that:

- ◇ The set $\{t_{n+1}^i\}_{i=0}^{n+1}$ is obtained from the set $\{t_n^i\}_{i=0}^n$ by adding one element.
- ◇ For some sequence $\varepsilon_n \to 0+$, we have $t_n^i - t_n^{i-1} < \varepsilon_n$ for all i.

Inscribe in α a sequence of polygons P_n with vertices $\alpha(t_n^i)$. Apply the claim above, to get a sequence of majorizing maps $F_n\colon R_n \to \mathcal{U}$. Note that for all $m > n$

- ◇ $\operatorname{Im} F_m$ lies in an ε_n-neighborhood of $\operatorname{Im} F_n$
- ◇ $\operatorname{Im} F_m \setminus \operatorname{Im} F_n$ lies in an ε_n-neighborhood of α.

It follows that the set

$$K = \alpha \cup \left(\bigcup_n \operatorname{Im} F_n\right)$$

is compact. Therefore the sequence (F_n) has a partial limit as $n \to \infty$; say F. Clearly F is a majorizing map for α. □

The following exercise is the rigidity case of the majorization theorem.

8.12.7. Exercise. *Let $\mathcal{U}$ be a ϖ^κ-geodesic* CAT(κ) *space and $\alpha\colon [0,\ell] \to \mathcal{U}$ be a closed curve with arclength parametrization. Assume that $\ell < 2\cdot\varpi^\kappa$ and there is a closed convex curve $\tilde\alpha\colon [0,\ell] \to \mathbb{M}^2(\kappa)$ such that*

$$|\alpha(t_0) - \alpha(t_1)|_{\mathcal{U}} = |\tilde\alpha(t_0) - \tilde\alpha(t_1)|_{\mathbb{M}^2(\kappa)}$$

for any t_0 and t_1. Then there is a distance preserving map $F\colon \operatorname{Conv}\tilde\alpha \to \mathcal{U}$ such that $F\colon \tilde\alpha(t) \mapsto \alpha(t)$ for any t.

8.12.8. Exercise. *Assume a closed curve α of length $< 2\cdot\varpi^\kappa$ in a ϖ^κ-geodesic* CAT(κ) *space is not a triangle. Show that exactly one of these statements holds:*

a) α is majorized by non-isometric convex regions in $\mathbb{M}^2(\kappa)$;

b) the majorizing map for α is distance preserving.

For $n = 1$, the following lemma states that in a CAT(κ) space, a sharp triangle comparison implies the presence of an isometric copy of the convex hull of the model triangle. This rigidity statement was proved by Alexandrov [16]. An analogue for CBB(κ) spaces fails, see Exercise 9.4.7.

8.12.9. Arm lemma. *Let $\mathcal{U}$ be a ϖ^κ-geodesic* $\mathrm{CAT}(\kappa)$ *space, and $P = [x^0x^1\dots x^{n+1}]$ be a polygon of length $< 2\cdot\varpi^\kappa$ in $\mathcal{U}$. Suppose $\tilde P = [\tilde x^0\tilde x^1\dots\tilde x^{n+1}]$ is a convex polygon in $\mathbb{M}^2(\kappa)$ such that*

❷ $$|\tilde x^i - \tilde x^{i-1}|_{\mathbb{M}^2(\kappa)} = |x^i - x^{i-1}|_{\mathcal{U}} \quad \text{and} \quad \measuredangle[x^i\,{}^{x^{i-1}}_{x^{i+1}}] \geqslant \measuredangle[\tilde x^i\,{}^{\tilde x^{i-1}}_{\tilde x^{i+1}}]$$

for all i. Then

a) $|\tilde x^0 - \tilde x^{n+1}|_{\mathbb{M}^2(\kappa)} \leqslant |x^0 - x^{n+1}|_{\mathcal{U}}$.

b) Equality holds in (a) if and only if the map $\tilde x^i \mapsto x^i$ can be extended to a distance-preserving map of $\mathrm{Conv}(\tilde x^0, \tilde x^1\dots$ $\dots\tilde x^{n+1})$ *onto* $\mathrm{Conv}(x^0, x^1\dots x^{n+1})$.

Proof. a). By majorization (8.12.4), P is majorized by a convex region $\tilde D$ in $\mathbb{M}^2(\kappa)$. By Proposition 8.12.2 and the definition of angle, $\tilde D$ is bounded by a convex polygon $\tilde P_R = [\,y^0\tilde y^1\dots\tilde y^{n+1}]$ that satisfies

$$|\tilde y^i - \tilde y^{i\pm1}|_{\mathbb{M}^2(\kappa)} = |x^i - x^{i\pm1}|_{\mathcal{U}}, \qquad |\tilde y^0 - \tilde y^{n+1}|_{\mathbb{M}^2(\kappa)} = |x^0 - x^{n+1}|_{\mathcal{U}},$$
$$\measuredangle[\tilde y^i\,{}^{\tilde y^{i-1}}_{\tilde y^{i+1}}] \geqslant \measuredangle[x^i\,{}^{x^{i-1}}_{x^{i+1}}] \geqslant \measuredangle[\tilde x^i\,{}^{\tilde x^{i-1}}_{\tilde x^{i+1}}],$$

for $1 \leqslant i \leqslant n$, where the final inequality is by ❷.

By the arm lemma (8.12.9), $|\tilde x^0 - \tilde x^{n+1}| \leqslant |\tilde y^0 - \tilde y^{n+1}|$. Since $|\tilde y^0 - \tilde y^{n+1}| = |x^0 - x^{n+1}|$, the part (a) follows.

b). Suppose equality holds in (a). Then angles at the j-th vertex of $\tilde P$, P and $\tilde P_R$ are equal for $1 \leqslant j \leqslant n$, and we may take $\tilde P = \tilde P_R$.

Let $F\colon \tilde D \to \mathcal{U}$ be the majorizing map for P, where $\tilde D$ is the convex region bounded by $\tilde P$, and $F|_{\tilde P}$ is length-preserving.

❸ *Let $\tilde x, \tilde y, \tilde z$ be three vertices of $\tilde P$, and x, y, z be the corresponding vertices of P. If $|\tilde x - \tilde y| = |x - y|$, $|\tilde y - \tilde z| = |x - z|$ and $\measuredangle[\tilde y\,{}^{\tilde x}_{\tilde z}] = \measuredangle[y\,{}^{x}_{z}]$, then $F|_{\mathrm{Conv}(\tilde x,\tilde y,\tilde z)}$ is distance-preserving.*

Since F is majorizing, F restricts to distance-preserving maps from $[\tilde x\tilde y]$ to $[xy]$ and $[\tilde y\tilde z]$ to $[yz]$. Suppose $\tilde p \in [\tilde x\tilde y]$ and $\tilde q \in [\tilde y\tilde z]$. Then

❹ $$|\tilde p - \tilde q|_{\mathbb{M}^2(\kappa)} = |F(\tilde p) - F(\tilde q)|_{\mathcal{U}}.$$

Indeed, inequality holds in one direction by majorization, and in the other direction by angle comparison (8.3.1c). By first variation formula (8.14.4), it follows that each pair of corresponding angles of triangles $[\tilde x\tilde y\tilde z]$ and $[xyz]$ are equal. But then ❹ holds for p, q on any two sides of these triangles, so F is distance-preserving on every geodesic of $\mathrm{Conv}(\tilde p, \tilde x, \tilde y)$. Hence the claim.

❺ *Suppose $F|_{\mathrm{Conv}(\tilde x^{n+1},\tilde x^0,\tilde x^1\dots\tilde x^k)}$ is distance-preserving for some k, $1 \leqslant k \leqslant n-1$. Then $F|\,\mathrm{Conv}(\tilde x^{n+1}, \tilde x^0, \tilde x^1\dots\tilde x^{k+1})$ is distance-preserving.*

Set $\tilde p = [\tilde x^{k-1}\tilde x^{k+1}] \cap [\tilde x^k \tilde x^{n+1}]$ and $p = [x^{k-1}x^{k+1}] \cap [x^k x^{n+1}]$.

The following maps are distance-preserving:

(i) $F|_{\mathrm{Conv}(\tilde x^{k-1},\tilde x^k,\tilde x^{k+1})}$,

(ii) $F|_{\mathrm{Conv}(\tilde x^{k+1},\tilde x^{k-1},\tilde x^{n+1})}$,

(iii) $F|_{\mathrm{Conv}(\tilde x^{n+1},\tilde x^k,\tilde x^{k+1})}$.

Indeed, (i) follows from ❸. Therefore $|\tilde x^{k-1} - \tilde x^{k+1}| = |x^{k-1} - x^{k+1}|$, and so F restricts to a distance-preserving map from $[\tilde x^{k-1}\tilde x^{k+1}]$ onto $[x^{k-1}x^{k+1}]$. With the induction hypothesis, it follows that $F(\tilde p) = p$, hence

❻ $$\measuredangle[\tilde x^{k-1}\,{}^{\tilde x^{k+1}}_{\tilde x^{n+1}}] = \measuredangle[x^{n+1}\,{}^{x^{k-1}}_{x^{k+1}}].$$

Then (ii) follows from ❻ and ❸. Since $|\tilde x^k - \tilde x^{n+1}| = |x^k - x^{n+1}|$, (iii) follows from ❻ and (i).

Let $\tilde\gamma$ be a geodesic of $\mathrm{Conv}(\tilde x^{n+1}, \tilde x^0, \tilde x^1 \dots \tilde x^{k+1})$. Then length $\tilde\gamma < \varpi^\kappa$. If $\tilde\gamma$ does not pass the point $[\tilde x^{k-1}\tilde x^{k+1}] \cap [\tilde x^k \tilde x^{n+1}]$, it follows from the induction hypothesis and (i) (ii), (iii) that $\gamma = F\circ\tilde\gamma$ is a local geodesic of length $< \varpi^\kappa$. By 8.4.4, γ is a geodesic. By continuity, $F\circ\tilde\gamma$ is a geodesic for all $\tilde\gamma$. ❺ follows and hence the lemma. □

8.12.10. Exercise. *Let $\mathcal U$ be a complete length* CAT(0) *space and for 4 points $x_1, x_2, x_3, x_4 \in \mathcal U$ there is a convex quadrilateral $[\tilde x_1\tilde x_2\tilde x_3\tilde x_4]$ in $\mathbb E^2$ such that*

$$|x_i - x_j|_{\mathcal U} = |\tilde x_i - \tilde x_j|_{\mathbb E^2}$$

for all i and j. Show that $\mathcal U$ contains an isometric copy of the solid quadrilateral *$\tilde x_1\tilde x_2\tilde x_3\tilde x_4$; that is, the convex hull of $\tilde x_1, \tilde x_2, \tilde x_3, \tilde x_4$ in $\mathbb E^2$*

8.13 Hadamard–Cartan theorem

The development of Alexandrov geomtry was greatly influenced by the Hadamard–Cartan theorem. Its original formulation states that if M is a complete Riemannian manifold with nonpositive sectional curvature, then the exponential map at any point $p \in M$ is a covering; in particular it implies that the universal cover of M is diffeomorphic to the Euclidean space of the same dimension.

In this generality, the theorem appeared in the lectures of Cartan [30]. For surfaces in the Euclidean plane, the theorem was proved by Hans von Mangoldt [31], and few years later independently by Hadamard [32].

Formulations for metric spaces of different generality were proved by Busemann [15], Rinow [33], and Gromov [34, p. 119]. A detailed proof of Gromov's statement when $\mathcal U$ is proper was given by Werner

Ballmann [35], using Birkhoff's curve-shortening. A proof in the non-proper geodesic case was given by the first author and Richard Bishop [27]. This proof applies more generally, to *convex spaces* (see Exercise 8.13.7). It was pointed out by Bruce Kleiner [29] and independently by Martin Bridson and André Haefliger [26] that this proof extends to length spaces as well as geodesic spaces, giving:

8.13.1. Hadamard–Cartan theorem. *Let $\mathcal{U}$ be a complete, simply connected length locally* CAT(0) *space. Then $\mathcal{U}$ is* CAT(0).

Proof. Since $\varpi^\kappa = \infty$, the map $\Phi\colon \mathcal{B} \to \mathcal{U}$ in Theorem (8.11.1) is a metric covering. Since $\mathcal{U}$ is simply connected, $\Phi\colon \mathcal{B} \to \mathcal{U}$ is an isometry. □

To formulate the generalized Hadamard–Cartan theorem, we need the following definition.

8.13.2. Definition. *Given $\ell \in (0,\infty]$, a metric space $\mathcal{X}$ is called ℓ-simply connected if it is connected and any closed curve of length $< \ell$ is null-homotopic in the class of curves of length $< \ell$ in $\mathcal{X}$.*

Note that there is a subtle difference between simply connected and ∞-simply connected spaces; the first states that any closed curve is null-homotopic while the second means that any rectifiable curve is null-homotopic in the class of rectifiable curves. However, as follows from Proposition 8.13.5, for locally CAT(κ) spaces these two definitions are equivalent. This fact makes it possible to deduce the Hadamard–Cartan theorem directly from the generalized Hadamard–Cartan theorem.

8.13.3. Generalized Hadamard–Cartan theorem. *A complete length space $\mathcal{U}$ is* CAT(κ) *if and only if $\mathcal{U}$ is $2{\cdot}\varpi^\kappa$-simply connected and $\mathcal{U}$ is locally* CAT(κ).

For proper spaces, the generalized Hadamard–Cartan theorem was proved by Brian Bowditch [36]. In the proof we need the following lemma.

8.13.4. Lemma.

Assume $\mathcal{U}$ is a complete length locally CAT(κ) *space, $\varepsilon > 0$, and $\gamma_1, \gamma_2\colon \mathbb{S}^1 \to \mathcal{U}$ are two closed curves. Assume*

a) $\operatorname{length}\gamma_1, \operatorname{length}\gamma_2 < 2{\cdot}\varpi^\kappa - 4{\cdot}\varepsilon$*;*

b) $|\gamma_1(x) - \gamma_2(x)| < \varepsilon$ *for any $x \in \mathbb{S}^1$, and the geodesic $[\gamma_1(x)\gamma_2(x)]$ is uniquely defined and depends continuously on x;*

c) γ_1 *is majorized by a convex region in $\mathbb{M}^2(\kappa)$.*

Then γ_2 is majorized by a convex region in $\mathbb{M}^2(\kappa)$.

Proof. Let D be a convex region in $\mathbb{M}^2(\kappa)$ that majorizes γ_1 under the map $F\colon D \to \mathcal{U}$ (see Definition 8.12.1). Denote by $\tilde\gamma_1$ the curve bounding D such that $F\circ\tilde\gamma_1 = \gamma_1$. Since

$$\begin{aligned}\operatorname{length}\tilde\gamma_1 = \operatorname{length}\gamma_1 &< \\ &< 2\cdot\varpi^\kappa - 4\cdot\varepsilon,\end{aligned}$$

there is a point $\tilde p \in D$ such that $|\tilde p - \tilde\gamma(x)|_{\mathbb{M}^2(\kappa)} < \frac{\varpi^\kappa}{2} - \varepsilon$ for any $x \in \mathbb{S}^1$. Denote by α_x the joint of the paths $F\circ\operatorname{path}_{[p\tilde\gamma_1(x)]_{\mathbb{M}^2(\kappa)}}$ and $\operatorname{path}_{[\gamma_1(x)\gamma_2(x)]}$ in $\mathcal{U}$. Note that α_x depends continuously on x and

$$\operatorname{length}\alpha_x < \tfrac{\varpi^\kappa}{2} \quad\text{and}\quad \alpha_x(1) = \gamma_2(x)$$

for any x.

Let us apply the lifting globalization theorem (8.11.1) for $p = F(\tilde p)$. We get a ϖ^κ-geodesic CAT(κ) space $\mathcal{B}$ and a locally isometric map $\Phi\colon \mathcal{B}\to\mathcal{U}$ with $\Phi(\hat p) = p$ for some $\hat p \in \mathcal{B}$, and with the lifting property for the curves starting at p with length $< \varpi^\kappa/2$. Applying the lifting property for α_x, we get existence of a curve $\hat\gamma_2\colon \mathbb{S}^1 \to \mathcal{B}$ such that

$$\gamma_2 = \Phi\circ\hat\gamma_2.$$

Since $\mathcal{B}$ is a geodesic CAT(κ) space, we can apply the majorization theorem (8.12.4) for $\hat\gamma_2$. The composition of the obtained majorization with Φ is a majorization of γ_2. □

Proof. The "only if" part follows from the Reshetnyak majorization theorem (8.12.4).

Let γ_t, $t\in[0,1]$ be a null-homotopy of curves in $\mathcal{U}$; that is, $\gamma_0(x) = p$ for some $p\in\mathcal{U}$ and any $x\in\mathbb{S}^1$. Assume further that $\operatorname{length}\gamma_t < 2\cdot\varpi^\kappa$ for any t. To prove the "only if" part, it is sufficient to show that γ_1 is majorized by a convex region in $\mathbb{M}^2(\kappa)$.

By semicontinuity of length (2.3.2), we can choose $\varepsilon > 0$ sufficiently small that

$$\operatorname{length}\gamma_t < 2\cdot\varpi^\kappa - 4\cdot\varepsilon$$

for all t.

By Corollary 8.11.2, we may assume in addition that $\mathrm{B}(\gamma_t(x),\varepsilon)$ is CAT(κ) for any t and x.

Choose a partition $0 = t_0 < t_1 < \ldots < t_n = 1$ so that $|\gamma_{t_i}(x) - \gamma_{t_{i-1}}(x)| < \varepsilon$ for any i and x. According to 8.2.1, for any i, the geodesic $[\gamma_{t_i}(x)\gamma_{t_{i-1}}(x)]$ depends continuously on x.

Note that $\gamma_0 = \gamma_{t_0}$ is majorized by a convex region in $\mathbb{M}^2(\kappa)$. Applying the lemma n times, we see that the same holds for $\gamma_1 = \gamma_{t_n}$. $\square$

8.13.5. Proposition. *Let $\mathcal{U}$ be a complete length locally* CAT(κ) *space. Then $\mathcal{U}$ is simply connected if and only if it is ∞-simply connected.*

Proof; "if" part. It is sufficient to show that any closed curve in $\mathcal{U}$ is homotopic to a closed broken geodesic.

Let γ_0 be a closed curve in $\mathcal{U}$. According to Corollary 8.11.2, there is $\varepsilon > 0$ such that $\mathrm{B}(\gamma(x), \varepsilon)$ is CAT(κ) for any x.

Choose a broken geodesic γ_1 such that $|\gamma_0(x) - \gamma_1(x)| < \varepsilon$ for any x. By 8.2.1, $\mathrm{path}_{[\gamma_0(x)\gamma_1(x)]}$ is uniquely defined and depends continuously on x.

Hence $\gamma_t(x) = \mathrm{path}_{[\gamma_0(x)\gamma_1(x)]}(t)$ gives a homotopy from γ_0 to γ_1.

"Only if" part. The proof is similar.

Assume γ_t is a homotopy between two rectifiable curves γ_0 and γ_1. Fix $\varepsilon > 0$ so that the ball $\mathrm{B}(\gamma_t(x), \varepsilon)$ is CAT(κ) for any t and x. Choose a partition $0 = t_0 < t_1 < \ldots < t_n = 1$ so that

$$|\gamma_{t_{i-1}}(x) - \gamma_{t_i}(x)| < \tfrac{\varepsilon}{10}$$

for any i and x. Set $\hat\gamma_{t_0} = \gamma_0$, $\hat\gamma_{t_n} = \gamma_{t_n}$. For each $0 < i < n$, approximate γ_{t_i} by a closed broken geodesic $\hat\gamma_i$.

Construct the homotopy from $\hat\gamma_{t_{i-1}}$ to $\hat\gamma_{t_i}$ setting

$$\hat\gamma_t = \mathrm{path}_{[\hat\gamma_{t_{i-1}}(x)\hat\gamma_{t_i}(x)]}(t).$$

Since ε is sufficiently small, by 8.2.6, we get that

$$\operatorname{length}\hat\gamma_t < 10\cdot(\operatorname{length}\hat\gamma_{t_{i-1}} + \operatorname{length}\hat\gamma_{t_i}).$$

In particular, $\hat\gamma_t$ is rectifiable for all t.

Joining the obtained homotopies for all i we obtain a homotoply from γ_0 to γ_1 in the class of rectifiable curves. $\square$

8.13.6. Exercise. *Let $\mathcal{X}$ be a double cover of $\mathbb{E}^3$ that branches along two distinct lines ℓ and m. Show that $\mathcal{X}$ is* CAT(0) *if and only if ℓ intersects m at a right angle.*

About convex spaces. A *convex space* $\mathcal{X}$ is a geodesic space such that the function $t \mapsto |\gamma(t) - \sigma(t)|$ is convex for any two geodesic paths $\gamma, \sigma : [0,1] \to \mathcal{X}$. A *locally convex space* is a length space in which

every point has a neighborhood that is a convex space in the restricted metric.

8.13.7. Exercise. *Assume $\mathcal{X}$ is a convex space such that the angle of any hinge is defined. Show that $\mathcal{X}$ is* CAT(0).

The following exercise gives an analog of Hadamard–Cartan theorem for locally convex spaces; see also [27].

8.13.8. Exercise. *Show that a complete, simply connected, locally convex space is a convex space.*

8.14 Angles

Recall that ω denotes a selective nonpriciple ultrafilter on $\mathbb{N}$, see Section 3.2.

8.14.1. Angle semicontinuity. *Let $\mathcal{U}_n$ be a ϖ^κ-geodesic* CAT(κ) *space for each $n \in \mathbb{N}$ and $\mathcal{U}_n \to \mathcal{U}_\omega$ as $n \to \omega$. Assume that a sequence of hinges $[p_n\,{}^{x_n}_{y_n}]$ in $\mathcal{U}_n$ converges to a hinge $[p_\omega\,{}^{x_\omega}_{y_\omega}]$ in $\mathcal{U}_\omega$. Then*

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] \geqslant \lim_{n\to\omega} \measuredangle[p_n\,{}^{x_n}_{y_n}].$$

Proof. By angle-sidelength monotonicity (8.3.2),

$$\measuredangle[p_\omega\,{}^{x_\omega}_{y_\omega}] = \inf\left\{\, \tilde{\measuredangle}^\kappa(p_\omega\,{}^{\bar x_\omega}_{\bar y_\omega}) : \bar x_\omega \in\,]p_\omega x_\omega],\ \bar y_\omega \in\,]p_\omega y_\omega]\,\right\}.$$

For fixed $\bar x_\omega \in\,]p_\omega x_\omega]$ and $\bar y_\omega \in\,]p_\omega x_\omega]$, choose $\bar x_n \in\,]p x_n]$ and $\bar y_n \in\,]p y_n]$ so that $\bar x_n \to \bar x_\omega$ and $\bar y_n \to \bar y_\omega$ as $n \to \omega$. Clearly

$$\tilde{\measuredangle}^\kappa(p_n\,{}^{\bar x_n}_{\bar y_n}) \to \tilde{\measuredangle}^\kappa(p_\omega\,{}^{\bar x_\omega}_{\bar y_\omega})$$

as $n \to \omega$.

By angle comparison (8.3.1c), $\measuredangle[p_n\,{}^{x_n}_{y_n}] \leqslant \tilde{\measuredangle}^\kappa(p_n\,{}^{\bar x_n}_{\bar y_n})$. Hence the result. □

Now we verify that the first variation formula holds in the CAT setting. Compare it to the first variation inequality (5.3.4) which holds for general metric spaces and to the strong angle lemma (7.7.6) for CBB spaces.

8.14.2. Strong angle lemma. *Let $\mathcal{U}$ be a ϖ^κ-geodesic* CAT(κ) *space. Then for any hinge $[p\,{}^{q}_{x}]$ in $\mathcal{U}$, we have*

❶ $$\measuredangle[p\,{}^{q}_{x}] = \lim_{\bar x\to p} \left\{\, \tilde{\measuredangle}^\kappa(p\,{}^{q}_{\bar x}) : \bar x \in\,]px]\,\right\}$$

for any $\kappa \in \mathbb{R}$ *such that* $|p-q| < \varpi^\kappa$.

Proof. By angle-sidelength monotonicity (8.3.2), the righthand side is defined and at least equal to the lefthand side.

By Lemma 5.3.1, we may take $\kappa = 0$ in ❶. By the cosine law and the first variation inequality (5.3.4), the righthand side is at most equal to the lefthand side. □

8.14.3. First variation. *Let* $\mathcal{U}$ *be a* ϖ^κ*-geodesic* $\mathrm{CAT}(\kappa)$ *space. For any nontrivial geodesic* $[py]$ *in* $\mathcal{U}$ *and point* $q \neq p$ *such that* $|p-q| < \varpi^\kappa$, *we have*

$$|q - \mathrm{geod}_{[py]}(t)| = |q-p| - t\cdot \cos\measuredangle[p\,{}^q_y] + o(t).$$

Proof. The first variation equation is equivalent to the strong angle lemma (8.14.2), as follows from the Euclidean cosine law. □

8.14.4. First variation (both-endpoints version). *Let* $\mathcal{U}$ *be a* ϖ^κ*-geodesic* $\mathrm{CAT}(\kappa)$ *space. For any nontrivial geodesics* $[py]$ *and* $[qz]$ *in* $\mathcal{U}$ *such that* $p \neq q$ *and* $|p-q| < \varpi^\kappa$, *we have*

$$|\mathrm{geod}_{[py]}(t) - \mathrm{geod}_{[qz]}(\tau)| = |q-p| - t\cdot \cos\measuredangle[p\,{}^q_y] - \tau\cdot \cos\measuredangle[q\,{}^p_z] + o(t+\tau).$$

Proof. By the first variation equation (8.14.3),

$$\begin{aligned}
&|\mathrm{geod}_{[py]}(t) - \mathrm{geod}_{[qz]}(\tau)| = \\
&= |q - \mathrm{geod}_{[py]}(t)| - \tau\cdot \cos\measuredangle[q\,{}^{\mathrm{geod}_{[py]}(t)}_z] + o(\tau) \\
&= |q-p| - t\cdot \cos\measuredangle[p\,{}^q_y] + o(t) - \tau\cdot \cos\measuredangle[q\,{}^{\mathrm{geod}_{[py]}(t)}_z] + o(\tau) = \\
&= |q-p| - t\cdot \cos\measuredangle[p\,{}^q_y] - \tau\cdot \cos\measuredangle[q\,{}^p_z] + o(t+\tau).
\end{aligned}$$

Here the final equality follows from

❷ $$\lim_{t\to 0} \measuredangle[q\,{}^{\mathrm{geod}_{[py]}(t)}_z] = \measuredangle[q\,{}^p_z].$$

"$\leqslant$" in ❷ holds by angle semicontinuity (8.14.1). "$\geqslant$" holds by the triangle inequality for angles, since angle comparison (8.3.1c) gives

$$\lim_{t\to 0} \measuredangle[q\,{}^p_{\mathrm{geod}_{[py]}(t)}] = 0.$$

□

8.15 Remarks and open problems

The following question was known in folklore in the 80's, but it seems that in print it was first mentioned in [37]. We do not see any reason why it should be true, but we also cannot construct a counterexample.

8.15.1. Open question. *Let $\mathcal{U}$ be a complete length* CAT(0) *space and $K \subset \mathcal{U}$ be a compact set. Is it true that K lies in a convex compact set $\bar{K} \subset \mathcal{U}$?*

The question can easily be reduced to the case when K is finite; so far it is not even known if any three points in a complete length CAT(0) space lie in a compact convex set.

Chapter 9

Kirszbraun revisited

This chapter is based on our paper [38] and earlier paper of Urs Lang and Viktor Schroeder [39].

9.1 Short map extension definitions.

9.1.1. Theorem. *Acomplete length space $\mathcal{L}$ is* $\mathrm{CBB}(\kappa)$ *if and only if for any 3-point set V_3 and any 4-point set $V_4 \supset V_3$ in $\mathcal{L}$, any short map $f\colon V_3 \to \mathbb{M}^2(\kappa)$ can be extended to a short map $F\colon V_4 \to \mathbb{M}^2(\kappa)$ (so $f = F|_{V_3}$).*

The proof of the "only if" part of Theorem 9.1.1 can be obtained as a corollary of Kirszbraun's theorem (9.4.1). We present another, more elementary proof; it use the following analog of Alexandrov lemma (5.2.1).

We say that two triangles with a common vertex *do not overlap* if their convex hulls intersect only at the common vertex.

9.1.2. Overlap lemma. *Let $[\tilde{x}^1\tilde{x}^2\tilde{x}^3]$ be a triangle in $\mathbb{M}^2(\kappa)$. Let $\tilde{p}^1, \tilde{p}^2, \tilde{p}^3$ be points such that, for any permutation $\{i,j,k\}$ of $\{1,2,3\}$, we have*

(i) $|\tilde{p}^i - \tilde{x}^k| = |\tilde{p}^j - \tilde{x}^k|$,

(ii) $\tilde{p}^i$ *and* $\tilde{x}^i$ *lie in the same closed halfspace determined by* $[\tilde{x}^j\tilde{x}^k]$,

Assume no pair of triangles $[\tilde{p}^i\tilde{x}^j\tilde{x}^k]$ overlap then

$$\measuredangle\tilde{p}^1 + \measuredangle\tilde{p}^2 + \measuredangle\tilde{p}^3 > 2\cdot\pi,$$

where $\measuredangle\tilde{p}^i$ denotes $\measuredangle[\tilde{p}^i\,{}^{\tilde{x}^k}_{\tilde{x}^j}]$ for a permutation $\{i,j,k\}$ of $\{1,2,3\}$.



Remarks. If $\kappa \leqslant 0$, then overlap lemma can be proved without using condition (i). This follows immediately from the formula that relates the sum of angles for the hexagon $[\tilde{p}^1\tilde{x}^2\tilde{p}^3\tilde{x}^1\tilde{p}^2\tilde{x}^3]$ and its area:

$$\measuredangle\tilde{p}^1 - \measuredangle\tilde{x}^2 + \measuredangle\tilde{p}^3 - \measuredangle\tilde{x}^1 + \measuredangle\tilde{p}^2 - \measuredangle\tilde{x}^3 = 2\cdot\pi - \kappa\cdot\text{area}.$$

The diagram shows that condition (i) is essential in case $\kappa > 0$.

Proof. Rotate the triangle $[\tilde{p}^3\tilde{x}^1\tilde{x}^2]$ around $\tilde{x}^1$ to make $[\tilde{x}^1\tilde{p}^3]$ coincide with $[\tilde{x}^1\tilde{p}^2]$. Let $\dot{x}^2$ denote the image of $\tilde{x}^2$ after rotation. Note that

$$\measuredangle[\tilde{x}^1\,{}^{\tilde{x}^3}_{\dot{x}^2}] = \min\{\,\measuredangle[\tilde{x}^1\,{}^{\tilde{x}^2}_{\tilde{p}^3}] + \measuredangle[\tilde{x}^1\,{}^{\tilde{p}^2}_{\tilde{x}^3}],\ 2\cdot\pi - (\measuredangle[\tilde{x}^1\,{}^{\tilde{x}^2}_{\tilde{p}^3}] + \measuredangle[\tilde{x}^1\,{}^{\tilde{p}^2}_{\tilde{x}^3}])\,\}.$$

By (ii), the triangles $[\tilde{p}^3\tilde{x}^1\tilde{x}^2]$ and $[\tilde{p}^2\tilde{x}^3\tilde{x}^1]$ do not overlap if and only if

❶ $$2\cdot\pi > \measuredangle[\tilde{x}^1\,{}^{\tilde{x}^2}_{\tilde{p}^3}] + \measuredangle[\tilde{x}^1\,{}^{\tilde{p}^2}_{\tilde{x}^3}] + \measuredangle[\tilde{x}^1\,{}^{\tilde{x}^2}_{\tilde{x}^3}]$$

and

❷ $$\measuredangle[\tilde{x}^1\,{}^{\tilde{x}^3}_{\tilde{x}^2}] > \measuredangle[\tilde{x}^1\,{}^{\tilde{x}^3}_{\dot{x}^2}].$$

The condition ❷ holds if and only if $|\tilde{x}^2 - \tilde{x}^3| > |\dot{x}^2 - \tilde{x}^3|$, which in turn holds if and only if

❸ $$\begin{aligned}\measuredangle\tilde{p}^1 &> \measuredangle[\tilde{p}^2\,{}^{\tilde{x}^3}_{\dot{x}^2}]\\ &= \min\{\measuredangle\tilde{p}^3 + \measuredangle\tilde{p}^2, 2\cdot\pi - (\measuredangle\tilde{p}^3 + \measuredangle\tilde{p}^2)\}.\end{aligned}$$

The inequality follows since the corresponding hinges have the same pairs of sidelengths. (The two pictures show that both possibilities for the minimum can occur.)



Now assume $\measuredangle\tilde{p}^1 + \measuredangle\tilde{p}^2 + \measuredangle\tilde{p}^3 \leqslant 2\cdot\pi$. Then ❸ implies

$$\measuredangle\tilde{p}^i > \measuredangle\tilde{p}^j + \measuredangle\tilde{p}^k.$$

Since no pair of triangles overlap, the same holds for any permutation (i,j,k) of $(1,2,3)$. Therefore

$$\measuredangle\tilde p^1+\measuredangle\tilde p^2+\measuredangle\tilde p^3>2\cdot(\measuredangle\tilde p^1+\measuredangle\tilde p^2+\measuredangle\tilde p^3),$$

a contradiction. □

Proof of 9.1.1; "if" part. Assume $\mathcal{L}$ is geodesic. Take $x^1,x^2,x^3\in\mathcal{L}$ so the model triangle $[\tilde x^1\tilde x^2\tilde x^3]=\tilde\triangle^\kappa(x^1x^2x^3)$ is defined. Choose $p\in\in\,]x^1x^2[$; apply Kirszbrun property for $V_3=\{x^1,x^2,x^3\}$ and $V_4=\{x^1,x^2,x^3,p\}$ and the map $f(x^i)=\tilde x^i$. You obtain point-on-side comparison (7.3.1*b*).

In case $\mathcal{L}$ is not geodesic, pass to its ultrapower $\mathcal{L}^\omega$. Note that the short map extension property survives for $\mathcal{L}^\omega$ and recall that $\mathcal{L}^\omega$ is geodesic (see 3.5.2). Thus, from above, $\mathcal{L}^\omega$ is a complete length CBB(κ) space. By Proposition 7.1.7, $\mathcal{L}$ is a complete length CBB(κ) space.

"Only if" part. Assume the contrary; that is, $x^1,x^2,x^3,p\in\mathcal{L}$, and $\tilde x^1,\tilde x^2,\tilde x^3\in\mathbb{M}^2(\kappa)$ are such that $|\tilde x^i-\tilde x^j|\leqslant|x^i-x^j|$ for all i,j and there is no point $\tilde p\in\mathbb{M}^2(\kappa)$ such that $|\tilde p-\tilde x^i|\leqslant|p-x^i|$ for all i.

Note that in this case all comparison triangles $\tilde\triangle^\kappa(px^ix^j)$ are defined. That is always true if $\kappa\leqslant0$. If $\kappa>0$, and say $\tilde\triangle^\kappa(px^1x^2)$ is undefined, then

$$\begin{aligned}|p-x^1|+|p-x^2|&\geqslant2\cdot\varpi^\kappa-|x^1-x^2|\geqslant\\&\geqslant2\cdot\varpi^\kappa-|\tilde x^1-\tilde x^2|\geqslant\\&\geqslant|\tilde x^1-\tilde x^3|+|\tilde x^2-\tilde x^3|.\end{aligned}$$

It follows that we can take $\tilde p$ on $[\tilde x^1\tilde x^3]$ or $[\tilde x^2\tilde x^3]$.

For each $i\in\{1,2,3\}$, consider a point $\tilde p^i\in\mathbb{M}^2(\kappa)$ such that $|\tilde p^i-\tilde x^i|$ is minimal among points satisfying $|\tilde p^i-\tilde x^j|\leqslant|p-x^j|$ for all $j\neq i$. Clearly, every $\tilde p^i$ is inside the triangle $[\tilde x^1\tilde x^2\tilde x^3]$ (that is, in $\operatorname{Conv}(\tilde x^1,\tilde x^2,\tilde x^3)$), and $|\tilde p^i-\tilde x^i|>|p-x^i|$ for each i. Since the function $x\mapsto\tilde\measuredangle^\kappa\{x;a,b\}$ is increasing, it follows that

(i) $|\tilde p^i-\tilde x^j|=|p-x^j|$ for $i\neq j$;

(ii) no pair of triangles from $[\tilde p^1\tilde x^2\tilde x^3]$, $[\tilde p^2\tilde x^3\tilde x^1]$, $[\tilde p^3\tilde x^1\tilde x^2]$ overlap in $[\tilde x^1\tilde x^2\tilde x^3]$.

As follows from the overlap lemma (9.1.2), in this case

$$\measuredangle[\tilde p^1\,{}^{\tilde x^2}_{\tilde x^3}]+\measuredangle[\tilde p^2\,{}^{\tilde x^3}_{\tilde x^1}]+\measuredangle[\tilde p^3\,{}^{\tilde x^1}_{\tilde x^2}]>2\cdot\pi.$$

Since $|\tilde x^i-\tilde x^j|\leqslant|x^i-x^j|$ we get

$$\measuredangle[\tilde p^k\,{}^{\tilde x^i}_{\tilde x^j}]\leqslant\tilde\measuredangle^\kappa(p\,{}^{x^i}_{x^j})$$

if (i,j,k) is a permutation of $(1,2,3)$. Therefore

$$\measuredangle^{\kappa}(p\,{}^{x^1}_{x^2}) + \measuredangle^{\kappa}(p\,{}^{x^2}_{x^3}) + \measuredangle^{\kappa}(p\,{}^{x^3}_{x^1}) > 2\cdot\pi;$$

the later contradicts CBB(κ) comparison (7.1.2). $\square$

9.1.3. Theorem. *Assume any pair of points at distance $< \varpi^{\kappa}$ in the metric space $\mathcal{U}$ are joined by a unique geodesic. Then $\mathcal{U}$ is* CAT(κ) *if and only if for any 3-point set V_3 with perimeter $< 2\cdot\varpi^{\kappa}$ and any 4-point set $V_4 \supset V_3$ in $\mathbb{M}^2(\kappa)$, any short map $f\colon V_3 \to \mathcal{U}$ can be extended to a short map $F\colon V_4 \to \mathcal{U}$.*

Note that the "only if" part of Theorem 9.1.3 does not follow directly from Kirszbraun's theorem, since the desired extension is in $\mathcal{U}$ — not its completion.

9.1.4. Lemma. *Let $x^1, x^2, x^3, y^1, y^2, y^3 \in \mathbb{M}(\kappa)$ be points such that $|x^i - x^j| \geqslant |y^i - y^j|$ for all i,j. Then there is a short map $\Phi\colon \mathbb{M}(\kappa) \to \mathbb{M}(\kappa)$ such that $\Phi(x^i) = y^i$ for all i; moreover, one can choose Φ so that*

$$\operatorname{Im}\Phi \subset \operatorname{Conv}(y^1, y^2, y^3).$$

We only give an idea of the proof of this lemma; alternatively, one can get the result as a corollary of Kirszbraun's theorem (9.4.1)

Idea of the proof. The map Φ can be constructed as a composition of an isometry of $\mathbb{M}(\kappa)$ and the following folding maps: Given a halfspace H in $\mathbb{M}(\kappa)$, consider the map $\mathbb{M}(\kappa) \to H$, that is identity on H and reflects all points outside of H into H. This map is a path isometry, in particular it is short.

The last part of the lemma can be proved by composing the constructed map with foldings along the sides of triangle $[y^1y^2y^3]$ and passing to a partial limit. $\square$

Proof of 9.1.3; "if" part. The point-on-side comparison (8.3.1b) follows by taking $V_3 = \{\tilde{x}, \tilde{y}, \tilde{p}\}$ and $V_4 = \{\tilde{x}, \tilde{y}, \tilde{p}, \tilde{z}\}$ where $z \in\,]xy[$. It is only necessary to observe that $F(\tilde{z}) = z$ by uniqueness of $[xy]$.

"Only if" part. Let $V_3 = \{\tilde{x}^1, \tilde{x}^2, \tilde{x}^3\}$ and $V_4 = \{\tilde{x}^1, \tilde{x}^2, \tilde{x}^3, \tilde{p}\}$.

Set $y^i = f(\tilde{x}^i)$ for all i; we need to find a point $q \in \mathcal{U}$ such that $|y^i - q| \leqslant |\tilde{x}^i - \tilde{p}|$ for all i.

Let D be the convex set in $\mathbb{M}^2(\kappa)$ bounded by the model triangle $[\tilde{y}^1\tilde{y}^2\tilde{y}^3] = \tilde{\triangle}^{\kappa}y^1y^2y^3$; that is, $D = \operatorname{Conv}(\tilde{y}^1, \tilde{y}^2, \tilde{y}^3)$.

Note that $|\tilde{y}^i - \tilde{y}^j| = |y^i - y^j| \leqslant |\tilde{x}^i - \tilde{x}^j|$ for all i,j. Applying Lemma 9.1.4, we get a short map $\Phi\colon \mathbb{M}(\kappa) \to D$ such that $\Phi\colon \tilde{x}^i \mapsto \tilde{y}^i$.

Further, by the majorization theorem (8.12.4), there is a short map $F\colon D \to \mathcal{U}$ such that $\tilde{y}^i \mapsto y^i$ for all i.

Thus one can take $q = F \circ \Phi(\tilde{p})$. $\square$

9.1.5. Exercise. *Assume $\mathcal{X}$ is a complete length space that satisfies the following condition: any 4-point subset can be isometrically embedded in Euclidean 3-space.*

Prove that $\mathcal{X}$ is isometric to a closed convex substet of a Hilbert space.

9.1.6. Exercise. *Let $\mathcal{F}_s$ be the metric on the 5 point set $\{p, q, x, y, z\}$ such that $|p - q| = s$ and all the remaining distances are equal 1. Assume $\mathcal{F}_s$ admits a distance preserving map into a complete length* CAT(0) *space. Show that $\mathcal{F}_s$ admits a distance preserving map into a complete length* CBB(0) *space.*

9.2 (1+n)-point comparison

The following theorem gives a more sensitive analog of CBB(κ) comparison (7.1.2).

9.2.1. (1+n)-point comparison. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space. Then for any array of points $p, x^1, \dots, x^n \in \mathcal{L}$ there is a model array $\tilde{p}, \tilde{x}^1, \dots, \tilde{x}^n \in \mathbb{M}^n(\kappa)$ such that*

a) $|\tilde{p} - \tilde{x}^i| = |p - x^i|$ for all i.

b) $|\tilde{x}^i - \tilde{x}^j| \geqslant |x^i - x^j|$ for all i, j.

Proof. It is sufficient to show that given $\varepsilon > 0$ there is a configuration $\tilde{p}, \tilde{x}^1, \dots, \tilde{x}^n \in \mathbb{M}^n(\kappa)$ such that

$$|\tilde{x}^i - \tilde{x}^j| \geqslant |x^i - x^j| \quad \text{and} \quad \big||\tilde{p} - \tilde{x}^i| - |p - x^i|\big| \leqslant \varepsilon.$$

Then one can pass to a limit configuration for $\varepsilon \to 0+$.

According to 7.2.4, the set $\mathrm{Str}(x^1, \dots, x^n)$ is dense in $\mathcal{L}$. Thus, there is a point $p' \in \mathrm{Str}(\tilde{x}^1, \dots, \tilde{x}^n)$ such that $|p' - p| \leqslant \varepsilon$. According to Corollary 11.6.5, $\mathrm{T}_{p'}$ contains a subcone E isometric to a Euclidean space that contains all vectors $\log[p'x^i]$. Passing to subspace if necessary, we can assume that $\dim E \leqslant n$.

Mark a point $\tilde{p} \in \mathbb{M}^n(\kappa)$ and choose a distance preserving map $\iota\colon E \to \mathrm{T}_{\tilde{p}}\mathbb{M}^n(\kappa)$. Set

$$\tilde{x}^i = \exp_{\tilde{p}} \circ \iota(\log[p'x^i]).$$

Thus $|\tilde{p} - \tilde{x}^i| = |p' - x^i|$ and therefore

$$\big||\tilde{p} - \tilde{x}^i| - |p - x^i|\big| \leqslant |p - p'| \leqslant \varepsilon.$$

From the hinge comparison (7.3.1c) we have

$$\measuredangle^{\kappa}(\tilde p\,{}^{\tilde x^i}_{\tilde x^j}) = \measuredangle[\tilde p\,{}^{\tilde x^i}_{\tilde x^j}] = \measuredangle[p'\,{}^{x^i}_{x^j}] \geqslant \measuredangle^{\kappa}(p'\,{}^{x^i}_{x^j}),$$

thus

$$|\tilde x^i - \tilde x^j| \geqslant |x^i - x^j|.$$

□

9.2.2. Exercise. *Let $p, x_1, \dots, x_n$ be a point array $p, x_1, \dots, x_n$ in a* CBB(0) *space. Consider the $n{\times}n$-matrix M with the components*

$$m_{i,j} = \tfrac{1}{2}\cdot(|x_i - p|^2 + |x_j - p|^2 - |x_i - x_j|^2).$$

Show that

❶ $$\boldsymbol{s}\cdot M\cdot \boldsymbol{s}^\top \geqslant 0$$

for any vector $\boldsymbol{s} = (s_1, \dots, s_n)$ with nonnegative components.

The inequality in the above exercise is discovered by Karl-Theodor Sturm [40]. It turns out to be weaker than (1+n)-point comparison. An example can be constructed by perturbing the 6-point metric isometric to a regular pentagon with its center by making its sides slightly longer and diagonals slightly shorter [41].



9.2.3. Exercise. *Give an example of metric on a finite set, that satisfies the comparison inequality*

$$\measuredangle^0(p\,{}^{x_1}_{x_2}) + \measuredangle^0(p\,{}^{x_2}_{x_3}) + \measuredangle^0(p\,{}^{x_3}_{x_1}) \leqslant 2\cdot\pi$$

for any quadruple of points (p, x_1, x_2, x_3), but is not isometric to a subset of Alexandrov space with curvature $\geqslant 0$.

9.2.4. Exercise. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space. Assume that a point array $(a^0, a^1, \dots, a^k)$ in $\mathcal{L}$ is* κ-strutting *for a point $p \in \mathcal{L}$. Show that there are point $\tilde p, \tilde a^0, \dots, \tilde a^m \in \mathbb{M}^{m+1}(\kappa)$ such that*

$$|\tilde p - \tilde a^i| = |p - a^i| \text{ and } |\tilde a^i - \tilde a^j| = |a^i - a^j|$$

for all i and j.

9.3 Helly's theorem

9.3.1. Helly's theorem. *Let $\mathcal{U}$ be a complete length* CAT(0) *space and $\{K_\alpha\}_{\alpha\in\mathcal{A}}$ be an arbitrary collection of closed bounded convex subsets of $\mathcal{U}$.*

If

$$\bigcap_{\alpha\in\mathcal{A}} K_\alpha = \varnothing$$

then there is an index array $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{A}$ such that

$$\bigcap_{i=1}^{n} K_{\alpha_i} = \varnothing.$$

Remarks.

(i) In general, none of K_α might be compact; otherwise the statement is trivial.

(ii) If $\mathcal{U}$ is a Hilbert space (not necessarily separable), then the above result is equivalent to the following statement: if a convex bounded set is closed in ordinary topology then it compact in the weak topology. One can define *weak topology* on arbitrary metric space, by taking exteriors of closed ball as its prebase. Then the result above implies analogous statement for complete length CAT(0) spaces (compare to [42]).

We present the original proof of Urs Lang and Viktor Schroeder from [39].

Proof of 9.3.1. Assume the contrary. Then for any finite set $F \subset \mathcal{A}$

$$K_F := \bigcap_{\alpha\in F} K_\alpha \neq \varnothing,$$

we will construct point z such that $z \in K_\alpha$ for each α. Thus we will arrive to contradiction since

$$\bigcap_{\alpha\in\mathcal{A}} K_\alpha = \varnothing.$$

Choose a point $p \in \mathcal{U}$ and set $r = \sup|K_F - p|$ where F runs all finite subsets of $\mathcal{A}$. Set p_F^* to be the closest point on K_F from p; according to closest-point projection lemma (8.6.3), p_F^* exits and is unique.

Take a nested sequence of finite subsets $F_1 \subset F_2 \subset \dots$ of $\mathcal{A}$, such that $|K_{F_n} - p| \to r$.

Let us show that the sequence $(p^*_{F_n})$ converges in itself. Indeed, if not, then for some fixed $\varepsilon > 0$, we can choose two subsequences (y'_n) and (y''_n) of $(p^*_{F_n})$ such that $|y'_n - y''_n| \geqslant \varepsilon$. Set z_n to be midpoint of $[y'_n y''_n]$. From point-on-side comparison (7.3.1*b*), there is $\delta > 0$ such that

$$|p - z_n| \leqslant \max\{|p - y'_n|, |p - y''_n|\} - \delta.$$

Thus

$$\varlimsup_{n\to\infty} |p - z_n| < r.$$

On the other hand, from convexity, each F_n contains all $z_{\bar{k}}$ with sufficiently large $\bar{k}$, a contradiction.

Thus, $p^*_{F_n}$ converges and we can set $z = \lim_n p^*_{F_n}$. Clearly

$$|p - z| = r.$$

Repeat the above arguments for the sequence $F'_n = F_n \cup \{\alpha\}$. As a result, we get another point z' such that $|p - z| = |p - z'| = r$ and $z, z' \in K_{F_n}$ for all n. Thus, if $z \neq z'$ the midpoint $\hat{z}$ of $[zz']$ would belong to all K_{F_n} and from comparison we would have $|p - \hat{z}| < r$, a contradiction.

Thus, $z' = z$; in particular $z \in K_\alpha$ for each $\alpha \in \mathcal{A}$. □

9.4 Kirszbraun's theorem

A slightly weaker version of the following theorem was proved by Urs Lang and Viktor Schroeder [39].

9.4.1. Kirszbraun's theorem. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space, $\mathcal{U}$ be a complete length* $\mathrm{CAT}(\kappa)$ *space, $Q \subset \mathcal{L}$ be arbitrary subset and $f\colon Q \to \mathcal{U}$ be a short map. Assume that there is $z \in \mathcal{U}$ such that $f(Q) \subset \mathrm{B}[z, \frac{\varpi^\kappa}{2}]_{\mathcal{U}}$. Then $f\colon Q \to \mathcal{U}$ can be extended to a short map $F\colon \mathcal{L} \to \mathcal{U}$ (that is, there is a short map $F\colon \mathcal{L} \to \mathcal{U}$ such that $F|_Q = f$.)*

The condition $f(Q) \subset \mathrm{B}[z, \frac{\varpi^\kappa}{2}]$ trivially holds for any $\kappa \leqslant 0$ since in this case $\varpi^\kappa = \infty$. The following example shows that this condition is needed for $\kappa > 0$.

The Conjecture 9.6.2 (if true) gives an equivalent condition for the existence of a short extension; roughly it states the following example is the only obstacle.

9.4.2. Example. *Let $\mathbb{S}^m_+$ be a closed m-dimensional unit hemisphere. Denote its boundary, which is isometric to $\mathbb{S}^{m-1}$, by $\partial\mathbb{S}^m_+$. Clearly,*

$\mathbb{S}^m_+$ is CBB(1) *and $\partial\mathbb{S}^m_+$ is* CAT(1)*, but the identity map $\partial\mathbb{S}^m_+ \to \partial\mathbb{S}^m_+$ cannot be extended to a short map $\mathbb{S}^m_+ \to \partial\mathbb{S}^m_+$ (there is no place for the pole).*

There is also a direct generalization of this example to a hemisphere in a Hilbert space of arbitrary cardinal dimension.

First we prove this theorem in the case $\kappa \leqslant 0$ (9.4.4). In the proof of the more complicated case $\kappa > 0$, we use the case $\kappa = 0$. The following lemma is the main ingredient in the proof.

9.4.3. Finite+one lemma. *Let $\kappa \leqslant 0$, $\mathcal{L}$ be a complete length* CBB(κ) *space, and $\mathcal{U}$ be a complete length* CAT(κ) *space. Suppose $x^1, x^2, \dots, x^n \in \mathcal{L}$ and $y^1, y^2, \dots, y^n \in \mathcal{U}$ are such that $|x^i - x^j| \geqslant |y^i - y^j|$ for all i, j.*

Then for any $p \in \mathcal{L}$, there is $q \in \mathcal{U}$ such that $|y^i - q| \leqslant |x^i - p|$ for each i.

Proof. It is sufficient to prove the lemma only for $\kappa = 0$ and -1. The proofs of these two cases are identical, only the formulas differ. In the proof, we assume $\kappa = 0$ and provide the formulas for $\kappa = -1$ in the footnotes.

From $(1+n)$-point comparison (9.2.1), there is a model configuration $\tilde{p}, \tilde{x}^1, \tilde{x}^2, \dots, \tilde{x}^n \in \mathbb{M}^n(\kappa)$ such that $|\tilde{p} - \tilde{x}^i| = |p - x^i|$ and $|\tilde{x}^i - \tilde{x}^j| \geqslant |x^i - x^j|$ for all i, j. It follows that we can assume that $\mathcal{L} = \mathbb{M}^n(\kappa)$.

For each i, consider functions $f^i\colon \mathcal{U} \to \mathbb{R}$ and $\tilde{f}^i\colon \mathbb{M}^n(\kappa) \to \mathbb{R}$ defined as follows[1]

$$(A)^0 \qquad f^i = \tfrac{1}{2}\cdot \operatorname{dist}^2_{y^i}\,, \quad \tilde{f}^i = \tfrac{1}{2}\cdot \operatorname{dist}^2_{\tilde{x}^i}\,.$$

Set $\boldsymbol{f} = (f^1, f^2, \dots, f^n)\colon \mathcal{U} \to \mathbb{R}^n$ and $\tilde{\boldsymbol{f}} = (\tilde{f}^1, \tilde{f}^2, \dots, \tilde{f}^n)\colon \mathbb{M}^n(\kappa) \to \mathbb{R}^n$.

Recall that the set $\operatorname{Up}\boldsymbol{f}(\mathcal{U}) \subset \mathbb{R}^n$ is defined on page 167. Note that it is sufficient to prove that $\tilde{\boldsymbol{f}}(\tilde{p}) \in \operatorname{Up}\boldsymbol{f}(\mathcal{U})$.

Clearly, $(f^i)'' \geqslant 1$. Thus, by the theorem on barycentric simplex (12.1.3a), the set $\operatorname{Up}\boldsymbol{f}(\mathcal{U}) \subset \mathbb{R}^n$ is convex.

Arguing by contradiction, let us assume that $\tilde{\boldsymbol{f}}(\tilde{p}) \notin \operatorname{Up}\boldsymbol{f}(\mathcal{U})$.

Then there exists a supporting hyperplane $\alpha_1\cdot x_1 + \dots \alpha_n\cdot x_n = c$ to $\operatorname{Up}\boldsymbol{f}(\mathcal{U})$, separating it from $\tilde{\boldsymbol{f}}(\tilde{p})$. According to Lemma 12.1.6b, $\alpha_i \geqslant 0$ for each i. So we can assume that $(\alpha_1, \alpha_2, \dots, \alpha_n) \in \Delta^{n-1}$

[1] In case $\kappa = -1$,

$$(A)^- \qquad f^i = \cosh\circ\operatorname{dist}_{y^i}\,, \quad \tilde{f}^i = \cosh\circ\operatorname{dist}_{\tilde{x}^i}\,.$$

(that is, $\alpha_i \geqslant 0$ for each i and $\sum \alpha_i = 1$) and

$$\sum_i \alpha_i \cdot \tilde{f}^i(\tilde{p}) < \inf \left\{ \sum_i \alpha_i \cdot f^i(q) : q \in \mathcal{U} \right\}.$$

The latter contradicts the following claim.

❶ *Given* $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \Delta^{n-1}$, *set*

$$h = \sum_i \alpha_i \cdot f^i \qquad h \colon \mathcal{U} \to \mathbb{R} \qquad z = \operatorname{MinPoint} h \in \mathcal{U}$$

$$\tilde{h} = \sum_i \alpha_i \cdot \tilde{f}^i \qquad \tilde{h} \colon \mathbb{M}^n(\kappa) \to \mathbb{R} \qquad \tilde{z} = \operatorname{MinPoint} \tilde{h} \in \mathbb{M}^n(\kappa)$$

Then $h(z) \leqslant \tilde{h}(\tilde{z})$.

Proof of the claim. Note that $\boldsymbol{d}_z h \geqslant 0$. Thus, for each i, we have[2]

$$(B)^0 \qquad \begin{aligned} 0 &\leqslant (\boldsymbol{d}_z h)(\uparrow_{[zy^i]}) = \\ &= -\sum_j \alpha_j \cdot |z - y^j| \cdot \cos \measuredangle[z\,{}^{y^i}_{y^j}] \leqslant \\ &\leqslant -\sum_j \alpha_j \cdot |z - y^j| \cdot \cos \tilde{\measuredangle}^0(z\,{}^{y^i}_{y^j}) = \\ &= -\tfrac{1}{2 \cdot |z - y^i|} \cdot \sum_j \alpha_j \cdot \left[|z - y^i|^2 + |z - y^j|^2 - |y^i - y^j|^2 \right]. \end{aligned}$$

In particular[3],

$$(C)^0 \qquad \sum_i \alpha_i \cdot \left[\sum_j \alpha_j \cdot \left[|z - y^i|^2 + |z - y^j|^2 - |y^i - y^j|^2 \right] \right] \leqslant 0,$$

[2] In case $\kappa = -1$, the same calculations give

$$(B)^- \qquad 0 \leqslant \ldots \leqslant -\tfrac{1}{\sinh |z - y^i|} \cdot \sum_j \alpha_j \cdot \left[\cosh |z - y^i| \cdot \cosh |z - y^j| - \cosh |y^i - y^j| \right].$$

[3] In case $\kappa = -1$, the same calculations give

$$(C)^- \qquad \sum_i \alpha_i \cdot \left[\sum_j \alpha_j \cdot \left[\cosh |z - y^i| \cdot \cosh |z - y^j| - \cosh |y^i - y^j| \right] \right] \leqslant 0.$$

or[4]

$$(D)^0 \qquad 2\cdot h(z) \leqslant \sum_{i,j} \alpha_i\cdot\alpha_j\cdot|y^i-y^j|^2.$$

Note, that if $\mathcal{U} \overset{iso}{=\!=} \mathbb{M}^n(\kappa)$, then all inequalities in (B,C,D) are sharp. Thus the same argument as above, repeated for $\tilde{x}^1,\tilde{x}^2,\dots,\tilde{x}^n \in$ $\in \mathbb{M}^n(\kappa)$ gives[5]

$$(E)^0 \qquad 2\cdot\tilde{h}(\tilde{z}) = \sum_{i,j} \alpha_i\cdot\alpha_j\cdot|\tilde{x}^i-\tilde{x}^j|^2.$$

Note that

$$|\tilde{x}^i-\tilde{x}^j| \geqslant |x^i-x^j| \geqslant |y^i-y^j|$$

for all i, j. Thus, (D) and (E) imply the claim. $\triangle\square$

9.4.4. Kirszbraun's theorem for nonpositive bound. *Let* $\kappa \leqslant$ $\leqslant 0$, $\mathcal{L}$ *be a complete length* CBB(κ) *space,* $\mathcal{U}$ *be a complete length* CAT(κ) *space,* $Q\subset\mathcal{L}$ *be arbitrary subset and* $f\colon Q\to\mathcal{U}$ *be a short map. Then there is a short extension* $F\colon\mathcal{L}\to\mathcal{U}$ *of* f*; that is, there is a short map* $F\colon\mathcal{L}\to\mathcal{U}$ *such that* $F|_Q = f$.

Remark. If $\mathcal{U}$ is proper, then we do not need Helly's theorem (9.3.1) — compactness of closed balls in $\mathcal{U}$ is sufficient in this case.

Proof of 9.4.4. By Zorn's lemma, we can assume that $Q\subset\mathcal{L}$ is a maximal set; that is, $f\colon Q\to\mathcal{U}$ does not admits a short extension to any larger set $Q'\supset Q$.

Let us argue by contradiction. Assume that $Q\neq\mathcal{L}$; choose $p\in$ $\in\mathcal{L}\backslash Q$. Then

$$\bigcap_{x\in Q}\overline{\mathrm{B}}[f(x),|p-x|]=\varnothing.$$

Since $\kappa\leqslant 0$, the balls are convex; thus, by Helly's theorem (9.3.1), one can choose a point array $x^1,x^2,\dots,x^n\in Q$ such that

$$❷ \qquad \bigcap_{i=1}^{n}\overline{\mathrm{B}}[y^i,|x^i-p|]=\varnothing,$$

---

[4]In case $\kappa=-1$,

$$(D)^- \qquad (h(z))^2 \leqslant \sum_{i,j} \alpha_i\cdot\alpha_j\cdot\cosh|y^i-y^j|.$$

[5]In case $\kappa=-1$,

$$(E)^- \qquad (\tilde{h}(\tilde{z}))^2 = \sum_{i,j} \alpha_i\cdot\alpha_j\cdot\cosh|\tilde{x}^i-\tilde{x}^j|.$$

where $y^i = f(x^i)$. Finally note that ❷ contradicts the Finite+one lemma (9.4.3) $\square$

Proof of Kirszbraun's theorem (9.4.1). The case $\kappa \leqslant 0$ is already proved in 9.4.4. Thus it remains to prove the theorem only in case $\kappa > 0$. After rescaling we may assume that $\kappa = 1$ and therefore $\varpi^\kappa = \pi$.

Since $\overline{\mathrm{B}}[z, \pi/2]_{\mathcal{U}}$ is a complete length CAT(κ) space, we can assume $\mathcal{U} = \overline{\mathrm{B}}[z, \pi/2]_{\mathcal{U}}$. In particular, $\operatorname{diam}\mathcal{U} \leqslant \pi$.

Further, any two points $x, y \in \mathcal{U}$ such that $|x - y| < \pi$ are joined by a unique geodesic; if $|x - y| = \pi$, then the concatenation of $[xz]$ and $[zy]$ as a geodesic from x to y. Hence $\mathcal{U}$ is geodesic.

We can also assume that $\operatorname{diam}\mathcal{L} \leqslant \pi$. Otherwise $\mathcal{L}$ is one-dimensional (see 7.8.1); in this case the result follows since $\mathcal{U}$ is geodesic.

Assume the theorem is false. Then there is a set $Q \subset \mathcal{L}$, a short map $f\colon Q \to \mathcal{U}$ and $p \in \mathcal{L}\backslash Q$ such that

❸ $$\bigcap_{x\in Q} \overline{\mathrm{B}}[f(x), |x - p|] = \varnothing.$$

We will apply 9.4.4 for $\kappa = 0$ to Euclidean cones $\mathring{\mathcal{L}} = \operatorname{Cone}\mathcal{L}$ and $\mathring{\mathcal{U}} = \operatorname{Cone}\mathcal{U}$. Note that

- ⋄ $\mathring{\mathcal{U}}$ is a complete length CAT(0) space,
- ⋄ since $\operatorname{diam}\mathcal{L} \leqslant \pi$ we have $\mathring{\mathcal{L}}$ is CBB(0).

Further, we will view spaces $\mathcal{L}$ and $\mathcal{U}$ as unit spheres in $\mathring{\mathcal{L}}$ and $\mathring{\mathcal{U}}$ respectively. In the cones $\mathring{\mathcal{L}}$ and $\mathring{\mathcal{U}}$ we will use "$|{*}|$" for distance to the vertex, denoted by 0, "$\cdot$" for cone multiplication, "$\measuredangle(x, y)$" for $\measuredangle[o\,{}^x_y]$ and "$\langle x, y\rangle$" for $|x|\cdot|y|\cdot\cos\measuredangle[o\,{}^x_y]$. In particular,

- ⋄ $|x - y|_{\mathcal{L}} = \measuredangle(x, y)$ for any $x, y \in \mathcal{L}$,
- ⋄ $|x - y|_{\mathcal{U}} = \measuredangle(x, y)$ for any $x, y \in \mathcal{U}$,
- ⋄ for any $y \in \mathcal{U}$, we have

❹ $$\measuredangle(z, y) \leqslant \tfrac{\pi}{2}.$$

Set $\mathring{Q} = \operatorname{Cone} Q \subset \mathring{\mathcal{L}}$ and let $\mathring{f}\colon \mathring{Q} \to \mathring{\mathcal{U}}$ be the natural cone extension of f; that is, $y = f(x) \Rightarrow t{\cdot}y = \mathring{f}(t{\cdot}x)$ for $t \geqslant 0$. Clearly $\mathring{f}$ is short.

Applying 9.4.4 for $\mathring{f}$, we get a short extension map $\mathring{F}\colon \mathring{\mathcal{L}} \to \mathring{\mathcal{U}}$. Set $s = \mathring{F}(p)$. Thus,

❺ $$|s - \mathring{f}(w)| \leqslant |p - w|$$

for any $w \in \mathring{Q}$. In particular, $|s| \leqslant 1$. Applying ❺ for $w = t{\cdot}x$ and $t \to \infty$ we get

❻ $$\langle f(x), s\rangle \geqslant \cos\measuredangle(p,x)$$

for any $x \in Q$.

By comparison, the geodesics $\mathrm{geod}_{[s\ t\cdot z]}$ converge as $t \to \infty$ and its limit is a ray; denote it by $\alpha\colon [0,\infty) \to \mathring{\mathcal{U}}$. From ❹, we have that the function $t \mapsto \langle f(x), \alpha(t)\rangle$ is nondecreasing. From ❻, for the necessarily unique point $\bar s$ on the ray α such that $|\bar s| = 1$ we also have



$$\langle f(x), \bar s\rangle \geqslant \cos\measuredangle(p,x)$$

or

$$\measuredangle(\bar s, f(x)) \leqslant \measuredangle(p, f(x))$$

for any $x \in Q$. The latter contradicts ❸. □

9.4.5. Exercise. *Let $\mathcal{U}$ is* CAT(0) *Assume there are two point arrays $x^0, x^1, \dots, x^k \in \mathcal{U}$ and $\tilde x^0, \tilde x^1, \dots, \tilde x^k \in \mathbb{E}^m$ such that $|x^i - x^j|_{\mathcal{U}} = |\tilde x^i - \tilde x^j|_{\mathbb{E}^m}$ for each i, j and for any point $z_0 \in \mathcal{U}$ there is $i > 0$ such that have $|z_0 - x_i| \geqslant |x_0 - x_i|$.*

Prove that there is a subset $Q \subset \mathcal{L}$ isometric to a convex set in $\mathbb{E}^m$ and containing all points x^i.

9.4.6. Exercise. *Let $\mathcal{L}$ be a complete length* CBB(0) *space, $x^0, x^1, \dots, x^k \in \mathcal{L}$ and $\tilde x^0, \tilde x^1, \dots, \tilde x^k \in \mathbb{E}^m$ be two point arrays such that $|x^i - x^j|_{\mathcal{L}} = |\tilde x^i - \tilde x^j|_{\mathbb{E}^m}$ for each i, j. Assume $\tilde x^0$ lies in the interior of* $\mathrm{Conv}(\tilde x^1, \dots, \tilde x^k)$.

Prove that there is a subset $Q \subset \mathcal{L}$ isometric to a convex set in $\mathbb{E}^m$ containing all points x^i.

9.4.7. Exercise. *Construct a three-dimensional complete length* CBB(0) *space $\mathcal{L}$ with a triangle $[xyz]$ such that all angles in $[xyz]$ are equal to the corresponding angles of its model triangle, but $[xyz]$ can not be filled by an isometric copy of the model solid triangle.*

The following statement we call $(2n{+}2)$-point comparison.

9.4.8. Exercise. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space. Consider $x, y \in \mathcal{U}$ and an array of pairs of points (p^1, q^1), $(p^2, q^2), \dots, (p^n, q^n)$ in $\mathcal{U}$, such that there is a model configuration $\tilde x$, $\tilde y$ and array of pairs $(\tilde p^1, \tilde q^1)$, $(\tilde p^2, \tilde q^2), \dots, (\tilde p^n, \tilde q^n)$ in $\mathbb{M}^3(\kappa)$ with the following properties:*

a) $[\tilde x \tilde p^1 \tilde q^1] = \tilde\triangle^\kappa x p^1 q^1$ *and* $[\tilde y \tilde p^n \tilde q^n] = \tilde\triangle^\kappa y p^n q^n$*;*

b) The simplex $\tilde p^i\tilde p^{i+1}\tilde q^i\tilde q^{i+1}$ is a model simplex[6] *of $p^ip^{i+1}q^iq^{i+1}$ for all i.*

Then for any choice of n points $\tilde z^i\in[\tilde p^i\tilde q^i]$, we have

$$|\tilde x-\tilde z^1|+|\tilde z^1-\tilde z^2|+\ldots+|\tilde z^{n-1}-\tilde z^n|+|\tilde z^n-\tilde y|\geqslant|x-y|.$$



9.4.9. Exercise. *Assume that a Riemannian manifold $\mathcal R$ satisfies the following condition: for an arbitrary subset $Q\subset\mathcal R$, any short map $Q\to\mathcal R$ can be extended to a short map $\mathcal R\to\mathcal R$. Show that $\mathcal R$ has constant sectional curvature.*

9.5 Curvature free

In this section we present a collection of exercises on a curvature-free analogs of Kirszbraun theorem.

The following exercise gives an analog of finite+one lemma (9.4.3) discovered by the second author and Stephan Stadler [43].

9.5.1. Exercise. *Let X and Y be metric spaces, $A\subset X$ and $f\colon A\to Y$ a short map. Assume Y is compact and for any finite set $V\subset X$ there is a short map $V\to Y$ that agrees with f in $V\cap A$. Then there is a short map $X\to Y$ that agrees with f in A.*

The following statement was first observed by John Isbell [44].

9.5.2. Exercise. *We say that a space $\mathcal X$ is* injective metric space *if for an arbitrary metric space $\mathcal Z$ and a subset $Q\subset\mathcal Z$, any short map $Q\to\mathcal X$ can be extended as a short map $\mathcal Z\to\mathcal X$.*

1. *Prove that any metric space $\mathcal X$ can be isometrically embedded into an injective metric space.*

[6] that is, perimeter of each triple in p^i,p^{i+1},q^i and q^{i+1} is $<2\cdot\pi$ and $|\tilde p^i-\tilde q^i|=|p^i-q^i|$, $|\tilde p^i-\tilde p^{i+1}|=|p^i-p^{i+1}|$, $|\tilde q^i-\tilde q^{i+1}|=|q^i-q^{i+1}|$, $|\tilde p^i-\tilde q^{i+1}|=|p^i-q^{i+1}|$ and $|\tilde p^{i+1}-\tilde q^i|=|p^{i+1}-q^i|$.

2. *Use it to construct an analog of convex hull in the category of metric spaces; this is called* injective hull.

9.5.3. Exercise. *We say that a compact space $\mathcal{X}$ is a* Kirszbrun source *if for an arbitrary complete metric space $\mathcal{Z}$ and subset $Q \subset \mathcal{X}$, any short map $Q \to \mathcal{Z}$ can be extended to a short map $\mathcal{X} \to \mathcal{Z}$. Prove that a metric space $\mathcal{X}$ is a Kirszbrun source if and only if it satisfies ultratriangle inequality for all triples of points; that is,*

$$|x - z| \leqslant \max\{|x - y|, |y - z|\}$$

for any $x, y, z \in \mathcal{X}$.

9.6 Remarks and open problems

9.6.1. Open problem. *Find a necessary and sufficient condition for a finite metric space to be isometrically embeddable into*

a) some length CBB(κ) *space.*

b) some length CAT(κ) *space.*

A metric on a finite set $\{a^1, a^2, \dots, a^n\}$, can be described by the matrix with components

$$s^{ij} = |a^i - a^j|^2,$$

which we will call the *decrypting matrix*. The set of decrypting matrices of all metrics that admit a distance preserving map into a CBB(0) or a CAT(0) space form a convex cone. The latter follows since the rescalings and products of CBB(0) (or CAT(0)) spaces are CBB(0) (or CAT(0) correspondingly). This convexity gives hope that the cone admits an explicit description. At the moment only some necessary and sufficient conditions are known; for more on this see our paper [38] and the paper of Nina Lebedeva, Vladimir Zolotov and the second author [41].

The following conjecture (if true) would give the right generality for Kirszbraun's theorem (9.4.1). Roughly it states that the example 9.4.2, is the only obstacle for extending short map.

9.6.2. Conjecture. *Assume $\mathcal{L}$ is a complete length* CBB(1) *space, $\mathcal{U}$ is a complete length* CAT(1) *space, $Q \subset \mathcal{L}$ is a proper subset and $f\colon Q \to \mathcal{U}$ be a short map that does not admit a short extension to any bigger set $Q' \supset Q$. Then*

a) Q is isometric to a sphere in a Hilbert space (of finite or cardinal dimension). Moreover, there is a point $p \in \mathcal{L}$ such that $|p - q| = \frac{\pi}{2}$ for any $q \in Q$.

b) The map $f\colon Q \to \mathcal{U}$ is a distance preserving map and there is no point $p' \in \mathcal{U}$ such that $|p'-q'| = \frac{\pi}{2}$ for any $q' \in f(Q)$.

Chapter 10

Warped products

The warped product is a construction that produces a new metric space, denoted by $\mathcal{B} \times_f \mathcal{F}$, from two metric spaces $\mathcal{B}$, $\mathcal{F}$ and a function $f\colon \mathcal{B} \to \mathbb{R}_{\geqslant 0}$.

Many important constructions such as direct product, cone, spherical suspension and join can be defined using warped products.

10.1 Definition

First we define the warped product for length spaces and then we expand the definition to allow arbitrary metric spaces $\mathcal{F}$.

Let $\mathcal{B}$ and $\mathcal{F}$ be length spaces and $f\colon \mathcal{B} \to [0,\infty)$ be a continuous function.

For any path $\gamma\colon [0,1] \to \mathcal{B} \times \mathcal{F}$, we write $\gamma = (\gamma_{\mathcal{B}}, \gamma_{\mathcal{F}})$ where $\gamma_{\mathcal{B}}$ is the projection of γ to $\mathcal{B}$, and $\gamma_{\mathcal{F}}$ is the projection to $\mathcal{F}$. If $\gamma_{\mathcal{B}}$ and $\gamma_{\mathcal{F}}$ are Lipschitz, set

❶ $$\operatorname{length}_f \gamma := \int\limits_0^1 \sqrt{v_{\mathcal{B}}^2 + (f \circ \gamma_{\mathcal{B}})^2 \cdot v_{\mathcal{F}}^2} \cdot dt,$$

where $\int$ is Lebesgue integral, $v_{\mathcal{B}}$ and $v_{\mathcal{F}}$ are the speeds of $\gamma_{\mathcal{B}}$ and $\gamma_{\mathcal{F}}$ respectively.

Note that $\operatorname{length}_f \gamma \geqslant \operatorname{length} \gamma_{\mathcal{B}}$. The integral in ❶ can be broken into the sum of two parts: one for the restriction of $\operatorname{length}_f \gamma$ as in ❶ to the nonzero set of $f \circ \gamma_{\mathcal{B}}$, and the other for the length of $\gamma_{\mathcal{B}}$ restricted to the zero set.

Consider the pseudometric on $\mathcal{B} \times_f \mathcal{F}$ defined by

$$|x - y| := \inf \left\{ \operatorname{length}_f \gamma : \gamma(0) = x, \gamma(1) = y \right\}$$

where the exact lower bound is taken for all Lipschitz paths $\gamma\colon [0,1] \to \to \mathcal{B}\times\mathcal{F}$. The corresponding metric space is called the *warped product with base $\mathcal{B}$, fiber $\mathcal{F}$ and warping function f*; it will be denoted by $\mathcal{B} \times_f \mathcal{F}$.

The points in $\mathcal{B} \times_f \mathcal{F}$ can be described by corresponding pairs $(p,\varphi) \in \mathcal{B} \times \mathcal{F}$; note that if $f(p) = 0$ for some $p \in \mathcal{B}$, then $(p,\varphi) = = (p,\psi)$ for any $\varphi,\psi \in \mathcal{F}$.

Note we are not claiming that every Lipschitz curve in $\mathcal{B} \times_f \mathcal{F}$ may be reparametrized as the image of a Lipschitz curve in $\mathcal{B} \times \mathcal{F}$; in fact it is not true.

10.1.1. Proposition. *The warped product $\mathcal{B} \times_f \mathcal{F}$ satisfies:*

a) The projection $(p,\varphi_0) \mapsto p$ is a submetry which when restricted to any horizontal leaf $\mathcal{B} \times \{\varphi_0\}$ is an isometry to $\mathcal{B}$.

b) If $f(p_0) \neq 0$, the projection $(p_0,\varphi) \mapsto \varphi$ of the vertical leaf $\{p_0\} \times \mathcal{F}$, with its length metric, is a homothety onto $\mathcal{F}$ with multiplier $\frac{1}{f(p_0)}$.

c) If f achieves its (local) minimum at p_0, then the vertical leaf $\{p_0\} \times \mathcal{F}$, with its length metric, is (locally) isometrically embedded in $\mathcal{B} \times_f \mathcal{F}$.

Proof. Claim (b) follows from the f-length formula ❶.

Also by ❶, the projection of $\mathcal{B} \times_f \mathcal{F}$ onto $\mathcal{B} \times \{\varphi_0\}$ given by $(p,\varphi) \mapsto (p,\varphi_0)$ is length-nonincreasing; hence (a).

The projection $(p,\varphi) \mapsto (p_0,\varphi)$ of a neighborhood of the vertical leaf $\{p_0\} \times \mathcal{F}$ to $\{p_0\} \times \mathcal{F}$ is length-nonincreasing if p_0 is a local minimum point of f. If p_0 is a global minimum point of f, then the same holds for the projection of whole space. Hence (c). □

Note that any horizontal leaf is weakly convex, but does not have to be convex even if $\mathcal{B} \times_f \mathcal{F}$ is a geodesic space, since vanishing of the warping function f allows geodesics to bifurcate into distinct horizontal leaves. For instance, if there is a geodesic with the ends in the zero set

$$Z = \{\, (p,\varphi) \in \mathcal{B} \times_f \mathcal{F}) : f(p) = 0 \,\},$$

then there is a geodesic with the same ends in each horizontal leaf.

10.1.2. Proposition. *Suppose $\mathcal{B}$ and $\mathcal{F}$ are length spaces and $f\colon \mathcal{B} \to \to [0,\infty)$ is a continuous function. Then the warped product $\mathcal{B} \times_f \mathcal{F}$ is a length space.*

Proof. It is sufficient to show that for any $\alpha\colon [0,1] \to \mathcal{B} \times_f \mathcal{F}$ there is a path $\beta\colon [0,1] \to \mathcal{B} \times \mathcal{F}$ such that

$$\operatorname{length} \alpha \geqslant \operatorname{length}_f \beta.$$

If $f\circ\alpha_{\mathcal{B}}(t)>0$ for any t, then the vertical projection $\alpha_{\mathcal{F}}$ is defined. In this case, let $\beta(t)=(\alpha_{\mathcal{B}}(t),\alpha_{\mathcal{F}}(t))\in\mathcal{B}\times\mathcal{F}$; clearly

$$\operatorname{length}\alpha=\operatorname{length}_f\beta.$$

If $f\circ\alpha_{\mathcal{B}}(t_0)=0$ for some t_0, let β be the concatenation of three curves in $\mathcal{B}\times\mathcal{F}$; namely: (1) the horizontal curve $(\alpha_{\mathcal{B}}(t),\varphi)$ for $t\leqslant t_0$, (2) a vertical path in form (s,φ) to (s,ψ) and (3) the horizontal curve $(\alpha_{\mathcal{B}}(t),\psi)$ for $t\geqslant t_0$. By ❶, the f-length of the middle path in the concatenation is vanishing; therefore the f-length of α is can not be smaller than length of $\alpha_{\mathcal{B}}$; that is,

$$\operatorname{length}_f\alpha\geqslant\operatorname{length}\alpha_{\mathcal{B}}=\operatorname{length}_f\beta.$$

The statement follows. □

Distance in a warped product is fiber-independent, in the sense that distances may be calculated by substituting for $\mathcal{F}$ a different length space:

10.1.3. Fiber-independence theorem. *Consider length spaces $\mathcal{B}$, $\mathcal{F}$ and $\check{\mathcal{F}}$, and a locally Lipschitz function $f:\mathcal{B}\to\mathbb{R}_{\geqslant0}$. Assume $p,q\in\mathcal{B}$, $\varphi,\psi\in\mathcal{F}$ and $\check\varphi,\check\psi\in\check{\mathcal{F}}$: Then*

$$|\varphi-\psi|_{\mathcal{F}}\geqslant|\check\varphi-\check\psi|_{\check{\mathcal{F}}}$$
$$\Downarrow$$
$$|(p,\varphi)-(q,\psi)|_{\mathcal{B}\times_f\mathcal{F}}\geqslant|(p,\check\varphi)-(q,\check\psi)|_{\mathcal{B}\times_f\check{\mathcal{F}}}$$

In particular,

$$|(p,\varphi)-(q,\psi)|_{\mathcal{B}\times_f\mathcal{F}}=|(p,0)-(q,\ell)|_{\mathcal{B}\times_f\mathbb{R}},$$

were $\ell=|\varphi-\psi|_{\mathcal{F}}$.

Proof. Let γ be a path in $(\mathcal{B}\times\mathcal{F})$.

Since $|\varphi-\psi|_{\mathcal{F}}\geqslant|\check\varphi-\check\psi|_{\check{\mathcal{F}}}$, there is a Lipschitz path $\gamma_{\check{\mathcal{F}}}$ from $\check\varphi$ to $\check\psi$ in $\check{\mathcal{F}}$ such that

$$(\operatorname{speed}\gamma_{\mathcal{F}})(t)\geqslant(\operatorname{speed}\gamma_{\check{\mathcal{F}}})(t)$$

for almost all $t\in[0,1]$. Consider the path $\check\gamma=(\gamma_{\mathcal{B}},\gamma_{\check{\mathcal{F}}})$ from $(p,\check\varphi)$ to $(q,\check\psi)$ in $\mathcal{B}\times_f\check{\mathcal{F}}$. Clearly

$$\operatorname{length}_f\gamma\geqslant\operatorname{length}_f\check\gamma.$$

□

10.1.4. Exercise. *Let $\mathcal{B}$ and $\mathcal{F}$ be two length spaces and $f,g\colon\mathcal{B}\to\mathbb{R}_{\geqslant}$ be two locally Lipschitz nonnegative functions. Assume $f(b)\leqslant g(b)$ for any $b\in\mathcal{B}$. Show that $\mathcal{B}\times_f\mathcal{F}\leqslant\mathcal{B}\times_g\mathcal{F}$; that is, there is a distance noncontracting map $\mathcal{B}\times_f\mathcal{F}\to\mathcal{B}\times_g\mathcal{F}$.*

10.2 Extended definition

The fiber-independence theorem gives an explicit formula:

$$|(p,\varphi)-(q,\psi)|_{\mathcal{B}\times_f\mathcal{F}}=|(p,0)-(q,|\varphi-\psi|_{\mathcal{F}})|_{\mathcal{B}\times_f\mathbb{R}}$$

for any $(p,\varphi),(q,\psi)\in\mathcal{B}\times\mathcal{F}$. It implies that if $\iota\colon A\to\check{A}$ is an isometry between two subsets $A\subset\mathcal{F}$ and $\check{A}\subset\check{\mathcal{F}}$ in length spaces $\mathcal{F}$ and $\check{\mathcal{F}}$, and $\mathcal{B}$ is a length space, then for any warping function $f\colon\mathcal{B}\to\mathbb{R}_{\geqslant 0}$, the map ι induces an isometry between the sets $\mathcal{B}\times_f A\subset\mathcal{B}\times_f\mathcal{F}$ and $\mathcal{B}\times_f\check{A}\subset\mathcal{B}\times_f\check{\mathcal{F}}$.

The latter observation allows us to define the warped product $\mathcal{B}\times_f\times_f\mathcal{F}$ where the fiber $\mathcal{F}$ does not carry its length metric. Indeed we can use Kuratowsky embedding to realize $\mathcal{F}$ as a subspace in a length space, say $\mathcal{F}'$. Therefore we can take the warped product $\mathcal{B}\times_f\mathcal{F}'$ and identify $\mathcal{B}\times_f\mathcal{F}$ with its subspace consisting of all pairs (b,φ) such that $\varphi\in\mathcal{F}$. According to the Fiber-independence theorem 10.1.3, the resulting space does not depend on the choice of $\mathcal{F}'$.

10.3 Examples

Direct product. The simplest example is the *direct product* $\mathcal{B}\times\times\mathcal{F}$, which could be also written as the warped product $\mathcal{B}\times_1\mathcal{F}$. For $p,q\in\mathcal{B}$ and $\varphi,\psi\in\mathcal{F}$, the direct product metric simplifies to

$$|(p,\varphi)-(q,\psi)|=\sqrt{|p-q|^2+|\varphi-\psi|^2}.$$

This is taken as the defining formula for the direct product of two arbitrary metric spaces $\mathcal{B}$ and $\mathcal{F}$.

Cones. The *Euclidean cone* $\operatorname{Cone}\mathcal{F}$ over a metric space $\mathcal{F}$ can be defined as the warped product $[0,\infty)\times_{\mathrm{id}}\mathcal{F}$. For $s,t\in[0,\infty)$ and $\varphi,\psi\in\mathcal{F}$, the cone metric is given by the cosine rule

$$|(s,\varphi)-(t,\psi)|=\sqrt{s^2+t^2-2\cdot s\cdot t\cdot\cos\alpha},$$

where $\alpha=\max\{\pi,|\varphi-\psi|\}$.

Instead of the Euclidean cosine rule, we may use the cosine rule in $\mathbb{M}^2(\kappa)$:

$$|(s,\varphi)-(t,\psi)|=\tilde{\Upsilon}^{\kappa}\{\alpha;s,t\}.$$

In this way we get κ-*cones* over $\mathcal{F}$, denoted by $\operatorname{Cone}^{\kappa}\mathcal{F}=[0,\infty)\times_{\mathrm{sn}^{\kappa}}\mathcal{F}$ for $\kappa\leqslant 0$ and $\operatorname{Cone}^{\kappa}\mathcal{F}=[0,\varpi^{\kappa}]\times_{\mathrm{sn}^{\kappa}}\mathcal{F}$ for $\kappa>0$.

The 1-cone $\operatorname{Cone}^1 \mathcal{F}$ is also called the *spherical suspension* over $\mathcal{F}$, and is also denoted by $\operatorname{Susp} \mathcal{F}$. That is,

$$\operatorname{Susp} \mathcal{F} = [0, \pi] \times_{\sin} \mathcal{F}.$$

10.3.1. Exercise. *Let $\mathcal{F}$ be a length space and $A \subset \mathcal{F}$. Show that $\operatorname{Cone}^\kappa A$ is convex in $\operatorname{Cone}^\kappa \mathcal{F}$ if and only if A is π-convex in $\mathcal{F}$.*

The elements of the Euclidean cone $\mathcal{K} = \operatorname{Cone} \mathcal{F}$ will often be referred to as *vectors*. The tip of $\mathcal{K}$ is usually denoted by 0 or $0_{\mathcal{K}}$. The *absolute value* $|w|$ of the vector w is defined as $|0 - w|_{\mathcal{K}}$; that is, the distance from w to the tip. The *scalar product* $\langle v, w \rangle$ of two vectors $v, w \in \mathcal{K}$ is defined by

$$\langle v, w \rangle := \big(|v - w|^2_{\mathcal{K}} - |v|^2 - |w|^2\big)/2.$$

Doubling. The doubling space $\mathcal{W}$ of a metric space $\mathcal{V}$ on a closed subset $A \subset \mathcal{V}$ can be also defined as a special type of warped product. Consider the fiber $\mathbb{S}^0$ consisting of two points with distance 2 from each other. Then $\mathcal{W}$ is isometric to the warped product with base $\mathcal{V}$, fiber $\mathbb{S}^0$ and warping function dist_A; that is,

$$\mathcal{W} \overset{iso}{=\!=} \mathcal{V} \times_{\operatorname{dist}_A} \mathbb{S}^0.$$

10.4 1-dimensional base

The following theorems provide conditions for the spaces and functions in a warped product with 1-dimensional base to have curvature bounds. These theorems are baby cases of the characterization of curvature bounds in warped products given in [45, 46].

10.4.1. Theorem.

a) If $\mathcal{L}$ is a complete length CBB(1) *space and* $\operatorname{diam} \mathcal{L} \leqslant \pi$ *then*

$$\begin{aligned} \operatorname{Susp} \mathcal{L} &= [0, \pi] \times_{\sin} \mathcal{L} \quad \textit{is } \mathrm{CBB}(1), \\ \operatorname{Cone} \mathcal{L} &= [0, \infty) \times_{\mathrm{id}} \mathcal{L} \quad \textit{is } \mathrm{CBB}(0), \\ \operatorname{Cone}^{-1} \mathcal{L} &= [0, \infty) \times_{\sinh} \mathcal{L} \quad \textit{is } \mathrm{CBB}(-1). \end{aligned}$$

Moreover the converse also holds in each of the three cases.

b) If $\mathcal{L}$ is a complete length CBB(0) *space then*

$$\begin{aligned} \mathbb{R} \times \mathcal{L} \quad &\textit{is a complete length } \mathrm{CBB}(0) \textit{ space}, \\ \mathbb{R} \times_{\exp} \mathcal{L} \quad &\textit{is a complete length } \mathrm{CBB}(-1) \textit{ space}. \end{aligned}$$

Moreover the converse also holds in each of the two cases.

c) If $\mathcal{L}$ is a complete length CBB(-1) *space, then* $\mathbb{R} \times_{\cosh} \mathcal{L}$ *is a complete length* CBB(-1) *space. Moreover the converse also holds.*

10.4.2. Theorem. *Let $\mathcal{L}$ be a metric space.*

a) If $\mathcal{L}$ is CAT(1) *then*

$$\begin{aligned} \operatorname{Susp}\mathcal{L} &= [0,\pi] \times_{\sin} \mathcal{L} \quad \textit{is } \mathrm{CAT}\textit{(1)},\\ \operatorname{Cone}\mathcal{L} &= [0,\infty) \times_{\mathrm{id}} \mathcal{L} \quad \textit{is } \mathrm{CAT}\textit{(0)},\\ \operatorname{Cone}^{-1}\mathcal{L} &= [0,\infty) \times_{\sinh} \mathcal{L} \quad \textit{is } \mathrm{CAT}\textit{(-1)}. \end{aligned}$$

Moreover, the converse also holds in each of the three cases.

b) If $\mathcal{L}$ is a complete length CAT(0) *space then $\mathbb{R} \times \mathcal{L}$ is* CAT(0) *and $\mathbb{R} \times_{\exp} \mathcal{L}$ is* CAT(-1). *Moreover the converse also holds in each of the two cases.*

c) If $\mathcal{L}$ is CAT(-1) *then $\mathbb{R} \times_{\cosh} \mathcal{L}$ is* CAT(-1). *Moreover the converse also holds.*

In the proof of the above two theorems we will use the following proposition.

10.4.3. Proposition.

a)

$$\begin{aligned} \operatorname{Susp}\mathbb{S}^{m-1} &= [0,\pi] \times_{\sin} \mathbb{S}^{m-1} \overset{iso}{=\!=} \mathbb{S}^m,\\ \operatorname{Cone}\mathbb{S}^{m-1} &= [0,\infty) \times_{\mathrm{id}} \mathbb{S}^{m-1} \overset{iso}{=\!=} \mathbb{E}^m,\\ \operatorname{Cone}^{-1}\mathbb{S}^{m-1} &= [0,\infty) \times_{\sinh} \mathbb{S}^{m-1} \overset{iso}{=\!=} \mathbb{M}^m(-1). \end{aligned}$$

b)

$$\begin{aligned} \mathbb{R}\times\mathbb{E}^{m-1} &\overset{iso}{=\!=} \mathbb{E}^m,\\ \mathbb{R} \times_{\exp} \mathbb{E}^{m-1} &\overset{iso}{=\!=} \mathbb{M}^m(-1). \end{aligned}$$

c)

$$\mathbb{R} \times_{\cosh} \mathbb{M}^{m-1}(-1) \overset{iso}{=\!=} \mathbb{M}^m(-1).$$

The proof is left to the reader.

Proof of 10.4.1. Let us prove the last statement in (a); the remaining statements are similar. Each proof is based on the Fiber-independence theorem 10.1.3 and the corresponding statement in Proposition 10.4.3.

Choose an arbitrary quadruple of points

$$(s,\varphi),(t^1,\varphi^1),(t^2,\varphi^2),(t^3,\varphi^3) \in [0,\infty) \times_{\sinh} \mathcal{L}.$$

Since $\operatorname{diam}\mathcal{L}\leqslant\pi$, $(1{+}n)$-point comparison (9.2.1) provides a quadruple of points $\psi,\psi^1,\psi^2,\psi^3\in\mathbb{S}^3$ such that

$$|\psi-\psi^i|_{\mathbb{S}^3}=|\varphi-\varphi^i|_{\mathcal{L}}$$

and

$$|\psi^i-\psi^j|_{\mathbb{S}^3}\geqslant|\varphi^i-\varphi^j|_{\mathcal{L}}$$

for all i and j.

According to Proposition 10.4.3a,

$$\operatorname{Cone}^{-1}\mathbb{S}^3=[0,\infty)\times_{\sinh}\mathbb{S}^3\overset{iso}{=\!=}\mathbb{M}^4(-1).$$

Consider the quadruple of points

$$(s,\psi),(t^1,\psi^1),(t^2,\psi^2),(t^3,\psi^3)\in\operatorname{Cone}^{-1}\mathbb{S}^3=\mathbb{M}^4(-1).$$

By the Fiber-independence theorem 10.1.3,

$$|(s,\psi)-(t^i,\psi^i)|_{[0,\infty)\times_{\sinh}\mathbb{S}^3}=|(s,\varphi)-(t^i,\varphi^i)|_{[0,\infty)\times_{\sinh}\mathcal{L}}$$

and

$$|(t^i,\psi^i)-(t^j,\psi^j)|_{[0,\infty)\times_{\sinh}\mathbb{S}^3}\geqslant|(t^i\varphi^i)-(t^j,\varphi^j)|_{[0,\infty)\times_{\sinh}\mathcal{L}}$$

for all i and j. Since four points of $\mathbb{M}^4(-1)$ lie in an isometrically embedded copy of $\mathbb{M}^3(-1)$, it remains to apply Exercise 7.1.3. □

10.5 General case

In this section we formulate general results on curvature bounds of warped products proved by the first author and Richard Bishop [45]

10.5.1. Theorem. *Let $\mathcal{B}$ be a complete finite-dimensional* CBB(κ) *length space, and the function $f\colon\mathcal{B}\to\mathbb{R}_{\geqslant}$ satisfy $f''+\kappa\cdot f\leqslant 0$. Denote by $Z\subset\mathcal{B}$ the zero set of f. Suppose $\mathcal{F}$ is a complete finite-dimensional* CBB(κ') *space. Then the warped product $\mathcal{B}\times_f\mathcal{F}$ is* CBB(κ) *in the following two cases:*

a) If $Z=\varnothing$ and

$$\kappa'\geqslant\kappa\cdot f^2(b)$$

for any $b\in\mathcal{B}$.

b) If $Z\neq\varnothing$ and

$$|d_zf|^2\leqslant\kappa'$$

for any $z\in Z$.

10.5.2. Theorem. *Let $\mathcal{B}$ be a complete* $\mathrm{CAT}(\kappa)$ *length space, and the function $f\colon \mathcal{B} \to \mathbb{R}_{\geqslant}$ satisfy $f'' + \kappa\cdot f \geqslant 0$, where f is Lipschitz on bounded sets or B is locally compact. Denote by $Z \subset \mathcal{B}$ the zero set of f. Suppose $\mathcal{F}$ is a complete* $\mathrm{CAT}(\kappa')$ *space. Then the warped product $\mathcal{B} \times_f \mathcal{F}$ is* $\mathrm{CAT}(\kappa)$ *in the following two cases:*

a) If $Z = \varnothing$ and

$$\kappa' \leqslant \kappa\cdot f^2(b)$$

for any $b \in \mathcal{B}$.

b) If $Z \neq \varnothing$,

$$[(d_z f)\uparrow_{[zb]}]^2 \geqslant \kappa'$$

for any minimizing geodesic $[zb]$ from Z to a point $b \in \mathcal{B}$ and

$$\kappa' \leqslant \kappa\cdot f^2(b)$$

for any $b \in \mathcal{B}$ such that $\operatorname{dist}_X b \geqslant \frac{\varpi^\kappa}{2}$.

The proofs are coming.

Part III

Structure and tools

Chapter 11

First order differentiation

11.1 Intrinsic property of tangent space

11.1.1. Theorem. *Let $\mathcal{U}$ be a complete length* $\mathrm{CAT}(\kappa)$ *space and $p \in \mathcal{U}$. Then $\mathrm{T}_p\mathcal{U}$ is a length space.*

Proof. Since $\mathrm{T}_p = \operatorname{Cone}\Sigma_p$, it is sufficient to show that for any hinge $[p\,{}^{x}_{y}]$ such that $\measuredangle[p\,{}^{x}_{y}] < \pi$ and any $\varepsilon > 0$, there is $z \in \mathcal{U}$ such that

❶ $$\measuredangle[p\,{}^{x}_{z}] < \tfrac{1}{2}\cdot\measuredangle[p\,{}^{x}_{y}] + \varepsilon, \quad \measuredangle[p\,{}^{y}_{z}] < \tfrac{1}{2}\cdot\measuredangle[p\,{}^{x}_{y}] + \varepsilon.$$

Fix a small positive number $\delta \ll \varepsilon$. Let $\bar{x} \in\,]px]$ and $\bar{y} \in\,]py]$ denote the points such that $|p - \bar{x}| = |p - \bar{y}| = \delta$. Let z denote the midpoint of $\bar{x}$ and $\bar{y}$.

Since $\delta \ll \varepsilon$ we can assume that

$$\tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{x}}_{\bar{y}}) < \measuredangle[p\,{}^{x}_{y}] + \varepsilon.$$

By Alexandrov's lemma (5.2.1), we have

$$\tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{x}}_{z}) + \tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{y}}_{z}) < \tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{x}}_{\bar{y}}).$$

By construction,

$$\tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{x}}_{z}) = \tilde{\measuredangle}^{\kappa}(p\,{}^{\bar{y}}_{z}).$$

Applying the angle comparison (8.3.1c), we get ❶. □

The following example was constructed by Stephanie Halbeisen [47]. It shows that an analogous statement does not hold for CBB

spaces. If the dimension is finite, such examples do not exist; for proper spaces the question is open, see 11.7.1.

11.1.2. Example. *There is a complete length* CBB *space* $\check{\mathcal{L}}$ *with a point* $p \in \check{\mathcal{L}}$ *such that the space of directions* $\Sigma_p\check{\mathcal{L}}$ *is not a* π*-length space, and therefore the tangent space* $\mathrm{T}_p\check{\mathcal{L}}$ *is not a length space.*

Construction. Let $\mathbb{H}$ be a Hilbert space formed by infinite sequences of real numbers $\boldsymbol{x} = (x_0, x_1, \dots)$ with norm $|\boldsymbol{x}|^2 = \sum_i (x_i)^2$. Fix $\varepsilon = 0.001$ and consider two functions $f, \check{f} : \mathbb{H} \to \mathbb{R}$

$$f(\boldsymbol{x}) = |\boldsymbol{x}|,$$

$$\check{f}(\boldsymbol{x}) = \max\left\{ |\boldsymbol{x}|, \max_{n\geqslant 1}\{(1+\varepsilon)\cdot x_n - \tfrac{1}{n}\}\right\}.$$

Both of these functions are convex and Lipschitz, therefore their graphs in $\mathbb{H}\times\mathbb{R}$ equipped with its length-metric form infinite dimensional Alexandrov spaces, say $\mathcal{L}$ and $\check{\mathcal{L}}$ (this is proved formally in 11.1.3). Set p to be the origin of $\mathbb{H}\times\mathbb{R}$.

Note that $\check{\mathcal{L}}\cap\mathcal{L}$ is starshaped in $\mathbb{H}$ with center at p. Further $\check{\mathcal{L}}\backslash\mathcal{L}$ consist of a countable number of disjoint sets

$$\Omega_n = \left\{ (\boldsymbol{x}, \check{f}(\boldsymbol{x})) \in \check{\mathcal{L}} : (1+\varepsilon)\cdot x_n - \tfrac{1}{n} > |\boldsymbol{x}| \right\}.$$

Note that $|\Omega_n - p| > \frac{1}{n}$ for each n. It follows that for any geodesic $[pq]$ in $\check{\mathcal{L}}$, a small subinterval $[p\bar{q}] \subset [pq]$ coincides with a straight segment in $\mathbb{H}\times\mathbb{R}$, which is also a geodesic in $\mathcal{L}$. Thus we can treat $\Sigma_p\mathcal{L}$ and $\Sigma_p\check{\mathcal{L}}$ as one set, with two angle metrics $\measuredangle$ and $\check{\measuredangle}$; let us denote by $\measuredangle_{\mathbb{H}\times\mathbb{R}}$ the angle in $\mathbb{H}\times\mathbb{R}$.

The space $\mathcal{L}$ is isometric to the Euclidean cone over $\Sigma_p\mathcal{L}$ with vertex at p; $\Sigma_p\mathcal{L}$ is isometric to a sphere in Hilbert space with radius $\frac{1}{\sqrt{2}}$. In particular, $\measuredangle$ is the length-metric of $\measuredangle_{\mathbb{H}\times\mathbb{R}}$ on $\Sigma_p\mathcal{L}$.

Therefore in order to show that $\check{\measuredangle}$ does not define a length-metric on $\Sigma_p\mathcal{L}$ it is sufficient to construct a pair of directions (ξ_+, ξ_-) such that

$$\check{\measuredangle}(\xi_+, \xi_-) < \measuredangle(\xi_+, \xi_-).$$

Set $\boldsymbol{e}_0 = (1, 0, 0, \dots)$, $\boldsymbol{e}_1 = (0, 1, 0, \dots), \dots \in \mathbb{H}$. Consider the following two rays in $\mathbb{H}\times\mathbb{R}$:

$$\gamma_+(t) = \tfrac{t}{\sqrt{2}}\cdot(\boldsymbol{e}_0, 1) \quad\text{and}\quad \gamma_-(t) = \tfrac{t}{\sqrt{2}}\cdot(-\boldsymbol{e}_0, 1), \quad t\in[0,+\infty).$$

They form unit-speed geodesics in both $\mathcal{L}$ and $\check{\mathcal{L}}$. Let $\xi_\pm$ be the directions of $\gamma_\pm$ at p. Denote by σ_n the half-planes in $\mathbb{H}$ spanned by $\boldsymbol{e}_0$ and $\boldsymbol{e}_n$; that is, $\sigma_n = \{\, x\cdot\boldsymbol{e}_0 + y\cdot\boldsymbol{e}_n : y \geqslant 0 \,\}$. Consider a sequence

of 2-dimensional sectors $Q_n = \check{\mathcal{L}} \cap (\sigma_n \times \mathbb{R})$. For each n, the sector Q_n intersects Ω_n and is bounded by two geodesic rays $\gamma_\pm$. Note that $Q_n \xrightarrow[\text{GH}]{} Q$, where Q is a solid Euclidean angle in $\mathbb{E}^2$ with angle measure $\beta < \measuredangle(\xi_+, \xi_-) = \frac{\pi}{\sqrt{2}}$. Indeed, Q_n is isometric to the subset of $\mathbb{E}^3$ described by

$$y \geqslant 0 \quad \text{and} \quad z = \max\left\{\sqrt{x^2+y^2}, (1+\varepsilon)\cdot y - \tfrac{1}{n}\right\}$$

with length-metric. Thus, its limit Q is isometric to the subset of $\mathbb{E}^3$ described by

$$y \geqslant 0 \quad \text{and} \quad z = \max\left\{\sqrt{x^2+y^2}, (1+\varepsilon)\cdot y\right\}$$

with length-metric. In particular, for any $t, \tau \geqslant 0$,

$$\begin{aligned}|\gamma_+(t) - \gamma_-(\tau)|_{\check{\mathcal{L}}} \leqslant \lim_{n\to\infty} |\gamma_+(t) - \gamma_-(\tau)|_{Q_n} = \\ = \tilde{\Upsilon}^0\{\beta; t, \tau\}.\end{aligned}$$

That is, $\check{\measuredangle}(\xi_+, \xi_-) \leqslant \beta < \measuredangle(\xi_+, \xi_-)$. □

11.1.3. Lemma. *Let $\mathbb{H}$ be a Hilbert space, $f\colon \mathbb{H} \to \mathbb{R}$ be a convex Lipschitz function and $S \subset \mathbb{H} \times \mathbb{R}$ be the graph of f equipped with the length-metric. Then S is* CBB(0).

Proof. Recall that for a subset $X \subset \mathbb{H} \times \mathbb{R}$, we will denote by $|* - *|_X$ the length-metric on X.

By Buyalo's theorem, any convex hypersurface in a Euclidean space, equipped with the length-metric, is non-negatively curved. Thus it is sufficient to show that for any 4-point set $\{x_0, x_1, x_2, x_3\} \subset S$, there is a finite-dimensional subspace $E \subset \mathbb{H} \times \mathbb{R}$ such that $\{x_i\} \in E$ and $|x_i - x_j|_{S\cap E}$ is arbitrary close to $|x_i - x_j|_S$.

Clearly $|x_i - x_j|_{S\cap E} \geqslant |x_i - x_j|_S$; thus it is sufficient to show that for given $\varepsilon > 0$ one can choose E, so that

❷. $$|x_i - x_j|_{S\cap E} < |x_i - x_j|_S + \varepsilon$$

For each pair (x_i, x_j), choose a broken line β_{ij} connecting x_i, x_j that lies under S (that is, outside of $\operatorname{Conv} S$) in $\mathbb{H} \times \mathbb{R}$ and has length at most $|x_i - x_j|_S + \varepsilon$. Take E to be the affine hull of all the vertexes in all β_{ij}. Thus,

$$|x_i - x_j|_{S\cap E} \leqslant \operatorname{length} \beta_{ij}$$

and ❷ follows. □

11.1.4. Exercise. *Construct a non-compact complete geodesic* CBB(0) *space that contains no rays.*

11.2 Differential

11.2.1. Exercise. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space and $p, q \in \mathcal{U}$. Assume $|p-q| < \varpi^\kappa$. Show that*

$$(\boldsymbol{d}_q \operatorname{dist}_p)(v) = -\langle \uparrow_{[qp]}, v\rangle.$$

11.2.2. Exercise. *Let $\mathcal{L}$ be a length* CBB(κ) *space and $p, q \in \mathcal{L}$ be distinct points. Assume $q \in \operatorname{Str}(p)$ or $p \in \operatorname{Str}(q)$ Show that*

$$(\boldsymbol{d}_q \operatorname{dist}_p)(v) = -\langle \uparrow_{[qp]}, v\rangle.$$

As it shown in Halbeisen's example (see section 11.1), a CBB space might have a tangent spaces that are not length spaces; thus concavity of the differential $\boldsymbol{d}_p f$ of a semiconcave function f is meaningless. Nevertheless, as the following lemma shows, differential $\boldsymbol{d}_p f$ of a semiconcave function allways satisfy the following weaker property similar to concavity (compare [12, p. 136], [48, p. 4.2]). In the finite dimensional case $\boldsymbol{d}_p f$ is concave.

11.2.3. Lemma. *Let $\mathcal{L}$ be a complete length* CBB *space $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be a locally Lipschitz semiconcave subfunction and $p \in \operatorname{Dom} f$. Then*

$$\sup\{\,(\boldsymbol{d}_p f)(\xi)\,:\,\xi \in \Sigma_p\,\} \geqslant \frac{(\boldsymbol{d}_p f)(u) + (\boldsymbol{d}_p f)(v)}{\sqrt{|u|^2 + 2\cdot\langle u, v\rangle + |v|^2}}$$

for any $u, v \in \mathrm{T}_p$.

Proof of 11.2.3. We can assume that $\alpha = \measuredangle(u,v) > 0$, otherwise the statement is trivial. Moreover, since $\mathrm{T}'_p = \operatorname{Cone}(\Sigma'_p)$ is dense in T_p and $\boldsymbol{d}_p f\colon \mathrm{T}_p \to \mathbb{R}$ is Lipschitz, we can assume that $u, v \in \mathrm{T}'_p$; that is, $\exp_p(t\cdot u)$ and $\exp_p(t\cdot v)$ are defined for all small $t > 0$.

Prepare a model configuration of five points: $\tilde p, \tilde u, \tilde v, \tilde v', \tilde w \in \mathbb{E}^2$ such that



- ◇ $\measuredangle[\tilde p\,{}^{\tilde u}_{\tilde v}] = \alpha$,
- ◇ $|\tilde p - \tilde u| = |u|$,
- ◇ $|\tilde p - \tilde v| = |v|$,
- ◇ $\tilde v'$ lies on the extension of $[\tilde p \tilde v]$ so that $\tilde v$ is midpoint of $[\tilde p \tilde v']$,
- ◇ $\tilde w$ is the midpoint for $\tilde u$ and $\tilde v$.

Note that

$$|\tilde p - \tilde w| = \tfrac{1}{2}\cdot\sqrt{|u|^2 + 2\cdot\langle u, v\rangle + |v|^2}.$$

Assume that $\mathcal{L}$ is geodesic.

For all small $t > 0$, construct points $u_t, v_t, v'_t, w_t \in \mathcal{L}$ the following way:

(a) $v_t = \exp_p(t{\cdot}v)$, $v'_t = \exp_p(t{\cdot}v')$
(b) $u_t = \exp_p(t{\cdot}u)$.
(c) w_t is the midpoint of $[u_t v_t]$.

Clearly $|p-u_t| = t{\cdot}|u|$, $|p-v_t| = t{\cdot}|v|$, $|p-v'_t| = t{\cdot}|v'|$. Since $\measuredangle(u,v)$ is defined, we have $|u_t-v_t| = t{\cdot}|\tilde u-\tilde v|+o(t)$ and $|u_t-v'_t| = t{\cdot}|\tilde u-\tilde v'|+o(t)$ (see Theorem 7.3.1c and Section 7.1).

From point-on-side and hinge comparison (7.3.1b+c), we get that

$$\measuredangle^{\kappa}(v_t\,{}^{p}_{w_t}) \geqslant \measuredangle^{\kappa}(v_t\,{}^{p}_{u_t}) \geqslant \measuredangle[\tilde v\,{}^{\tilde p}_{\tilde u}] + \tfrac{o(t)}{t}$$

and

$$\measuredangle^{\kappa}(v_t\,{}^{v'_t}_{w_t}) \geqslant \measuredangle^{\kappa}(v_t\,{}^{v'_t}_{u_t}) \geqslant \measuredangle[\tilde v\,{}^{\tilde v'}_{\tilde u}] + \tfrac{o(t)}{t}.$$

Clearly, $\measuredangle[\tilde v\,{}^{\tilde p}_{\tilde u}]+\measuredangle[\tilde v\,{}^{\tilde x}_{\tilde u'}] = \pi$. From adjacent angle comparison (7.3.1a), $\measuredangle^{\kappa}(v_t\,{}^{p}_{v_t}) + \measuredangle^{\kappa}(v_t\,{}^{u_t}_{v'_t}) \leqslant \pi$. Hence $\measuredangle^{\kappa}(v_t\,{}^{p}_{w_t}) \to \measuredangle[\tilde v\,{}^{\tilde p}_{\tilde v}]$ as $t\to 0+$ and thus

$$|p-w_t| = t|\tilde p-\tilde w| + o(t).$$

Since f is λ-concave we have

$$\begin{aligned}2{\cdot}f(w_t) &\geqslant f(u_t)+f(v_t)+\tfrac{\lambda}{4}{\cdot}|u_t-v_t|^2 = \\ &= 2{\cdot}f(p)+t{\cdot}[(\boldsymbol{d}_pf)(u)+(\boldsymbol{d}_pf)(v)]+o(t).\end{aligned}$$

Applying λ-concavity of f, we get

$$\text{❶}\qquad (\boldsymbol{d}_pf)(\uparrow_{[pw_t]}) \geqslant \frac{t{\cdot}[(\boldsymbol{d}_pf)(u)+(\boldsymbol{d}_pf)(v)]+o(t)}{2{\cdot}t{\cdot}|\tilde p-\tilde w|+o(t)}$$

and the lemma follows.

Finally, if $\mathcal{L}$ is not geodesic one needs to make two adjustments in the above constructions. Namely:

(i) For geodesic $[u_tv_t]$ to be defined, one has to take in (b) $u_t\in\operatorname{Str}(v_t)$, $u_t\approx\exp_p(t{\cdot}u)$. More precicely,

$$|u_t-\exp_p(t{\cdot}u)| = o(t).$$

Thus instead of $|p-u_t| = t{\cdot}|u|$ we have

$$|p-u_t| = t|u|+o(t),$$

but it is sufficient for the rest of proof.

(ii) The direction $\uparrow_{[pw_t]}$ might be undefined. Thus, in the estimate ❶, instead of $\uparrow_{[pw_t]}$, one should take $\uparrow_{[pw'_t]}$ for some point $w'_t\in\operatorname{Str}(p)$ near w_t (that is, $|w_t-w'_t| = o(t)$) □

11.3 Ultratangent space

The following theorem is often used together with the observation that ultralimit of any sequence of length spaces is geodesic (see 3.3.2).

11.3.1. Theorem.

a) If $\mathcal{L}$ is a CBB(κ) *space and $p \in \mathcal{L}$, then T_p^ω is* CBB(0).

b) If $\mathcal{U}$ is a CAT(κ) *space and $p \in \mathcal{U}$, then T_p^ω is* CAT(0).

The proofs of both parts are nearly identical.

Proof; (a). Since $\mathcal{L}$ is a complete length CBB(κ) space, then for its blowup $n{\cdot}\mathcal{L}$ is a complete length CBB(κ/n^2) space. By Proposition 7.1.7, $\omega{\cdot}\mathcal{L}$ is CBB(0) and so is T_p^ω as a metric component of $\omega{\cdot}\mathcal{L}$.

(b). Since $\mathcal{U}$ is a complete length CAT(κ) space, its blowup $n{\cdot}\mathcal{U}$ is CAT(κ/n^2). By Proposition 8.1.6, $\omega{\cdot}\mathcal{U}$ is CAT(0) and so is T_p^ω as a metric component of $\omega{\cdot}\mathcal{U}$. □

Recall that the tangent space T_p can be considered as a subset of T_p^ω (see 3.4.1). Therefore we get the following:

11.3.2. Corollary.

a) If $\mathcal{L}$ is a CBB(κ) *space and $p \in \mathcal{L}$, then T_p is* CBB(0). *Moreover, T_p satisfies (1+n)-point comparison (9.2.1).*

b) If $\mathcal{U}$ is a CAT(κ) *space and $p \in \mathcal{U}$, then T_p is* CAT(0). *Moreover, T_p satisfies (2n+2)-comparison (9.4.8).*

11.3.3. Proposition. *Assume $\mathcal{X}$ is a complete length* CBB *space or* CAT *space and $f\colon \mathcal{X} \looparrowright \mathbb{R}$ be a semiconcave locally Lipscitz subfunction. Then for any $p \in \operatorname{Dom} f$, the ultradifferential $\boldsymbol{d}_p^\omega\colon \mathrm{T}_p^\omega \to \mathbb{R}$ is a concave function.*

Proof. Fix a geodesic $[x^\omega y^\omega]$ in T_p^ω. It is sufficient to show that for any subarc $[\bar{x}^\omega \bar{y}^\omega]$ of $[x^\omega y^\omega]$ that does not contains the ends there is a sequence of geodesics $[\bar{x}^n \bar{y}^n]$ in $n{\cdot}\mathcal{X}$ converging to $[\bar{x}^\omega \bar{y}^\omega]$.

Choose any sequences $\bar{x}^n, \bar{y}^n \in n{\cdot}\mathcal{X}$ such that $\bar{x}^n \to \bar{x}^\omega$ and $\bar{y}^n \to$ $\to \bar{y}^\omega$ as $n \to \omega$. Note taht $[\bar{x}^n \bar{y}^n]$ converges to $[\bar{x}^\omega \bar{y}^\omega]$ as $n \to \omega$. The latter holds tivially in case CAT and in CBB case follows from 7.7.2. □

11.4 Gradient

11.4.1. Definition of gradient. *Let $\mathcal{L}$ be a complete length* CBB *space, $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be a subfunction and for a point $p \in \operatorname{Dom} f$ the differential $\boldsymbol{d}_p f\colon \mathrm{T}_p \to \mathbb{R}$ is defined.*

A tangent vector $g \in \mathrm{T}_p$ is called a gradient of f at p *(briefly, $g = \nabla_p f$) if*

a) $(\boldsymbol{d}_p f)(w) \leqslant \langle g, w\rangle$ *for any* $w \in \mathrm{T}_p$, *and*

b) $(\boldsymbol{d}_p f)(g) = \langle g, g\rangle$.

11.4.2. Existence and uniqueness of the gradient. *Let $\mathcal{L}$ be a complete* CBB(κ) *space and $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be locally Lipschitz and semi-concave subfunction. Then for any point $p \in \operatorname{Dom} f$, there is unique gradient $\nabla_p f \in \mathrm{T}_p$.*

Proof; uniqueness. If $g, g' \in \mathrm{T}_p$ are two gradients of f then

$$\langle g, g\rangle = (\boldsymbol{d}_p f)(g) \leqslant \langle g, g'\rangle, \qquad \langle g', g'\rangle = (\boldsymbol{d}_p f)(g') \leqslant \langle g, g'\rangle.$$

Therefore,

$$|g - g'|^2 = \langle g, g\rangle - 2\cdot\langle g, g'\rangle + \langle g', g'\rangle = 0;$$

that is, $g = g'$.

Existence. Note first that if $\boldsymbol{d}_p f \leqslant 0$, then one can take $\nabla_p f = 0$.

Otherwise, if $s = \sup\{\,(\boldsymbol{d}_p f)(\xi) : \xi \in \Sigma_p\,\} > 0$, it is sufficient to show that there is $\overline{\xi} \in \Sigma_p$ such that

❶ $$(\boldsymbol{d}_p f)\left(\overline{\xi}\right) = s.$$

Indeed, if $\overline{\xi}$ exists, then applying Lemma 11.2.3 for $u = \overline{\xi}$, $v = \varepsilon\cdot w$ with $\varepsilon \to 0+$, we get

$$(\boldsymbol{d}_p f)(w) \leqslant \langle w, s\cdot\overline{\xi}\rangle$$

for any $w \in \mathrm{T}_p$; that is, $s\cdot\overline{\xi}$ is the gradient at p.

Take a sequence of directions $\xi_n \in \Sigma_p$, such that $(\boldsymbol{d}_p f)(\xi_n) \to s$. Yet once applying Lemma 11.2.3 for $u = \xi_n$, $v = \xi_m$, we get

$$s \geqslant \frac{(\boldsymbol{d}_p f)(\xi_n) + (\boldsymbol{d}_p f)(\xi_m)}{\sqrt{2 + 2\cdot\cos\measuredangle(\xi_n, \xi_m)}}.$$

Therefore $\measuredangle(\xi_n, \xi_m) \to 0$ as $n, m \to \infty$; that is, (ξ_n) converges in itself. Clearly $\overline{\xi} = \lim_n \xi_n$ is satisfies ❶. □

Calculus

The next lemma roughly states that the gradient points in the direction of maximal slope; moreover if the slope in the given direction is almost maximal, then it is almost direction of the gradient.

11.4.3. Lemma. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $f\colon \mathcal{L} \to\!\!\!\!\to \mathbb{R}$ be locally Lipschitz and semiconcave and $p \in \operatorname{Dom} f$.*

Assume $|\nabla_p f| > 0$; set $\overline{\xi} = \frac{1}{|\nabla_p f|}\cdot\nabla_p f$. Then

a) If for some $v \in \mathrm{T}_p$, we have

$$|v| \leqslant 1+\varepsilon \quad \text{and} \quad (\boldsymbol{d}_p f)(v) > |\nabla_p f|\cdot(1-\varepsilon),$$

then

$$|\overline{\xi} - v| < 100\cdot\sqrt{\varepsilon}.$$

b) If $v_n \in \mathrm{T}_p$ be a sequence of vectors such that

$$\varlimsup_{n\to\infty} |v_n| \leqslant 1 \quad \text{and} \quad \varliminf_{n\to\infty} (\boldsymbol{d}_p f)(v_n) \geqslant |\nabla_p f|$$

then

$$\lim_{n\to\infty} v_n = \overline{\xi}.$$

c) $\overline{\xi}$ is the unique maximum direction for the restriction $\boldsymbol{d}_p f|_{\Sigma_p}$. In particular,

$$|\nabla_p f| = \sup\{\, \boldsymbol{d}_p f \,:\, \xi \in \Sigma_p f \,\}.$$

Proof. According to definition of gradient,

$$\begin{aligned}|\nabla_p f|\cdot(1-\varepsilon) < (\boldsymbol{d}_p f)(v) &\leqslant \\ &\leqslant \langle v, \nabla_p f\rangle = \\ &= |v|\cdot|\nabla_p f|\cdot\cos\measuredangle(\nabla_p f, v).\end{aligned}$$

Thus $|v| > 1-\varepsilon$ and $\cos\measuredangle(\nabla_p f, v) > \frac{1-\varepsilon}{1+\varepsilon}$. Hence (a).

Statements (b) and (c) follow directly from (a). □

As a corollary of the above lemma and Proposition 4.6.2 we get the following:

11.4.4. Chain rule. *Let $\mathcal{L}$ be a complete length* CBB *space, $f\colon \mathcal{L} \to\!\!\!\!\to \mathbb{R}$ be a semiconcave function and $\varphi\colon \mathbb{R} \to \mathbb{R}$ be a non-decreasing semiconcave function. Then $\varphi\circ f$ is semiconcave and $\nabla_x(\varphi\circ f) = \varphi^+(f(x))\cdot\nabla_x f$ for any $x \in \operatorname{Dom} f$.*

The following inequalities describe an important property of the "gradient vector field".



11.4.5. Lemma. *Let $\mathcal{L}$ be a complete length* CBB *space, $f\colon \mathcal{L} \to\!\!\!\!\to \mathbb{R}$ satisfies $f'' + \kappa\cdot f \leqslant \lambda$ for some $\kappa, \lambda \in \mathbb{R}$, $[pq] \subset \operatorname{Dom} f$ and $\ell = |p-q|$. Then*

$$\langle \uparrow_{[pq]}, \nabla_p f\rangle \geqslant \frac{f(q) - f(p)\cdot\mathrm{cs}^\kappa\ell - \lambda\cdot\mathrm{md}^\kappa\ell}{\mathrm{sn}^\kappa\ell}.$$

In particular,

a) if $\kappa = 0$,

$$\langle \uparrow_{[pq]}, \nabla_p f\rangle \geqslant \big(f(q) - f(p) - \tfrac{\lambda}{2}\cdot\ell^2\big)/\ell;$$

b) if $\kappa = 1$, $\lambda = 0$ *we have*

$$\langle \uparrow_{[pq]}, \nabla_p f\rangle \geqslant (f(q) - f(p)\cdot \cos\ell)\,/\,\sin\ell;$$

c) if $\kappa = -1$, $\lambda = 0$ *we have*

$$\langle \uparrow_{[pq]}, \nabla_p f\rangle \geqslant (f(q) - f(p)\cdot \cosh\ell)\,/\,\sinh\ell;$$

Proof of 11.4.5. Note that $\operatorname{geod}_{[pq]}(0) = p$, $\operatorname{geod}_{[pq]}(\ell) = q$ and $(\operatorname{geod}_{[pq]})^+(0) = \uparrow_{[pq]}$. Thus,

$$\begin{aligned}\langle \uparrow_{[pq]}, \nabla_p f\rangle &\geqslant d_p f(\uparrow_{[pq]}) = \\ &= (f\circ \operatorname{geod}_{[pq]})^+(0) \geqslant \\ &\geqslant \frac{f(q) - f(p)\cdot \operatorname{cs}^\kappa\ell - \lambda\cdot \operatorname{md}^\kappa\ell}{\operatorname{sn}^\kappa\ell}.\end{aligned}$$

□

The following corollary states that gradient vector field is monotonic in the sense similar to definition of *monotone operators*; see for example [49].

11.4.6. Monotonicity of gradient. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space,* $f\colon \mathcal{L} \to \mathbb{R}$ *be locally Lipschitz and* λ*-concave and* $[pq] \subset \subset \operatorname{Dom} f$. *Then*

$$\langle \uparrow_{[pq]}, \nabla_p f\rangle + \langle \uparrow_{[qp]}, \nabla_q f\rangle \geqslant -\lambda\cdot|p-q|.$$

Proof. Add two inequalities from 11.4.5*a*. □

11.4.7. Lemma. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space,* $f, g\colon \mathcal{L} \to \mathbb{R}$ *and* $p \in \operatorname{Dom} f \cap \operatorname{Dom} g$.

Then

$$|\nabla_p f - \nabla_p g|^2_{\mathrm{T}_p} \leqslant (|\nabla_p f| + |\nabla_p g|)\cdot \sup\{\, |(\boldsymbol{d}_p f)(\xi) - (\boldsymbol{d}_p g)(\xi)| : \xi \in \Sigma_p \,\}.$$

In particular, if $f_n\colon \mathcal{L} \to \mathbb{R}$ *is a sequence of locally Lipschitz and semiconcave subfunctions,* $p \in \operatorname{Dom} f_n$ *for each* n *and* $\boldsymbol{d}_p f_n$ *converges uniformly on* Σ_p *then sequence* $\nabla_p f_n \in \mathrm{T}_p$ *converges.*

Proof. Set

$$s = \sup\{\, |(\boldsymbol{d}_p f)(\xi) - (\boldsymbol{d}_p g)(\xi)| : \xi \in \Sigma_p \,\}.$$

Clearly for any $v \in \mathrm{T}_p$, we have

$$|(\boldsymbol{d}_p f)(v) - (\boldsymbol{d}_p g)(v)| \leqslant s\cdot|v|.$$

From the definition of gradient (11.4.1) we have:

$$\begin{aligned}(\boldsymbol{d}_p f)(\nabla_p g) &\leqslant \langle \nabla_p f, \nabla_p g\rangle, & (\boldsymbol{d}_p g)(\nabla_p f) &\leqslant \langle \nabla_p f, \nabla_p g\rangle,\\ (\boldsymbol{d}_p f)(\nabla_p f) &= \langle \nabla_p f, \nabla_p f\rangle, & (\boldsymbol{d}_p g)(\nabla_p g) &= \langle \nabla_p g, \nabla_p g\rangle.\end{aligned}$$

Therefore,

$$\begin{aligned}|\nabla_p f - \nabla_p g| &= \langle \nabla_p f, \nabla_p f\rangle + \langle \nabla_p g, \nabla_p g\rangle - 2\cdot\langle \nabla_p f, \nabla_p g\rangle \leqslant\\ &\leqslant (\boldsymbol{d}_p f)(\nabla_p f) + (\boldsymbol{d}_p g)(\nabla_p g) -\\ &\quad - (\boldsymbol{d}_p f)(\nabla_p g) - (\boldsymbol{d}_p g)(\nabla_p f) \leqslant\\ &\leqslant s\cdot(|\nabla_p f| + |\nabla_p g|).\end{aligned}$$

□

11.4.8. Exercise. *Let $\mathcal{L}$ be an m-dimensional complete length* $\mathrm{CBB}(\kappa)$ *space and a, b, p are mutually distinct points in $\mathcal{L}$. Prove that*

$$(d_p \operatorname{dist}_a)(\nabla_p \operatorname{dist}_b) + \cos \measuredangle^{\kappa}(p\,{}^a_b) \leqslant 0.$$

11.4.9. Exercise. *Let $\mathcal{L}$ be an m-dimensional complete length* $\mathrm{CBB}(\kappa)$ *space, the function $f\colon \mathcal{L} \to \mathbb{R}$ be semiconcave and locally Lipschitz and $\alpha\colon \mathbb{I} \to \mathcal{L}$ be a Lipschitz curve. Show that*

$$\langle \nabla_{\alpha(t)} f, \alpha^+(t)\rangle = (\boldsymbol{d}_{\alpha(t)} f)(\alpha^+(t))$$

for almost all $t \in \mathbb{I}$.

Semicontinuity

In this section we collect number of corollaries of the following lemma.

11.4.10. Ultralimit of |gradient|. *Assume*

- $(\mathcal{L}_n)$ *is a sequence of complete* $\mathrm{CBB}(\kappa)$ *spaces and* $(\mathcal{L}_n, p_n) \to$ $\to (\mathcal{L}_\omega, p_\omega)$ *as* $n \to \omega$*;*
- $f_n\colon \mathcal{L}_n \looparrowright \mathbb{R}$ *and* $f_\omega\colon \mathcal{L}_\omega \looparrowright \mathbb{R}$ *are locally Lipschitz and* λ*-concave and* $f_n \to f_\omega$ *as* $n \to \omega$*;*
- $x_n \in \operatorname{Dom} f_n$ *and* $x_n \to x_\omega \in \operatorname{Dom} f_\omega$ *as* $n \to \omega$.

Then

$$|\nabla_{x_\omega} f_\omega| \leqslant \lim_{n\to\omega} |\nabla_{x_n} f_n|.$$

Remarks. In order to construct an example such that $|\nabla_{x_\omega} f_\omega| < < \lim_{n\to\omega} |\nabla_{x_n} f_n|$; one can take $\mathcal{L}_n = \mathcal{L}_\omega = \mathbb{R}$, $f_n(x) = f_\omega(x) = -|x|$ and $x_n \to 0+$.

From the convergence of gradient curves (proved later in 14.2.10), one can deduce the following slightly stronger statement.

11.4.11. Proposition. *Assume*

- $\mathcal{L}_n$ *is a sequence of complete* CBB(κ) *spaces and* $(\mathcal{L}_n, p_n) \to \to (\mathcal{L}_\omega, p_\omega)$ *as* $n \to \omega$*;*
- $f_n\colon \mathcal{L}_n \looparrowright \mathbb{R}$ *and* $f_\omega\colon \mathcal{L}_\omega \looparrowright \mathbb{R}$ *are locally Lipschitz and* λ*-concave and* $f_n \to f_\omega$ *as* $n \to \omega$*.*

Then

$$|\nabla_{x_\omega} f_\omega| = \inf\{\lim_{n\to\omega} |\nabla_{x_n} f_n|\},$$

where infimum is taken for all sequences $x_n \in \operatorname{Dom} f_n$ *such that* $x_n \to \to x_\omega \in \operatorname{Dom} f_\omega$ *as* $n \to \omega$*.*

Proof of 11.4.10. Fix an $\varepsilon > 0$ and choose $y_\omega \in \operatorname{Dom} f_\omega$ sufficiently close to x_ω such that

$$|\nabla_{x_\omega} f_\omega| - \varepsilon < \frac{f_\omega(y_\omega) - f_\omega(x_\omega)}{|x_\omega - y_\omega|}.$$

Choose $y_n \in \mathcal{L}_n$ such that $y_n \to y_\omega$ as $n \to \omega$. Sinse $|x_\omega - y_\omega|$ is sufficiently small, the λ-concavity of f_n implies that

$$|\nabla_{x_\omega} f_\omega| - 2\cdot\varepsilon < (\boldsymbol{d}_{x_n} f_n)(\uparrow_{[x_n y_n]}),$$

for ω-almost all n. Hence,

$$|\nabla_{x_\omega} f_\omega| - 2\cdot\varepsilon \leqslant \lim_{n\to\omega} |\nabla_{x_n} f_n|.$$

Sinse $\varepsilon > 0$ is arbitrary, the proposition follows. $\square$

Note that the distance preserving map $\iota\colon \mathcal{L} \hookrightarrow \mathcal{L}^\omega$ induces an embedding

$$\boldsymbol{d}_p\iota\colon \mathrm{T}_p\mathcal{L} \hookrightarrow \mathrm{T}_p\mathcal{L}^\omega.$$

Thus, we can (and will) consider $\mathrm{T}_p\mathcal{L}$ as a subcone of $\mathrm{T}_p\mathcal{L}^\omega$.

11.4.12. Corollary. *Let* $\mathcal{L}$ *be a complete length* CBB *space and* $f\colon \mathcal{L} \looparrowright \mathbb{R}$ *be locally Lipschitz semiconcave subfunction. Then for any point* $p \in \operatorname{Dom} f$ *we have*

$$\nabla_p f = \nabla_p f^\omega.$$

Proof. Note that $\mathcal{L} \supset \operatorname{Dom} f \subset \operatorname{Dom} f^\omega \subset \mathcal{L}^\omega$. Applying 11.4.10 for $\mathcal{L}_n = \mathcal{L}$ and $x_n = x$, we get that $|\nabla_x f| \geqslant |\nabla_x f^\omega|$.

On the other hand, $f = f^\omega|_{\mathcal{L}}$, hence $\boldsymbol{d}_p f = \boldsymbol{d}_p f^\omega|\mathrm{T}_p\mathcal{L}$. Thus, from 11.4.3*c*, $|\nabla_x f| \leqslant |\nabla_x f^\omega|$. Therefore

❷ $$|\nabla_x f| = |\nabla_x f^\omega|.$$

Further,

$$\begin{aligned}|\nabla_x f|^2 &= (\boldsymbol{d}_p f)(\nabla_x f)\\ &= \boldsymbol{d}_p f^\omega(\nabla_x f) \leqslant\\ &\leqslant \langle \nabla_x f^\omega, \nabla_x f\rangle =\\ &= |\nabla_x f^\omega|\cdot|\nabla_x f|\cdot \cos\measuredangle(\nabla_x f^\omega, \nabla_x f).\end{aligned}$$

Together with ❷, this implies $\measuredangle(\nabla_x f^\omega, \nabla_x f) = 0$ and the statement follows. □

11.4.13. Semicontinuity of |gradient|. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be locally Lipschitz semiconcave subfunction. Then the function $x \mapsto |\nabla_x f|$ is lower-continuous; that is for any sequence $x_n \to x \in \operatorname{Dom} f$, we have*

$$|\nabla_x f| \leqslant \varliminf_{n\to\infty} |\nabla_{x_n} f|.$$

Proof. According to 11.4.12, $|\nabla_x f| = |\nabla_x f^\omega|$. Applying 11.4.10 for $x_n \to x$, we get that

$$\lim_{n\to\omega} |\nabla_{x_n} f| \geqslant |\nabla_x f^\omega| = |\nabla_x f|.$$

Passing to arbitrary subsequence of (x_n) we obtain the result. □

11.5 Polar vectors

Here we give a corollary of Lemma 11.4.7. It will be used to prove basic properties of tangent space.

11.5.1. Anti-sum lemma. *Let $\mathcal{L}$ be a complete length* CBB *space and $p \in \mathcal{L}$.*

Given two vectors $u, v \in \mathrm{T}_p$ there is unique vector $w \in \mathrm{T}_p$ such that

$$\langle u, x\rangle + \langle v, x\rangle + \langle w, x\rangle \geqslant 0$$

for any $x \in \mathrm{T}_p$ and

$$\langle u, w\rangle + \langle v, w\rangle + \langle w, w\rangle = 0.$$

If T_p would be a length space, then the lemma would follow from the existence of the gradient (11.4.2) applied to the function $\mathrm{T}_p \to \mathbb{R}$ defined by $x \mapsto -(\langle u, x\rangle + \langle v, x\rangle)$. However, tangent space T_p might be not a length space; see Halbeisen's example 11.1.2.

Applying the above lemma for $u = v$, we get the following statement.

11.5.2. Existence of polar vector. *Let $\mathcal{L}$ be a complete length* CBB *space and $p \in \mathcal{L}$. Given a vector $u \in \mathrm{T}_p$ there is unique vector $u^* \in \mathrm{T}_p$ such that $\langle u^*, u^*\rangle + \langle u, u^*\rangle = 0$ and u^* is* polar *to u; that is, $\langle u^*, x\rangle + \langle u, x\rangle \geqslant 0$ for any $x \in \mathrm{T}_p$.*

In particular, for any vector $u \in \mathrm{T}_p$ there is a polar vector $u^* \in \mathrm{T}_p$ such that $|u^*| \leqslant |u|$.

The Milka's lemma provides a refinement of this statement; it states that in the finite-dimensional case, we can assume that $|u^*| = |u|$.

Proof of 11.5.1. Choose two sequences of points $a_n, b_n \in \mathrm{Str}(p)$ such that $\uparrow_{[pa_n]} \to u/|u|$ and $\uparrow_{[pb_n]} \to v/|v|$. Consider sequence of functions

$$f_n = |u|\cdot \mathrm{dist}_{a_n} + |v|\cdot \mathrm{dist}_{b_n}\,.$$

According to Exercise 11.2.2,

$$(\boldsymbol{d}_p f_n)(x) = -|u|\cdot\langle\uparrow_{[pa_n]}, x\rangle - |v|\cdot\langle\uparrow_{[pb_n]}, x\rangle.$$

Thus, we have the following uniform convergence for all $x \in \Sigma_p$

$$(\boldsymbol{d}_p f_n)(x) \xrightarrow[n\to\infty]{} -\langle u, x\rangle - \langle v, x\rangle.$$

According to Lemma 11.4.7, sequence $\nabla_p f_n$ converges. Set

$$w = \lim_n \nabla_p f_n.$$

From definition of gradient

$$\begin{aligned}\langle w, w\rangle &= \lim_{n\to\infty}\langle\nabla_p f_n, \nabla_p f_n\rangle = \\ &= \lim_{n\to\infty}(\boldsymbol{d}_p f_n)(\nabla_p f_n) = \\ &= -\langle u, w\rangle - \langle v, w\rangle,\end{aligned} \qquad \begin{aligned}\langle w, x\rangle &= \lim_{n\to\infty}\langle\nabla_p f_n, x\rangle \geqslant \\ &\geqslant \lim_{n\to\infty}(\boldsymbol{d}_p f_n)(x) = \\ &= -\langle u, x\rangle - \langle v, x\rangle.\end{aligned}$$

□

11.6 Linear subspace of tangent space

11.6.1. Definition. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $p \in \mathcal{L}$ and $u, v \in \mathrm{T}_p$. We say that vectors u and v are* opposite *to each other, (briefly, $u + v = 0$) if $|u| = |v| = 0$ or $\measuredangle(u, v) = \pi$ and $|u| = |v|$.*

The subcone

$$\mathrm{Lin}_p = \{\, v \in \mathrm{T}_p : \exists\, w \in \mathrm{T}_p \ \ \textit{such that} \ \ w + v = 0 \,\}$$

will be called linear subcone *of* T_p.

The reason for the name "linear" will become evident in Theorem 11.6.4.

11.6.2. Proposition. *Let* $\mathcal{L}$ *be a complete length* CBB *space and* $p \in \mathcal{L}$. *Given two vectors* $u, v \in \mathrm{T}_p$, *the following statements are equivalent:*

a) $u + v = 0$;
b) $\langle u, x\rangle + \langle v, x\rangle = 0$ *for any* $x \in \mathrm{T}_p$;
c) $\langle u, \xi\rangle + \langle v, \xi\rangle = 0$ *for any* $\xi \in \Sigma_p$.

Proof. The condition $u + v = 0$ is equivalent to

$$\langle u, u\rangle = -\langle u, v\rangle = \langle v, v\rangle;$$

thus $(b)\Rightarrow(a)$. Since T_p is isometric to a subset of T_p^ω, the splitting theorem (14.4.1) applied for T_p^ω gives $(a)\Rightarrow(b)$.

The equivalence $(b)\Leftrightarrow(c)$ is trivial. □

11.6.3. Proposition. *Let* $\mathcal{L}$ *be a complete length* CBB *space and* $p \in \mathcal{L}$. *Then for any three vectors* $u, v, w \in \mathrm{T}_p$, $u + v = 0$ *and* $u + w = 0$ *implies* $v = w$.

Proof. From Proposition 11.6.2 both v and w satisfy the condition in corollary 11.5.2. Hence the result. □

Let $u \in \mathrm{Lin}_p$; that is $u + v = 0$ for some $v \in \mathrm{T}_p$. Given $s < 0$, set

$$s{\cdot}u := (-s){\cdot}v.$$

This way we define multiplication of any vector in Lin_p by any real number (positive and negative). Proposition 11.6.3 implies that such multiplication is uniquely defined.

11.6.4. Theorem. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space and* $p \in \mathcal{L}$. *Then* Lin_p *is a subcone of* T_p *isometric to a Hilbert space.*

Before comming to the proof of the theorem, let us give its corollary.

11.6.5. Corollary. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space and* $p \in \mathrm{Str}(x_1, x_2, \dots, x_n)$. *Then there is a subcone* $E \subset \mathrm{T}_p$ *that isometric to a Euclidean space such that* $\log[px_i] \in E$ *for every* i.

Proof. By the definition of Str (7.2.3), $\log[px_i] \in \mathrm{Lin}_p$ for each i. It remains to apply Theorem 11.6.4. $\square$

The main difficulty in the proof of Theorem 11.6.4 comes from the fact that in general T_p is not a length space; see Habeisen's example (11.1.2). Othervise, if the tangent space is a length space, the statement would follow directly from the splitting theorem (14.4.1). In fact the proof of this theorem is a far walk around — we use construction of gradient, as well as the splitting theorem, namely its corollary (14.4.2). Thus in order to understand our proof one needs to read most of Chapter 14.

Proof of 11.6.4. First we show that Lin_p is a complete geodesic CBB(0) space.

Recall that T_p^ω is a complete geodesic CBB(0) space (see 3.3.2 and 11.3.1*a*) and Lin_p is a closed subset of T_p^ω. Thus, it is sufficient to show that the metric on Lin_p inherited from T_p^ω is a length-metric.

Fix two vectors $x, y \in \mathrm{Lin}_p$. Let u and v be such that $u + \frac{1}{2}\cdot x = 0$ and $v + \frac{1}{2}\cdot y = 0$. Apply Lemma 11.5.1 to the vectors u and v; let $w \in \mathrm{T}_p$ denotes the obtained tangent vector.

❶ *w is a midpoint of $[xy]$.*

Indeed, according to Lemma 11.5.1,

$$\begin{aligned}|w|^2 &= -\langle w,u\rangle - \langle w,v\rangle = \\ &= \tfrac{1}{2}\cdot\langle w,x\rangle + \tfrac{1}{2}\cdot\langle w,y\rangle.\end{aligned}$$

Therefore,

$$\begin{aligned}|x-w|^2 + |w-y|^2 &= 2\cdot|w|^2 + |x|^2 + |y|^2 - 2\cdot\langle w,x\rangle - 2\cdot\langle w,y\rangle = \\ &= |x|^2 + |y|^2 - \langle w,x\rangle - \langle w,y\rangle \leqslant \\ &\leqslant |x|^2 + |y|^2 + \langle u,x\rangle + \langle v,x\rangle + \langle u,y\rangle + \langle v,y\rangle = \\ &= \tfrac{1}{2}\cdot|x|^2 + \tfrac{1}{2}\cdot|y|^2 - \langle x,y\rangle = \\ &= \tfrac{1}{2}\cdot|x-y|^2.\end{aligned}$$

Thus $|x-w| = |w-y| = \frac{1}{2}\cdot|x-y|$ and ❶ follows. $\triangle$

Note that for any $v \in \mathrm{Lin}_p$ there is a line ℓ that contains v and 0, thus applying 14.4.2, we get that Lin_p is isometric to a Hilbert space. $\square$

11.7 Comments

11.7.1. Open question. *Let $\mathcal{L}$ be a proper length* CBB(κ) *space. Is it true that for any $p \in \mathcal{L}$, the tangent space T_p is a length space?*

Chapter 12

Dimension of CAT spaces

In this chapter we discuss the constructions introduced by Bruce Kleiner in [13].

The material of this chapter is used mostly for CAT spaces, but the results in section 12.1 find some applications for finite dimensional CBB spaces as well.

12.1 The case of complete geodesic spaces

The following construction gives a $\bar{k}$-dimensional submanifold for a given "nondegenerate" array of $\bar{k}+1$ strongly convex functions.

12.1.1. Definition. *For two real arrays $\boldsymbol{v}$, $\boldsymbol{w} \in \mathbb{R}^{\bar{k}+1}$, $\boldsymbol{v} = (v^0, v^1, \dots$ $\dots, v^{\bar{k}})$ and $\boldsymbol{w} = (w^0, w^1, \dots, w^{\bar{k}})$, we will write $\boldsymbol{v} \succcurlyeq \boldsymbol{w}$ if $v^i \geqslant w^i$ for each i.*

Given a subset $Q \subset \mathbb{R}^{\bar{k}+1}$, denote by $\operatorname{Up} Q$, the smallest upper set containing Q and $\operatorname{Min} Q$, the set of minimal elements of Q with respect to $\succcurlyeq$; that is,

$$\begin{aligned} \operatorname{Up} Q &= \left\{ \boldsymbol{v} \in \mathbb{R}^{\bar{k}+1} : \exists\, \boldsymbol{w} \in Q \text{ such that } \boldsymbol{v} \succcurlyeq \boldsymbol{w} \right\}, \\ \operatorname{Min} Q &= \left\{ \boldsymbol{v} \in Q : \text{if } \boldsymbol{v} \succcurlyeq \boldsymbol{w} \in Q \text{ then } \boldsymbol{w} = \boldsymbol{v} \right\}. \end{aligned}$$

12.1.2. Definition. *Let $\mathcal{X}$ be a metric space and $\boldsymbol{f} = (f^0, f^1, \dots$ $\dots, f^{\bar{k}}) \colon \mathcal{X} \to \mathbb{R}^{\bar{k}+1}$ be a function array. The set*

$$\operatorname{Web} \boldsymbol{f} := \boldsymbol{f}^{-1} \left[\operatorname{Min} \boldsymbol{f}(\mathcal{X}) \right] \subset \mathcal{X}$$

will be called the web *on* $\boldsymbol{f}$.

Given an array $\boldsymbol{f} = (f^0, f^1, \dots, f^k)$, we denote by $\boldsymbol{f}^{-i}$ the subarray of $\boldsymbol{f}$ with f^i removed; that is,

$$\boldsymbol{f}^{-i} := (f^0, \dots, f^{i-1}, f^{i+1}, \dots, f^k).$$

Clearly $\operatorname{Web} \boldsymbol{f}^{-i} \subset \operatorname{Web} \boldsymbol{f}$. Define the *inner web* of $\boldsymbol{f}$ as

$$\operatorname{InWeb} \boldsymbol{f} = \operatorname{Web} \boldsymbol{f} \setminus \left(\bigcup_i \operatorname{Web} \boldsymbol{f}^{-i} \right).$$

We say that a function array is *nondegenerate* if $\operatorname{InWeb} \boldsymbol{f} \neq \varnothing$.

Example. If $\mathcal{X}$ is a geodesic space, then $\operatorname{Web}(\operatorname{dist}_x, \operatorname{dist}_y)$ is the union of all geodesics from x to y and

$$\operatorname{InWeb}(\operatorname{dist}_x, \operatorname{dist}_y) = \operatorname{Web}(\operatorname{dist}_x, \operatorname{dist}_y) \setminus \{x, y\}.$$

Barycenters. Let us denote by $\Delta^k \subset \mathbb{R}^{k+1}$ the *standard k-simplex*; that is, $\boldsymbol{x} = (x^0, x^1, \dots, x^k) \in \Delta^k$ if $\sum_{i=0}^k x^i = 1$ and $x^i \geqslant 0$ for all i.

Let $\mathcal{X}$ be a metric space and $\boldsymbol{f} = (f^0, f^1, \dots, f^k) \colon \mathcal{X} \to \mathbb{R}^{k+1}$ be a function array. Consider the map $\mathfrak{S}_{\boldsymbol{f}} \colon \Delta^k \to \mathcal{X}$, defined by

$$\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x}) = \operatorname{MinPoint} \sum_{i=0}^k x^i \cdot f^i,$$

where $\operatorname{MinPoint} f$ denotes a point of minimum of f. The map $\mathfrak{S}_{\boldsymbol{f}}$ will be called a *barycentric simplex* of $\boldsymbol{f}$. Note that for general function array $\boldsymbol{f}$, the value $\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x})$ might be undefined or nonuniquely defined.

It is clear from the definition, that $\mathfrak{S}_{\boldsymbol{f}^{-i}}$ coincides with the restriction of $\mathfrak{S}_{\boldsymbol{f}}$ to the corresponding face of Δ^k.

12.1.3. Theorem. *Let $\mathcal{X}$ be a complete geodesic space and $\boldsymbol{f} = (f^0, f^1, \dots, f^k) \colon \mathcal{X} \to \mathbb{R}^{k+1}$ be an array of strongly convex and locally Lipschitz functions. Then $\boldsymbol{f}$ defines a $C^{\frac{1}{2}}$-embedding* $\operatorname{Web} \boldsymbol{f} \hookrightarrow \mathbb{R}^{k+1}$.

Moreover,

a) $W = \operatorname{Up}[\boldsymbol{f}(\mathcal{X})]$ is a convex closed subset of $\mathbb{R}^{k+1}$ and $S = \partial_{\mathbb{R}^{k+1}} W$ is a convex hypersurface in $\mathbb{R}^{k+1}$.

b)

$$\boldsymbol{f}(\operatorname{Web} \boldsymbol{f}) = \operatorname{Min} W \subset S$$

and

$$\boldsymbol{f}(\operatorname{InWeb} \boldsymbol{f}) = \operatorname{Int}_S(\operatorname{Min} W).$$

c) *The barycentric simplex* $\mathfrak{S}_{\boldsymbol{f}}\colon \Delta^{\bar{k}} \to \mathcal{X}$ *is a uniquely defined Lipshitz map and* $\operatorname{Im}\mathfrak{S}_{\boldsymbol{f}} = \operatorname{Web}\boldsymbol{f}$. *In particular* $\operatorname{Web}\boldsymbol{f}$ *is compact.*

d) *Let us equip* $\Delta^{\bar{k}}$ *with metric induced by* ℓ^1*-norm on* $\mathbb{R}^{\bar{k}+1}$. *Then the Lipschitz constant of* $\mathfrak{S}_{\boldsymbol{f}}\colon \Delta^{\bar{k}} \to \mathcal{U}$ *can be estimated in terms of positive lower bounds on* $(f^i)''$ *and Lipshitz constants of* f^i *in a neigborhood of* $\operatorname{Web}\boldsymbol{f}$ *for all* i.

In particular, from (a) and (b), it follows that $\operatorname{InWeb}\boldsymbol{f}$ *is* $C^{\frac{1}{2}}$*-homeomorphic to an open set of* $\mathbb{R}^{\bar{k}}$.

The proof is preceded by few preliminary statements.

12.1.4. Lemma. *Suppose* $\mathcal{X}$ *is a complete geodesic space and let* $f\colon \mathcal{X} \to \mathbb{R}$ *be a locally Lipschitz, strongly convex function. Then the minimum point of* f *is uniquely defined.*

Proof. Without loss of generality, we can assume that f is 1-convex; in particular the following claim holds.

❶ *if* z *is a midpoint of the geodesic* $[xy]$, *then*

$$s \leqslant f(z) \leqslant \tfrac{1}{2}\cdot f(x) + \tfrac{1}{2}\cdot f(y) - \tfrac{1}{8}\cdot|x-y|^2.$$

Uniqueness. Assume that x and y are distinct minimum points of f. From ❶, we get

$$f(z) < f(x) = f(y),$$

a contradiction.

Existence. Fix a point $p \in \mathcal{X}$; let $\pounds \in \mathbb{R}$ be a Lipschitz constant of f in a neighborhood of p.

Consider function $\varphi(t) = f \circ \operatorname{geod}_{[px]}(t)$. Clearly φ is 1-convex and $\varphi^+(0) \geqslant -\pounds$. Setting $\ell = |p-x|$, we get

$$\begin{aligned} f(x) = \varphi(\ell) &\geqslant \\ &\geqslant f(p) - \pounds\cdot\ell + \tfrac{1}{2}\cdot\ell^2 \geqslant \\ &\geqslant f(p) - \tfrac{1}{2}\cdot\pounds^2. \end{aligned}$$

In particular,

$$\begin{aligned} s := \inf\{\, f(x) : x \in \mathcal{X} \,\} &\geqslant \\ \geqslant f(p) - \tfrac{1}{2}\cdot\pounds^2. \end{aligned}$$

Choose a sequence of points $p_n \in \mathcal{X}$ such that $f(p_n) \to s$. Applying ❶, for $x = p_n$, $y = p_m$, we get that (p_n) is converging in itself. Hence, p_n converges to a minimum point of f. □

12.1.5. Definition. *Let Q be a closed subset of $\mathbb{R}^{k+1}$. A vector $\boldsymbol{x} = (x^0, x^1, \dots, x^k) \in \mathbb{R}^{k+1}$ is* subnormal *to Q at point $\boldsymbol{v} \in Q$ if*

$$\langle \boldsymbol{x}, \boldsymbol{w} - \boldsymbol{v} \rangle := \sum_i x^i \cdot (w^i - v^i) \geqslant 0$$

for any $\boldsymbol{w} \in Q$.

12.1.6. Lemma. *Let $\mathcal{X}$ be a complete geodesic space and $\boldsymbol{f} = (f^0, f^1, \dots, f^k)\colon \mathcal{X} \to \mathbb{R}^{k+1}$ be an array of strongly convex and locally Lipschitz functions. Set $W = \operatorname{Up} \boldsymbol{f}(\mathcal{X})$. Then:*

a) W is a closed convex set bounded below with respect to $\succcurlyeq$.

b) If $\boldsymbol{x}$ is a subnormal vector to W, then $\boldsymbol{x} \succcurlyeq \mathbf{0}$.

c) $S = \partial_{\mathbb{R}^{k+1}} W$ is a complete convex hypersurface in $\mathbb{R}^{k+1}$.

Proof. Denote by $\bar{W}$ the closure of W.

Convexity of f^i implies that for any two points $p, q \in \mathcal{X}$ and $t \in [0, 1]$ we have

❷ $$(1-t) \cdot \boldsymbol{f}(p) + t \cdot \boldsymbol{f}(q) \succcurlyeq \boldsymbol{f} \circ \operatorname{path}_{[pq]}(t).$$

where $\operatorname{path}_{[pq]}$ denotes a geodesic path from p to q. Therefore W as well as $\bar{W}$ are convex sets in $\mathbb{R}^{k+1}$.

Set

$$w^i = \min \left\{ f^i(x) : x \in \mathcal{X} \right\}.$$

By Lemma 12.1.4, w^i is finite for each i. Clearly $\boldsymbol{w} = (w^0, w^1, \dots, w^k)$ is a lower bound of $\bar{W}$ with respect to $\succcurlyeq$.

It is clear that W has nonempty interior and $W \neq \mathbb{R}^{k+1}$ since it is bounded below. Therefore $S = \partial_{\mathbb{R}^{k+1}} W = \partial_{\mathbb{R}^{k+1}} \bar{W}$ is a complete convex hypersurface in $\mathbb{R}^{k+1}$.

Since $\bar{W}$ is closed and bounded below, we also have

❸ $$\bar{W} = \operatorname{Up}[\operatorname{Min} \bar{W}].$$

Choose arbitrary $\boldsymbol{v} \in S$. Let $\boldsymbol{x} \in \mathbb{R}^{k+1}$ be a subnormal vector to $\bar{W}$ at $\boldsymbol{v}$. In particular, $\langle \boldsymbol{x}, \boldsymbol{y} \rangle \geqslant 0$ for any $\boldsymbol{y} \succcurlyeq \mathbf{0}$; that is, $\boldsymbol{x} \succcurlyeq \mathbf{0}$.

Further, according to Lemma 12.1.4, the function $\sum_i x^i \cdot f^i$ has uniquely defined minimum point, say p. Clearly

❹ $$\boldsymbol{v} \succcurlyeq \boldsymbol{f}(p) \quad \text{and} \quad \boldsymbol{f}(p) \in \operatorname{Min} W.$$

Note that for any $\boldsymbol{u} \in \bar{W}$ there is $\boldsymbol{v} \in S$ such that $\boldsymbol{u} \succcurlyeq \boldsymbol{v}$. Therefore ❹ implies that

$$\bar{W} \subset \operatorname{Up}[\operatorname{Min} W] \subset W.$$

Hence $\bar{W} = W$; that is, W is closed. $\square$

Proof of 12.1.3. Without loss of generality we may assume that all f^i are 1-convex.

Given $\boldsymbol{v} = (v^0, v^1, \dots, v^k) \in \mathbb{R}^{k+1}$, consider the function $h_{\boldsymbol{v}} \colon \mathcal{X} \to \mathbb{R}$ defined as

$$h_{\boldsymbol{v}}(p) = \max_i \{f^i(p) - v^i\}.$$

Note that $h_{\boldsymbol{v}}$ is 1-convex. Set

$$\Phi(\boldsymbol{v}) := \operatorname{MinPoint} h_{\boldsymbol{v}}.$$

According to Lemma 12.1.4, $\Phi(\boldsymbol{v})$ is uniquely defined.

From the definition of web (12.1.2) we have $\Phi \circ \boldsymbol{f}(p) = p$ for any $p \in \operatorname{Web} \boldsymbol{f}$; that is, Φ is a left inverse to the restriction $\boldsymbol{f}|_{\operatorname{Web} \boldsymbol{f}}$. In particular,

❺ $$\operatorname{Web} \boldsymbol{f} = \operatorname{Im} \Phi.$$

Given $\boldsymbol{v}, \boldsymbol{w} \in \mathbb{R}^{k+1}$, set $p = \Phi(\boldsymbol{v})$ and $q = \Phi(\boldsymbol{w})$. Since $h_{\boldsymbol{v}}$ and $h_{\boldsymbol{w}}$ are 1-convex, we have

$$h_{\boldsymbol{v}}(q) \geqslant h_{\boldsymbol{v}}(p) + \tfrac{1}{2} \cdot |p - q|^2, \qquad h_{\boldsymbol{w}}(p) \geqslant h_{\boldsymbol{w}}(q) + \tfrac{1}{2} \cdot |p - q|^2.$$

Therefore,

$$\begin{aligned} |p - q|^2 &\leqslant 2 \cdot \sup_{x \in \mathcal{X}} \{|h_{\boldsymbol{v}}(x) - h_{\boldsymbol{w}}(x)|\} \leqslant \\ &\leqslant 2 \cdot \max_i \{|v^i - w^i|\}. \end{aligned}$$

In particular, Φ is $C^{\frac{1}{2}}$-continuous, or $\boldsymbol{f}|_{\operatorname{Web} \boldsymbol{f}}$ is a $C^{\frac{1}{2}}$-embedding.

As in Lemma 12.1.6, set $W = \operatorname{Up} \boldsymbol{f}(\mathcal{X})$ and $S = \partial_{\mathbb{R}^{k+1}} W$. Recall that S is a convex hypersurface in $\mathbb{R}^{k+1}$. Clearly $\boldsymbol{f}(\operatorname{Web} \boldsymbol{f}) = \operatorname{Min} W \subset S$. From the definition of inner web we have $\boldsymbol{v} \in \boldsymbol{f}(\operatorname{InWeb} \boldsymbol{f})$ if and only if $\boldsymbol{v} \in S$ and for any i there is $\boldsymbol{w} = (w^0, w^1, \dots, w^k) \in W$ such that $w^j < v^j$ for all $j \neq i$. Thus, $\boldsymbol{f}(\operatorname{InWeb} \boldsymbol{f})$ is open in S. That is, $\operatorname{InWeb} \boldsymbol{f}$ is $C^{\frac{1}{2}}$-homeomorphic to an open set in a convex hypersurface $S \subset \mathbb{R}^{k+1}$.

(c)+(d). Since f^i is 1-convex, for any $\boldsymbol{x} = (x^0, x^1, \dots, x^k) \in \Delta^k$, the convex combination

$$\left(\sum_i x^i \cdot f^i \right) : \mathcal{X} \to \mathbb{R}$$

is also 1-convex. Therefore, according to Lemma 12.1.4, $\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x})$ is defined for any $\boldsymbol{x} \in \Delta^k$.

For $\boldsymbol{x}, \boldsymbol{y} \in \Delta^{\bar{k}}$, set

$$f_{\boldsymbol{x}} = \sum_i x^i \cdot f^i, \qquad f_{\boldsymbol{y}} = \sum_i y^i \cdot f^i,$$
$$p = \mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x}), \qquad q = \mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{y}),$$
$$\ell = |p - q|.$$

Note the following:

- ⋄ The function $\varphi(t) = f_{\boldsymbol{x}} \circ \operatorname{geod}_{[pq]}(t)$ has minimum at 0. Therefore $\varphi^+(0) \geqslant 0$
- ⋄ The function $\psi(t) = f_{\boldsymbol{y}} \circ \operatorname{geod}_{[pq]}(t)$ has minimum at ℓ. Therefore $\psi^-(\ell) \geqslant 0$.

From 1-convexity of $f_{\boldsymbol{y}}$, we have $\psi^+(0) + \psi^-(\ell) + \ell \leqslant 0$.

Let $\pounds$ be a Lipschitz constant for all f^i in a neighborhood $\Omega \ni p$. Then

$$\psi^+(0) \leqslant \varphi^+(0) + \pounds \cdot \|\boldsymbol{x} - \boldsymbol{y}\|_1,$$

where $\|\boldsymbol{x} - \boldsymbol{y}\|_1 = \sum_{i=0}^{\bar{k}} |x^i - y^i|$. That is, given $\boldsymbol{x} \in \Delta^{\bar{k}}$, there is a constant $\pounds$ such that

$$|\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x}) - \mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{y})| = \ell \leqslant$$
$$\leqslant \pounds \cdot \|\boldsymbol{x} - \boldsymbol{y}\|_1$$

for any $\boldsymbol{y} \in \Delta^{\bar{k}}$. In particular, there is $\varepsilon > 0$ such that if $\|\boldsymbol{x} - \boldsymbol{y}\|_1$, $\|\boldsymbol{x} - \boldsymbol{z}\|_1 < \varepsilon$, then $\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{y})$, $\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{z}) \in \Omega$. Thus, the same argument as above implies

$$|\mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{y}) - \mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{z})| = \ell \leqslant \pounds \cdot \|\boldsymbol{y} - \boldsymbol{z}\|_1$$

for any $\boldsymbol{y}$ and $\boldsymbol{z}$ sufficiently close to $\boldsymbol{x}$; that is, $\mathfrak{S}_{\boldsymbol{f}}$ is locally Lipschitz. Since $\Delta^{\bar{k}}$ is compact, $\mathfrak{S}_{\boldsymbol{f}}$ is Lipschitz.

Clearly $\mathfrak{S}_{\boldsymbol{f}}(\Delta^{\bar{k}}) \subset \operatorname{Web} \boldsymbol{f}$; it remains to show that $\mathfrak{S}_{\boldsymbol{f}}(\Delta^{\bar{k}}) \supset \supset \operatorname{Web} \boldsymbol{f}$. According to Lemma 12.1.6, $W = \operatorname{Up} \boldsymbol{f}(\mathcal{X})$ is a closed convex set in $\mathbb{R}^{\bar{k}+1}$. Let $p \in \operatorname{Web} \boldsymbol{f}$; clearly $\boldsymbol{f}(p) \in \operatorname{Min} W \subset S = \partial_{\mathbb{R}^{\bar{k}+1}} W$. Let $\boldsymbol{x}$ be a subnormal vector to W at $\boldsymbol{f}(p)$. According to Lemma 12.1.6, $\boldsymbol{x} \succcurlyeq \boldsymbol{0}$. Without loss of generality we may assume that $\sum_i x^i = 1$; that is, $\boldsymbol{x} \in \Delta^{\bar{k}}$. By Lemma 12.1.4, p is the unique minimum point of $\sum_i x^i \cdot f^i$; that is, $p = \mathfrak{S}_{\boldsymbol{f}}(\boldsymbol{x})$. □

12.2 The case of CAT spaces

Let $\boldsymbol{a} = (a^0, a^1, \dots, a^{\bar{k}})$ be a point array in a metric space $\mathcal{U}$. Recall that $\operatorname{dist}_{\boldsymbol{a}}$ denotes the distance map

$$(\operatorname{dist}_{a^0}, \operatorname{dist}_{a^1}, \dots, \operatorname{dist}_{a^{\bar{k}}}) \colon \mathcal{U} \to \mathbb{R}^{\bar{k}+1}$$

which can be also regarded as a function array. The *radius of the point array* $\boldsymbol{a}$ is defined to be the radius of the set $\{a^0, a^1, \dots, a^{\bar{k}}\}$; that is,

$$\operatorname{rad} \boldsymbol{a} = \inf \left\{ r > 0 : a^i \in \mathrm{B}(z, r) \text{ for all } i \text{ and some } z \in \mathcal{U} \right\}.$$

Fix $\kappa \in \mathbb{R}$. Let $\boldsymbol{a} = (a^0, a^1, \dots, a^{\bar{k}})$ be a point array of radius $< \frac{\varpi^\kappa}{2}$ in a metric space $\mathcal{U}$. Consider function array $\boldsymbol{f} = (f^0, f^1, \dots$ $\dots, f^{\bar{k}})$ where

$$f^i(x) = \mathrm{md}^\kappa |a^i - x|.$$

Assuming the barycentric simplex $\mathfrak{S}_{\boldsymbol{f}}$ is defined, then $\mathfrak{S}_{\boldsymbol{f}}$ is called the κ-*barycentric simplex* for the point array $\boldsymbol{a}$; it will be denoted by $\mathfrak{S}^\kappa_{\boldsymbol{a}}$. The points $a^0, a^1, \dots, a^{\bar{k}}$ are called *vertexes of* κ-*barycentric simplex*. Note that once we say the κ-barycentric simplex is defined, we automatically assume that $\operatorname{rad} \boldsymbol{a} < \frac{\varpi^\kappa}{2}$.

12.2.1. Theorem. *Let $\mathcal{U}$ be a complete length* $\mathrm{CAT}(\kappa)$ *space and $\boldsymbol{a} = (a^0, a^1, \dots a^{\bar{k}})$ be a point array with radius $< \frac{\varpi^\kappa}{2}$. Then:*

a) The κ-barycentric simplex $\mathfrak{S}^\kappa_{\boldsymbol{a}} \colon \Delta^{\bar{k}} \to \mathcal{U}$ is defined. Moreover, $\mathfrak{S}^\kappa_{\boldsymbol{a}}$ is a Lipschitz map, and if $\Delta^{\bar{k}}$ is equipped with the ℓ^1-metric, then its Lipschitz contant can be estimated in terms of κ and the radius of $\boldsymbol{a}$ (in particular it does not depend on $\bar{k}$).

b) $\mathrm{Web}(\mathrm{dist}_{\boldsymbol{a}}) = \operatorname{Im} \mathfrak{S}^\kappa_{\boldsymbol{a}}$. *Moreover, if a closed convex set K contains all a^i, then* $\mathrm{Web}(\mathrm{dist}_{\boldsymbol{a}}) \subset K$.

c) The restriction[1] $\mathrm{dist}_{\boldsymbol{a}^{-0}} |_{\mathrm{InWeb}(\mathrm{dist}_{\boldsymbol{a}})}$ *is an open $C^{\frac{1}{2}}$-embedding in $\mathbb{R}^{\bar{k}}$. Thus there is an inverse of* $\mathrm{dist}_{\boldsymbol{a}^{-0}} |_{\mathrm{InWeb}(\mathrm{dist}_{\boldsymbol{a}})}$*, say* $\Phi \colon \mathbb{R}^{\bar{k}} \hookrightarrow \mathcal{U}$.

The subfunction $f = \mathrm{dist}_{a^0} \circ \Phi$ is semiconvex and locally Lipschitz. Moreover, if $\kappa \leqslant 0$, then f is convex.

In particular, $\mathrm{Web}(\mathrm{dist}_{\boldsymbol{a}})$ *is compact and* $\mathrm{InWeb}(\mathrm{dist}_{\boldsymbol{a}})$ *is $C^{\frac{1}{2}}$-homeomorphic to an open subset of $\mathbb{R}^{\bar{k}}$.*

12.2.2. Definition. *The submap $\Phi \colon \mathbb{R}^{\bar{k}} \hookrightarrow \mathcal{X}$ as in Theorem 12.2.1c will be called the* $\mathrm{dist}_{\boldsymbol{a}}$-web embedding *with* brace dist_{a^0} .

12.2.3. Definition. *Let $\mathcal{U}$ be a complete length* $\mathrm{CAT}(\kappa)$ *space and $\boldsymbol{a} = (a^0, a^1, \dots a^{\bar{k}})$ be a point array with radius $< \frac{\varpi^\kappa}{2}$. If* $\mathrm{InWeb}(\mathrm{dist}_{\boldsymbol{a}})$ *is nonempty, then the point array $\boldsymbol{a}$ is called* nondegenerate.

Lemma 12.2.5 will provide examples of nondegenerate point arrays, which can be used in Theorem 12.2.1*c*.

[1] Recall that $\mathrm{dist}_{\boldsymbol{a}^{-0}}$ denotes array $(\mathrm{dist}_{a^1}, \dots, \mathrm{dist}_{a^{\bar{k}}})$.

12.2.4. Corollary. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space,* $\boldsymbol{a} = (a^0, a^1, \dots a^m)$ *be a nondegenerate point array of radius* $< \frac{\varpi^\kappa}{2}$ *in* $\mathcal{U}$ *and* $\sigma = \mathfrak{S}^\kappa_{\boldsymbol{a}}$ *be corresponding* κ*-baricentric simplex. Then for some* $\boldsymbol{x} \in \Delta^m$ *the differential* $\boldsymbol{d}_{\boldsymbol{x}}\sigma$ *is linear and the image* $\operatorname{Im} \boldsymbol{d}_{\boldsymbol{x}}\sigma$ *forms a subcone isometric to an* m*-dimensional Euclidean space in the tangent cone* $\mathrm{T}_{\sigma(\boldsymbol{x})}$.

Proof. Denote by $\tau\colon \mathcal{U} \to \mathbb{R}^m$ the distance map $\operatorname{dist}_{\boldsymbol{a}^{-0}}$.

According to Theorem 12.2.1, σ is Lipschitz and the distance map τ, gives an open embedding of $\operatorname{InWeb}(\operatorname{dist}_{\boldsymbol{a}}) = \sigma(\Delta^m)\backslash\sigma(\partial\Delta^m)$. Note that τ is Lipschitz. According to Rademacher's theorem the differential $\boldsymbol{d}_{\boldsymbol{x}}(\tau\circ\sigma)$ is linear for almost all $\boldsymbol{x} \in \Delta^m$. Further since $\operatorname{InWeb}(\operatorname{dist}_{\boldsymbol{a}}) \neq \varnothing$, the area formula implies that $\boldsymbol{d}_{\boldsymbol{x}}(\tau\circ\sigma)$ is surjective on a set of positive masure of points $\boldsymbol{x} \in \Delta^m$.

Note that $\boldsymbol{d}_{\boldsymbol{x}}(\tau\circ\sigma) = (\boldsymbol{d}_{\sigma(\boldsymbol{x})}\tau)\circ(\boldsymbol{d}_{\boldsymbol{x}}\sigma)$. Applying Rademacher's theorem again, we get that $\boldsymbol{d}_{\boldsymbol{x}}\sigma$ is linear for almost all $\boldsymbol{x} \in \Delta^m$, at these points $\operatorname{Im} \boldsymbol{d}_{\boldsymbol{x}}\sigma$ forms a subcone isometric to a Euclidean space in $\mathrm{T}_{\sigma(\boldsymbol{x})}$. Clearly the dimension of $\operatorname{Im} \boldsymbol{d}_{\boldsymbol{x}}(\tau\circ\sigma)$ is at least as big as dimension of $\operatorname{Im} \boldsymbol{d}_{\boldsymbol{x}}\sigma$. Hence the result. $\square$

Proof of 12.2.1. Fix $z \in \mathcal{U}$ and $r < \frac{\varpi^\kappa}{2}$ such that $|z - a^i| < r$ for all i. Note that the set $K \cap \overline{\mathrm{B}}[z, r]$ is convex, closed, and contains all a^i. Applying the theorem on short retract (Exercise 8.6.5), we get the second part of (b).

The rest of the statements are proved first in case $\kappa \leqslant 0$ and then the remaining case $\kappa > 0$ is reduced to the case $\kappa = 0$.

Case $\kappa \leqslant 0$. Consider function array $f^i = \operatorname{md}^\kappa \circ \operatorname{dist}_{a^i}$. From the definition of web (12.1.2), it is clear that $\operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}}) = \operatorname{Web} \boldsymbol{f}$. Further, from the definition of κ-barycentric simplex, $\mathfrak{S}^\kappa_{\boldsymbol{a}} = \mathfrak{S}_{\boldsymbol{f}}$.

All the functions f^i are strongly convex (see 8.5.1b). Therefore (12.2.1a), (12.2.1b) and the first statements in (12.2.1b) follow from Theorem 12.1.3.

Case $\kappa > 0$. Applying rescaling, we can assume $\kappa = 1$; so $\varpi^\kappa = \varpi^1 = \pi$.

Set $\mathring{\mathcal{U}} = \operatorname{Cone}\mathcal{U}$; by 10.4.2$a$, $\mathring{\mathcal{U}}$ is CAT(0). Let us denote by ι the natural embedding of $\mathcal{U}$ as the unit sphere in $\mathring{\mathcal{U}}$ and $\operatorname{proj}\colon \mathring{\mathcal{U}} \looparrowright \mathcal{U}$ the submap, defined as $\operatorname{proj}(v) = \iota^{-1}(v/|v|)$ for all $v \neq 0$. Note that there is $z \in \mathcal{U}$ and $\varepsilon > 0$ such that the set

$$K_\varepsilon = \left\{ v \in \mathring{\mathcal{U}} : \langle \iota(z), v\rangle \geqslant \varepsilon \right\}$$

contains all $\iota(a^i)$. Note that $0 \notin K_\varepsilon$ and the set K_ε is closed and convex. The later follows from Exercise 8.5.2, since $v \mapsto -\langle \iota(z), v\rangle$ is a Busemann function.

Denote by $\iota(\boldsymbol{a})$ the point array $(\iota(a^0), \iota(a^1), \dots, \iota(a^{\bar{k}}))$ in $\mathring{\mathcal{U}}$. From the case $\kappa = 0$, we get that $\operatorname{Im} \mathfrak{S}^0_{\iota(\boldsymbol{a})} \subset K_\varepsilon$. In particular $\operatorname{Im} \mathfrak{S}^0_{\iota(\boldsymbol{a})} \not\ni 0$ and thus $\operatorname{proj} \circ \mathfrak{S}^0_{\iota(\boldsymbol{a})}$ is defined. Direct calculations show

$$\mathfrak{S}^1_{\boldsymbol{a}} = \operatorname{proj} \circ \mathfrak{S}^0_{\iota(\boldsymbol{a})} \quad \text{and} \quad \operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}}) = \operatorname{proj}[\operatorname{Web}(\operatorname{dist}_{\iota(\boldsymbol{a})})].$$

It reduces the case $\kappa = 1$ of the theorem to the case $\kappa = 0$, which is proved already. $\square$

12.2.5. Lemma. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space, $\boldsymbol{a} = (a^0, a^1, \dots a^{\bar{k}})$ be an array of radius $< \frac{\varpi^\kappa}{2}$ and $B^i = \overline{\mathrm{B}}[a^i, r^i]$ for some array of positive reals $(r^0, r^1, \dots, r^{\bar{k}})$. Assume that $\bigcap_i B^i = \varnothing$, but $\bigcap_{i \neq j} B^i \neq \varnothing$ for any j. Then $\boldsymbol{a}$ is nondegenerate.*

Proof. Without loss of generality we can assume that $\mathcal{U}$ is geodesic and $\operatorname{diam} \mathcal{U} < \varpi^\kappa$. If not, choose a point $z \in \mathcal{U}$ and $r < \frac{\varpi^\kappa}{2}$ such that $|z - a^i| \leqslant r$ for each i and consider $\overline{\mathrm{B}}[z, r]$ instead of $\mathcal{U}$. The later can be done since $\overline{\mathrm{B}}[z, r]$ is convex and closed, thus $\overline{\mathrm{B}}[z, r]$ is a complete length CAT(κ) space and $\operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}}) \subset \overline{\mathrm{B}}[z, r]$; see 8.6.1 and 12.2.1*b*.

By Theorem 12.2.1, $\operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}})$ is a compact set, therefore there is a point $p \in \operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}})$ minimizing the function

$$f(x) = \max_i \{\operatorname{dist}_{B^i} x\} = \max\{0, |a^0 - x| - r^0, \dots, |a^{\bar{k}} - x| - r^{\bar{k}}\}.$$

By the definition of web (12.1.2), p is also the minimum point of f on $\mathcal{U}$. Let us show that

❶ *$p \notin B^j$ for any j.*

Indeed, assume the contrary; that is,

❷ $$p \in B^j$$

for some j. Then p is a point of local minimum for the function

$$h^j(x) = \max_{i \neq j} \{\operatorname{dist}_{B^i} x\}.$$

Hence

$$\max_{i \neq j} \{\measuredangle[p\,{}^{x}_{a^i}]\} \geqslant \tfrac{\pi}{2}$$

for any $x \in \mathcal{U}$. From angle comparison (8.3.1*c*), it follows that p is a global minimum of h^j and hence

$$p \in \bigcap_{i \neq j} B^i.$$

The latter and ❷ contradict $\bigcap_i B^i = \varnothing$. $\triangle$

From the definition of web, it also follows that

$$\mathrm{Web}(\mathrm{dist}_{\boldsymbol{a}^{-j}}) \subset \bigcup_{i\neq j} B^i.$$

Therefore, the claim implies that $p \notin \mathrm{Web}(\mathrm{dist}_{\boldsymbol{a}^{-j}})$ for each j; that is, $p \in \mathrm{InWeb}(\mathrm{dist}_{\boldsymbol{a}})$. $\square$

12.3 Dimension of CBA spaces

We start with two examples.

The first example shows that the dimension of a complete length CAT spaces is not local; that is, such spaces might have open sets with different linear dimension.

Such an example can be constructed by gluing at one point two Euclidean spaces of different dimensions. According to Reshetnyak's gluing theorem (8.9.1), this construction gives a CAT(0) spaces.

The second example provides a complete length CAT space with topological dimension one and arbitrary large Hausdorff dimension. Thus, for complete length CAT spaces, one should not expect any relations between topological and Hausdorff dimensions except for the one provided by Szpilrajn's theorem (6.1.5).

To construct the second type of examples, note that the completion of any metric tree has topological dimension one and is CAT(κ) for any κ. Start with a binary tree Γ, and a sequence $\varepsilon_n > 0$ such that $\sum_n \varepsilon_n < \infty$. Define the metric on Γ by prescribing length of edge from level n to the level $n+1$ to be ε_n. For an appropriately chosen sequence ε_n, the completion of Γ will contain a Cantor set of arbitrary large dimension.

The following is a version of the theorem proved by Bruce Kleiner in [13] with an improvement made by Alexander Lytchak in [50].

12.3.1. Theorem. *For any complete length* CAT(κ) *space* $\mathcal{U}$*, the following statements are equivalent:*

a) $\mathrm{LinDim}\,\mathcal{U} \geqslant m$*;*

b) For some $z \in \mathcal{U}$ *there is an array of* $m+1$ *balls* $B^i = \mathrm{B}(a^i, r^i)$ *with* $a^0, a^1, \dots, a^m \in \mathrm{B}(z, \frac{\varpi^\kappa}{2})$ *such that*

$$\bigcap_i B^i = \varnothing \quad \textit{and} \quad \bigcap_{i\neq j} B^i \neq \varnothing \quad \textit{for each } j.$$

c) There is a $C^{\frac{1}{2}}$-embedding $\Phi\colon \overline{\mathrm{B}}[1]_{\mathbb{E}^m} \hookrightarrow \mathcal{U}$; that is, Φ is bi-Hölder with exponent $\frac{1}{2}$.

d) There is a closed separable set $K \subset \mathcal{U}$ such that

$$\operatorname{TopDim} K \geqslant m.$$

Remarks. Theorem 12.4.1 gives a stronger version of part (c) in finite dimensional case. Namely, a complete length CAT space with linear dimension m admits a bi-Lipschitz embedding Φ of an open set of $\mathbb{R}^m$. Moreover, the Lipschitz constants of Φ can be made arbitrary close to 1.

12.3.2. Corollary. *For any separable complete length* CAT *space $\mathcal{U}$, we have*

$$\operatorname{TopDim}\mathcal{U} = \operatorname{LinDim}\mathcal{U}.$$

Any simplicial complex can be equipped with a length-metric such that each k-simplex is isometric to the standard simplex

$$\Delta^k = \left\{ (x_0, \dots, x_k) \in \mathbb{R}^{k+1} : x_i \geqslant 0, \quad x_0 + \ldots + x_k = 1 \right\}$$

with the metric induced by the ℓ^1-norm on $\mathbb{R}^{k+1}$. This metric will be called the *ℓ^1-metric* on the simplicial complex.

12.3.3. Lemma. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space and $\rho\colon \mathcal{U} \to \mathbb{R}$ be a continuous positive function. Then there is a simplicial complex $\mathcal{N}$ equiped with ℓ^1-metric, a locally Lipschitz map $\Phi\colon \mathcal{U} \to \mathcal{N}$ and a Lipschitz map $\Psi\colon \mathcal{N} \to \mathcal{U}$ such that*

a) The displacement *of composition $\Psi \circ \Phi\colon \mathcal{U} \to \mathcal{U}$ is bounded by ρ; that is,*

$$|x - \Psi \circ \Phi(x)| < \rho(x)$$

for any $x \in \mathcal{U}$

b) If $\operatorname{LinDim}\mathcal{U} \leqslant m$ then the Ψ-image of any closed simplex in $\mathcal{N}$ coincides with the image of its m-skeleton.

Proof. Without loss of generality, we can assume that for any x we have $\rho(x) < \rho_0$ for some fixed $\rho_0 < \frac{\varpi^\kappa}{2}$.

By Stone's theorem, any metric space is paracompact. Thus, we can choose a locally finite covering $\{\, \Omega_\alpha : \alpha \in \mathcal{A} \,\}$ of $\mathcal{U}$ such that $\Omega_\alpha \subset \subset \mathrm{B}(x, \frac{1}{3}\cdot\rho(x))$ for any $x \in \Omega_\alpha$.

Denote by $\mathcal{N}$ the nerve of the covering $\{\Omega_\alpha\}$; that is, $\mathcal{N}$ is an abstract simplicial complex with set of vertexes formed by $\mathcal{A}$ such that $\{\alpha^0, \alpha^1, \dots, \alpha^n\} \subset \mathcal{A}$ are vertexes of a simplex if and only if $\Omega_{\alpha^0} \cap \Omega_{\alpha^1} \cap \ldots \cap \Omega_{\alpha^n} \neq \varnothing$.

Fix a Lipschitz partition of unity $\varphi_\alpha\colon \mathcal{U} \to [0,1]$, subordinate to $\{\Omega_\alpha\}$. Consider the map $\Phi\colon \mathcal{U} \to \mathcal{N}$ such that the barycentric coordinate of $\Phi(p)$ is $\varphi_\alpha(p)$. Note that Φ is locally Lipschitz. Clearly Φ-preimage of any open simplex in $\mathcal{N}$ lies in Ω_α for some $\alpha \in \mathcal{A}$.

For each $\alpha \in \mathcal{A}$, choose $x_\alpha \in \Omega_\alpha$. Let us extend the map $\alpha \mapsto \mapsto x_\alpha$ to a map $\Psi\colon \mathcal{N} \to \mathcal{U}$ that is κ-barycentric on each simplex; according to Theorem 12.2.1a, this extension exists, Ψ is Lipschtz and its Lipschitz constant depends only on ρ_0 and κ.

(a) Fix $x \in \mathcal{U}$. Denote by Δ the minimal simplex that contains $\Phi(x)$; and let $(\alpha^0, \alpha^1, \dots, \alpha^n)$ be the vertexes of Δ. Note that α is a vertex of Δ if and only if $\varphi_\alpha(x) > 0$. Thus,

$$|x - x_{\alpha^i}| < \tfrac{1}{3}\cdot\rho(x)$$

for any i. Therefore

$$\operatorname{diam}\Psi(\Delta) \leqslant \max_{i,j}\{|x_{\alpha^i} - x_{\alpha^j}|\} < \tfrac{2}{3}\cdot\rho(x).$$

In particular

$$|x - \Psi\circ\Phi(x)| \leqslant |x - x_{\alpha^0}| + \operatorname{diam}\Psi(\Delta) < \rho(x).$$

(b) Assume the contrary; that is, $\Psi(\mathcal{N})$ is not inculded in the Ψ-image of m-skeleton of $\mathcal{N}$. Then for some $\bar{k} > m$, there is a $\bar{k}$-simplex $\Delta^{\bar{k}}$ in $\mathcal{N}$, such that the baricentric simplex $\sigma = \Psi|_{\Delta^{\bar{k}}}$ is nondegenrate; that is,

$$W = \Psi(\Delta^{\bar{k}})\backslash\Psi(\partial\Delta^{\bar{k}}) \neq \varnothing.$$

Applying Corollary 12.2.4, we get that $\operatorname{LinDim}\mathcal{U} \geqslant \bar{k}$, a contradiction. □

Proof of 12.3.1. Note that

- ⋄ The implication $(b)\Rightarrow(c)$ follows directly from Lemma 12.2.5 and Theorem 12.2.1c.
- ⋄ The implication $(c)\Rightarrow(d)$ is trivial.

(d)⇒(a). According to Theorem 6.1.7, there is a continuous map $f\colon K \to \mathbb{R}^m$ with a stable value. By Tietze extension theorem, it is possible to extend f to a continuous map $F\colon \mathcal{U} \to \mathbb{R}^m$.

Fix $\varepsilon > 0$. Since F is continuous, there is a continuous positive function ρ defined on $\mathcal{U}$ such that

$$|x - y| < \rho(x) \quad\Rightarrow\quad |F(x) - F(y)| < \tfrac{1}{3}\cdot\varepsilon.$$

Apply Lemma 12.3.3 for the function ρ. For the obtained simplicial complex $\mathcal{N}$, and maps $\Phi\colon \mathcal{U} \to \mathcal{N}$, $\Psi\colon \mathcal{N} \to \mathcal{U}$, we have

$$|F\circ\Psi\circ\Phi(x) - F(x)| < \tfrac{1}{3}\cdot\varepsilon$$

for any $x \in \mathcal{U}$.

According to Lemma 4.2.5, there is a locally Lipschitz map $F_\varepsilon : \mathcal{U} \to \to \mathbb{R}^{m+1}$ such that $|F_\varepsilon(x) - F(x)| < \frac{1}{3}\cdot\varepsilon$ for any $x \in \mathcal{U}$.

Note that $\Phi(K)$ is contained in a countable subcomplex of $\mathcal{N}$, say $\mathcal{N}'$. Indeed, since K is separable, there is a countable dense collection of points $\{x_n\}$ in K. Denote by Δ_n the minimal simplex of $\mathcal{N}$ that contains $\Phi(x_n)$. Then $\Phi(K) \subset \bigcup_i \Delta_n$.

Arguing by contradiction, assume $\operatorname{LinDim}\mathcal{U} < m$. By 12.3.3*b*, the image $F_\varepsilon \circ \Psi \circ \Phi(K)$ lies in the F_ε-image of $(m-1)$-skeliton of $\mathcal{N}'$; in particular it can be covered by countable collection of Lipschitz images of $(m-1)$-simplexes. Hence $\mathbf{0} \in \mathbb{R}^m$ is not a stable value of the restriction $F_\varepsilon \circ \Psi \circ \Phi|_K$. Since $\varepsilon > 0$ is arbitrary, we get that $\mathbf{0} \in \mathbb{R}^m$ is not a stable value of $f = F|_K$, a contradiction.

(a)⇒(b). The following claim follows from the definition of tangent space.

❶ *Let $q \in \mathcal{U}$ and $\dot x^1, \dot x^2, \dots, \dot x^n \in \mathrm{T}_q$. Then given $\delta > 0$, there is a point array $x^1, x^2, \dots, x^n$ in $\mathcal{U}$ such that*

$$\measuredangle(\dot x^i, \log[q x^i]) < \delta$$

and for some fixed $\mathfrak{z} > 0$ we have

$$\tfrac{1}{\mathfrak{z}}\cdot|q - x^i| = |\dot x^i| \quad \text{and} \quad |\tfrac{1}{\mathfrak{z}}\cdot|x^i - x^j| - |\dot x^i - \dot x^j|| < \delta$$

for all i and j.

Moreover the value $\mathfrak{z}$ can be taken arbitrary small.

Proof of the claim. For each i choose a geodesic γ^i from q that goes almost in the directions of $\dot x^i$. Then take the point x^i on γ^i on the distance $\mathfrak{z}\cdot|\dot x^i|$ from q. △

Choose $q \in \mathcal{U}$ such that T_q contains a subcone E isometric to m-dimensional Euclidean space. Note that one can choose $\varepsilon > 0$ and a point arrray $(\dot a^0, \dot a^1, \dots, \dot a^m)$ in $E \subset \mathrm{T}_q$ such that $\bigcap_i \overline{\mathrm{B}}[\dot a^i, 1+\varepsilon] = \varnothing$ and $\bigcap_{i\ne j} \overline{\mathrm{B}}[\dot a^i, 1-\varepsilon] \ne \varnothing$ for each j.

Applying Claim ❶, we get a point array $(a^0, a^1, \dots, a^m)$ in $\mathcal{U}$ such that $\bigcap_i \overline{\mathrm{B}}[a^i, \mathfrak{z}] = \varnothing$ and $\bigcap_{i\ne j} \overline{\mathrm{B}}[a^i, \mathfrak{z}] \ne \varnothing$ for each j. Since $\mathfrak{z} > 0$ can be chousen arbitrary small, it implies (b). □

12.4 Finite-dimensional CAT spaces

Recall that web embedding and its brace is defined in 12.2.2.

12.4.1. Theorem. *Let $\mathcal{U}$ be a complete length* CAT(κ) *space,* $\operatorname{LinDim}\mathcal{U} = m$, *and $\boldsymbol{a} = (a^0, a^1, \dots a^m)$ be a point array in $\mathcal{U}$ with*

radius $< \frac{\varpi^\kappa}{2}$. *Then the* $\operatorname{dist}_{\boldsymbol{a}}$*-web embedding* $\Phi\colon \mathbb{R}^m \hookrightarrow \mathcal{U}$ *with brace* $\operatorname{dist}_{a^0}$ *is locally Lipschitz.*

Note that if $\boldsymbol{a}$ is degenerate, that is, if $\operatorname{InWeb}(\operatorname{dist}_{\boldsymbol{a}}) = \varnothing$, then the domain of the web embedding Φ above is empty and hence the conclusion of the theorem trivially holds.

12.4.2. Lemma. *Let* $\mathcal{U}$ *be a complete length* $\mathrm{CAT}(\kappa)$ *space, and* $\boldsymbol{a} = (a^0, a^1, \dots a^{\bar{k}})$ *be a point array with radius* $< \frac{\varpi^\kappa}{2}$. *Then for any* $p \in \operatorname{InWeb}(\operatorname{dist}_{\boldsymbol{a}})$, *there is* $\varepsilon > 0$ *such that if for some* $q \in \operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}})$ *and* $b \in \mathcal{U}$ *we have*

$$|p - q| < \varepsilon, \quad |p - b| < \varepsilon \quad \textit{and} \quad \measuredangle[q\,{}^{b}_{a^i}] < \tfrac{\pi}{2} + \varepsilon$$

for each i, *then the array* $b, a^0, a^1, \dots, a^m$ *is nondegenerate.*

Proof. Without loss of generality, we may assume that $\mathcal{U}$ is geodesic and $\operatorname{diam}\mathcal{U} < \varpi^\kappa$. If not, consider instead of $\mathcal{U}$, a ball $\overline{\mathrm{B}}[z, r] \subset \mathcal{U}$ for some $z \in \mathcal{U}$ and $r < \frac{\varpi^\kappa}{2}$ such that $|z - a^i| \leqslant r$ for each i.

From angle comparison (8.3.1c), it follows that $p \in \operatorname{InWeb}\boldsymbol{a}$ if and only if both of the following conditions hold:

1. $\max_i\{\measuredangle[p\,{}^{a^i}_{u}]\} \geqslant \frac{\pi}{2}$ for any $u \in \mathcal{U}$,
2. for each i there is $u^i \in \mathcal{U}$ such that $\measuredangle[p\,{}^{a^j}_{u^i}] < \frac{\pi}{2}$ for all $j \neq i$.

Due to the semicontinuity of angles (8.14.1), there is $\varepsilon > 0$ such that for any $x \in \mathrm{B}(p, 10\cdot\varepsilon)$ we have

❶ $$\measuredangle[x\,{}^{a^j}_{u^i}] < \tfrac{\pi}{2} - 10\cdot\varepsilon \quad \text{for all} \quad j \neq i.$$

Now assume for sufficiently small $\varepsilon > 0$ there are points $b \in \mathcal{U}$ and $q \in \operatorname{Web}(\operatorname{dist}_{\boldsymbol{a}})$ such that

❷ $$|p - q| < \varepsilon, \quad |p - b| < \varepsilon, \quad \measuredangle[q\,{}^{b}_{a^i}] < \tfrac{\pi}{2} + \varepsilon \quad \text{for all} \quad i.$$

Acording to Theorem 12.2.1b, for all small $\varepsilon > 0$, we have

$$\operatorname{rad}\{b, a^0, a^1, \dots, a^{\bar{k}}\} < \tfrac{\varpi^\kappa}{2}.$$

Fix sufficiently small $\delta > 0$ and set

$$v^i = \operatorname{geod}_{[qu^i]}(\tfrac{1}{3}\cdot\delta) \quad \text{and} \quad w^i = \operatorname{geod}_{[v^i b]}(\tfrac{2}{3}\cdot\delta).$$

Clearly,

$$\begin{aligned}|b - w^i| &= |b - v^i| - \tfrac{2}{3}\cdot\delta \leqslant \\ &\leqslant |b - q| - \tfrac{1}{3}\cdot\delta.\end{aligned}$$

Further, the inequalities ❶ and ❷ imply

$$\begin{aligned}|a^j - w^i| &< |a^j - v^i| + \tfrac{2}{3}\cdot\varepsilon\cdot\delta < \\ &< |a^i - q| - \varepsilon\cdot\delta < \\ &< |a^i - q|\end{aligned}$$

for all $i \neq j$.

Set $B^i = \overline{\mathrm{B}}[a^i, |a^i - q|]$ and $B^{m+1} = \overline{\mathrm{B}}[b, |a^i - q| - \frac{1}{3}\cdot\delta]$. Clearly

$$\bigcap_{i\neq m+1} B^i = \{q\},$$

$$\bigcap_{i\neq j} B^i \ni w^j \quad \text{for} \quad j \neq m+1.$$

$$\bigcap_{i} B^i = \{q\} \cap B^{m+1} = \varnothing.$$

Lemma 12.2.5 finishes the proof. □

Proof of 12.4.1. From 12.2.2, it is sufficient to show that the Φ is locally Lipshitz. Assume the contrary; that is, there are sequences $\boldsymbol{y}_n, \boldsymbol{z}_n \to \boldsymbol{x} \in \operatorname{Dom}\Phi$ such that

❸ $$\frac{|\Phi(\boldsymbol{y}_n) - \Phi(\boldsymbol{z}_n|)}{|\boldsymbol{y}_n - \boldsymbol{z}_n|} \to \infty \quad \text{as} \quad n \to \infty.$$

Set $p = \Phi(\boldsymbol{x})$, $q_n = \Phi(\boldsymbol{y}_n)$, and $b_n = \Phi(\boldsymbol{z}_n)$. From 12.2.2, p, q_n, $b_n \in$ $\in \operatorname{InWeb}(\operatorname{dist}_{\boldsymbol{a}})$ and $q_n, b_n \to p$ as $n \to \infty$. Fix arbitrary $\varepsilon > 0$. Note that ❸ implies

$$\measuredangle[q_n\, {}^{a^i}_{b_n}] < \tfrac{\pi}{2} + \varepsilon$$

for all $i > 0$ and all large n. Further, according to 12.2.2, the subfunction $(\operatorname{dist}_{\boldsymbol{a}^0}) \circ \Phi$ is locally Lipschitz. Therefore we also have

$$\measuredangle[q_n\, {}^{a^0}_{b_n}] < \tfrac{\pi}{2} + \varepsilon$$

for all large n. According to Lemma 12.4.2, the point array $b_n, a^0, \dots$ $\dots, a^k$ for large n is nondegenerate.

Applying Corollary 12.2.4, we get a contradiction □

12.5 Remarks and open problems

The following conjecture (in an equivalent form) appers in [13], see also [51, p. 133].

12.5.1. Conjecture. *For any complete length* CAT *space* $\mathcal{U}$*, we have*

$$\operatorname{TopDim}\mathcal{U} = \operatorname{LinDim}\mathcal{U}.$$

By Corollary 12.3.2 it holds for separable spaces

Chapter 13

Dimension of CBB spaces

In sections 13.3 and 13.4, we prove equivalence of some dimension-like invariants for CBB spaces. In Section 12.3 we discuss the dimension of CAT spaces.

13.1 Struts and rank

Our definitions of strut and distant chart differ from the one given by Burago, Gromov and Perelman in [3]; it is closer to Perelman's definitions [52, 53].

The term "strut" seems to have the closest meaning to the original Russian term used in [3]. In the official translation, it appears under name "burst" and in the authors translation it was "strainer". It seems that both are not good, so we decided to switch to "strut".

13.1.1. Definition of struts. *Let $\mathcal{L}$ be a complete length* CBB *space. We say that a point array $(a^0, a^1, \dots, a^k)$ in $\mathcal{L}$ is* κ-strutting *for a point $p \in \mathcal{L}$ if $\measuredangle^\kappa(p\,{}^{a^i}_{a^j}) > \frac{\pi}{2}$ for all $i \neq j$.*

13.1.2. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and $p \in$ $\in \mathcal{L}$. Let us define rank of $\mathcal{L}$ at p as*

$$\operatorname{rank}_p = \operatorname{rank}_p \mathcal{L} := \operatorname{pack}_{\pi/2} \Sigma_p - 1.$$

Thus rank takes values in $\mathbb{Z}_{\geqslant 0} \cup \{\infty\}$.

13.1.3. Proposition. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $p \in \mathcal{L}$. Then the following conditions are equivalent:*

a) $\operatorname{rank}_p \geqslant \bar{k}$

b) There is a κ*-strutting at* p *point array* $(a^0, a^1, \dots, a^{\bar{k}})$.

Proof of 13.1.3, (b)⇒(a). For each i, choose a point $\acute{a}^i \in \operatorname{Str}(p)$ sufficiently close to a^i (so $[p\acute{a}^i]$ exists for each i). One can choose $\acute{a}^i$ on such a way that we still have $\measuredangle^\kappa(p\,{}^{\acute{a}^i}_{\acute{a}^j}) > \frac{\pi}{2}$ for all $i \neq j$.

From hinge comparison (7.3.1*c*) we have

$$\measuredangle(\uparrow_{[p\acute{a}^j]}, \uparrow_{[p\acute{a}^j]}) \geqslant \measuredangle^\kappa(p\,{}^{\acute{a}^i}_{\acute{a}^j}) > \tfrac{\pi}{2}$$

for all $i \neq j$. In particular $\operatorname{pack}_{\pi/2} \Sigma_p \geqslant \bar{k} + 1$.

(a)⇒(b). Assume $\xi^0, \xi^1, \dots, \xi^{\bar{k}}$ is an array of directions in Σ_p, such that $\measuredangle(\xi^i, \xi^j) > \frac{\pi}{2}$ if $i \neq j$.

Without loss of generality, we may assume that each direction ξ^i is geodesic; that is, for each i there is a geodesic γ^i in $\mathcal{L}$ such that $\gamma^i(0) = p$ and $\xi^i = (\gamma^i)^+(0)$. From the definition of angle, it follows that for sufficiently small $\varepsilon > 0$ the array of points $a^i = \gamma^i(\varepsilon)$ satisfies (*b*). □

13.1.4. Corollary. *Let* $\mathcal{L}$ *be a complete length* CBB *space and* $\bar{k} \in \mathbb{Z}_{\geqslant 0}$. *Then the set of all points in* $\mathcal{L}$ *with rank* $\geqslant \bar{k}$ *is open.*

Proof. Given an array of points $\boldsymbol{a} = (a^0, \dots, a^{\bar{k}})$ in $\mathcal{L}$ consider the set $\Omega_{\boldsymbol{a}}$ of all points $p \in \mathcal{L}$ such that array $\boldsymbol{a}$ is κ-strutting for a point p. Cleary $\Omega_{\boldsymbol{a}}$ is open.

According to Proposition 13.1.3, the set of points in $\mathcal{L}$ with rank $\geqslant \bar{k}$ can be presented as

$$\bigcup_{\boldsymbol{a}} \Omega_{\boldsymbol{a}},$$

where the union is taken for all $\bar{k}$-array $\boldsymbol{a}$ of points in $\mathcal{L}$, hence the result. □

13.2 Right-inverse theorem

13.2.1. Right-inverse theorem. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space,* $p, b \in \mathcal{L}$ *and* $\boldsymbol{a} = (a^1, \dots, a^{\bar{k}})$ *be a point array in* $\mathcal{L}$.

Assume that $(b, a^1, a^2, \dots, a^{\bar{k}})$ *is* κ*-strutting for* p. *Then the distance map* $\operatorname{dist}_{\boldsymbol{a}} : \mathcal{L} \to \mathbb{R}^{\bar{k}}$ *has a right inverse defined in a neighborhood of* $\operatorname{dist}_{\boldsymbol{a}} p \in \mathbb{R}^{\bar{k}}$; *that is, there is a submap* $\Phi\colon \mathbb{R}^{\bar{k}} \hookrightarrow \mathcal{L}$ *such that* $\operatorname{Dom}\Phi \ni \operatorname{dist}_{\boldsymbol{a}} p$ *and* $\operatorname{dist}_{\boldsymbol{a}}[\Phi(\boldsymbol{x})] = \boldsymbol{x}$ *for any* $\boldsymbol{x} \in \operatorname{Dom}\Phi$. *Moreover*

a) The map Φ can be chosen to be $C^{\frac{1}{2}}$-continuous (that is, Hölder continuous with exponent $\frac{1}{2}$) and such that

$$\Phi(\operatorname{dist}_{\boldsymbol{a}} p) = p.$$

b) The distance map $\operatorname{dist}_{\boldsymbol{a}} : \mathcal{L} \to \mathbb{R}^{\bar{k}}$ is locally co-Lipschitz (in particular, open) in a neighborhood of p.

Part b of the theorem is closely related to [3, Theorem 5.4] by Burago, Gromov and Perelman, but the proof presented here is different; yet alternative proof can be build on [54, Proposition 4.3] by Lytchak.

Proof. Fix some $\varepsilon, r, \lambda > 0$ such that the following conditions hold:

(i) Each distance function $\operatorname{dist}_{a^i}$ and dist_b is $\frac{\lambda}{2}$-concave in $\mathrm{B}(p,r)$.

(ii) For any $q \in \mathrm{B}(p,r)$, we have $\measuredangle^{\kappa}(q\,{}^{a^i}_{a^j}) > \frac{\pi}{2} + \varepsilon$ for all $i \neq j$ and $\measuredangle^{\kappa}(q\,{}^{b}_{a^i}) > \frac{\pi}{2} + \varepsilon$ for all i. In addition $\varepsilon < \frac{1}{10}$.

Given $\boldsymbol{x} = (x^1, x^2, \dots, x^{\bar{k}}) \in \mathbb{R}^{\bar{k}}$ consider the function $f_{\boldsymbol{x}} \colon \mathcal{L} \to \mathbb{R}$ defined as

$$f_{\boldsymbol{x}} = \min_i\{h^i_{\boldsymbol{x}}\} + \varepsilon \cdot \operatorname{dist}_b\,,$$

where $h^i_{\boldsymbol{x}}(q) = \min\{0, |a^i - q| - x^i\}$. Note that for any $\boldsymbol{x} \in \mathbb{R}^{\bar{k}}$, the function $f_{\boldsymbol{x}}$ is $(1+\varepsilon)$-Lipschitz and λ-concave in $\mathrm{B}(p,r)$. Denote by $\alpha_{\boldsymbol{x}}(t)$ the $f_{\boldsymbol{x}}$-gradient curve that starts at p.

❶ *If for some $\boldsymbol{x} \in \mathbb{R}^{\bar{k}}$ and $t_0 \leqslant \frac{r}{2}$, we have $|\operatorname{dist}_{\boldsymbol{a}} p - \boldsymbol{x}| \leqslant \frac{\varepsilon^2}{10} \cdot t_0$ then $\operatorname{dist}_{\boldsymbol{a}}[\alpha_{\boldsymbol{x}}(t_0)] = \boldsymbol{x}$.*

First note that Claim ❶ follows if for any $q \in \mathrm{B}(p,r)$, we have

(i) $(\boldsymbol{d}_q \operatorname{dist}_{a^i})(\nabla_q f_{\boldsymbol{x}}) < -\frac{1}{10} \cdot \varepsilon^2$ if $|a^i - q| > x^i$ and

(ii) $(\boldsymbol{d}_q \operatorname{dist}_{a^i})(\nabla_q f_{\boldsymbol{x}}) > \frac{1}{10} \cdot \varepsilon^2$ if $|a^i - q| - x^i = \min_j\{|a^j - q| - x^j\} < 0$.

Indeed, since $t_0 \leqslant \frac{r}{2}$, $\alpha_{\boldsymbol{x}}(t) \in \mathrm{B}(p,r)$ for all $t \in [0, t_0]$. Consider the following real-to-real functions:

$$\begin{aligned} \varphi(t) &:= \max_i\{|a^i - \alpha_{\boldsymbol{x}}(t)| - x^i\}, \\ \psi(t) &:= \min_i\{|a^i - \alpha_{\boldsymbol{x}}(t)| - x^i\}. \end{aligned} \qquad ❷$$

Then from (i), we have $\varphi^+ < -\frac{1}{10} \cdot \varepsilon^2$ if $\varphi > 0$ and $t \in [0, t_0]$. The same way, from (ii), we have $\psi^+ > \frac{1}{10} \cdot \varepsilon^2$ if $\psi < 0$ and $t \in [0, t_0]$. Since $|\operatorname{dist}_{\boldsymbol{a}} p - \boldsymbol{x}| \leqslant \frac{\varepsilon^2}{10} \cdot t_0$, it follows that $\varphi(0) \leqslant \frac{\varepsilon^2}{10} \cdot t_0$ and $\psi(0) \geqslant -\frac{\varepsilon^2}{10} \cdot t_0$. Thus $\varphi(t_0) \leqslant 0$ and $\psi(t_0) \geqslant 0$. On the other hand, from ❷ we have $\varphi(t_0) \geqslant \psi(t_0)$. That is, $\varphi(t_0) = \psi(t_0) = 0$; hence Claim ❶ follows.

Thus, to prove Calim ❶, it remains to prove (i) and (ii). First let us prove it assuming that $\mathcal{L}$ is geodesic.

Note that

❸ $$(\boldsymbol{d}_q \operatorname{dist}_b)(\uparrow_{[qa^i]}) \leqslant \cos \measuredangle^{\kappa}(q\,{}^{b}_{a^j}) < -\tfrac{\varepsilon}{2},$$

for all i and

❹ $$(\boldsymbol{d}_q \operatorname{dist}_{a^j})(\uparrow_{[qa^i]}) \leqslant \cos \measuredangle^{\kappa}(q\,{}^{a^i}_{a^j}) < -\tfrac{\varepsilon}{2}$$

for all $j \neq i$. Further, ❹ implies

❺ $$(\boldsymbol{d}_q h^j_{\boldsymbol{x}})(\uparrow_{[qa^i]}) \leqslant 0.$$

for all $i \neq j$. The assumption in (i), implies

$$\boldsymbol{d}_q f_{\boldsymbol{x}} = \min_{j\neq i}\{\boldsymbol{d}_q h^j_{\boldsymbol{x}}\} + \varepsilon\cdot(\boldsymbol{d}_q \operatorname{dist}_b).$$

Thus,

$$\begin{aligned}-(\boldsymbol{d}_q \operatorname{dist}_{a^i})(\nabla_q f_{\boldsymbol{x}}) &\geqslant \langle \uparrow_{[qa^i]}, \nabla_q f_{\boldsymbol{x}}\rangle \geqslant \\ &\geqslant (\boldsymbol{d}_q f_{\boldsymbol{x}})(\uparrow_{[qa^i]}) = \\ &= \min_{i\neq j}\{(\boldsymbol{d}_q h^i_{\boldsymbol{x}})(\uparrow_{[qa^i]})\} + \varepsilon\cdot(\boldsymbol{d}_q \operatorname{dist}_b)(\uparrow_{[qa^i]}).\end{aligned}$$

Therefore (i) follows from ❸ and ❺.

The assumption in (ii) implies that $f_{\boldsymbol{x}}(q) = h^i_{\boldsymbol{x}}(q) + \varepsilon\cdot \operatorname{dist}_b$ and

$$\boldsymbol{d}_q f_{\boldsymbol{x}} \leqslant \boldsymbol{d}_q \operatorname{dist}_{a^i} + \varepsilon\cdot(\boldsymbol{d}_p \operatorname{dist}_b).$$

Therefore,

$$\begin{aligned}(\boldsymbol{d}_q \operatorname{dist}_{a^i})(\nabla_q f_{\boldsymbol{x}}) &\geqslant \boldsymbol{d}_q f_{\boldsymbol{x}}(\nabla_q f_{\boldsymbol{x}}) \geqslant \\ &\geqslant \Big[(\boldsymbol{d}_q f_{\boldsymbol{x}})(\uparrow_{[qb]})\Big]^2 \geqslant \\ &\geqslant \Big[\min_i\{\cos \measuredangle^{\kappa}(q\,{}^{b}_{a^i})\} - \varepsilon^2\Big]^2.\end{aligned}$$

Thus, (ii) follows from ❸ since $\varepsilon < \frac{1}{10}$.

That finishes the proof of ❶ in case if $\mathcal{L}$ is geodesic. If $\mathcal{L}$ is not geodesic, perform the above estimate in $\mathcal{L}^\omega$, the ultrapower of $\mathcal{L}$. (Recall that according to 3.5.2, $\mathcal{L}^\omega$ is geodesic.) △

Set $t_0(\boldsymbol{x}) = \frac{10}{\varepsilon^2}\cdot|\operatorname{dist}_{\boldsymbol{a}} p - \boldsymbol{x}|$; so we have equality in ❶. Define the submap

$$\Phi\colon \boldsymbol{x} \mapsto \boldsymbol{\alpha}_{\boldsymbol{x}} \circ t_0(\boldsymbol{x}), \quad \operatorname{Dom}\Phi = \mathrm{B}(\operatorname{dist}_{\boldsymbol{a}} p, \tfrac{\varepsilon^2\cdot r}{20}) \subset \mathbb{R}^{\bar{k}}.$$

It follows from Claim ❶, that $\operatorname{dist}_{\boldsymbol{a}}[\Phi(\boldsymbol{x})] = \boldsymbol{x}$ for any $\boldsymbol{x} \in \operatorname{Dom}\Phi$.

Clearly $t_0(p) = 0$; thus $\Phi(\operatorname{dist}_{\boldsymbol{a}} p) = p$. Further, by construction of $f_{\boldsymbol{x}}$,

$$|f_{\boldsymbol{x}}(q) - f_{\boldsymbol{y}}(q)| \leqslant |\boldsymbol{x} - \boldsymbol{y}|,$$

for any $q \in \mathcal{L}$. Therefore, according to Lemma 14.2.7, Φ is $C^{\frac{1}{2}}$-continuous. Thus, we proved (a).

Further, note that

❻ $$|p - \Phi(\boldsymbol{x})| \leqslant (1+\varepsilon)\cdot t_0(\boldsymbol{x}) \leqslant \tfrac{11}{\varepsilon^2}\cdot|\operatorname{dist}_{\boldsymbol{a}} p - \boldsymbol{x}|.$$

Note that one can repeat the above construction for any $p' \in \mathrm{B}(p, \frac{r}{4})$, $\varepsilon' = \varepsilon$ and $r' = \frac{r}{2}$. The inequality ❻, for the obtained map Φ', implies that for any $p', q \in \mathrm{B}(p, \frac{r}{4})$ there is $q' \in \mathcal{L}$ such that $\Phi'(q) = \Phi'(q')$ and

$$|p' - q'| \leqslant \tfrac{11}{\varepsilon^2}\cdot|\operatorname{dist}_{\boldsymbol{a}} p' - \boldsymbol{x}|.$$

That is, the distance map $\operatorname{dist}_{\boldsymbol{a}}$ is locally $\frac{11}{\varepsilon^2}$-co-Lipschitz in $\mathrm{B}(p, \frac{r}{4})$. □

13.3 Dimension theorem for CBB

As the main dimension-like invariant, we will use the linear dimension LinDim; see Definition 6.2.1, in other words by default *dimension* means *linear dimension.*

The following theorem is the main result of this section.

13.3.1. Theorem. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space, $q \in \mathcal{L}$, $R > 0$ and $m \in \mathbb{Z}_{\geqslant 0}$. Then the following statements are equivalent:*

A) $\operatorname{LinDim}\mathcal{L} \geqslant m$.

B) There is a point $p \in \mathcal{L}$ that admits a κ-strutting array $(b, a^1, \dots, a^m) \in \mathcal{L}^{m+1}$.

C) Let Euk^m be the set of all points $p \in \mathcal{L}$ such that there is an isometric cone embedding $\mathbb{E}^m \hookrightarrow \mathrm{T}_p$. Then Euk^m contains a dense G-delta set in $\mathcal{L}$.

D) There is a $C^{\frac{1}{2}}$-embedding; that is, a bi-Hölder embedding with exponent $\frac{1}{2}$,

$$\overline{\mathrm{B}}[1]_{\mathbb{E}^m} \hookrightarrow \mathrm{B}(q, R).$$

.

E)

$$\operatorname{pack}_\varepsilon \mathrm{B}(q, R) > \frac{c}{\varepsilon^m}$$

for some fixed $c > 0$ and any $\varepsilon > 0$.

In particular:

(i) If $\operatorname{LinDim}\mathcal{L}=\infty$*, then all the statements (C), (D) and (E) are satisfied for all* $m\in\mathbb{Z}_{\geqslant 0}$*.*

(ii) If the statement (D) or (E) is satisfied for some choice of $q\in\mathcal{L}$ *and* $R>0$*, then it also is satisfied for any other choice of* q *and* R*.*

For finite-dimensional spaces, Theorem 13.4.1 gives a stronger version of the theorem above.

The proof of the above theorem with exception of statement (D) was given in [20]. At that time, it was not known whether for any complete length $\mathrm{CBB}(\kappa)$ space $\mathcal{L}$,

$$\operatorname{LinDim}\mathcal{L}=\infty \quad\Rightarrow\quad \operatorname{TopDim}\mathcal{L}=\infty.$$

The latter implication was proved by Grigory Perelman [55]; it was done by combining an idea of Conrad Plaut with the technique of gradient flow. The statement 13.3.1D is somewhat stronger.

To prove Theorem 13.3.1 we will need the following three propositions.

13.3.2. Proposition. *Let* $\mathcal{L}$ *be a complete length* $\mathrm{CBB}(\kappa)$ *space and* $p\in\mathcal{L}$*. Assume there is an isometric cone embedding* $\iota\colon\mathbb{E}^m\hookrightarrow\mathrm{T}_p\mathcal{L}$*. Then either*

a) $\operatorname{Im}\iota=\mathrm{T}_p\mathcal{L}$*, or*

b) there is a point p' *arbitrarily close to* p *such that there is an isometric cone embedding* $\iota'\colon\mathbb{E}^{m+1}\hookrightarrow\mathrm{T}_{p'}\mathcal{L}$*.*

Proof. Assume $\iota(\mathbb{E}^m)$ is a proper subset of $\mathrm{T}_p\mathcal{L}$. Equivalently, there is a direction $\xi\in\Sigma_p\backslash\iota(\mathbb{S}^{m-1})$, where $\mathbb{S}^{m-1}\subset\mathbb{E}^m$ is the unit sphere.

Fix $\varepsilon>0$ so that $\measuredangle(\xi,\sigma)>\varepsilon$ for any $\sigma\in\iota(\mathbb{S}^{m-1})$. Choose a *maximal* ε*-packing* in $\iota(\mathbb{S}^{m-1})$; that is, an array of directions $\zeta^1,\zeta^2,\dots$ $\dots,\zeta^n\in\iota(\mathbb{S}^{m-1})$ so that $n=\operatorname{pack}_\varepsilon\mathbb{S}^{m-1}$ and $\measuredangle(\zeta^i,\zeta^j)>\varepsilon$ for any $i\neq j$.

Choose an array of points $x,z^1,z^2,\dots,z^n\in\mathcal{L}$ so that $\uparrow_{[px]}\approx\xi$, $\uparrow_{[pz^i]}\approx\zeta^i$; here write "$\approx$" for "sufficiently close". We can choose this array on such a way that $\measuredangle^\kappa(p\,{}^{x}_{z^i})>\varepsilon$ for all i and $\measuredangle^\kappa(p\,{}^{z^i}_{z^j})>\varepsilon$ for all $i\neq j$. Applying Corollary 11.6.5, we can find a point p' arbitrary close to p so that all directions $\uparrow_{[p'x]},\uparrow_{[p'z^1]},\uparrow_{[p'z^2]},\dots,\uparrow_{[p'z^n]}$ belong to an isometric copy of $\mathbb{S}^{\bar k-1}$ in $\Sigma_{p'}$. In addition, we can assume that $\measuredangle^\kappa(p'\,{}^{x}_{z^i})>\varepsilon$ and $\measuredangle^\kappa(p'\,{}^{z^i}_{z^j})>\varepsilon$. From hinge comparison (7.3.1c), $\measuredangle(\uparrow_{[p'x]},\uparrow_{[p'z^i]})>\varepsilon$ and $\measuredangle(\uparrow_{[p'z^i]},\uparrow_{[p'z^j]})>\varepsilon$; that is,

$$\operatorname{pack}_\varepsilon\mathbb{S}^{\bar k-1}\geqslant n+1>\operatorname{pack}_\varepsilon\mathbb{S}^{m-1}.$$

Hence $\bar{k} > m$. □

13.3.3. Proposition. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space, then for any two poins $p, \bar{p} \in \mathcal{L}$ and any $R, \bar{R} > 0$ there is a contsant $\delta = \delta(\kappa, R, \bar{R}, |p - \bar{p}|) > 0$ such that*

$$\operatorname{pack}_{\delta\cdot\varepsilon} \mathrm{B}(\bar{p}, \bar{R}) \geqslant \operatorname{pack}_{\varepsilon} \mathrm{B}(p, R).$$



Proof. According to 7.6.4, we can assume that $\kappa \leqslant 0$.

Let $n = \operatorname{pack}_{\varepsilon} \mathrm{B}(p, R)$ and $x^1, x^2, \dots, x^n \in \mathrm{B}(p, R)$ be a maximal ε-packing; that is, $|x^i - x^j| > \varepsilon$ for all $i \neq j$. Without loss of generality we can assume that $x^1, x^2, \dots, x^n \in \mathrm{Str}(\bar{p})$. Thus, for each i there is a unique geodesic $[\bar{p}x^i]$ (see 7.2.4). Choose factor $s > 0$, so that $\bar{r} > s\cdot(|p - \bar{p}| + R)$. For each i, take $\bar{x}^i \in [\bar{p}x^i]$ so that $|\bar{p} - \bar{x}^i| = s\cdot(|p - x^i|)$. From 7.3.2$a$,

$$\measuredangle^{\kappa}\left(\bar{p}\,{}^{\bar{x}^i}_{\bar{x}^j}\right) \geqslant \measuredangle^{\kappa}(\bar{p}\,{}^{x^i}_{x^j}).$$

Applying cosine rule, we get a constant $\delta = \delta(\kappa, R, \bar{R}, |p - \bar{p}|) > 0$ such that

$$|\bar{x}^i - \bar{x}^j| > \delta\cdot(|x^i - x^j|) > \delta\cdot\varepsilon$$

for all $i \neq j$. Hence the statement follows. □

13.3.4. Proposition. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space, $r < \varpi^{\kappa}$ and $p \in \mathcal{L}$. Assume that*

❶ $$\operatorname{pack}_{\varepsilon} \mathrm{B}(p, r) > \operatorname{pack}_{\varepsilon} \overline{\mathrm{B}}[r]_{\mathbb{M}^m(\kappa)}$$

for some $\varepsilon > 0$. Then there is a G-delta set $A \subset \mathcal{L}$, that is dense in a neighborhood of p and such that $\dim \mathrm{Lin}_q > m$ *for any $q \in A$.*

Proof. Choose a maximal ε-packing in $\mathrm{B}(p, r)$; that is, an array of points $x^1, x^2, \dots, x^n \in \mathrm{B}(p, r)$ so that $n = \operatorname{pack}_{\varepsilon} \mathrm{B}(p, r)$ and $|x^i - x^j| > \varepsilon$ for any $i \neq j$. Choose a neighborhood $\Omega \ni p$, such that $|q - x^i| < r$ for any $q \in \Omega$ and all i. Set

$$A = \Omega \cap \mathrm{Str}(x^1, x^2, \dots, x^n).$$

According to Theorem 7.2.4, A is a G-delta set taht is dense in Ω.

Assume $\bar{k} = \dim \mathrm{Lin}_q \leqslant m$ for some $q \in A$. Consider an array of vectors $v^1, v^2, \dots, v^n \in \mathrm{Lin}_q$, $v^i = \log[qx^i]$. Clearly

$$|v^i| = |q - x^i| < r$$

and from hinge comparison (7.3.1c) we have

$$\tilde{\Upsilon}^{\kappa}[o\,{}^{v^i}_{v^j}] \geqslant |x^i - x^j| > \varepsilon.$$

Note that the ball $\mathrm{B}(0,r)_{\mathrm{Lin}_q}$ equipped with the metric $\rho(v,w) = \tilde{\Upsilon}^{\kappa}[o\,{}^{v}_{w}]$ is isometric to $\overline{\mathrm{B}}[r]_{\mathbb{M}^{\bar{k}}(\kappa)}$. Thus,

$$\mathrm{pack}_\varepsilon\, \overline{\mathrm{B}}[r]_{\mathbb{M}^{\bar{k}}(\kappa)} \geqslant \mathrm{pack}_\varepsilon\, \mathrm{B}(p,r),$$

which contradicts $\bar{k} \leqslant m$ and ❶. □

The proof is essentially done in 13.3.2, 13.3.3, 13.3.4, 13.2.1; here we only assemble the proof from these parts.

We will prove implications

$$(C) \Rightarrow (A) \Rightarrow (B) \Rightarrow (E) \Rightarrow (C) \Rightarrow (D) \Rightarrow (E).$$

Proof of 13.3.1. The implication $(C)\Rightarrow(A)$ is trivial. The proof of $(D)\Rightarrow(E)$ is valid for general metric spaces; it is based on general relations between topological dimension, Hausdorff measure and $\mathrm{pack}_\varepsilon$.

(A)⇒(B). Choose a point $p \in \mathcal{L}$ such that $\dim \mathrm{Lin}_p \geqslant m$. Clearly one can choose an array of directions $\xi^0, \xi^1, \dots, \xi^m \in \mathrm{Lin}_p$ so that $\measuredangle(\xi^i, \xi^j) > \frac{\pi}{2}$ for all $i \neq j$. Choose an array of points $x^0, x^1, \dots, x^m \in \mathcal{L}$ so that each $\uparrow_{[px^i]}$ is sufficiently close to ξ^i; in particular, we have $\measuredangle[p\,{}^{x^i}_{x^j}] > \frac{\pi}{2}$. Choose points $a^i \in\,]px^i]$ sufficiently close to p. One can do it so that each $\measuredangle^{\kappa}(p\,{}^{a^i}_{a^j})$ is arbitrary close to $\measuredangle[p\,{}^{a^i}_{a^j}]$, in particular, $\measuredangle^{\kappa}(p\,{}^{a^i}_{a^j}) > \frac{\pi}{2}$. Finally, set $b = a^0$.

(B)⇒(E). Let $p \in \mathcal{L}$ be a point that admits a κ-strutting array $b, a^1, \dots, a^m \in \mathcal{L}$. The right-inverse mapping map theorem (13.2.1b) implies that the distance map $\mathrm{dist}_{\boldsymbol{a}} : \mathcal{L} \to \mathbb{R}^m$,

$$\mathrm{dist}_{\boldsymbol{a}} : x \mapsto (|a^1 - x|, |a^2 - x|, \dots, |a^n - x|)$$

is open in a neighborhood of p. Since the distance map $\mathrm{dist}_{\boldsymbol{a}}$ is Lipschitz, for any $r > 0$, there is $c > 0$ such that

$$\mathrm{pack}_\varepsilon\, \mathrm{B}(p,r) > \frac{c}{\varepsilon^m}.$$

Applying 13.3.3, we get similar inequality for any other ball in $\mathcal{L}$; that is, for any point $q \in \mathcal{L}$, $R > 0$ there is $c' > 0$ such that

$$\mathrm{pack}_\varepsilon\, \mathrm{B}(q,R) > \frac{c'}{\varepsilon^m}.$$

(E)⇒(C). Note that for any $q' \in \mathcal{L}$ and $R' > |q - q'| + R$ we have

$$\begin{aligned}\operatorname{pack}_\varepsilon \mathrm{B}(q', R') &\geqslant \operatorname{pack}_\varepsilon \mathrm{B}(q, R) \geqslant \\ &\geqslant \frac{c}{\varepsilon^m} > \\ &> \operatorname{pack}_\varepsilon \overline{\mathrm{B}}[R']_{\mathbb{M}^{m-1}(\kappa)}.\end{aligned}$$

for all sufficiently small $\varepsilon > 0$. Applying 13.3.4, we get that

$$\begin{gathered}\mathrm{Euk}^m = \\ \{\, p \in \mathcal{L} : \text{there is an isometric cone embedding } \iota : \mathbb{E}^m \hookrightarrow \mathrm{T}_p \,\}\end{gathered}$$

contains a G-delta set that is dense in a neighborhood of any point $q' \in \mathcal{L}$.

(C)⇒(D). Since Euk^m contains a dense G-delta set in $\mathcal{L}$, we can choose $p \in \mathrm{B}(q, R)$ with an isometric cone embedding $\iota\colon \mathbb{E}^m \hookrightarrow \mathrm{T}_p$.

Repeating the construction in (A)⇒(B), we get a κ-strutting array $b, a^1, \dots, a^m \in \mathcal{L}$ for p.

Applying the right-inverse theorem (13.2.1), we obtain a $C^{\frac{1}{2}}$-submap

$$\Phi\colon \mathbb{R}^m \looparrowright \mathrm{B}(q, R)$$

that is a right inverse for $\operatorname{dist}_{\boldsymbol{a}} : \mathcal{L} \to \mathbb{R}^m$ such that $\Phi(\operatorname{dist}_{\boldsymbol{a}} p) = p$. In particular, Φ is a $C^{\frac{1}{2}}$-embedding of $\operatorname{Dom}\Phi$.

(D)⇒(E). Let $W \subset \mathrm{B}(q, R)$ be the image of the embedding. Since $\operatorname{TopDim} W = m$, Szpilrajn's theorem (6.1.5) implies that

$$\operatorname{HausMes}_m W > 0.$$

Given $\varepsilon > 0$, consider a maximal ε-packing of W; that is, an array of points $(x^1, x^2, \dots, x^n)$ in W such that $n = \operatorname{pack}_\varepsilon W$ and $|x^i - x^j| > \varepsilon$ for all $i \neq j$. Note that W is covered by balls $\mathrm{B}(x^i, 2\cdot\varepsilon)$.

By the definition of Hausdorff measure,

$$\operatorname{pack}_\varepsilon W \geqslant \frac{c}{\varepsilon^m} \cdot \operatorname{HausMes}_m W$$

for a fixed constant $c > 0$ and all small $\varepsilon > 0$. Hence (E) follows. □

13.4 Finite dimensional CBB spaces

In this section we will show that all reasonable notions of dimension coincide on the class of Alexandrov spaces with curvature bounded below.

First we will prove a stronger version of Theorem 13.3.1 for the finite-dimensional case.

13.4.1. Theorem. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, m be a nonnegative integer, $0 < R \leqslant \varpi^\kappa$ and $q \in \mathcal{L}$. Then the following statements are equivalent:*

a) $\operatorname{LinDim}\mathcal{L} = m$*;*

b) m is the maximal integer, such that there is a point $p \in \mathcal{L}$ that admits a κ-strutting array $b, a^1, \dots, a^m$.

c) $\mathrm{T}_p \overset{iso}{=\!=} \mathbb{E}^m$ *for any point p in a dense G-delta set of $\mathcal{L}$.*

d) There is an open bi-Lipschitz embedding

$$\overline{\mathrm{B}}[1]_{\mathbb{E}^m} \hookrightarrow \mathrm{B}(q,R) \subset \mathcal{L};$$

e) For any $\varepsilon > 0$,

$$\operatorname{pack}_\varepsilon \overline{\mathrm{B}}[R]_{\mathbb{M}^m(\kappa)} \geqslant \operatorname{pack}_\varepsilon \mathrm{B}(q,R),$$

moreover, there is $c = c(q,R) > 0$ such that

$$\operatorname{pack}_\varepsilon \mathrm{B}(q,R) > \frac{c}{\varepsilon^m}.$$

The above theorem was essentially proved in [3].

Using theorems 13.3.1 and 13.4.1, one can show that linear dimension is equal to many different types of dimensions, such as *small* and *big inductive dimension* and *upper* and *lower box counting dimensions* (also known as *Minkowski dimensions*), *homological dimension* and so on.

The following two corollaries follow from 13.4.1*e*.

13.4.2. Corollary. *Any m-dimensional complete length* CBB *space is proper and geodesic.*

13.4.3. Corollary. *Let $(\mathcal{L}_n)$ be a sequence of length* CBB(κ) *spaces and $\mathcal{L}_n \to \mathcal{L}_\omega$ as $n \to \omega$. Assume* $\operatorname{LinDim} L_n \leqslant m$ *for all n. Then* $\operatorname{LinDim} L_\omega \leqslant m$.

13.4.4. Corollary. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space. Then for any open $\Omega \subset \mathcal{L}$, we have*

$$\operatorname{LinDim}\mathcal{L} = \operatorname{LinDim}\Omega = \operatorname{TopDim}\Omega = \operatorname{HausDim}\Omega,$$

where TopDim *and* HausDim *denotes topological dimension (6.1.2) and Hausdorff dimension (6.1.1) correspondingly.*

In particular, $\mathcal{L}$ is dimension homogeneous; that is, any open set has the same linear dimension.

Proof of 13.4.4. The equality

$$\operatorname{LinDim}\mathcal{L} = \operatorname{LinDim}\Omega$$

follows from 13.3.1A&C.

If $\operatorname{LinDim}\mathcal{L} = \infty$, then applying 13.3.1$D$, for $\mathrm{B}(q,R) \subset \Omega$, we get that there is a compact subset $K \subset \Omega$ with arbitrary large $\operatorname{TopDim} K$ therefore

$$\operatorname{TopDim}\Omega = \infty.$$

By Szpilrajn's theorem (6.1.5), $\operatorname{HausDim} K \geqslant \operatorname{TopDim} K$. Thus, we also have

$$\operatorname{HausDim}\Omega = \infty.$$

If $\operatorname{LinDim}\mathcal{L} = m < \infty$, then first inequality in 13.4.1e implies that

$$\operatorname{HausDim}\mathrm{B}(q,R) \leqslant m.$$

According to Corollary 13.4.2, $\mathcal{L}$ is proper and in particular it has countable base. Thus applying Szpilrajn theorem again, we get

$$\operatorname{TopDim}\Omega \leqslant \operatorname{HausDim}\Omega \leqslant m.$$

Finally, 13.4.1d implies that $m \leqslant \operatorname{TopDim}\Omega$. □

Proof of 13.4.1. The equivalence $(a)\Leftrightarrow(b)$ follows from 13.3.1.

(a)⇒(c). If $\operatorname{LinDim}\mathcal{L} = m$, then from Theorem 13.3.1, we get that Euk^m contains a dense G-delta set in $\mathcal{L}$. From 13.3.2, it follows that T_p is isometric to $\mathbb{E}^m$ for any $p \in \mathrm{Euk}^m$.

(c)⇒(d). It is proved exactly the same way as implication *(C)⇒(D)* of theorem 13.3.1, but we have to apply existence of distance chart instead of the right-inverse theorem.

(d)⇒(e). From (d), it follows that, there is a point $p \in \mathrm{B}(q,R)$ and $r > 0$ such that $\mathrm{B}(p,r) \subset \mathcal{L}$ is bi-Lipschit homeomorphic to a bounded open set of $\mathbb{E}^m$. Thus, there is $c > 0$ such that

❶ $$\operatorname{pack}_\varepsilon \mathrm{B}(p,r) > \frac{c}{\varepsilon^m}.$$

Applying 13.3.3, we get that inequality ❶, with different constants, holds for any other ball, in particular for $\mathrm{B}(q,R)$.

Applying 13.3.4, we get the first inequality.

(e)⇒(a). From theorem 13.3.1, we have $\operatorname{LinDim}\mathcal{L} \geqslant m$. Applying theorem 13.3.1 again, we get that if $\operatorname{LinDim}\mathcal{L} \geqslant m+1$, then for some $c > 0$ and any $\varepsilon > 0$

$$\operatorname{pack}_\varepsilon \mathrm{B}(q,R) \geqslant \frac{c}{\varepsilon^{m+1}},$$

but

$$\frac{c'}{\varepsilon^m} \geqslant \operatorname{pack}_\varepsilon \mathrm{B}(q,R),$$

for any $\varepsilon > 0$, a contradiction. □

The following exercise was suggested by Alexander Lytchak.

13.4.5. Exercise. *Let $\mathcal{L}$ be a complete length* CBB *space and $\Sigma_p\mathcal{L}$ is compact for any $p \in \mathcal{L}$. Prove that $\mathcal{L}$ is finite dimensional.*

13.5 One-dimensional CBB spaces

13.5.1. Theorem. *Let $\mathcal{L}$ be an one-dimensional complete length* CBB(κ) *space. Then $\mathcal{L}$ is isometric to a connected complete Riemannian* 1*-dimensional manifold with possibly non-empty boundary.*

Proof. Clearly $\mathcal{L}$ is connected. It remains to show the following.

❶ *For any point $p \in \mathcal{L}$ there is $\varepsilon > 0$ such that $\mathrm{B}(p,\varepsilon)$ is isometric to either $[0,\varepsilon)$ or $(-\varepsilon,\varepsilon)$.*

First let us show that

❷ *If $p \in\,]xy[$ for some x, $y \in \mathcal{L}$ and $\varepsilon < \min\{|p-x|, |p-y|\}$ then $\mathrm{B}(p,\varepsilon) \subset\,]xy[$. In particular $\mathrm{B}(p,\varepsilon) \overset{iso}{=\!=} (-\varepsilon,\varepsilon)$.*

Assume the contrary; that is, there is

$$z \in \mathrm{B}(p,\varepsilon)\backslash\,]xy[.$$



Consider a geodesic $[pz]$, let $q \in [pz]\cap[xy]$ be the point that maximizes the distance $|p-q|$. At q, we have three distinct directions: to x, to y and to z. Moreover, $\measuredangle[q\,{}^{x}_{y}] = \pi$. Thus, according to Proposition 13.3.2, $\operatorname{LinDim}\mathcal{L} > 1$, a contradiction.

Now assume there is no geodesic passing thru p. Since $\operatorname{LinDim}\mathcal{L} = 1$ there is a point $x \neq p$.

Fix a positive value $\varepsilon < |p-x|$. Let us show that

❸ $\mathrm{B}(p,\varepsilon)\subset[px]$; *in particular* $\mathrm{B}(p,\varepsilon)\overset{iso}{=\!=}[0,\varepsilon)$.



Assume the contrary; let $z\in\mathrm{B}(p,\varepsilon)\backslash[py]$. Choose a point $w\in\,]py[$ such that

$$|p-w|+|p-z|<\varepsilon.$$

Consider geodesic $[wz]$, let $q\in[py]\cap[wz]$ be the point that maximizes the distance $|w-q|$. Since no geodesics pass thru p, we have $p\neq q$. As above, $\measuredangle[q\,{}^{p}_{y}]=\pi$ and $\uparrow_{[qz]}$ is distinct from $\uparrow_{[qp]}$ and $\uparrow_{[qp]}$. Thus, according to Proposition 13.3.2, $\operatorname{LinDim}\mathcal{L}>1$, a contradiction.

Clearly ❷ + ❸ ⇒ ❶; hence the result. □

Chapter 14

Gradient flow

Gradient flow provides a tool in Alexandrov's geometry, which we use everywhere in the book.

The technique of gradient flow takes its roots in the so called *Sharafutdinov's retraction*, introduced by Vladimir Sharafutdinov in [56]. It was used widely in comparison geometry since then. In CBB spaces, it was first used by Grigory Perelman and third author in [55]. Bit later, independently Jürgen Jost and Uwe Mayer [57, 58] used the gradient flow in CAT spaces. Later, Alexander Lytchak unified and generalized these two approaches to a wide class of metric spaces in [54]. It was yet developed further by Shin-ichi Ohta in [48] and by Giuseppe Sevaré in [59].

Gradient flow could be considered as a nonsmooth version of first order ordinary differential equations. The following exercise is a stripped version of Sharfutdinov's retraction; it gives the idea behind gradient flow.

14.0.1. Exercise. *Assume that a one parameter family of convex sets $K_t \subset \mathbb{E}^m$ is nested; that is, $K_{t_1} \supset K_{t_2}$ if $t_1 \leqslant t_2$. Show that there is a family of short maps $\varphi_t\colon \mathbb{E}^m \to K_t$ such that $\varphi_t|_{K_t} = \mathrm{id}$ and $\varphi_{t_2} \circ \varphi_{t_1} = \varphi_{t_2}$ for all $t_1 \leqslant t_2$.*

14.1 Gradient-like curves

Gradient-like curves will be used later in the construction of gradient curves. The latter appear to be a special reparametrization of gradient-like curves.

14.1.1. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be locally Lipschitz semiconcave subfunction.*

A Lipschitz curve $\hat\alpha\colon [s_{\min}, s_{\max}) \to \operatorname{Dom} f$ *will be called* f-gradient-like curve *if*

$$\hat\alpha^+ = \tfrac{1}{|\nabla_{\hat\alpha} f|}\cdot\nabla_{\hat\alpha} f;$$

that is, for any $s \in [s_{\min}, s_{\max})$, $\hat\alpha^+(s)$ *is defined and*

$$\hat\alpha^+(s) = \tfrac{1}{|\nabla_{\hat\alpha(s)} f|}\cdot\nabla_{\hat\alpha(s)} f.$$

Note that in particular this definition implies that $|\nabla_p f| > 0$ for any point p on $\hat\alpha$. The next theorem shows that a weaker condition is equivalent.

14.1.2. Theorem. *Let* $\mathcal{L}$ *be a complete length* CBB *space,* $f\colon \mathcal{L} \to\!\!\to \mathbb{R}$ *be locally Lipschitz semiconcave subfunction and* $|\nabla_p f| > 0$ *for any* $p \in \operatorname{Dom} f$.

A curve $\hat\alpha\colon [s_{\min}, s_{\max}) \to \operatorname{Dom} f$ *is an* f*-gradient-like curve if and only if it is* 1*-Lipschitz and*

$$\lim_{s\to s_0+} \frac{f\circ\hat\alpha(s) - f\circ\hat\alpha(s_0)}{s - s_0} \geqslant |\nabla_{\hat\alpha(s_0)} f| \qquad ❶$$

for almost all $s_0 \in [s_{\min}, s_{\max})$.

Proof. The "only if" part follows directly from definition. To prove the "if" part, note that for any $s_0 \in [s_{\min}, s_{\max})$ we have

$$\begin{aligned}\lim_{s\to s_0+} \frac{f\circ\hat\alpha(s) - f\circ\hat\alpha(s_0)}{s - s_0} &\geqslant \lim_{s\to s_0+} \oint_{s_0}^{s} |\nabla_{\hat\alpha(\mathsf{s})} f|\cdot \boldsymbol{d}\mathsf{s} \geqslant \\ &\geqslant |\nabla_{\hat\alpha(s_0)} f|;\end{aligned}$$

the first inequality follows from ❶ and the second from lower semicontinuity of the function $x \mapsto |\nabla_x f|$, see 11.4.13. Form 11.4.3, we have

$$\hat\alpha^+(s_0) = \tfrac{1}{|\nabla_{\hat\alpha(s_0)} f|}\cdot\nabla_{\hat\alpha(s_0)} f;$$

hence the result. □

14.1.3. Theorem. *Let* $\mathcal{L}$ *be a complete length* CBB *space and* $f\colon \mathcal{L} \to\!\!\to \mathbb{R}$ *be locally Lipschitz and* λ*-concave. Assume* $\hat\alpha\colon [0, s_{\max}) \to\!\!\to \operatorname{Dom} f$ *is an* f*-gradient-like curve. Then*

$$(f\circ\hat\alpha)'' \leqslant \lambda$$

everywhere on $[0, s_{\max})$.

Closely related statements were proved independently by Uwe Mayer and Shin-ichi Ohta [58, p. 2.36] and [48, p. 5.7].

14.1.4. Corollary. *Let $\mathcal{L}$ be a complete length* CBB *space, $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be a locally Lipschitz and semiconcave function and $\hat\alpha\colon [0, s_{\max}) \to \to \operatorname{Dom} f$ be an f-gradient-like curve. Then function $s \mapsto |\nabla_{\hat\alpha(s)} f|$ is right-continuous; that is, for any $s_0 \in [0, s_{\max})$ we have*

$$|\nabla_{\hat\alpha(s_0)} f| = \lim_{s\to s_0+} |\nabla_{\hat\alpha(s)} f|.$$

Proof. Applying 14.1.3 locally, we have that $f\circ\hat\alpha(s)$ is semiconcave. The statement follows since

$$(f\circ\hat\alpha)^+(s) = (\boldsymbol{d}_p f)\left(\tfrac{1}{|\nabla_{\hat\alpha(s)} f|}\cdot\nabla_{\hat\alpha(s)} f\right) = |\nabla_{\hat\alpha(s)} f|.$$

□

Proof of 14.1.3. For any $s > s_0$,

$$\begin{aligned}(f\circ\hat\alpha)^+(s_0) &= |\nabla_{\hat\alpha(s_0)} f| \geqslant \\ &\geqslant (d_{\hat\alpha(s_0)} f)(\uparrow_{[\hat\alpha(s_0)\hat\alpha(s)]}) \geqslant \\ &\geqslant \frac{f\circ\hat\alpha(s) - f\circ\hat\alpha(s_0)}{|\hat\alpha(s) - \hat\alpha(s_0)|} - \tfrac{\lambda}{2}\cdot|\hat\alpha(s) - \hat\alpha(s_0)|.\end{aligned}$$

Set $\lambda_+ = \max\{0, \lambda\}$. Since $s - s_0 \geqslant |\hat\alpha(s) - \hat\alpha(s_0)|$, for any $s > s_0$ we have

❷ $$(f\circ\hat\alpha)^+(s_0) \geqslant \frac{f\circ\hat\alpha(s) - f\circ\hat\alpha(s_0)}{s - s_0} - \tfrac{\lambda_+}{2}\cdot(s - s_0).$$

Thus $f\circ\hat\alpha$ is λ_+-concave. That finishes the proof for $\lambda \geqslant 0$. For $\lambda < 0$ we get only that $f\circ\hat\alpha$ is 0-concave.

Note that $|\hat\alpha(s) - \hat\alpha(s_0)| = s - s_0 - o(s - s_0)$, thus

❸ $$(f\circ\hat\alpha)^+(s_0) \geqslant \frac{f\circ\hat\alpha(s) - f\circ\hat\alpha(s_0)}{s - s_0} - \tfrac{\lambda}{2}\cdot(s - s_0) + o(s - s_0).$$

Together, ❷ and ❸ imply that $f\circ\hat\alpha$ is λ-concave. □

14.1.5. Proposition. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $p, q \in \mathcal{L}$. Assume $\hat\alpha\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ be a dist_p-gradient-like curve such that $\hat\alpha(s) \to z \in\,]pq[$ as $s \to s_{\max}+$ then α is a unit-speed geodesic that lies in $[pq]$.*

Proof. Clearly,

❹ $$\tfrac{d^+}{dt}|q - \hat\alpha(t)| \geqslant -1$$

On the other hand,

❺
$$\begin{aligned}\tfrac{d^+}{dt}|p-\hat\alpha(t)| &\geqslant (\boldsymbol{d}_{\hat\alpha(t)}\operatorname{dist}_p)(\uparrow_{[\hat\alpha(t)q]}) \geqslant\\ &\geqslant -\cos\measuredangle^\kappa(\hat\alpha(t)\,{}^p_q).\end{aligned}$$

Inequalities ❹ and ❺ imply that the function $t\mapsto \measuredangle^\kappa(q\,{}^{\hat\alpha(t)}_p)$ is nondecreasing. Hence the result. □

14.2 Gradient curves

In this section we define gradient curves and tie them tightly to gradient-like curves which were introduced in Section 14.1.

14.2.1. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L}\to\mathbb{R}$ be a locally Lipschitz and semiconcave function.*

A locally Lipschitz curve $\alpha\colon[t_{\min},t_{\max})\to\operatorname{Dom}f$ will be called f-gradient curve *if*

$$\alpha^+=\nabla_\alpha f;$$

that is, for any $t\in[t_{\min},t_{\max})$, $\alpha^+(t)$ is defined and $\alpha^+(t)=\nabla_{\alpha(t)}f$.

The following exercise describe a global geometric property of gradient curves without direct reference of its function. It use the notion of self-contracting curves which was introduced by Aris Daniilidis, Olivier Ley, Stéphane Sabourau [60].

14.2.2. Exercise. *Let $\mathcal{L}$ be a complete length* CBB *space $f\colon\mathcal{L}\to\mathbb{R}$ be a concave locally Lipschitz function and $\alpha\colon\mathbb{I}\to\mathcal{L}$ is an f-gradient curve. Show that α is* self-contracting*; that is,*

$$|\alpha(t_1)-\alpha(t_3)|\geqslant|\alpha(t_2)-\alpha(t_3)|$$

if $t_1\leqslant t_2\leqslant t_3$.

The next lemma states that gradient and gradient-like curves are special reparametrizations of each-other.

14.2.3. Lemma. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $f\colon\mathcal{L}\to\mathbb{R}$ be a locally Lipschitz semiconcave subfunction such that $|\nabla_p f|>0$ for any $p\in\operatorname{Dom}f$.*

Assume $\alpha\colon[0,t_{\max})\to\operatorname{Dom}f$ be a locally Lipschitz curve and $\hat\alpha\colon[0,s_{\max})\to\operatorname{Dom}f$ be its reparametrization by arc-length, so $\alpha=$

$\hat\alpha\circ\sigma$ *for some homeomorphism* $\sigma\colon [0,t_{\max})\to[0,s_{\max})$*, then*

$$\alpha^+ = \nabla_\alpha f$$
$$\Updownarrow$$
$$\hat\alpha^+ = \frac{1}{|\nabla_{\hat\alpha} f|}\cdot\nabla_{\hat\alpha} f \quad \textit{and} \quad \sigma^{-1}(s) = \int\limits_0^s \frac{\boldsymbol{d}\mathfrak{s}}{(f\circ\hat\alpha)'(\mathfrak{s})}.$$

Proof; ($\Rightarrow$). According to 4.4.1,

❶
$$\begin{aligned}\sigma'(t) &\overset{a.e.}{=\!=} |\alpha^+(t)| = \\ &= |\nabla_{\alpha(t)} f|.\end{aligned}$$

Note that

$$\begin{aligned}(f\circ\alpha)'(t) &\overset{a.e.}{=\!=} (f\circ\alpha)^+(t) = \\ &= |\nabla_{\alpha(t)} f|^2.\end{aligned}$$

Setting $s=\sigma(t)$, we have

$$\begin{aligned}(f\circ\hat\alpha)'(s) &\overset{a.e.}{=\!=} \frac{(f\circ\alpha)'(t)}{\sigma'(t)} \overset{a.e.}{=\!=} \\ &\overset{a.e.}{=\!=} |\nabla_{\alpha(t)} f| = \\ &= |\nabla_{\hat\alpha(s)} f|.\end{aligned}$$

From 14.1.2, it follows that $\hat\alpha(t)$ is an f-gradient-like curve; that is,

$$\hat\alpha^+ = \frac{1}{|\nabla_{\hat\alpha} f|}\cdot\nabla_{\hat\alpha} f.$$

In particular, $(f\circ\hat\alpha)^+(s) = |\nabla_{\hat\alpha^+(s)} f|$ and from ❶,

$$\begin{aligned}\sigma^{-1}(s) &= \int\limits_0^s \frac{1}{|\nabla_{\hat\alpha(\mathfrak{s})} f|}\cdot\boldsymbol{d}\mathfrak{s} = \\ &= \int\limits_0^s \frac{1}{(f\circ\hat\alpha)'(\mathfrak{s})}\cdot\boldsymbol{d}\mathfrak{s}.\end{aligned}$$

($\Leftarrow$). Clearly,

$$\begin{aligned}\sigma(t) &= \int\limits_0^t (f\circ\hat\alpha)^+(\sigma(\mathfrak{t}))\cdot\boldsymbol{d}\mathfrak{t} = \\ &= \int\limits_0^t |\nabla_{\alpha(\mathfrak{t})} f|\cdot\boldsymbol{d}\mathfrak{t}.\end{aligned}$$

According to 14.1.4, the function $s \mapsto |\nabla_{\hat\alpha(s)} f|$ is right-continuous. Therefore so is the function $t \mapsto |\nabla_{\hat\alpha\circ\sigma(t)} f| = |\nabla_{\alpha(t)} f|$. Hence, for any $t_0 \in [0, t_{\max})$ we have

$$\begin{aligned}\sigma^+(t_0) &= \lim_{t\to t_0+} \oint_{t_0}^{t} |\nabla_{\alpha(\mathfrak{t})} f| \cdot \boldsymbol{d}\mathfrak{t} = \\ &= |\nabla_{\alpha(t_0)} f|.\end{aligned}$$

Thus, we have

$$\begin{aligned}\alpha^+(t_0) &= \sigma^+(t_0)\cdot\hat\alpha^+(\sigma(t_0)) = \\ &= \nabla_{\alpha(t_0)} f.\end{aligned}$$

□

14.2.4. Exercise. *Let $\mathcal{L}$ be a complete length* CBB *space, $f\colon \mathcal{L} \to \mathbb{R}$ be a locally Lipschitz semiconcave function. Assume $\alpha\colon \mathbb{I} \to \mathcal{L}$ is a Lipschitz curve such that*

$$\begin{aligned}\alpha^+(t) &\leqslant |\nabla_{\alpha(t)} f|, \\ (f\circ\alpha)^+(t) &\geqslant |\nabla_{\alpha(t)} f|\end{aligned}$$

for almost all t. Show that α is an f-gradient curve.

14.2.5. Exercise. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L} \to \mathbb{R}$ be a concave locally Lipschitz function. Show that $\alpha\colon \mathbb{R} \to \mathcal{L}$ is an f-gradient curve if and only if*

$$|x - \alpha(t_1)|^2_{\mathcal{L}} - |x - \alpha(t_0)|^2_{\mathcal{L}} \leqslant 2\cdot(t_1 - t_0)\cdot(f\circ\alpha(t_1) - f(x))$$

for any $t_1 > t_0$ and $x \in \mathcal{L}$.

Distance estimates

14.2.6. First distance estimate. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $f\colon \mathcal{L} \to \mathbb{R}$ be a locally Lipschitz and λ-concave function. Assume $\alpha, \beta\colon [0, t_{\max}) \to \mathcal{L}$ be two f-gradient. Then for any $t \in [0, t_{\max})$*

$$|\alpha(t) - \beta(t)| \leqslant e^{\lambda\cdot t}\cdot|\alpha(0) - \beta(0)|.$$

Moreover, the same conclusion holds for a locally Lipschitz and λ-concave subfunction $f\colon \mathcal{L} \looparrowright \mathbb{R}$ if for any $t \in [0, t_{\max})$ there is a geodesic $[\alpha(t)\,\beta(t)]$ in $\operatorname{Dom} f$.

Proof. If $\mathcal{L}$ is not geodesic space, let us pass to its ultrapower $\mathcal{L}^\omega$.

Fix a choice of geodesic $[\alpha(t)\,\beta(t)]$ for each t.

Set $\ell(t) = |\alpha(t) - \beta(t)|$, from the first variation inequality (5.3.4) and the estimate in 11.4.6 we get

$$\ell^+(t) \leqslant -\langle\uparrow_{[\alpha(t)\beta(t)]}, \nabla_{\alpha(t)} f\rangle - \langle\uparrow_{[\beta(t)\alpha(t)]}, \nabla_{\beta(t)} f\rangle \leqslant \lambda\cdot\ell(t).$$

Hence the result. □

14.2.7. Second distance estimate. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $\varepsilon > 0$ and $f, g\colon \mathcal{L} \to \mathbb{R}$ be two λ-concave locally Lipschitz function such that $|f - g| < \varepsilon$. Assume $\alpha, \beta\colon [0, t_{\max}) \to \mathcal{L}$ are correspondingly f- and g-gradient curves such that $\alpha(0) = \beta(0)$. Then*

$$|\alpha(t) - \beta(t)| \leqslant \sqrt{\tfrac{1}{2\cdot\varepsilon\cdot\lambda}\cdot\left(e^{\frac{t\cdot\lambda}{\varepsilon}} - 1\right)}$$

for any $t \in [0, t_{\max})$. In particular, if $t_{\max} < \infty$, then

$$|\alpha(t) - \beta(t)| \leqslant c\cdot\sqrt{\varepsilon\cdot t}$$

for some constant $c = c(t_{\max}, \lambda)$.

Moreover, the same conclusion holds for locally Lipschitz and λ-concave subfunctions $f, g\colon \mathcal{L} \looparrowright \mathbb{R}$ if for any $t \in [0, t_{\max})$ there is a geodesic $[\alpha(t)\,\beta(t)]$ in $\operatorname{Dom} f \cap \operatorname{Dom} g$.

Proof. Set $\ell = \ell(t) = |\alpha(t) - \beta(t)|$. Fix t, set $p = \alpha(t)$ and $q = \beta(t)$. From the first variation formula and 11.4.5,

$$\begin{aligned}\ell^+ &\leqslant -\langle\uparrow_{[pq]}, \nabla_p f\rangle - \langle\uparrow_{[qp]}, \nabla_q g\rangle \leqslant \\ &\leqslant -\Big(f(q) - f(p) - \lambda\cdot\tfrac{\ell^2}{2}\Big)/\ell - \Big(g(p) - g(q) - \lambda\cdot\tfrac{\ell^2}{2}\Big)/\ell \leqslant \\ &\leqslant \lambda\cdot\ell + \tfrac{2\cdot\varepsilon}{\ell}.\end{aligned}$$

Integrating the above estimate, we get

$$\ell(t) \leqslant \sqrt{\tfrac{1}{2\cdot\varepsilon\cdot\lambda}\cdot\left(e^{\frac{t\cdot\lambda}{\varepsilon}} - 1\right)}.$$

□

Existence, uniqueness, convergence

In general, "past" of gradient curves can not be determined by present. For example, consider concave function $f\colon \mathbb{R} \to \mathbb{R}$, $f(x) = -|x|$; two

curves $\alpha(t) = \min\{0, t\}$ and $\beta(t) = 0$ are f-gradient and $\alpha(t) = \beta(t) = 0$ for all $t \geqslant 0$, however $\alpha(t) \neq \beta(t)$ for all $t < 0$.

The next theorem shows that "future" gradient curve is unique.

14.2.8. Picard's theorem. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be semiconcave subfunction. Assume $\alpha, \beta\colon [0, t_{\max}) \to \to \operatorname{Dom} f$ be two f-gradient curves such that $\alpha(0) = \beta(0)$. Then $\alpha(t) = \beta(t)$ for any $t \in [0, t_{\max})$.*

Proof. Follows from the first distance estimate (14.2.6). □

14.2.9. Local existence. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $f\colon \mathcal{L} \looparrowright \mathbb{R}$ be locally Lipschitz λ-concave subfunction. Then for any $p \in \operatorname{Dom} f$*

a) if $|\nabla_p f| > 0$, then for some $\varepsilon > 0$, there is an f-gradient-like curve $\hat\alpha\colon [0, \varepsilon) \to \mathcal{L}$ that starts at p (that is, $\hat\alpha(0) = p$);

b) for some $\delta > 0$, there is an f-gradient curve $\alpha\colon [0, \delta) \to \mathcal{L}$ that starts at p (that is $\alpha(0) = p$).

This theorem was proved by Grigory Perelman and the third author in [55]; here we present a simplified proof given by Alexander Lytchak in [54].

Proof. Note that if $|\nabla_p f| = 0$, then one can take constant curve $\alpha(t) = p$. Otherwise, take $\varepsilon > 0$, such that $\mathrm{B}(p, \varepsilon) \subset \operatorname{Dom} f$, the restriction $f|_{\mathrm{B}(p,\varepsilon)}$ is Lipschitz and $|\nabla_x f| > \varepsilon$ for all $x \in \mathrm{B}(p, \varepsilon)$; the latter is possible due to semicontinuity of |gradient| (11.4.13).

The curves $\hat\alpha$ and α will be constructed in the following three steps. First we construct an f^ω-gradient-like curve $\hat\alpha_\omega\colon [0, \varepsilon) \to \mathcal{L}^\omega$ as an ω-limit of certain sequence of broken geodesics in $\mathcal{L}$. Second, we parametrize $\hat\alpha_\omega$ as in 14.2.3, to obtain an f^ω-gradient curve α_ω in $\mathcal{L}^\omega$. Third, applying Picard's theorem (14.2.8) together with Lemma 3.2.3, we obtain that α_ω lies in $\mathcal{L} \subset \mathcal{L}^\omega$ and therefore one can take $\alpha = \alpha_\omega$ and $\hat\alpha = \hat\alpha_\omega$.

Note that if $\mathcal{L}$ is proper, then $\mathcal{L} = \mathcal{L}^\omega$ and $f^\omega = f$. Thus, in this case, the third step is not necessary.

Step 1. Given $n \in \mathbb{N}$, by open-close argument, we can construct a unit-speed curve $\hat\alpha_n\colon [0, \varepsilon] \to \mathcal{L}$ starting at p, with a partition of $[0, \varepsilon)$ into countable number of half-open intervals $[\sigma_i, \bar\sigma_i)$ so that for each i we have

(i) $\hat\alpha_n([\sigma_i, \bar\sigma_i])$ is a geodesic and $\bar\sigma_i - \sigma_i < \frac{1}{n}$.

(ii) $f \circ \hat\alpha_n(\bar\sigma_i) - f \circ \hat\alpha_n(\sigma_i) > (\bar\sigma_i - \sigma_i)\cdot(|\nabla_{\hat\alpha_n(\sigma_i)} f| - \frac{1}{n})$.

Pass to a subsequence of $(\hat\alpha_n)$ such that $f \circ \hat\alpha_n$ uniformly converges; set

$$h(s) = \lim_{n\to\infty} f \circ \hat\alpha_n(s).$$

Set $\hat\alpha_\omega = \lim_{n\to\omega}\hat\alpha_n$, it is a curve in $\mathcal{L}^\omega$ that starts at $p\in\mathcal{L}\subset\mathcal{L}^\omega$. Clearly $\hat\alpha_\omega$ is 1-Lipschitz. From (ii) and 11.4.10, we get that

$$(f^\omega\circ\hat\alpha_\omega)^+(\sigma)\geqslant|\nabla_{\hat\alpha_\omega(\sigma)}f^\omega|.$$

According to 14.1.2, $\hat\alpha_\omega\colon[0,\varepsilon)\to\mathcal{L}^\omega$ is an f^ω-gradient-like curve.

Step 2. Clearly $h(s)=f^\omega\circ\alpha_\omega$. Therefore, according to 14.1.3, h is λ-concave. Thus, we can define a homeomorphism $\sigma\colon[0,\delta]\to[0,\varepsilon]$ by

❷ $$\sigma^{-1}(s)=\int\limits_0^s\frac{1}{h'(\mathring{s})}\cdot\boldsymbol{d}\mathring{s},$$

According to 14.2.3, $\alpha(t)=\hat\alpha\circ\sigma(t)$ is an f^ω-gradient curve in $\mathcal{L}^\omega$.

Step 3. Clearly, $\nabla_p f=\nabla_p f^\omega$ for any $p\in\mathcal{L}\subset\mathcal{L}^\omega$; more formally if $\iota\colon\mathcal{L}\hookrightarrow\mathcal{L}^\omega$ is the natural embedding, then $(\boldsymbol{d}_p\iota)(\nabla_p f)=\nabla_p f^\omega$. Thus, it is sufficient to show that $\alpha_\omega\subset\mathcal{L}$. Assume the contrary, then according to 3.2.3 there is a subsequence $\hat\alpha_{n_k}$ such that

$$\hat\alpha_\omega\neq\hat\alpha'_\omega:=\lim_{k\to\omega}\hat\alpha_{n_k}.$$

Clearly $h(s)=f^\omega\circ\hat\alpha_\omega=f^\omega\circ\hat\alpha'_\omega$. Thus, for $\sigma\colon[0,\delta]\to[0,\varepsilon]$ defined by ❷, we have that both curves $\hat\alpha_\omega\circ\sigma$ and $\hat\alpha'_\omega\circ\sigma$ are f^ω-gradient. Thus, from Picard's theorem (14.2.8), we get that $\hat\alpha_\omega\circ\sigma=\hat\alpha'_\omega\circ\sigma$ and therefore $\hat\alpha_\omega=\hat\alpha'_\omega$, a contradiction □

14.2.10. Ultralimit of gradient curves. *Assume*

- $(\mathcal{L}_n)$ *is a sequence of complete* CBB(κ) *spaces and* $(\mathcal{L}_n,p_n)\to\to(\mathcal{L}_\omega,p_\omega)$ *as* $n\to\omega$*;*
- $f_n\colon\mathcal{L}_n\to\mathbb{R}$ *are* $\pounds$*-Lipschitz and* λ*-concave* $f_n\to f_\omega$ *as* $n\to\omega$ *and* $p_\omega\in\operatorname{Dom}f_\omega$*.*

Then

a) f_ω *is* λ*-concave.*

b) If $|\nabla_{p_\omega}f_\omega|>0$*, then there is* $\varepsilon>0$ *such that, the* f_n*-gradient-like curves* $\hat\alpha_n\colon[0,\varepsilon)\to\mathcal{L}_n$ *are defined for* ω*-almost all* n*. And moreover, a curve* $\hat\alpha_\omega\colon[0,\varepsilon)\to\mathcal{L}_\omega$ *is a gradient-like curve that starts at* p_ω *if and only if* $\hat\alpha_n(s)\to\to\hat\alpha_\omega(s)$ *as* $n\to\omega$ *for all* $s\in[0,\varepsilon)$*.*

c) For some $\delta>0$*, the* f_n*-gradient curves* $\alpha_n\colon[0,\delta)\to\mathcal{L}_n$ *are defined for* ω*-almost all* n*. And moreover, a curve* $\alpha_\omega\colon[0,\delta)\to\mathcal{L}_\omega$ *is a gradient curve that starts at* p_ω *if and only if* $\alpha_n(t)\to\alpha_\omega(t)$ *as* $n\to\omega$ *for all* $t\in[0,\delta)$*.*

Note that according to Exercise 3.7.2, the part (a) does not hold for general metric spaces. The idea in the proof is the same as in the proof of local existence (14.2.9). We will need the following lemma

Proof; (a) Fix a geodesic $\gamma_\omega\colon \mathbb{I} \to \operatorname{Dom} f_\omega$; we need to show that the function

❸ $$t \mapsto f_\omega \circ \gamma_\omega(t) - \tfrac{\lambda}{2}\cdot t^2$$

is concave.

Since f_n are $\pounds$-Lipschitz, so is f_ω. Therefore it is sufficient to prove the concavity in the interior of $\mathbb{I}$; in other words we can assume that γ_ω connects points p_ω and q_ω and can be extended behind its ends as a minimizing geodesic. In the latter case, by Theorem 7.2.4, γ_ω is unique geodesic connecting p_ω to q_ω .

Construct two sequences of points $p_n, q_n \in \mathcal{L}_n$ such that $p_n \to p_\omega$ and $q_n \to q_\omega$ as $n \to \omega$. Since $\mathcal{L}_n$ is almost geodesic (7.2.4), we can assume that for each n there is a geodesic γ_n from p_n to q_n in $\mathcal{L}_n$.

Since f_n is λ concave, the function

$$t \mapsto f_n \circ \gamma_n(t) - \tfrac{\lambda}{2}\cdot t^2$$

is concave.

The ω-limit of sequence γ_n is a geodesic in $\mathcal{L}_\omega$ from p_ω to q_ω. By uniqueness of such geodesic we have that $\gamma_n \to \gamma_\omega$ as $n \to \omega$. Passing to the limit, we get ❸.

"if"-part of (b). Take $\varepsilon > 0$ so small that $\mathrm{B}(p_\omega, \varepsilon) \subset \operatorname{Dom} f_\omega$ and $|\nabla_{x_\omega} f_\omega| > 0$ for any $x_\omega \in \mathrm{B}(p_\omega, \varepsilon)$ (that is possible due to 11.4.13).

Clearly $\hat\alpha_\omega$ is 1-Lipschitz. From 11.4.10, we get $(f_\omega \circ \hat\alpha_\omega)^+(s) \geqslant$ $\geqslant |\nabla_{\hat\alpha_\omega(s)} f^\omega|$. According to 14.1.2, $\hat\alpha_\omega\colon [0,\varepsilon) \to \mathcal{L}^\omega$ is an f_ω-gradient-like curve.

"if"-part of (c). Assume first $|\nabla_{p_\omega} f_\omega| > 0$ — so we can apply "if"-part of (b). Let $h_n = f_n \circ \hat\alpha_n\colon [0,\varepsilon) \to \mathbb{R}$ and $h_\omega = f_\omega \circ \hat\alpha_\omega$. From 14.1.3, h_n are λ-concave and clearly $h_n \to h_\omega$ as $n \to \omega$. Let us define reparametrizations

$$\sigma_n^{-1}(s) = \int\limits_0^s \frac{1}{h_n'(\mathfrak{s})}\cdot \boldsymbol{d}\mathfrak{s}, \qquad \sigma_\omega^{-1}(s) = \int\limits_0^s \frac{1}{h_\omega'(\mathfrak{s})}\cdot \boldsymbol{d}\mathfrak{s}.$$

The λ-convexity of h_n implies that $\sigma_n \to \sigma_\omega$ as $n \to \omega$. By 14.2.3, $\alpha_n = \hat\alpha_n \circ \sigma_n$. Applying "if"-part of (b) together with Lemma 14.2.3, we get that $\alpha_\omega = \hat\alpha_\omega \circ \sigma_\omega$ is gradient curve.

The remaining case $|\nabla_{p_\omega} f_\omega| = 0$, can be reduced to the one above using the following trick. Consider sequence of spaces $\mathcal{L}_n^\times = \mathcal{L}_n \times \mathbb{R}$, with the sequence of subfunction $f_n^\times\colon \mathcal{L}_n^\times \to \mathbb{R}$, defined by

$$f_n^\times(p,t) = f_n(p) + t.$$

According to Theorem 10.4.1b, $\mathcal{L}_n^\times$ is a complete length CBB(κ_-) space for $\kappa_- = \min\{\kappa, 0\}$. Note that $f_n^\times$ are λ_+-concave, for $\lambda_+ = \max\{\lambda, 0\}$. Analogously define $\mathcal{L}_\omega^\times = \mathcal{L}_\omega \times \mathbb{R}$, and $f_\omega^\times(p,t) = f_\omega(p) + t$.

Clearly $\mathcal{L}_n^\times \to \mathcal{L}_\omega^\times$, $f_n^\times \to f_\omega^\times$ as $n \to \omega$ and $|\nabla_x f_\omega^\times| > 0$ for any $x \in \operatorname{Dom} f_\omega^\times$. Thus, for the sequence $f_n^\times \colon \mathcal{L}_n^\times \looparrowright \mathbb{R}$, we can apply "if"-part of (b). It remains to note that $\alpha_\omega^\times(t) = (\alpha_\omega(t), t)$ is a $f_\omega^\times$-gradient curve in $\mathcal{L}_\omega^\times$ if and only if $\alpha_\omega(t)$ is a f_ω-gradient curve.

"only if"-part of (c) and (b). The "only if"-part of (c) follows from the "if"-part of (c) and Picard's theorem (14.2.8). Applying Lemma 14.2.3, we get "only if"-part of (b). □

Directly form the local existence (14.2.9) and the distance estimates (14.2.6), we get the following:

14.2.11. Global existence. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $f \colon \mathcal{L} \looparrowright \mathbb{R}$ be locally Lipschitz and λ-concave subfunction. Then for any $p \in \operatorname{Dom} f$, there is $t_{\max} \in (0, \infty]$ such that there is an f-gradient curve $\alpha \colon [0, t_{\max}) \to \mathcal{L}$ with $\alpha(0) = p$. Moreover, for any sequence $t_n \to t_{\max}-$, the sequence $\alpha(t_n)$ does not have a limit point in* $\operatorname{Dom} f$.

The following theorem gurantees the existance of gradient curves for all times for the special type of semiconcave functions that play important role in the theory. It follows from 14.2.11, 14.1.3 and 14.2.3.

14.2.12. Theorem. *Let $\mathcal{L}$ be a complete length* CBB *space and $f \colon \mathcal{L} \to \mathbb{R}$ satisfies*

$$f'' + \kappa \cdot f \leqslant \lambda$$

for some real values κ and λ. Then f has complete gradient*; that is, for any $x \in \mathcal{L}$ there is a f-gradient curve $\alpha \colon [0, \infty) \to \mathcal{L}$ that starts at x.*

14.3 Gradient flow

In this section we define gradient flow for semiconcave subfunctions and reformulate theorems obtained earlier in this chapter using this new terminology.

Let $\mathcal{L}$ be a complete length CBB space and $f \colon \mathcal{L} \looparrowright \mathbb{R}$ be a locally Lipschitz semiconcave subfunctions. For any $t \geqslant 0$ write $\mathfrak{d}_f^t(x) = y$ if there is a f-gradient curve α such that $\alpha(0) = x$ and $\alpha(t) = y$. The defined map $\mathfrak{d}_f^t \colon \mathcal{L} \to \mathcal{L}$ is called *f-gradient flow* for time t.

From 14.2.7, it follows that for any $t \geqslant 0$, the domain of definition of $\mathfrak{d}_f^t$ is an open subset of $\mathcal{L}$; that is, $\mathfrak{d}_f^t$ is a submap. Moreover, if

f is defined on whole $\mathcal{L}$ and $f'' + \mathrm{K}\cdot f \leqslant \lambda$ for some $\mathrm{K}, \lambda \in \mathbb{R}$, then according to 14.2.12, $\mathfrak{d}_f^t(x)$ is defined for all pairs $(x,t) \in \mathcal{L} \times \mathbb{R}_{\geqslant 0}$.

Clearly $\mathfrak{d}_f^{t_1+t_2} = \mathfrak{d}_f^{t_1} \circ \mathfrak{d}_f^{t_2}$; in other words gradient flow is given by an action of semigroup $(\mathbb{R}_{\geqslant 0}, +)$.

From the first distance estimate 14.2.6, we get the following.

14.3.1. Proposition. *Let $\mathcal{L}$ be a complete length* CBB *space and $f\colon \mathcal{L} \to \mathbb{R}$ be a semiconcave functions. Then the map $x \mapsto \mathfrak{d}_f^t(x)$ is locally Lipschitz.*

Moreover, if f is λ-concave, then $\mathfrak{d}_f^t$ is $e^{\lambda\cdot t}$-Lipschitz.

Next proposition states that gradient flow is stable under Gromov–Hausdorff convergence. It follows directly from the proposition on ultralimit of gradient curves 14.2.10.

14.3.2. Proposition. *If $\mathcal{L}_n$ be an m-dimensional complete length* CBB(κ) *space, $\mathcal{L}_n \xrightarrow{\tau} \mathcal{L}$, $f_n\colon \mathcal{L}_n \to \mathbb{R}$ is a sequence of λ-concave functions that converges to $f\colon \mathcal{L} \to \mathbb{R}$, then $\mathfrak{d}_{f_n}^t\colon \mathcal{L}_n \to \mathcal{L}_n$ converges to $\mathfrak{d}_f^t\colon \mathcal{L} \to \mathcal{L}$.*

14.3.3. Exercise. *Let $\mathcal{L}$ be an m-dimensional complete length* CBB *space, $\partial\mathcal{L} = \varnothing$, $K \subset \mathcal{L}$ be a compact subset and $f\colon \mathcal{L} \to \mathbb{R}$ be semiconcave function. Assume that for some $t > 0$ the gradient flow $\mathfrak{d}_f^t$ is defined everywhere in K, prove that*

$$\partial_{\mathcal{L}}\mathfrak{d}_f^t K \subset \mathfrak{d}_f^t \partial K.$$

14.4 Line splitting theorem

Let $\mathcal{X}$ be a metric space and $A, B \subset \mathcal{X}$. We will write

$$\mathcal{X} = A \oplus B$$

If there projections $\mathrm{proj}_A\colon \mathcal{X} \to A$ and $\mathrm{proj}_B\colon \mathcal{X} \to B$ such that

$$|x-y|^2 = |\mathrm{proj}_A(x) - \mathrm{proj}_A(y)|^2 + |\mathrm{proj}_B(x) - \mathrm{proj}_B(y)|^2$$

for any two points $x, y \in \mathcal{X}$.

Note that if

$$\mathcal{X} = A \oplus B$$

then

- ⋄ A intersects B at single point,
- ⋄ both sets A and B are convex sets in $\mathcal{X}$.

Recall that a line in a metric space is a both side infinite geodesic; it has to minimze the length on each segment.

14.4.1. Line splitting theorem. *Let $\mathcal{L}$ be a complete length* CBB(0) *space and γ be a line in $\mathcal{L}$. Then*

$$\mathcal{L} = X \oplus \gamma(\mathbb{R})$$

for some subset $X \subset \mathcal{L}$.

For smooth 2-dimensional surfaces, this theorem was proved by Stefan Cohn-Vossen in [61]. For the Riemannian manifolds of higher dimensions it was proved by Victor Toponogov in [19]. Then it was generalized by Anatoliy Milka in [62] to Alexandrov's spaces, almost the same prove is given in [4, p. 1.5].

Further generalizations of the splitting theorem for Riemannian manifolds with nonnegative Ricci curvature were obtained by Jeff Cheeger and Detlef Gromoll [63]. Jost-Hinrich Eshenburg obtained an analogous result for the Lorentzian manifolds [64], that is, pseudo-Riemannian manifolds of signature $(1, n)$.

We present a proof that use gradient flow for Busemann's functions, it is close in the spirit to the proof given in [63].

Before goint into the proof, let us state few corollaries of the theorem.

14.4.2. Corollary. *Let $\mathcal{L}$ be a complete length* CBB(0) *space. Then there is an isometric splitting*

$$\mathcal{L} = \mathcal{L}' \oplus H$$

where $H \subset \mathcal{L}$ is a subset isometric to a Hilbert space and $\mathcal{L}' \subset \mathcal{L}$ be a convex subset that contains no lines.

14.4.3. Corollary. *Let $\mathcal{K}$ be an n-dimensional complete length* CBB(0) *cone and $v_+, v_- \in \mathcal{K}$ be a pair of opposite vectors (that is, $v_+ + v_- = 0$). Then there is an isometry $\iota\colon K \to K' \times \mathbb{R}$, where K' be a complete length* CBB(0) *space with a cone structure with vertex o' such that $\iota : v_\pm \mapsto (o', \pm|v_\pm|)$.*

14.4.4. Corollary. *Assume $\mathcal{L}$ is an m-dimensional complete length* CBB(1) *space, $m \geqslant 2$ and* $\operatorname{rad}\mathcal{L} = \pi$, *then*

$$\mathcal{L} \overset{iso}{=\!=} \mathbb{S}^m.$$

Proof of 14.4.1. Consider two Busemann's functions bus_+ and bus_- asociated with rays $\gamma : [0, \infty) \to \mathcal{L}$ and $\gamma : (-\infty, 0] \to \mathcal{L}$; that is,

$$\mathrm{bus}_\pm(x) = \lim_{t\to\infty} |\gamma(\pm t) - x| - t.$$

According to Exercise 7.4.2, both functions $\mathrm{bus}_\pm$ are concave.

Fix $x \in \mathcal{L}$. Note that since γ is a line, we have

$$\mathrm{bus}_+(x) + \mathrm{bus}_-(x) \geqslant 0.$$

On the other hand, by 7.4.1*b* $f(t) = \mathrm{dist}_x^2 \circ \gamma(t)$ is 2-concave. In particular, $f(t) \leqslant t^2 + at + b$ for some constants $a, b \in \mathbb{R}$. Passing to the limit as $t \to \pm\infty$, we get

$$\mathrm{bus}_+(x) + \mathrm{bus}_-(x) \leqslant 0.$$

Hence

$$\mathrm{bus}_+(x) + \mathrm{bus}_-(x) = 0$$

for any $x \in \mathcal{L}$. In particular the functions $\mathrm{bus}_\pm$ are affine; that is they are convex and concave at the same time.

It follows that for any x,

$$\begin{aligned} |\nabla_x \mathrm{bus}_\pm| &= \sup\{\, \boldsymbol{d}_x \mathrm{bus}_\pm(\xi) : \xi \in \Sigma_x \,\} = \\ &= \sup\{\, -\boldsymbol{d}_x \mathrm{bus}_\mp(\xi) : \xi \in \Sigma_x \,\} \equiv \\ &\equiv 1. \end{aligned}$$

By Exercise 14.2.4, 1-Lipschitz curve α, such that $\mathrm{bus}_\pm(\alpha(t)) = t + c$ is a $\mathrm{bus}_\pm$-gradient curve. In particular, $\alpha(t)$ is a bus_+-gradient curve if and only if $\alpha(-t)$ is a bus_--gradient curve. It follows that for any $t > 0$, the $\mathrm{bus}_\pm$-gradient flows commute; that is

$$\mathfrak{d}^t_{\mathrm{bus}_+} \circ \mathfrak{d}^t_{\mathrm{bus}_-} = \mathrm{id}_{\mathcal{L}}\,.$$

Set

$$\mathfrak{d}^t = \begin{cases} \mathfrak{d}^t_{\mathrm{bus}_+} & \text{if } t \geqslant 0 \\ \mathfrak{d}^t_{\mathrm{bus}_-} & \text{if } t < 0 \end{cases}$$

it defines an $\mathbb{R}$-action on $\mathcal{L}$.

Set $\mathcal{L}' = \mathrm{bus}_+^{-1}(0) = \mathrm{bus}_-^{-1}(0) \subset \mathcal{L}$, it is a closed convex set, therefore it forms an Alexandrov's space. Consider map $h\colon \mathcal{L}' \times \mathbb{R} \to \mathcal{L}$ defined by $h\colon (x,t) \mapsto \mathfrak{d}^t(x)$. Note that h is onto. Applying distance estimate 14.2.6 for $\mathfrak{d}^t_{\mathrm{bus}_+}$ and $\mathfrak{d}^t_{\mathrm{bus}_-}$, we get that h is a short and non-contracting at the same time; that is, h is an isometry. $\square$

14.5 Radial curves

The radial curves are specially reparametrized gradient curves for distance functions. This parametrization makes them to behave as unit-speed geodesics in a natural comparison sense.

14.5.1. Definition. *Assume $\mathcal{L}$ is a complete length* CBB *space,* $\kappa \in \mathbb{R}$*, and* $p \in \mathcal{L}$*. A curve*

$$\sigma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$$

is called (p,κ)-radial curve *if*

$$s_{\min} = |p - \sigma(s_{\min})| \in (0, \tfrac{\varpi^\kappa}{2}),$$

and it satisfies the following differential equation

❶ $$\sigma^+(s) = \frac{\operatorname{tg}\kappa|p-\sigma(s)|}{\operatorname{tg}\kappa s}\cdot\nabla_{\sigma(s)}\operatorname{dist}_p\,.$$

for any $s \in [s_{\min}, s_{\max})$*, here* $\operatorname{tg}\kappa x := \frac{\operatorname{sn}^\kappa x}{\operatorname{cs}^\kappa x}$*.*

If $x = \sigma(s_{\min})$*, we say that* σ starts in x*.*

Note that according to the definition $s_{\max} \leqslant \frac{\varpi^\kappa}{2}$.

Further you will see that the (p,κ)-radial curves work best for CBB(κ) spaces.

14.5.2. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and* $p \in \mathcal{L}$*. A unit-speed geodesic* $\gamma\colon \mathbb{I} \to \mathcal{L}$ *is called* p-radial geodesic *if* $|p - \gamma(s)| \equiv s$*.*

The proof of the following two propositions follows directly from the definitions:

14.5.3. Proposition. *Let $\mathcal{L}$ be a complete length* CBB *space and* $p \in \mathcal{L}$*. Assume* $\frac{\varpi^\kappa}{2} \geqslant s_{\max}$*. Then any* p*-radial geodesic* $\gamma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ *is a* (p,κ)*-radial curve.*

14.5.4. Proposition. *Let $\mathcal{L}$ be a complete length* CBB *space,* $p \in \mathcal{L}$ *and* $\sigma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ *be a* (p,κ)*-radial curve. Then for any* $s \in [s_{\min}, s_{\max})$*, we have* $|p - \sigma(s)| \leqslant s$*.*

Moreover, if for some s_0 *we have* $|p - \sigma(s_0)| = s_0$ *then the restriction* $\sigma|_{[s_{\min}, s_0]}$ *is a* p*-radial geodesic.*

14.5.5. Existence and uniqueness. *Let $\mathcal{L}$ be a complete length* CBB *space,* $\kappa \in \mathbb{R}$*,* $p \in \mathcal{L}$*, and* $x \in \mathcal{L}$*. Assume* $0 < |p - x| < \frac{\varpi^\kappa}{2}$*. Then there is unique* (p,κ)*-radial curve* $\sigma\colon [|p-x|, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ *that starts in* x*; that is,* $\sigma(|p-x|) = x$*.*

Proof; existence. Set

$$\operatorname{itg}^\kappa\colon [0, \tfrac{\varpi^\kappa}{2}) \to \mathbb{R}, \quad \operatorname{itg}^\kappa(t) = \int\limits_0^t \operatorname{tg}\kappa\mathfrak{t}\cdot\boldsymbol{d}\mathfrak{t}.$$

Clearly $\operatorname{itg}^\kappa$ is smooth and increasing. From 4.6.2 it follows that the composition

$$f = \operatorname{itg}^\kappa \circ \operatorname{dist}_p$$

is semiconcave in $\mathrm{B}(p, \frac{\varpi^\kappa}{2})$.

According to 14.2.9, there is an f-gradient curve $\alpha\colon [0, t_{\max}) \to \mathcal{L}$ defined on the maximal interval such that $\alpha(0) = x$.

Now consider solution of differential equation $\tau(t)$, $\tau' = (\operatorname{tg}\kappa\tau)^2$ and $\tau(0) = r$. Note that $\tau(t)$ is also a gradient curve for function $\operatorname{itg}^\kappa$ defined on $[0, \frac{\varpi^\kappa}{2})$. Direct calculations show that composition $\alpha \circ \tau^{-1}$ is a (p, κ)-radial curve.

Uniqueness. Assume σ^1, σ^2 be two (p, κ)-radial curves that starts in x. Then compositions $\sigma^i \circ \tau$ both give f-gradient curves. By Picard's theorem (14.2.8), we have $\sigma^1 \circ \tau \equiv \sigma^2 \circ \tau$. Therefore $\sigma^1(s) = \sigma^2(s)$ for any $s \geqslant r$ such that both sides are defined. □

14.6 Radial comparisons

In this section we show that radial curves in some comparison sense behave as unit-speed geodesics.

14.6.1. Radial monotonicity. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space and p, q be distinct points in $\mathcal{L}$. Assume $\sigma\colon [s_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ is a (p, κ)-radial curve. Then the function*

$$s \mapsto \mathcal{Z}^\kappa\{|q - \sigma(s)|; |p - q|, s\}$$

is nonincreasing in all the domain of definition.

From Radial monotonicity, by straightforward calculations one gets the following.

14.6.2. Corollary. *Let $\kappa \leqslant 0$, $\mathcal{L}$ is a complete* $\mathrm{CBB}(\kappa)$ *space and $p, q \in \mathcal{L}$. Assume $\sigma\colon [s_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ is a (p, κ)-radial curve. Then for any $w \geqslant 1$, the function*

$$s \mapsto \mathcal{Z}^\kappa\{|q - \sigma(s)|; |p - q|, w{\cdot}s\}$$

is non-increasing in whole domain of definition.

14.6.3. Radial comparison. *Let $\mathcal{L}$ be a complete length* $\mathrm{CBB}(\kappa)$ *space and $p \in \mathcal{L}$. Assume $\rho\colon [r_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ and $\sigma\colon [s_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ are two (p, κ)-radial curves. Set*

$$\varphi_{\min} = \mathcal{Z}^\kappa\left(p\,{}^{\rho(r_{\min})}_{\sigma(s_{\min})}\right).$$

Then for any $r \in [r_{\min}, \frac{\varpi^\kappa}{2})$ *and* $s \in [s_{\min}, \frac{\varpi^\kappa}{2})$, *we have*

$$\mathcal{Z}^\kappa\{|\rho(r) - \sigma(s)|; r, s\} \leqslant \varphi_{\min},$$

or equivalently,

$$|\rho(r) - \sigma(s)| \leqslant \tilde{\Upsilon}^\kappa\{\varphi_{\min}; r, s\}.$$

We prove both of the theorems simultaneously. The proof is an application of 11.4.5 plus trigonometric manipulations. We give a prove first in the simplest case $\kappa = 0$ and then the harder case $\kappa \neq 0$.

We proof case $\kappa = 0$ separately since it is easier to follow. The arguments for both cases are nearly the same, but the case $\kappa \neq 0$ require an extra idea.

Proof of 14.6.1 and 14.6.3 in case $\kappa = 0$. Set

$$\begin{aligned} R = R(r) &= |p - \rho(r)|, \\ S = S(s) &= |p - \sigma(s)|, \\ \ell = \ell(r, s) &= |\rho(r) - \sigma(s)|, \\ \varphi = \varphi(r, s) &= \mathcal{Z}^0\{\ell(r, s); r, s\}. \end{aligned}$$



It will be sufficient to prove the following two inequalities:

$$(*)^0_\varphi \qquad \frac{\partial^+\varphi}{\partial r}(s_{\min}, r) \leqslant 0, \qquad \frac{\partial^+\varphi}{\partial s}(s, r_{\min}) \leqslant 0$$

$$(**)^0_\varphi \qquad s\cdot\frac{\partial^+\varphi}{\partial s} + r\cdot\frac{\partial^+\varphi}{\partial r} \leqslant 0.$$

Once they are proved, the radial monotonicity follows from $(*)^0_\varphi$. The radial comparison follows from both $(*)^0_\varphi$ and $(**)^0_\varphi$. Indeed, one can connect $(s_{\min}, r_{\min})$ and (s_0, r_0) in $[s_{\min}, \infty) \times [r_{\min}, \infty)$ by a join of coordinate line and a segment defined by $r/s = r_0/s_0$. According



to $(*)^0_\varphi$ and $(**)^0_\varphi$, the value of φ does not increase while pair (r, s) moving along this join. Thus $\varphi(r_0, s_0) \leqslant \varphi(r_{\min}, s_{\min}) = \varphi_{\min}$.

It remains to show $(*)^0_\varphi$ and $(**)^0_\varphi$. First let us rewrite the inequalities $(*)^0_\varphi$ and $(**)^0_\varphi$ in an equivalent form:

$$(*)^0_\ell \qquad \begin{aligned} \frac{\partial^+}{\partial s}\ell(s, r_{\min}) &\leqslant \cos \measuredangle^0\{r_{\min}; s, \ell\}, \\ \frac{\partial^+}{\partial r}\ell(s_{\min}, r) &\leqslant \cos \measuredangle^0\{s_{\min}; r, \ell\}, \end{aligned}$$

$$(**)^0_\ell \qquad s\cdot\frac{\partial^+\ell}{\partial s} + r\cdot\frac{\partial^+\ell}{\partial r} \leqslant s\cdot \cos \measuredangle^0\{r; s, \ell\} + r\cdot \cos \measuredangle^0\{s; r, \ell\} = \ell.$$

Set

$$(A)^0 \qquad f = \tfrac{1}{2}\cdot \operatorname{dist}_p^2.$$

Clearly f is 1-concave and

$$(B)^0 \qquad \rho^+(r) = \tfrac{1}{r}\cdot\nabla_{\rho(r)} f \quad \text{and} \quad \sigma^+(s) = \tfrac{1}{s}\cdot\nabla_{\sigma(s)} f.$$

Thus from 11.4.5, we have

$$(C)^0 \qquad \frac{\partial^+\ell}{\partial r} = -\tfrac{1}{r}\cdot\langle\nabla_{\rho(r)} f, \uparrow_{[\rho(r)\sigma(s)]}\rangle \leqslant \frac{\ell^2 + R^2 - S^2}{2\cdot\ell\cdot r}.$$

Since $R(r) \leqslant r$ and $S(s_{\min}) = s_{\min}$, we get

$$(D)^0 \qquad \begin{aligned} \frac{\partial^+\ell}{\partial r}(r, s_{\min}) &\leqslant \frac{\ell^2 + r^2 - s_{\min}^2}{2\cdot\ell\cdot r} = \\ &= \cos \measuredangle^0\{s_{\min}; r, \ell\}, \end{aligned}$$

which is the first inequality in $(*)^0_\ell$. By switching places of ρ and σ we obtain the second inequality in $(*)^0_\ell$. Further, summing together $(C)^0$ with its mirror-inequality for $\frac{\partial^+\ell}{\partial s}$, we get

$$(E)^0 \qquad r\cdot\frac{\partial^+\ell}{\partial r}+s\cdot\frac{\partial^+\ell}{\partial s}\leqslant\frac{\ell^2+R^2-S^2}{2\cdot\ell}+\frac{\ell^2+S^2-R^2}{2\cdot\ell}=\ell$$

which is $(**)^0_\ell$. □

Proof of 14.6.1 and 14.6.3 in case $\kappa\neq 0$. Set as before

$$R=R(r)=|p-\rho(r)|,\qquad \ell=\ell(r,s)=|\rho(r)-\sigma(s)|$$
$$S=S(s)=|p-\sigma(s)|,\qquad \varphi=\varphi(r,s)=\measuredangle^\kappa\{\ell(r,s);r,s\}.$$

The statement follows from the following three inequalities:

$$(*)^\pm_\varphi \qquad \frac{\partial^+\varphi}{\partial r}(s_{\min},r)\leqslant 0\quad \frac{\partial^+\varphi}{\partial s}(s,r_{\min})\leqslant 0$$

$$(**)^\pm_\varphi \qquad \operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S\cdot\frac{\partial^+\varphi}{\partial s}+\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R\cdot\frac{\partial^+\varphi}{\partial r}\leqslant 0$$

Once they are proved, the radial monotonicity follows from $(*)^\pm_\varphi$. The radial comparison follows from both $(*)^0_\varphi$ and $(**)^\pm_\varphi$. Indeed, functions $s\mapsto\operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S$ and $r\mapsto\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R$ are Lipschitz. Thus there is a solution for differential equation

$$(r',s')=(\operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S,\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R)$$

with any initial data. $(r_0,s_0)\in[r_{\min},\frac{\varpi^\kappa}{2})\times[s_{\min},\frac{\varpi^\kappa}{2})$. (Unlike case $\kappa=0$ the solution can not be written explicitly.) Since $\operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S$, $\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R>0$, this solution $(r(t),s(t))$ must meet one of coordinate rays $\{r_{\min}\}\times[s_{\min},\frac{\varpi^\kappa}{2})$ or $[r_{\min},\frac{\varpi^\kappa}{2})\times\{s_{\min}\}$. That is, one can connect pair $(s_{\min},r_{\min})$ to (s_0,r_0) by a join of coordinate line and the solution $(r(t),s(t))$. According to $(*)^\pm_\varphi$ and $(**)^\pm_\varphi$, the value of φ does not increase while pair (r,s) moving along this join. Thus $\varphi(r_0,s_0)\leqslant$ $\leqslant\varphi(r_{\min},s_{\min})$.

As before we rewrite the inequalities $(*)^\pm_\varphi$ and $(**)^\pm_\varphi$ in terms of ℓ:

$$(*)^\pm_\ell \qquad \begin{aligned}&\frac{\partial^+\ell}{\partial s}(s,r_{\min})\leqslant\cos\measuredangle^\kappa\{r_{\min};s,\ell\},\\&\frac{\partial^+\ell}{\partial r}(s_{\min},r)\leqslant\cos\measuredangle^\kappa\{s_{\min};r,\ell\},\end{aligned}$$

$$(**)^\pm_\ell \qquad \begin{aligned}&\operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S\cdot\frac{\partial^+\ell}{\partial s}+\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R\cdot\frac{\partial^+\ell}{\partial r}\leqslant\\&\qquad\leqslant\operatorname{sn}^\kappa s\cdot\operatorname{cs}^\kappa S\cdot\cos\measuredangle^\kappa\{r;s,\ell\}+\operatorname{sn}^\kappa r\cdot\operatorname{cs}^\kappa R\cdot\cos\measuredangle^\kappa\{s;r,\ell\}\end{aligned}$$

Further, set

$$(A)^{\pm} \qquad f = -\tfrac{1}{\kappa}\cdot\mathrm{cs}^{\kappa}\circ\mathrm{dist}_p = \mathrm{md}^{\kappa}\circ\mathrm{dist}_p - \tfrac{1}{\kappa}.$$

Clearly $f'' + \kappa\cdot f \leqslant 0$ and

$$(B)^{\pm} \qquad \begin{aligned} \rho^{+}(r) &= \frac{1}{\operatorname{tg}\kappa r\cdot\mathrm{cs}^{\kappa}R}\cdot\nabla_{\rho(r)}f \\ \sigma^{+}(s) &= \frac{1}{\operatorname{tg}\kappa s\cdot\mathrm{cs}^{\kappa}S}\cdot\nabla_{\sigma(s)}f. \end{aligned}$$

Thus from 11.4.5, we have

$$(C)^{\pm} \qquad \begin{aligned} \frac{\partial^{+}\ell}{\partial r} &= -\frac{1}{\operatorname{tg}\kappa r\cdot\mathrm{cs}^{\kappa}R}\cdot\langle\nabla_{\rho(r)}f, \uparrow_{[\rho(r)\sigma(s)]}\rangle \leqslant \\ &\leqslant \frac{1}{\operatorname{tg}\kappa r\cdot\mathrm{cs}^{\kappa}R}\cdot\frac{\mathrm{cs}^{\kappa}S - \mathrm{cs}^{\kappa}R\cdot\mathrm{cs}^{\kappa}\ell}{\kappa\cdot\mathrm{sn}^{\kappa}\ell} = \\ &= \frac{\frac{\mathrm{cs}^{\kappa}S}{\mathrm{cs}^{\kappa}R} - \mathrm{cs}^{\kappa}\ell}{\kappa\cdot\operatorname{tg}\kappa r\cdot\mathrm{sn}^{\kappa}\ell}. \end{aligned}$$

Note that for all $\kappa \neq 0$, the function $x \mapsto \frac{1}{\kappa\cdot\mathrm{cs}^{\kappa}x}$ is increasing. Thus, since $R(r) \leqslant r$ and $S(s_{\min}) = s_{\min}$, we get

$$(D)^{\pm} \qquad \begin{aligned} \frac{\partial^{+}\ell}{\partial r}(r, s_{\min}) &\leqslant \frac{\frac{\mathrm{cs}^{\kappa}s_{\min}}{\mathrm{cs}^{\kappa}r} - \mathrm{cs}^{\kappa}\ell}{\kappa\cdot\operatorname{tg}\kappa r\cdot\mathrm{sn}^{\kappa}\ell} = \\ &= \frac{\mathrm{cs}^{\kappa}s_{\min} - \mathrm{cs}^{\kappa}\ell\cdot\mathrm{cs}^{\kappa}r}{\kappa\cdot\mathrm{sn}^{\kappa}r\cdot\mathrm{sn}^{\kappa}\ell} = \\ &= \cos\measuredangle^{\kappa}\{s_{\min}; r, \ell\}, \end{aligned}$$

which is the first inequality in $(*)^{\pm}_{\ell}$ for $\kappa \neq 0$. By switching places of ρ and σ we obtain the second inequality in $(*)^{\pm}_{\ell}$. Further, summing together $(C)^{\pm}$ with its mirror-inequality for $\frac{\partial^{+}\ell}{\partial s}$, we get

$$(E)^{\pm} \qquad \begin{aligned} \mathrm{sn}^{\kappa}r\cdot\mathrm{cs}^{\kappa}R\cdot\frac{\partial^{+}\ell}{\partial r} &+ \mathrm{sn}^{\kappa}s\cdot\mathrm{cs}^{\kappa}S\cdot\frac{\partial^{+}\ell}{\partial s} \leqslant \\ &\leqslant \frac{\mathrm{cs}^{\kappa}S\cdot\mathrm{cs}^{\kappa}r - \mathrm{cs}^{\kappa}\ell\cdot\mathrm{cs}^{\kappa}R\cdot\mathrm{cs}^{\kappa}r}{\kappa\cdot\mathrm{sn}^{\kappa}\ell} + \\ &\quad + \frac{\mathrm{cs}^{\kappa}R\cdot\mathrm{cs}^{\kappa}s - \mathrm{cs}^{\kappa}\ell\cdot\mathrm{cs}^{\kappa}S\cdot\mathrm{cs}^{\kappa}s}{\kappa\cdot\mathrm{sn}^{\kappa}\ell} = \\ &= \mathrm{sn}^{\kappa}r\cdot\mathrm{cs}^{\kappa}R\cdot\frac{\mathrm{cs}^{\kappa}s - \mathrm{cs}^{\kappa}\ell\cdot\mathrm{cs}^{\kappa}r}{\kappa\cdot\mathrm{sn}^{\kappa}r\cdot\mathrm{sn}^{\kappa}\ell} + \\ &\quad + \mathrm{sn}^{\kappa}s\cdot\mathrm{cs}^{\kappa}S\cdot\frac{\mathrm{cs}^{\kappa}r - \mathrm{cs}^{\kappa}\ell\cdot\mathrm{cs}^{\kappa}s}{\kappa\cdot\mathrm{sn}^{\kappa}s\cdot\mathrm{sn}^{\kappa}\ell} = \\ &= \mathrm{sn}^{\kappa}r\cdot\mathrm{cs}^{\kappa}R\cdot\cos\measuredangle^{\kappa}\{r; s, \ell\} + \\ &\quad + \mathrm{sn}^{\kappa}s\cdot\mathrm{cs}^{\kappa}S\cdot\cos\measuredangle^{\kappa}\{s; r, \ell\} \end{aligned}$$

which is $(**)^{\pm}_{\ell}$. $\square$

14.6.4. Exercise. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $x, y, z \in \mathcal{L}$. Assume $\measuredangle^{\kappa}(z\,{}^{x}_{y}) = \pi$. Show that there is a geodesic $[xy]$ that contains z. In particular, x can be connected to y by a minimizing geodesic.*

Compare to Exercise 9.4.6

14.7 Gradient exponent

Let $\mathcal{L}$ be a complete length CBB(κ) space, $p \in \mathcal{L}$ and $\xi \in \Sigma_p$. Consider a sequence of points $x_n \in \mathcal{L}$ so that $\uparrow_{[px_n]} \to \xi$. Set $r_n = |p - x_n|$; denote by $\sigma_n \colon [r_n, \frac{\varpi^{\kappa}}{2}) \to \mathcal{L}$ be the (p, κ)-radial curve that starts in x_n.

From the radial comparison (14.6.3), we have that $\sigma_n \colon [r_n, \frac{\varpi^{\kappa}}{2}) \to$ $\to \mathcal{L}$ converge to a curve $\sigma_\xi \colon (0, \frac{\varpi^{\kappa}}{2}) \to \mathcal{L}$ and this limit is independent from the choice of the sequence x_n. Set $\sigma_\xi(0) = p$ and if $\kappa > 0$ define

$$\sigma_\xi(\tfrac{\varpi^{\kappa}}{2}) = \lim_{t \to \frac{\varpi^{\kappa}}{2}} \sigma_\xi(t).$$

The obtained curve σ_ξ will be called (p, κ)-*radial curve in direction* ξ.

Let us define *gradient exponent* as

$$\operatorname{gexp}^{\kappa}_p \colon r \cdot \xi \mapsto \sigma_\xi(r) \colon \overline{\mathrm{B}}[0, \tfrac{\varpi^{\kappa}}{2}] \subset \mathrm{T}_p \to \mathcal{L}.$$

Here are properties of radial curves reformulated in terms of gradient exponent:

14.7.1. Theorem. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space. Then*

a) If $p, q \in \mathcal{L}$ be points such that $|p - q| \leqslant \frac{\varpi^{\kappa}}{2}$, then for any geodesic $[pq]$ in $\mathcal{L}$, we have

$$\operatorname{gexp}^{\kappa}_p(\log[pq]) = q$$

b) For any $v, w \in \overline{\mathrm{B}}[0, \frac{\varpi^{\kappa}}{2}] \subset \mathrm{T}_p$,

$$|\operatorname{gexp}^{\kappa}_p v - \operatorname{gexp}^{\kappa}_p w| \leqslant \tilde{\Upsilon}^{\kappa}[0\,{}^{v}_{w}].$$

In other words, if we denote by $\mathcal{T}^{\kappa}_p$ the set $\overline{\mathrm{B}}[0, \frac{\varpi^{\kappa}}{2}] \subset \mathrm{T}_p$ equipped with metric $|v - w|_{\mathcal{T}^{\kappa}_p} = \tilde{\Upsilon}^{\kappa}[0\,{}^{v}_{w}]$, then

$$\operatorname{gexp}^{\kappa}_p : \mathcal{T}^{\kappa}_p \to \mathcal{L}$$

is a short map.

c) Let $\mathcal{L}$ be a complete length $\mathrm{CBB}(\kappa)$ *space, $p, q \in \mathcal{L}$ and $|p-q| \leqslant \frac{\varpi^\kappa}{2}$. If $v \in \mathrm{T}_p$, $|v| \leqslant 1$ and*

$$\sigma(t) = \mathrm{gexp}_p^\kappa(t{\cdot}v),$$

then the function

$$s \mapsto \mathcal{Z}^\kappa(\sigma|_0^s, q) := \mathcal{Z}^\kappa\{|q-\sigma(s)|; |q-\sigma(0)|, s\}$$

is non-increasing in whole domain of definition.

Proof. Follows directly from construction of gexp_p^κ and the radial comparison (14.6.3). □

Applying the theorem above together with 13.4.1*c*, we obtain the following.

14.7.2. Corollary. *Let $\mathcal{L}$ be an m-dimensional complete length* $\mathrm{CBB}(\kappa)$ *space, $p \in \mathcal{L}$ and $0 < R \leqslant \frac{\varpi^\kappa}{2}$. Then there is a short map $f\colon \overline{\mathrm{B}}[R]_{\mathbb{M}^m(\kappa)} \to \mathcal{L}$ such that $\operatorname{Im} f = \overline{\mathrm{B}}[p, R] \subset \mathcal{L}$.*

14.7.3. Exercise. *Let $\mathcal{L} \subset \mathbb{E}^2$ be Euclidean halfplane. Clearly $\mathcal{L}$ is a two-dimensional complete length* $\mathrm{CBB}(0)$ *space. Given a point $x \in \mathbb{E}^2$ let us denote by* $\mathrm{proj}(x)$ *the closest point on $\mathcal{L}$.*

Apply the Radial comparison (14.6.3) to show that for any interior point $p \in \mathcal{L}$ we have

$$\mathrm{gexp}_p v = \mathrm{proj}(p+v).$$

14.7.4. Exercise. *Let $\mathcal{L}$ be an m-dimensional complete length* $\mathrm{CBB}(\kappa)$ *space and* $\operatorname{rad}\mathcal{L} = R$. *Prove that there is a* $(\mathrm{sn}^\kappa R)$*-Lipschitz map $\Phi\colon \mathbb{S}^{m-1} \to \mathcal{L}$ such that $\operatorname{Im}\Phi \supset \partial\mathcal{L}$.*

14.8 Remarks

Gradient flow on Riemannian manifolds

The gradient flow for general semiconcave functions on smooth Riemannian manifold can be introduced much cheaper. To do this note that the distance estimates proved in the Section 14.2 can be proved the same way for gradient curves of smooth semiconcave subfunctions. By Greene–Wu lemma [65], given a λ-concave function f, a compact set $K \subset \operatorname{Dom} f$ and $\varepsilon > 0$ there is a smooth $(\lambda - \varepsilon)$-concave function that is ε-close to f on K. Hence one can apply the smoothing and pass to the limit as $\varepsilon \to 0$. Note that by the second distance estimate the obtained limit curve does not depend on the smoothing.

Gradient curves of a family of functions

Gradient flow can be extended to a family of functions.

Chanyoung Jun used this flow to study Pursuit curves in CAT(0) spaces [66, 67]. Also the optimal transport plan, or equivalently geodesic in Wasserstein metric, can be described as gradient flow for a family of semiconcave functions. This observation was used by the third author to prove Alexandrov spaces with nonnegative curvature have nonnegative Ricci curvature in the sense of Lott–Villani–Sturm [68].

Gradient curves for non-Lipschitz functions

In this book, we only consider gradient curves for locally Lipschitz semiconcave subfunctions; it turns out to be sufficient for all our needs. However, instead of Lipschitz semi-concave subfunctions, it is more natural to consider upper semi-continuous semi-concave functions with target in $[-\infty,\infty)$ and to assume in addition that the function take finite values at a dense set in the domain of definition. The set of such subfunctions on a CBB space $\mathcal{L}$ will be denoted as $\mathrm{LCC}(\mathcal{L})$ for **l**ower semi-**c**ontinous and semi-**c**oncave.

In this section we describe the adjustments needed to construct gradient curves in for the subfunctions in $\mathrm{LCC}(\mathcal{L})$.

There is one place where this type of functions appears; it is the entropy and some other closely related functionals on the Wasserstein space over a CBB(0) space. The gradient flow for these function play an important role in the theory of optimal transport, see [69] and references there in.

Differential. First we need to extend the definition of differential (5.7.1) to LCC subfunctions.

Let $\mathcal{L}$ be a complete length CBB space and $f\in\mathrm{LCC}(\mathcal{L})$. Given a point $p\in\operatorname{Dom}f$, and a geodesic direction $\xi=\uparrow_{[pq]}$, set $\hat{\boldsymbol{d}}_pf(\xi)=(f\circ\circ\,\mathrm{geod}_{[pq]})^+(0)$. Since f is semiconcave the value $\hat{\boldsymbol{d}}_pf(\xi)$ is defined if $f\circ\mathrm{geod}_{[pq]}(t)$ is finite at all sufficiently small values $t>0$, but $\hat{\boldsymbol{d}}_pf(\xi)$ may take value ∞. Note that $\hat{\boldsymbol{d}}_pf$ is defined on a dense subset of Σ_p.

Set

$$\boldsymbol{d}_pf(\zeta)=\varlimsup_{\xi\to\zeta}\hat{\boldsymbol{d}}_pf(\xi)$$

and then set $\boldsymbol{d}_pf(v)=|v|\cdot\boldsymbol{d}_pf(\xi)$ if $v=|v|\cdot\xi$ for some $\xi\in\Sigma_p$.

In other words, we define differential as the smallest upper semi-continuous positive-homogeneous function $\boldsymbol{d}_pf\colon\mathrm{T}_p\to\mathbb{R}$ such that $\boldsymbol{d}_pf(\xi)\geqslant\hat{\boldsymbol{d}}_pf(\xi)$ if $\hat{\boldsymbol{d}}_pf(\xi)$ is defined.

Existence and uniqueness of the gradient. Note that in the proof of 11.4.2, we used Lipschitz condition just once, to show that

$$\begin{aligned} s &= \sup\{\,(\boldsymbol{d}_p f)(\xi) : \xi \in \Sigma_p\,\} = \\ &= \varlimsup_{x\to p} \frac{f(x)-f(p)}{|x-p|} < \\ &< \infty. \end{aligned}$$

The value s above will be denoted as $|\nabla|_p f$. Note that if the gradient $\nabla_p f$ is defined then $|\nabla|_p f = |\nabla_p f|$ and otherwise $|\nabla|_p f = \infty$.

Summarizing the discussion above, we get the following.

11.4.2′ Existence and uniqueness of the gradient. *Assume $\mathcal{L}$ a complete* CBB(κ) *space and $f \in \mathrm{LCC}(\mathcal{L})$. Then for any point $p \in \in \operatorname{Dom} f$ either there is unique gradient $\nabla_p f \in \mathrm{T}_p$ or $|\nabla|_p f = \infty$*

Further, in all the results of Section 11.4, we can only assume that $f \in \mathrm{LCC}(\mathcal{L})$ and its gradient is defined at the points in the consideration; the proofs are the same.

In sections 11.4–14.2 require almost no changes; mainly, where it is appropriate, one needs to exchange $|\nabla_p f|$ to $|\nabla|_p f$ and/or assume that the gradient is defined at the points of interest. Also one has to take ❶ in Theorem 14.1.2 as the definition of gradient-like curve. Then the theorem states that any gradient-like curve $\alpha\colon \mathbb{I} \to \mathcal{L}$ satisfies the definition 14.1.1 at $t \in \mathbb{I}$ if $\nabla_{\hat\alpha(s)} f$ is defined. Further in the Definition 14.2.1, should be changed to the following

14.2.1′. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and $f \in \mathrm{LCC}(\mathcal{L})$.*

*A curve $\alpha\colon [t_{\min}, t_{\max}) \to \operatorname{Dom} f$ will be called f-*gradient curve *if*

$$\alpha^+(t) = \nabla_{\alpha(t)} f$$

if $\nabla_{\alpha(t)} f$ is defined and

$$(f\circ\alpha)^+(t) = \infty$$

otherwise.

In the proof of Local existence (14.2.9), the condition (ii) has to be exchanged to the following condition

(ii)′ $f\circ\hat\alpha_n(\bar\sigma_i) - f\circ\hat\alpha_n(\sigma_i) > (\bar\sigma_i - \sigma_i)\cdot \max\{n, |\nabla|_{\hat\alpha_n(\sigma_i)} f - \frac{1}{n})\}$.

Any gradient curve $\alpha[0,\ell) \to \mathcal{L}$ for a subfunction $f \in \mathrm{LCC}(\mathcal{L})$ satisfy the equation

$$\alpha^+(t) = \nabla_{\alpha(t)} f$$

at all values t with possible exception $t = 0$. In particular, the gradient of f is defined at all points of any f-gradient curve with the exception for the initial point.

Slower radial curves

Let $\kappa \geqslant 0$. Assume that for some function ψ, the curves defined by equation

$$\sigma^+(s) = \psi(s, |p - \sigma(s)|) \cdot \nabla_{\sigma(s)} \operatorname{dist}_p$$

satisfy radial comparison 14.6.3. Then in fact $\sigma(s)$ are radial curves; that is,

$$\psi(s, |p - \sigma(s)|) = \frac{\operatorname{tg} \kappa |p - \sigma(s)|)}{\operatorname{tg} \kappa s},$$

see exercise 14.7.3.

In case $\kappa < 0$, such ψ is not unique. In particular one can take curves defined by simpler equation

$$\sigma^+(s) = \frac{\operatorname{sn}^\kappa |p - \sigma(s)|}{\operatorname{sn}^\kappa s} \cdot \nabla_{\sigma(s)} \operatorname{dist}_p = \frac{1}{\operatorname{sn}^\kappa s} \cdot \nabla_{\sigma(s)} (\operatorname{md}^\kappa \circ \operatorname{dist}_p).$$

Among all curves of that type, the radial curves for curvature κ as defined in 14.5.1 maximize the growth of $|p - \sigma(s)|$.

Radial curves for sets

Here we generalize the constructions of radial curves and gradient exponent. Roughly, we show that one can use a distance function dist_A to any closed set A instead of the distant function to one point. We only give the corresponding definitions and state the results, the proofs are straightforward generalization of corresponding one-point-set version.

First we give a more general form of the definition of radial curves (14.5.1) and the definition of radial geodesic (14.5.2):

14.8.1. Definition. *Assume $\mathcal{L}$ is a complete length* CBB *space, $\kappa \in \mathbb{R}$, and $A \subset \mathcal{L}$ be a closed subset. A curve $\sigma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ is called* (A, κ)-radial curve *if $s_{\min} = \operatorname{dist}_A \sigma(s_{\min}) \in (0, \frac{\varpi^\kappa}{2})$, and it satisfies the following differential equation*

$$\sigma^+(s) = \frac{\operatorname{tg} \kappa |p - \sigma(s)|}{\operatorname{tg} \kappa s} \cdot \nabla_{\sigma(s)} \operatorname{dist}_A .$$

for any $s \in [s_{\min}, s_{\max})$, here $\operatorname{tg} \kappa x = \frac{\operatorname{sn}^\kappa x}{\operatorname{cs}^\kappa x}$.

If $x = \sigma(s_{\min})$, we say that σ starts in *x.*

14.8.2. Definition. *Let $\mathcal{L}$ be a complete length* CBB *space and $A \subset \mathcal{L}$ be a closed subset. A unit-speed geodesic $\gamma\colon \mathbb{I} \to \mathcal{L}$ is called* A-radial geodesic *if $\operatorname{dist}_A \gamma(s) \equiv s$.*

The following propositions are analogous to the propositions 14.5.3 and 14.5.4. Their proofs follow directly from the definitions:

14.8.3. Proposition. *Let $\mathcal{L}$ be a complete length* CBB *space, $A \subset \mathcal{L}$ be a closed subset. Assume that $\frac{\varpi^\kappa}{2} \geqslant s_{\max}$. Then any* dist_A*-radial geodesic $\gamma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ is an (A,κ)-radial curve.*

14.8.4. Proposition. *Let $\mathcal{L}$ be a complete length* CBB *space, $A \subset \mathcal{L}$ be a closed subset and $\sigma\colon [s_{\min}, s_{\max}) \to \mathcal{L}$ be a (A,κ)-radial curve. Then for any $s \in [s_{\min}, s_{\max})$, we have $\operatorname{dist}_A \sigma(s) \leqslant s$.*

Moreover, if for some s_0 we have $\operatorname{dist}_A \sigma(s_0) = s_0$ then the restriction $\sigma|_{[s_{\min}, s_0]}$ is a A-radial geodesic.

Here is the corresponding generalization of existence and uniqueness for (A,κ)-radial curves; it can be proved the same way as 14.5.5

14.8.5. Existence and uniqueness. *Let $\mathcal{L}$ be a complete length* CBB *space, $\kappa \in \mathbb{R}$, $A \subset \mathcal{L}$ be a closed subset, and $x \in \mathcal{L}$. Assume $0 < \operatorname{dist}_A x < \frac{\varpi^\kappa}{2}$. Then there is unique (A,κ)-radial curve $\sigma\colon [\operatorname{dist}_A x, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ that starts in x.*

Next we formulate radial monotonicity and radial comparison for (A,κ)-radial curves. The proof of these two statements are almost exactly the same as the proof of 14.6.1 and 14.6.3.

14.8.6. Radial monotonicity. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space, $A \subset \mathcal{L}$ be a closed subset and $q \in \mathcal{L}\backslash A$. Assume $\sigma\colon [s_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ is an (A,κ)-radial curve. Then the function*

$$s \mapsto \measuredangle^\kappa\{|q - \sigma(s)|; \operatorname{dist}_A q, s\}$$

is nonincreasing in all the domain of definition.

To formulate generalized radial comparison, we need to introduce a short cut notation. Given a set A and two points x and y in a metric space define

$$\measuredangle^\kappa\left(A\,{}^{x}_{y}\right) := \measuredangle^\kappa\{|x - y|; \operatorname{dist}_A x, \operatorname{dist}_A y\}$$

Note that distances $|x - y|$, $\operatorname{dist}_A x$ and $\operatorname{dist}_A y$ might not satisfy the triangle inequality. Therefore the model angle $\measuredangle^\kappa\left(A\,{}^{x}_{y}\right)$ might be undefined even for $\kappa \leqslant 0$.

14.8.7. Radial comparison. *Let $\mathcal{L}$ be a complete length* CBB(κ) *space and $A \subset \mathcal{L}$ be a closed set. Assume $\rho\colon [r_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ and $\sigma\colon [s_{\min}, \frac{\varpi^\kappa}{2}) \to \mathcal{L}$ are two* dist_A*-radial curves for curvature κ. Assume further that*

$$\varphi_{\min} = \measuredangle^\kappa\left(A\,{}^{\rho(r_{\min})}_{\sigma(s_{\min})}\right)$$

is defined. Then for any $r \in [r_{\min}, \frac{\varpi^\kappa}{2})$ *and* $s \in [s_{\min}, \frac{\varpi^\kappa}{2})$*, we have*

$$|\rho(r) - \sigma(s)| \leqslant \tilde{\Upsilon}^\kappa\{\varphi_{\min}; r, s\}.$$

Finally, assume p be an isolated point of a closed set A in a complete length CBB space $\mathcal{L}$. Applying the same limiting procedure as in Section 14.7, for any $\xi \in \Sigma_p$ one can construct an (A, κ)-radial curve σ_ξ such that $\sigma_\xi(0) = p$ and $\sigma^+(0) = \xi$. This way we obtain a map $\operatorname{gexp}^\kappa_A \colon \mathrm{T}_p \looparrowright \mathcal{L}$, $r{\cdot}\xi \mapsto \sigma_\xi(r)$. For the constructed map, the following analog of 14.7.1 holds; the proof is straightforward.

14.8.8. Theorem. *Let* $\mathcal{L}$ *be a complete length* CBB(κ) *space and* $A \subset \mathcal{L}$ *is a closed subset with an isolated point* $p \in A$*. Then*

a) Assume $\operatorname{dist}_A q = |p - q| \leqslant \frac{\varpi^\kappa}{2}$ *be an* A*-radial geodesic. Then*

$$\operatorname{gexp}^\kappa_A(\log[pq]) = q$$

b) For any $v, w \in \overline{\mathrm{B}}[0, \frac{\varpi^\kappa}{2}] \subset \mathrm{T}_p$*,*

$$|\operatorname{gexp}^\kappa_p v - \operatorname{gexp}^\kappa_p w| \leqslant \tilde{\Upsilon}^\kappa[0\,{}^{v}_{w}].$$

In other words, if we denote by $\mathcal{T}^\kappa_p$ *the set* $\overline{\mathrm{B}}[0, \frac{\varpi^\kappa}{2}] \subset \mathrm{T}_p$ *equipped with metric* $|v - w|_{\mathcal{T}^\kappa_p} = \tilde{\Upsilon}^\kappa[0\,{}^{v}_{w}]$*, then*

$$\operatorname{gexp}^\kappa_p : \mathcal{T}^\kappa_p \to \mathcal{L}$$

is a short map.

c) Let $\mathcal{L}$ *be a complete length* CBB(κ) *space,* $p, q \in \mathcal{L}$ *and* $|p - q| \leqslant \frac{\varpi^\kappa}{2}$*. If* $v \in \mathrm{T}_p$*,* $|v| \leqslant 1$ *and*

$$\sigma(t) = \operatorname{gexp}^\kappa_p(t{\cdot}v),$$

then the function $s \mapsto \measuredangle^\kappa(\sigma|^s_0, q)$ *is non-increasing in whole domain of definition.*

Index

R-geodesic space, 12
R-length space, 12
$V^{\wedge n}$, 62
$[**]$, 3, 47
CAT(κ) comparison, 101
Δ^m, 282
$\mathbb{I}$, 47
LinDim, 69
$\pounds$-Lipschitz map, 38
$\pounds$-co-Lipshitz map, 38
$\mathbb{M}^2(\kappa)$, 3
Σ_∞, 326
$\mathrm{Str}(p)$, 76
T'_*, 202
T_∞, 326
MinPoint, 282
cs^κ, 4
$|* - *|$, 3
ℓ^1-metric, 291
ε-linear function, 64
ε-pack, 10
$\mathrm{geod}_{[**]}$, 47
$[*\,{}^*_*]$, 48
itg^κ, 255
κ-barycentric simplex, 287
κ-fat, 81
κ-monotonic curve, 309
κ-thin, 107
$\measuredangle$, 3
    $\measuredangle[*\,{}^*_*]$, 50
    $\measuredangle^{\mathrm{up}}$, 54
md^κ, 4
$\tilde\triangle^\kappa$
    $\tilde\triangle^\kappa(***)$, 48
    $\tilde\triangle^\kappa\{*,*,*\}$, 3
μ-measurable, 55
∇, 235
ω
    ω-limit, 30
ω-limit space, 32
$\tilde\Upsilon^\kappa$
    $\tilde\Upsilon^\kappa[*\,{}^*_*]$, 50
    $\tilde\Upsilon^\kappa\{*;*,*\}$, 4
sn^κ, 4
$\mathfrak{S}_{\boldsymbol{f}}$, 282
$\mathbb{Z}^\kappa$
    $\mathbb{Z}^\kappa(*\,{}^*_*)$, 49
    $\mathbb{Z}^\kappa(*|{}^*_*,*)$, 299
$[***]$, 48
$[***]$, 3
ϖ^κ, 3

absolute value, 163
Alexandrov's lemma, 49
angle, 50
    extended model angle, 100
    strong angle, 96
    upper angle, 54
arc-point model angle, 299

barrier, 43
barycentric simplex of function array, 282
base of development, 83
blowup, 204
both-side differentiable, 299
bounded set, 10
Busemann function, 48

Carathéodory's construction, 56
closed ball, 9
closed neighborhood, 10
co-Lipshitz map, 38
common space, 20
comparison

adjacent angle comparison, 79
adjacent-angles comparison, 105
development comparison, 84, 114
function comparison, 82
point-on-side comparison, 105
complete gradient, 251
cone, 52
cone at infinity, 326
conic map, 303
constantspeed geodesic, 47
continuous submap, 37
convex
κ-convex curve, 298
convex with respect to a point, 84
locally convex development with respect to a point, 84
convex set, 14
weakly convex set, 14
coordinate of the distant chart, 212
cubical analog, 190
cubical complex, 189
cubulation, 189
curve, 41

decomposed triangle, 108
decrypting matrix, 152
defined angles, 53
development, 83
Diameter, 10
differentiable
both-sided differentiable, 201
left differentiable, 201
right differentiable, 201
differential, 201
dimension
geometric dimension, 70
Hausdorff dimension, 67
Linear dimension, 69
local dimension, 70
topological dimension, 67
directionally limited, 61
disjoint union of metric spaces, 8
distance, 7
distance function, 8
distance preserving map, 39
distant chart, 212
doubling, 17
doubling measure, 59
doubling space, 58

Euclidean polyhedral space, 183
Euclidean volume, 58
extended model angle, 100
exterior power, 62
extremal set, 269, 270
extremal subset, 272

flag complex, 186
funny quasigeodesic, 321

G-delta geodesic space, 76
geodesic, 47
G-delta geodesic space, 76
local geodesic, 48
geodesic path, 48
geodesic space, 12
geodesic tangent vector, 53
geometric dimension, 70
gluing, 17
gradient, 235
gradient flow, 251
Gromov–Hausdorff convergence, 21
Gromov–Hausdorff distance, 25

Hamilton–Jacobi shifts, 307
Hausdorff measure, 56
hinge, 48
hinge comparison, 79
hyperbolic polyhedral space, 183

ideal boundary, 326
inner web, 282
isometric spaces, 40
isometry, 40
ε-siometry, 24

Jensen's inequality, 43

Kirszbraun target, 154
Kirszbrun source, 154
Kuratowski map, 18

lemma
    Alexandrov's lemma, 49
length space, 11
length-metric, 11
lifting of a distance function, 22
lifting of a point, 22
liftings of the distance chart, 214
line, 47
linear subspace, 208
Lipschitz curve, 41
Lipschitz map, 38
local dimension, 70
locally length space, 12
locally Lipschitz curve, 41
locally Lipschitz map, 38
long measure, 56

maximal ε-packing, 10, 219
metric component, 9
model angle
    arc-point model angle, 299
    extended model angle, 100
model triangle, 48

natural map, 107
negative critical point, 169
no-triangle condition, 186
nondegenerate function array, 282
nondegenerate point array, 287
nonprinciple ultrafilter, 29

open ball, 9
open set, 10
open tubular neighborhood, 10
opposite, 208
order, 67
orientable, 279
overlap, 141

plane
    κ-plane, 3
point-on-side comparison, 79
polar, 298
polar vector, 207
polyhedral spaces
    locally polyhedral spaces, 183
polytope, 175
positive critical point, 169
pre-quasigeodesic, 309
    Ω-pre-quasigeodesic, 314
primitive extremal set, 270
probation curve, 306
proper extremal subset, 272
proper space, 10

quasigeodesic
    κ-quasigeodesic, 298
quotient space, 16

radial geodesic, 255, 265
Radius, 10
radius of the point array, 287
ray, 47
real interval, 41
rectifiable set, 63
    countably rectifiable, 63
    locally rectifiable, 63
refinement, 67
reflection, 17
regular
    $(\bar{k}, \varepsilon)$-regular, 64
    $\bar{k}$-regular, 64
regular point of semiconvex function, 64

saddle function, 175
scalar product, 164
selective ultrafilter, 29
self-contracting curve, 244
short map, 38
simply connected space at infinity, 190
soul, 325
spherical measure, 56
spherical polyhedral space, 183
spherical polytope, 175
stable value, 68
standard simplex, 282
star of vertex, 188
straight
    $*$-straight point, 76
strong angle, 96
strongly two-convex set, 172
strut, 221
strut of the distant chart, 212

subfunction, 37
submap, 37
submetry, 40
subnormal vector, 284
subspace, 8

tangent vector, 53
tip of the cone, 52
totally convex, 14
totally convex set
    totally R-convex set, 15
triangle, 48
triangulation of a polyhedral space
        space, 183
turn to the point p, 321
two-convex set, 168

ultrafilter, 29
    nonprinciple ultrafilter, 29
    selective ultrafilter, 29
ultralimit, 30
unit-speed curve, 41
unit-speed geodesic, 47

vector, 163
vertexes of barycentric simplex, 287
Vitali covering, 61

weakly convex set, 14
    weakly R-convex set, 15
web, 282
web embedding, 287

Bibliography

[1] Bruce Kleiner and Bernhard Leeb. "Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings". *Inst. Hautes Études Sci. Publ. Math.* 86 (1997), 115–197 (1998).

[2] Walter Rudin. "Homogeneity problems in the theory of Čech compactifications". *Duke Math. J.* 23 (1956), pp. 409–419.

[3] Yu. Burago, M. Gromov, and G. Perelman. "A. D. Aleksandrov spaces with curvatures bounded below". *Russian Math. Surveys* 47.2 (1992), pp. 1–58.

[4] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry.* Vol. 33. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2001, pp. xiv+415.

[5] Bernd Kirchheim. "Rectifiable metric spaces: local structure and regularity of the Hausdorff measure". *Proc. Amer. Math. Soc.* 121.1 (1994), pp. 113–123.

[6] Witold Hurewicz and Henry Wallman. *Dimension Theory.* Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941, pp. vii+165.

[7] L. Pontrjagin. "Sur une hypothèse fondamentale de la théorie de la dimension." *C. R. Acad. Sci., Paris* 190 (1930), pp. 1105–1107.

[8] A. S. Besicovitch and P. A. P. Moran. "The measure of product and cylinder sets". *J. London Math. Soc.* 20 (1945), pp. 110–120.

[9] Jun-iti Nagata. *Modern dimension theory.* Revised. Vol. 2. Sigma Series in Pure Mathematics. Heldermann Verlag, Berlin, 1983, pp. ix+284.

[10] Ryszard Engelking. *Dimension theory.* North-Holland Publishing Co., Amsterdam-Oxford-New York; PWN—Polish Scientific Publishers, Warsaw, 1978, x+314 pp. (loose errata).

[11] Thomas Foertsch and Viktor Schroeder. "Minkowski versus Euclidean rank for products of metric spaces". *Adv. Geom.* 2.2 (2002), pp. 123–131.

[12] Conrad Plaut. "Metric spaces of curvature $\geq k$". *Handbook of geometric topology.* North-Holland, Amsterdam, 2002, pp. 819–898.

[13] Bruce Kleiner. "The local structure of length spaces with curvature bounded above". *Math. Z.* 231.3 (1999), pp. 409–456.

[14] Abraham Wald. „Begründung eiiner Koordinatenlosen Differentialgeometrie der Flächen". *Ergebnisse eines mathematischen Kolloquium* 6 (1935), S. 24–46.

[15] Herbert Busemann. "Spaces with non-positive curvature". *Acta Math.* 80 (1948), pp. 259–310.

[16] A. D. Alexandrow. "Über eine Verallgemeinerung der Riemannschen Geometrie". *Schr. Forschungsinst. Math.* 1 (1957), pp. 33–84.

[17] И. М. Либерман. «Геодезические линии на выпуклых поверхностях». *Докл. АН СССР* 32.2 (1941), с. 310—312.

[18] P. Pizzetti. "Paragone fra due triangoli a lati uguali". *Atti della Reale Accademia dei Lincei, Rendiconti (5). Classe di Scienze Fisiche, Matematiche e Naturali* 16.1 (1907), pp. 6–11.

[19] Виктор Андреевич Топоногов. «Римановы пространства кривизны, ограниченной снизу». *Успехи математических наук* 14.1 (85 (1959), с. 87—130.

[20] Conrad Plaut. “Spaces of Wald curvature bounded below”. *J. Geom. Anal.* 6.1 (1996), pp. 113–134.

[21] Katsuhiro Shiohama. *An introduction to the geometry of Alexandrov spaces.* Vol. 8. Lecture Notes Series. Seoul National University, Research Institute of Mathematics, Global Analysis Research Center, Seoul, 1993, pp. ii+78.

[22] Anton Petrunin. “A globalization for non-complete but geodesic spaces”. *Math. Ann.* 366.1-2 (2016), pp. 387–393.

[23] Urs Lang and Viktor Schroeder. “On Toponogov’s comparison theorem for Alexandrov spaces”. *Enseign. Math.* 59.3-4 (2013), pp. 325–336.

[24] Yu. G. Reshetnyak. “Inextensible mappings in a space of curvature no greater than K”. *Sibirsk. Mat. Ž.* 9.4 (1968), pp. 683–689.

[25] Sergei Ivanov. *To what extent is convexity a local property?* MathOverflow. (version: 2012-06-09). eprint: `https://mathoverflow.net/q/22091`.

[26] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature.* Vol. 319. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999, pp. xxii+643.

[27] Stephanie B. Alexander and Richard L. Bishop. “The Hadamard-Cartan theorem in locally convex metric spaces”. *Enseign. Math. (2)* 36.3-4 (1990), pp. 309–320.

[28] Stephanie B. Alexander and Richard L. Bishop. “Comparison theorems for curves of bounded geodesic curvature in metric spaces of curvature bounded above”. *Differential Geom. Appl.* 6.1 (1996), pp. 67–86.

[29] Werner Ballmann. *Lectures on spaces of nonpositive curvature.* Vol. 25. DMV Seminar. With an appendix by Misha Brin. Birkhäuser Verlag, Basel, 1995, pp. viii+112.

[30] É. Cartan. *Leçons sur la Géométrie des Espaces de Riemann.* Gauthier-Villars, Paris, 1928, pp. vi+273.

[31] Hans von Mangoldt. „Ueber diejenigen Punkte auf positiv gekrümmten Flächen, welche die Eigenschaft haben, dass die von ihnen ausgehenden geodätischen Linien nie aufhören, kürzeste Linien zu sein“. *J. Reine Angew. Math.* 91 (1881), S. 23–53.

[32] J. Hadamard. “Sur la forme des lignes géodésiques à l’infini et sur les géodésiques des surfaces réglées du second ordre”. *Bull. Soc. Math. France* 26 (1898), pp. 195–216.

[33] Willi Rinow. *Die innere Geometrie der metrischen Räume.* Die Grundlehren der mathematischen Wissenschaften, Bd. 105. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1961, pp. xv+520.

[34] M. Gromov. “Hyperbolic groups”. *Essays in group theory.* Vol. 8. Math. Sci. Res. Inst. Publ. Springer, New York, 1987, pp. 75–263.

[35] Werner Ballmann. “Singular spaces of nonpositive curvature”. *Sur les groupes hyperboliques d’après Mikhael Gromov (Bern, 1988).* Vol. 83. Progr. Math. Birkhäuser Boston, Boston, MA, 1990, pp. 189–201.

[36] B. H. Bowditch. “Notes on locally CAT(1) spaces”. *Geometric group theory (Columbus, OH, 1992).* Vol. 3. Ohio State Univ. Math. Res. Inst. Publ. de Gruyter, Berlin, 1995, pp. 1–48.

[37] Eva Kopecká and Simeon Reich. “Nonexpansive retracts in Banach spaces”. *Fixed point theory and its applications.* Vol. 77. Banach Center Publ. Polish Acad. Sci. Inst. Math., Warsaw, 2007, pp. 161–174.

[38] Stephanie Alexander, Vitali Kapovitch, and Anton Petrunin. “Alexandrov meets Kirszbraun”. *Proceedings of the Gökova Geometry-Topology Conference 2010.* Int. Press, Somerville, MA, 2011, pp. 88–109.

[39] U. Lang and V. Schroeder. “Kirszbraun’s theorem and metric spaces of bounded curvature”. *Geom. Funct. Anal.* 7.3 (1997), pp. 535–560.

[40] K. T. Sturm. "Metric spaces of lower bounded curvature". *Exposition. Math.* 17.1 (1999), pp. 35–47.

[41] Nina Lebedeva, Anton Petrunin, and Vladimir Zolotov. *Bipolar comparison.* arXiv: `1711.09423 [math.DG]`.

[42] Nicolas Monod. "Superrigidity for irreducible lattices and geometric splitting". *J. Amer. Math. Soc.* 19.4 (2006), pp. 781–814.

[43] Anton Petrunin and Stephan Stadler. *Metric minimizing surfaces revisited.* arXiv: `1707.09635 [math.DG]`.

[44] J. R. Isbell. "Six theorems about injective metric spaces". *Comment. Math. Helv.* 39 (1964), pp. 65–76.

[45] S. B. Alexander and R. L. Bishop. "Curvature bounds for warped products of metric spaces". *Geom. Funct. Anal.* 14.6 (2004), pp. 1143–1181.

[46] Stephanie B. Alexander and Richard L. Bishop. "Warped products admitting a curvature bound". *Adv. Math.* 303 (2016), pp. 88–122.

[47] Stephanie Halbeisen. "On tangent cones of Alexandrov spaces with curvature bounded below". *Manuscripta Math.* 103.2 (2000), pp. 169–182.

[48] Shin-ichi Ohta. "Gradient flows on Wasserstein spaces over compact Alexandrov spaces". *Amer. J. Math.* 131.2 (2009), pp. 475–516.

[49] Robert R. Phelps. *Convex functions, monotone operators and differentiability.* Second. Vol. 1364. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1993, pp. xii+117.

[50] A. Lytchak. "Differentiation in metric spaces". *St. Petersburg Math. J.* 16.6 (2005), pp. 1017–1041.

[51] M. Gromov. "Asymptotic invariants of infinite groups". *Geometric group theory, Vol. 2 (Sussex, 1991).* Vol. 182. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.

[52] G. Perelman. *Alexandrov paces with curvature bounded from below II.* Preprint LOMI, 1991. 35pp. eprint: `http://www.math.psu.edu/petrunin/`.

[53] G. Ya. Perelman. "Elements of Morse theory on Aleksandrov spaces". *St. Petersburg Math. J.* 5.1 (1994), pp. 205–213.

[54] A. Lytchak. "Open map theorem for metric spaces". *St. Petersburg Math. J.* 17.3 (2006), pp. 477–491.

[55] G. Perelman and A. Petrunin. *Quasigeodesics and Gradient curves in Alexandrov spaces.* eprint: `http://www.math.psu.edu/petrunin/`.

[56] В. А. Шарафутдинов. «Теорема Погорелова — Клингенберга для многообразий, гомеоморфных $\mathbb{R}^n$». *Сибирский математический журнал* 18.4 (1977), с. 915—925.

[57] Jürgen Jost. "Nonlinear Dirichlet forms". *New directions in Dirichlet forms.* Vol. 8. AMS/IP Stud. Adv. Math. Amer. Math. Soc., Providence, RI, 1998, pp. 1–47.

[58] Uwe F. Mayer. "Gradient flows on nonpositively curved metric spaces and harmonic maps". *Comm. Anal. Geom.* 6.2 (1998), pp. 199–253.

[59] Giuseppe Savaré. "Gradient flows and diffusion semigroups in metric spaces under lower curvature bounds". *C. R. Math. Acad. Sci. Paris* 345.3 (2007), pp. 151–154.

[60] Aris Daniilidis, Olivier Ley, and Stéphane Sabourau. "Asymptotic behaviour of self-contracted planar curves and gradient orbits of convex functions". *J. Math. Pures Appl. (9)* 94.2 (2010), pp. 183–199.

[61] Stefan Cohn-Vossen. „Totalkrümmung und geodätische Linien auf einfachzusammenhängenden offenen vollständigen Flächenstücken". *Mat. Sb.* 1(43).2 (1936).

[62] А. Д. Милка. «Метрическое строение одного класса пространств, содержащих прямые линии.«». *Укр. геометр, сб.* 4 (1967), с. 43—48.

[63] Jeff Cheeger and Detlef Gromoll. "The splitting theorem for manifolds of nonnegative Ricci curvature". *J. Differential Geometry* 6 (1971/72), pp. 119–128.

[64] J.-H. Eschenburg. "The splitting theorem for space-times with strong energy condition". *J. Differential Geom.* 27.3 (1988), pp. 477–491.

[65] R. E. Greene and H. Wu. "On the subharmonicity and plurisubharmonicity of geodesically convex functions". *Indiana Univ. Math. J.* 22 (1972/73), pp. 641–653.

[66] Chanyoung Jun. "Pursuit-evasion and time-dependent gradient flow in singular spaces". PhD thesis. University of Illinois at Urbana-Champaign, 2012.

[67] C. Jun. *Time-dependent gradient curves on CAT(0) spaces.* arXiv: `1801.04583 [math.MG]`.

[68] Anton Petrunin. "Alexandrov meets Lott-Villani-Sturm". *Münster J. Math.* 4 (2011), pp. 53–64.

[69] Cédric Villani. *Optimal transport.* Vol. 338. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Old and new. Springer-Verlag, Berlin, 2009, pp. xxii+973.