

An invitation to Alexandrov geometry:
spaces with curvature bounded below

Vitali Kapovitch and Anton Petrunin
with with an appendix by
Nina Lebedeva and Anton Petrunin

Contents

1 Preliminaries	1
A. Prerequisite 1 ; B. Notations 1 ; C. Length spaces 2 ; D. Geodesics 3 ; E. Menger's lemma 3 ; F. Triangles and model tangles 4 ; G. Hinges and their angle measure 5 ; H. Triangle inequality for angles 5 .	
2 Definitions	7
A. Four-point comparison 7 ; B. Alexandrov's lemma 8 ; C. Hinge comparison 10 ; D. Equivalent conditions 11 ; E. Thin and fat triangles 11 ; F. Function comparison 12 ; G. Comments 13 .	
3 Globalization	15
A. Globalization 15 ; B. On general curvature bound 18 ; C. Remarks 19 .	
4 Calculus	21
A. Space of directions 21 ; B. Tangent space 22 ; C. Semiconcave functions 23 ; D. Differential 23 ; E. Gradient 24 .	
5 Gradient flow	29
A. Velocity of curve 29 ; B. Gradient curves 30 ; C. Distance estimates 31 ; D. Gradient flow 33 ; E. Gradient exponent 34 ; F. Remarks 35 .	
6 Line splitting	37
A. Busemann function 37 ; B. Splitting theorem 38 ; C. Comments 40 .	
7 Dimension	41
A. Polar vectors 41 ; B. Linear subspace of tangent space 43 ; C. Linear dimension 45 ; D. Space of directions 46 ; E. Comments 48 .	
8 Limit spaces	49
A. Hausdorff convergence 49 ; B. Gromov–Hausdorff convergence 50 ; C. Survival of curvature bounds 51 ; D. Volume 52 ; E. Gromov's selection theorem 53 ; F. Comments 55 .	

9 Homotopy finiteness theorem	57
A. Controlled concavity 57 ; B. Liftings 58 ; C. Nerves 59 ; D. Homotopy stability 60 ; E. Comments 61 .	
10 Boundary	63
A. Definition 63 ; B. Conic neighborhoods 63 ; C. Topology 64 ; D. Tangent space 66 ; E. Doubling 66 ; F. Remarks 70 .	
11 Quotients	73
A. Quotient space 73 ; B. Generalizations 74 ; C. Hopf's conjecture 75 ; D. Erdős' problem rediscovered 77 ; E. Crystallographic actions 78 ; F. Remarks 79 .	
12 Surface of convex body	81
A. Surface of convex polyhedra 81 ; B. Curvature 83 ; C. Surface of convex body 84 ; D. Comments 84 .	
A Alexandrov's embedding theorem	85
A. Introduction 85 ; B. Space of polyhedrons and metrics 86 ; C. About the proof 87 .	
Semisolutions	91
Bibliography	107

Preface

As in our previous invitation [2], we try to demonstrate the beauty and power of Alexandrov geometry by reaching interesting applications and theorems with a minimum of preparation. This time we do spaces with curvature bounded below in the sense of Alexandrov.

This subject is more technical, this time we jumped over proofs of couple of technical results, namely existence part in generalized Picard's theorem (5.3) and Perelman's theorem about conic neighborhoods (10.3). The rest of our presentation is nearly rigorous.

In Lecture 1, we discuss necessary preliminaries and fix notations.

Lecture 2 introduces the main object of our study — spaces with curvature bounded below in the sense of Alexandrov.

In Lecture 3 we formulate and prove the globalization theorem — local Alexandrov condition implies global. To simplify the presentation we consider only compact case, but this case is leading.

In Lecture 4 we do beginning of calculus — tangent space and space of directions, differential, and gradient.

Lecture 5 introduces gradient flow — this is the main technical tool in the theory.

Lecture 6 proves the line splitting theorem. It provides the first application of gradient flow.

In Lecture 7 we introduce and discuss dimension of Alexandrov spaces.

Lecture 8 shows that lower curvature bound survives in the Gromov–Hausdorff limit. Further, we introduce volume, prove the Bishop–Gromov inequality and use it to prove Gromov's selection theorem.

Lecture 9 starts with Perelman's construction of concave functions. Further, we apply it with Gromov's selection theorem to prove the homotopy finiteness theorem. This proof illustrates the main source of applications of Alexandrov geometry.

In Lecture 10 we introduce boundary of finite-dimensional Alexandrov space and prove the doubling theorem.

Lecture 11 we show that quotient Alexandrov space by isometric group action is an Alexandrov space and give several applications of this statement. These proofs illustrate another source of applications of Alexandrov geometry.

In Lecture 12 we show that surface of a convex body in Euclidean space is an Alexandrov space. This is historically the first serious application of Alexandrov geometry.

Finally, Appendix A sketches Alexandrov embedding theorem of convex polyhedra. Historically, this theorem is the first remarkable result in Alexandrov geometry that dates back to 1941. The proof is very well written by Alexandrov, but we decided to include its sketch here due to its beauty and importance. This appendix was written by Nina Lebedeva and the second author for a book about .

Let us give a list of available texts on Alexandrov spaces with curvature bounded below:

- ◊ The first introduction to Alexandrov geometry is given in the original paper of Yuriy Burago, Michael Gromov, and Grigory Perelman [11] and its extension [45] written by Perelman.
- ◊ A brief and reader-friendly introduction was written by Katsuhiko Shiohama [59, Sections 1–8].
- ◊ Another reader-friendly introduction, written by Dmitri Burago, Yuriy Burago, and Sergei Ivanov [10, Chapter 10].
- ◊ Survey by Conrad Plaut [57].
- ◊ Survey by the second author [50].

Acknowledgments. Our notes were shaped in a number of lectures given by the authors at different occasions in Penn State, including the MASS program, at the Summer School “Algebra and Geometry” in Yaroslavl, at SPbSU, and University of Toronto. We want to thank these institution for hospitality and support.

We were partially supported by the following grants: Vitali Kapovitch — NSERC Discovery grants; Anton Petrunin — NSF grant DMS-2005279.

Lecture 1

Preliminaries

A Prerequisite

We assume that the reader is familiar with the following topics in metric geometry:

- ◇ Compactness and proper metric spaces; recall that a metric space is proper if all its closed balls (with finite radius) are compact.
- ◇ Complete metric spaces and completion.
- ◇ Curves, semicontinuity of length and rectifiability.
- ◇ Hausdorff and Gromov–Hausdorff convergence. These are defined in Lecture 8, but it is better to have prior acquaintance with these convergences.

These topics are treated in [10] and [54]. Occasionally, we use some topology; any introductory text in algebraic topology should be sufficient.

B Notations

The distance between two points x and y in a metric space \mathcal{X} will be denoted by $|x - y|$ or $|x - y|_{\mathcal{X}}$. The latter notation is used if we need to emphasize that the distance is taken in the space \mathcal{X} .

Given radius $r \in [0, \infty]$ and center $x \in \mathcal{X}$, the sets

$$\begin{aligned} B(x, r) &= \{ y \in \mathcal{X} : |x - y| < r \}, \\ \overline{B}[x, r] &= \{ y \in \mathcal{X} : |x - y| \leq r \} \end{aligned}$$

are called, respectively, the open and the closed balls. The notations $B(x, r)_{\mathcal{X}}$ and $\overline{B}[x, r]_{\mathcal{X}}$ might be used if we need to emphasize that these balls are taken in the metric space \mathcal{X} .

We will denote by \mathbb{S}^n , \mathbb{E}^n , and \mathbb{H}^n the n -dimensional sphere (with angle metric), Euclidean space, and Lobachevsky space respectively. More generally, $\mathbb{M}^n(\kappa)$ will denote the model n -space of curvature κ ; that is,

- ◇ if $\kappa > 0$, then $\mathbb{M}^n(\kappa)$ is the n -sphere of radius $\frac{1}{\sqrt{\kappa}}$, so $\mathbb{S}^n = \mathbb{M}^n(1)$
- ◇ $\mathbb{M}^n(0) = \mathbb{E}^n$,
- ◇ if $\kappa < 0$, then $\mathbb{M}^n(\kappa)$ is the Lobachevsky n -space \mathbb{H}^n rescaled by factor $\frac{1}{\sqrt{-\kappa}}$; in particular $\mathbb{M}^n(-1) = \mathbb{H}^n$.

C Length spaces

Let \mathcal{X} be a metric space. If for any $\varepsilon > 0$ and any pair of points $x, y \in \mathcal{X}$, there is a path α connecting x to y such that

$$\text{length } \alpha < |x - y| + \varepsilon,$$

then \mathcal{X} is called a length space and the metric on \mathcal{X} is called a length metric.

1.1. Exercise. *Let \mathcal{X} be a complete length space. Show that for any compact subset $K \subset \mathcal{X}$ there is a compact path-connected subset $K' \subset \mathcal{X}$ that contains K .*

Induced length metric. Directly from the definition, it follows that if $\alpha: [0, 1] \rightarrow \mathcal{X}$ is a path from x to y (that is, $\alpha(0) = x$ and $\alpha(1) = y$), then

$$\text{length } \alpha \geq |x - y|.$$

Set

$$\|x - y\| = \inf \{ \text{length } \alpha \}$$

where the greatest lower bound is taken for all paths from x to y . It is straightforward to check that $(x, y) \mapsto \|x - y\|$ is an ∞ -metric; that is, $(x, y) \mapsto \|x - y\|$ is a metric in the extended positive reals $[0, \infty]$. The metric $\|* - *\|$ is called the induced length metric.

1.2. Exercise. *Suppose $(\mathcal{X}, |* - *|)$ is a complete metric space. Show that $(\mathcal{X}, \|* - *\|)$ is complete; that is, any Cauchy sequence of points in $(\mathcal{X}, \|* - *\|)$ converges in $(\mathcal{X}, \|* - *\|)$.*

Let A be a subset of a metric space \mathcal{X} . Given two points $x, y \in A$, consider the value

$$|x - y|_A = \inf_{\alpha} \{ \text{length } \alpha \},$$

where the greatest lower bound is taken for all paths α from x to y in A . In other words, $|* - *|_A$ denotes the induced length metric on

the subspace A . (The notation $|\ast - \ast|_A$ conflicts with the previously defined notation for distance $|x - y|_{\mathcal{X}}$ in a metric space \mathcal{X} . However, most of the time we will work with ambient length spaces where the meaning will be unambiguous.)

D Geodesics

Let \mathcal{X} be a metric space and \mathbb{I} a real interval. A distance-preserving map $\gamma: \mathbb{I} \rightarrow \mathcal{X}$ is called a geodesic¹; in other words, $\gamma: \mathbb{I} \rightarrow \mathcal{X}$ is a geodesic if

$$|\gamma(s) - \gamma(t)|_{\mathcal{X}} = |s - t|$$

for any pair $s, t \in \mathbb{I}$.

If $\gamma: [a, b] \rightarrow \mathcal{X}$ is a geodesic such that $p = \gamma(a)$, $q = \gamma(b)$, then we say that γ is a geodesic from p to q . In this case, the image of γ is denoted by $[pq]$, and, with abuse of notations, we also call it a geodesic. We may write $[pq]_{\mathcal{X}}$ to emphasize that the geodesic $[pq]$ is in the space \mathcal{X} .

In general, a geodesic from p to q need not exist and if it exists, it need not be unique. However, once we write $[pq]$ we assume that we have chosen such geodesic.

A geodesic path is a geodesic with constant-speed parameterization by the unit interval $[0, 1]$.

A metric space is called geodesic if any pair of its points can be joined by a geodesic.

Evidently, any geodesic space is a length space.

1.3. Exercise. *Show that any proper length space is geodesic.*

E Menger's lemma

1.4. Lemma. *Let \mathcal{X} be a complete metric space. Assume that for any pair of points $x, y \in \mathcal{X}$, there is a midpoint z . Then \mathcal{X} is a geodesic space.*

This lemma is due to Karl Menger [40, Section 6].

Proof. Choose $x, y \in \mathcal{X}$; set $\alpha(0) = x$, and $\alpha(1) = y$.

Let $\alpha(\frac{1}{2})$ be a midpoint between $\alpha(0)$ and $\alpha(1)$. Further, let $\alpha(\frac{1}{4})$ and $\alpha(\frac{3}{4})$ be midpoints between the pairs $(\alpha(0), \alpha(\frac{1}{2}))$ and $(\alpha(\frac{1}{2}), \alpha(1))$ respectively. Applying the above procedure recursively, on the n -th

¹Others call it differently: *shortest path*, *minimizing geodesic*. Also, note that the meaning of the term *geodesic* is different from what is used in Riemannian geometry, altho they are closely related.

step we define $\alpha(\frac{k}{2^n})$, for every odd integer k such that $0 < \frac{k}{2^n} < 1$, as a midpoint of the already defined $\alpha(\frac{k-1}{2^n})$ and $\alpha(\frac{k+1}{2^n})$.

This way we define $\alpha(t)$ for all dyadic rationals t in $[0, 1]$. Moreover, α has Lipschitz constant $|x - y|$. Since \mathcal{X} is complete, the map α can be extended continuously to $[0, 1]$. Moreover,

$$\text{length } \alpha \leq |x - y|.$$

□

1.5. Exercise. *Let \mathcal{X} be a complete metric space. Assume that for any pair of points $x, y \in \mathcal{X}$, there is an almost midpoint; that is, given $\varepsilon > 0$, there is a point z such that*

$$|x - z| < \frac{1}{2} \cdot |x - y| + \varepsilon \quad \text{and} \quad |y - z| < \frac{1}{2} \cdot |x - y| + \varepsilon.$$

Show that \mathcal{X} is a length space.

F Triangles and model tangles

Triangles. Given a triple of points p, q, r in a metric space \mathcal{X} , a choice of geodesics $([qr], [rp], [pq])$ will be called a triangle; we will use the short notation $[pqr] = [pqr]_{\mathcal{X}} = ([qr], [rp], [pq])$.

Given a triple $p, q, r \in \mathcal{X}$ there may be no triangle $[pqr]$ simply because one of the pairs of these points cannot be joined by a geodesic. Also, many different triangles with these vertices may exist, any of which can be denoted by $[pqr]$. If we write $[pqr]$, it means that we have chosen such a triangle.

Model triangles. Given three points p, q, r in a metric space \mathcal{X} , let us define its model triangle $[\tilde{p}\tilde{q}\tilde{r}]$ (briefly, $[\tilde{p}\tilde{q}\tilde{r}] = \tilde{\Delta}(pqr)_{\mathbb{E}^2}$) to be a triangle in the Euclidean plane \mathbb{E}^2 such that

$$|\tilde{p} - \tilde{q}|_{\mathbb{E}^2} = |p - q|_{\mathcal{X}}, \quad |\tilde{q} - \tilde{r}|_{\mathbb{E}^2} = |q - r|_{\mathcal{X}}, \quad |\tilde{r} - \tilde{p}|_{\mathbb{E}^2} = |r - p|_{\mathcal{X}}.$$

The same way we can define the hyperbolic and the spherical model triangles $\tilde{\Delta}(pqr)_{\mathbb{H}^2}$, $\tilde{\Delta}(pqr)_{\mathbb{S}^2}$ in the Lobachevsky plane \mathbb{H}^2 and the unit sphere \mathbb{S}^2 . In the latter case, the model triangle is said to be defined if in addition

$$|p - q| + |q - r| + |r - p| < 2 \cdot \pi.$$

In this case, the model triangle again exists and is unique up to an isometry of \mathbb{S}^2 .

Model angles. If $[\tilde{p}\tilde{q}\tilde{r}] = \tilde{\Delta}(pqr)_{\mathbb{E}^2}$ and $|p - q|, |p - r| > 0$, the angle measure of $[\tilde{p}\tilde{q}\tilde{r}]$ at \tilde{p} will be called the model angle of the triple p, q, r and will be denoted by $\tilde{\angle}(p_r^q)_{\mathbb{E}^2}$.

The same way we define $\tilde{\angle}(p_r^q)_{\mathbb{M}^2(\kappa)}$; in particular, $\tilde{\angle}(p_r^q)_{\mathbb{H}^2}$ and $\tilde{\angle}(p_r^q)_{\mathbb{S}^2}$. We may use the notation $\tilde{\angle}(p_r^q)$ if it is evident which of the model spaces is meant.

1.6. Exercise. Show that for any triple of point p, q , and r , the function

$$\kappa \mapsto \tilde{\angle}(p_r^q)_{\mathbb{M}^2(\kappa)}$$

is nondecreasing in its domain of definition.

G Hinges and their angle measure

Hinges. Let $p, x, y \in \mathcal{X}$ be a triple of points such that p is distinct from x and y . A pair of geodesics $([px], [py])$ will be called a hinge and will be denoted by $[p_y^x] = ([px], [py])$.

Angles. The angle measure of a hinge $[p_y^x]$ is defined as the following limit

$$\angle[p_y^x] = \lim_{\bar{x}, \bar{y} \rightarrow p} \tilde{\angle}(p_{\bar{y}}^{\bar{x}}),$$

where $\bar{x} \in [px]$ and $\bar{y} \in [py]$.

Note that if $\angle[p_y^x]$ is defined, then

$$0 \leq \angle[p_y^x] \leq \pi.$$

1.7. Exercise. Suppose that in the above definition, one uses spherical or hyperbolic model angles instead of Euclidean. Show that it does not change the value $\angle[p_y^x]$.

1.8. Exercise. Give an example of a hinge $[p_y^x]$ in a metric space with an undefined angle measure $\angle[p_y^x]$.


H Triangle inequality for angles

1.9. Proposition. Let $[px_1], [px_2]$, and $[px_3]$ be three geodesics in a metric space. Suppose all the angle measures $\alpha_{ij} = \angle[p_{x_j}^{x_i}]$ are defined. Then

$$\alpha_{13} \leq \alpha_{12} + \alpha_{23}.$$

Proof. Since $\alpha_{13} \leq \pi$, we can assume that $\alpha_{12} + \alpha_{23} < \pi$. Denote by γ_i the unit-speed parametrization of $[px_i]$ from p to x_i . Given any $\varepsilon > 0$, for all sufficiently small $t, \tau, s \in \mathbb{R}_{\geq 0}$ we have

$$\begin{aligned} |\gamma_1(t) - \gamma_3(\tau)| &\leq |\gamma_1(t) - \gamma_2(s)| + |\gamma_2(s) - \gamma_3(\tau)| < \\ &< \sqrt{t^2 + s^2 - 2 \cdot t \cdot s \cdot \cos(\alpha_{12} + \varepsilon)} + \\ &\quad + \sqrt{s^2 + \tau^2 - 2 \cdot s \cdot \tau \cdot \cos(\alpha_{23} + \varepsilon)} \leq \end{aligned}$$



Below we define $s(t, \tau)$ so that for $s = s(t, \tau)$, this chain of inequalities can be continued as follows:

$$\leq \sqrt{t^2 + \tau^2 - 2 \cdot t \cdot \tau \cdot \cos(\alpha_{12} + \alpha_{23} + 2 \cdot \varepsilon)}.$$

Thus for any $\varepsilon > 0$,

$$\alpha_{13} \leq \alpha_{12} + \alpha_{23} + 2 \cdot \varepsilon.$$

Hence the result follows.

To define $s(t, \tau)$, consider three half-lines $\tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3$ on a Euclidean plane starting at one point, such that $\angle(\tilde{\gamma}_1, \tilde{\gamma}_2) = \alpha_{12} + \varepsilon$, $\angle(\tilde{\gamma}_2, \tilde{\gamma}_3) = \alpha_{23} + \varepsilon$, and $\angle(\tilde{\gamma}_1, \tilde{\gamma}_3) = \alpha_{12} + \alpha_{23} + 2 \cdot \varepsilon$. We parametrize each half-line by the distance from the starting point. Given two positive numbers $t, \tau \in \mathbb{R}_{\geq 0}$, let $s = s(t, \tau)$ be the number such that $\tilde{\gamma}_2(s) \in [\tilde{\gamma}_1(t), \tilde{\gamma}_3(\tau)]$. Clearly, $s \leq \max\{t, \tau\}$, so t, τ, s may be taken sufficiently small. \square

1.10. Exercise. Prove that the sum of adjacent angles is at least π .

More precisely: suppose two hinges $[p_z^x]$ and $[p_z^y]$ are adjacent; that is, they share side $[pz]$, and the union of two sides $[px]$ and $[py]$ form a geodesic $[xy]$. Show that

$$\angle[p_z^x] + \angle[p_z^y] \geq \pi$$

whenever each angle on the left-hand side is defined.

Give an example showing that the inequality can be strict.

1.11. Exercise. Assume that the angle measure of $[q_x^p]$ is defined. Let γ be the unit speed parametrization of $[qx]$ from q to x . Show that

$$|p - \gamma(t)| \leq |q - p| - t \cdot \cos(\angle[q_x^p]) + o(t).$$

Lecture 2

Definitions

A Four-point comparison

Let p, x, y, z be a quadruple of points in a metric space. If the inequality

$$\mathbf{1} \quad \tilde{\angle}(p_y^x)_{\mathbb{E}^2} + \tilde{\angle}(p_z^y)_{\mathbb{E}^2} + \tilde{\angle}(p_x^z)_{\mathbb{E}^2} \leq 2 \cdot \pi$$

holds, then we say that the quadruple meets CBB(0) comparison.

2.1. Definition. A metric space \mathcal{X} has nonnegative curvature in the sense of Alexandrov (briefly, $\mathcal{X} \in \text{CBB}(0)$) if CBB(0) comparison holds for any quadruple in \mathcal{X} such that each model angle in **1** is defined.

If instead of \mathbb{E}^2 , we use \mathbb{S}^2 or \mathbb{H}^2 , then we get the definition of CBB(1) and CBB(-1) comparisons. Note that $\tilde{\angle}(p_y^x)_{\mathbb{E}^2}$ and $\tilde{\angle}(p_y^x)_{\mathbb{H}^2}$ are defined if $p \neq x, p \neq y$, but for $\tilde{\angle}(p_y^x)_{\mathbb{S}^2}$ we require in addition that

$$|p - x| + |p - y| + |x - y| < 2 \cdot \pi;$$

if this inequality does not hold, then we assume that CBB(1) comparison holds for this quadruple.

More generally, one may apply this definition to $\mathbb{M}^2(\kappa)$. This way we define CBB(κ) comparison for any real κ . However, if you see CBB(κ) comparison, it is usually safe to assume that $\kappa = -1, 0$, or 1 (applying rescaling, the CBB(κ) comparison can be reduced to these three cases).

2.2. Exercise. Show that \mathbb{E}^n is CBB(0).

2.3. Exercise. Show that a metric space \mathcal{X} is CBB(0) if and only if for any quadruple of points $p, x_1, x_2, x_3 \in \mathcal{X}$ there is a quadruple of points $q, y_1, y_2, y_3 \in \mathbb{E}^3$ such that

$$|p - x_i|_{\mathcal{X}} \geq |q - y_i|_{\mathbb{E}^2} \quad \text{and} \quad |x_i - x_j|_{\mathcal{X}} \leq |y_i - y_j|_{\mathbb{E}^2}$$

for all i and j .

B Alexandrov's lemma

Recall that $[xy]$ denotes a geodesic from x to y ; set


$$]xy[= [xy] \setminus \{x\}, \quad]xy[= [xy] \setminus \{y\}, \quad]xy[= [xy] \setminus \{x, y\}.$$

2.4. Lemma. Let p, x, y, z be distinct points in a metric space such that $z \in]xy[$. Then the following expressions have the same sign:

- (a) $\tilde{\angle}(x_y^p) - \tilde{\angle}(x_z^p)$,
- (b) $\tilde{\angle}(z_x^p) + \tilde{\angle}(z_y^p) - \pi$.

The same holds for the hyperbolic and spherical model angles, but in the latter case, one has to assume in addition that

$$|p - z| + |p - y| + |x - y| < 2 \cdot \pi.$$



Proof. Consider the model triangle $[\tilde{x}\tilde{p}\tilde{z}] = \tilde{\Delta}(xpz)$. Take a point \tilde{y} on the extension of $[\tilde{x}\tilde{z}]$ beyond \tilde{z} so that $|\tilde{x} - \tilde{y}| = |x - y|$ (and therefore $|\tilde{x} - \tilde{z}| = |x - z|$).

Since increasing the opposite side in a plane triangle increases the corresponding angle, the following expressions have the same sign:

- (i) $\angle[\tilde{x}\tilde{p}\tilde{y}] - \tilde{\angle}(x_y^p)$,
- (ii) $|\tilde{p} - \tilde{y}| - |p - y|$,
- (iii) $\angle[\tilde{z}\tilde{p}\tilde{y}] - \tilde{\angle}(z_y^p)$.

Since


$$\angle[\tilde{x}\tilde{p}\tilde{y}] = \angle[\tilde{x}\tilde{p}\tilde{z}] = \tilde{\angle}(x_z^p)$$

and

$$\angle[\tilde{z}\tilde{p}\tilde{y}] = \pi - \angle[\tilde{z}\tilde{p}\tilde{x}] = \pi - \tilde{\angle}(z_x^p),$$

the statement follows.

The spherical and hyperbolic cases can be proved in the same way. \square



2.5. Exercise. Assume p, x, y, z are as in Alexandrov's lemma. Show that

$$\tilde{\angle}(p_y^x) \geq \tilde{\angle}(p_z^x) + \tilde{\angle}(p_y^z),$$

with equality if and only if the expressions in (a) and (b) vanish.

Note that

$$p \in]xy[\implies \tilde{\angle}(p_y^x) = \pi.$$

Applying it with Alexandrov's lemma and CBB(0) comparison, we get the following.


2.6. Claim. If p, x, y, z are points in a CBB(0) such that $p \in]xy[$, then

$$\tilde{\angle}(x_z^y) \leq \tilde{\angle}(x_z^p).$$

2.7. Exercise. Let $[p_y^x]$ be a hinge in a CBB(0) space. Consider the function

$$f: (|p - \bar{x}|, |p - \bar{y}|) \mapsto \tilde{\angle}(p_{\bar{y}}^{\bar{x}}),$$

where $\bar{x} \in]px]$ and $\bar{y} \in]py]$. Show that f is nonincreasing in each argument.



Note that 2.7 implies the following.

2.8. Claim. For any hinge $[p_y^x]$ in a CBB(0) space, the angle measure $\angle[p_y^x]$ is defined, and

$$\angle[p_y^x] \geq \tilde{\angle}(p_y^x).$$

2.9. Exercise. Let $[p_y^x]$ be a hinge in a CBB(0) space. Suppose $\angle[p_y^x] = 0$; show that $[px] \subset [py]$ or $[py] \subset [px]$.

2.10. Exercise. Let $[xy]$ be a geodesic in a CBB(0) space. Suppose $z \in]xy[$ show that there is a unique geodesic $[xz]$ and $[xz] \subset [xy]$.

Recall that adjacent hinges are defined in 1.10.

2.11. Exercise. Let $[p_z^x]$ and $[p_z^y]$ be adjacent hinges in a CBB(0) space. Show that

$$\angle[p_z^x] + \angle[p_z^y] = \pi.$$

2.12. Exercise. Show that

$$\tilde{\angle}(x_p^y) = \tilde{\angle}(x_p^v) \iff \tilde{\angle}(x_p^y) = \tilde{\angle}(x_p^w)$$

if p, x, y, v, w are points in a CBB(0) space such that $v, w \in]xy[$.

2.13. Exercise. Suppose hinges $[x_n y_n z_n]$ in a CBB(0) space converge to the hinge $[x_\infty y_\infty z_\infty]$; that is geodesics $[x_n y_n]$ and $[x_n z_n]$ converge to the geodesics $[x_\infty y_\infty]$ and $[x_\infty z_\infty]$ in the Hausdorff sense. Show that

$$\lim_{n \rightarrow \infty} \angle[x_n y_n z_n] \geq \angle[x_\infty y_\infty z_\infty].$$

C Hinge comparison

Let $[p_y^x]$ be a hinge in a CBB(0) space. By 2.9, the angle measure $\angle[p_y^x]$ is defined and

$$\angle[p_y^x] \geq \tilde{\angle}(p_y^x).$$

Further, according to 2.11, we have

$$\angle[p_z^x] + \angle[p_z^y] = \pi$$

for adjacent hinges $[p_z^x]$ and $[p_z^y]$ in a CBB(0) space.

The following theorem implies that a geodesic space is CBB(0) if the above conditions hold for all its hinges.

2.14. Theorem. A geodesic space \mathcal{L} is CBB(0) if the following conditions hold.

(a) For any hinge $[x_y^p]$ in \mathcal{L} , the angle $\angle[x_y^p]$ is defined and


$$\angle[x_y^p] \geq \tilde{\angle}(x_y^p).$$

(b) For any two adjacent hinges $[p_z^x]$ and $[p_z^y]$ in \mathcal{L} , we have

$$\angle[p_z^x] + \angle[p_z^y] \leq \pi.$$

Proof. Consider a point $w \in]pz[$ close to p . From (b), it follows that

$$\angle[w_z^x] + \angle[w_z^p] \leq \pi \quad \text{and} \quad \angle[w_z^y] + \angle[w_z^p] \leq \pi.$$



Since $\angle[w_y^x] \leq \angle[w_z^x] + \angle[w_z^y]$ (see 1.9), we get

$$\angle[w_z^x] + \angle[w_z^y] + \angle[w_z^p] \leq 2 \cdot \pi.$$

Applying (a),

$$\tilde{\angle}(w_z^x) + \tilde{\angle}(w_z^y) + \tilde{\angle}(w_z^p) \leq 2 \cdot \pi.$$

Passing to the limits $w \rightarrow p$, we have

$$\tilde{\angle}(p_z^x) + \tilde{\angle}(p_z^y) + \tilde{\angle}(p_z^p) \leq 2 \cdot \pi.$$

□

D Equivalent conditions

The following theorem summarizes 2.6, 2.8, 2.11, 2.14.

2.15. Theorem. *Let \mathcal{L} be a geodesic space. Then the following conditions are equivalent.*

- (a) \mathcal{L} is CBB(0).
- (b) (adjacent angle comparison) for any geodesic $[xy]$ and point $z \in]xy[, z \neq p$ in \mathcal{L} , we have

$$\tilde{\Delta}(z_x^p) + \tilde{\Delta}(z_y^p) \leq \pi.$$

- (c) (point-on-side comparison) for any geodesic $[xy]$ and $z \in]xy[$ in \mathcal{L} , we have

$$\tilde{\Delta}(x_y^p) \leq \tilde{\Delta}(x_z^p).$$

- (d) (hinge comparison) for any hinge $[x_y^p]$ in \mathcal{L} , the angle $\Delta[x_y^p]$ is defined and

$$\Delta[x_y^p] \geq \tilde{\Delta}(x_y^p).$$

Moreover,

$$\Delta[z_y^p] + \Delta[z_x^p] \leq \pi$$

for any adjacent hinges $[z_y^p]$ and $[z_x^p]$.

Moreover, the implications $(a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d)$ hold in any space, not necessarily geodesic.

2.16. Advanced Exercise. *Construct a geodesic space \mathcal{X} that is not CBB(0) and satisfies the following condition: for any three points $p, x, y \in \mathcal{X}$ there is a geodesic $[xy]$ such that for any $z \in]xy[$*

$$\tilde{\Delta}(z_x^p) + \tilde{\Delta}(z_y^p) \leq \pi.$$

E Thin and fat triangles

Consider the model triangle $[\tilde{x}\tilde{y}\tilde{z}] = \tilde{\Delta}(xyz)_{\mathbb{E}^2}$ in the Euclidean plane. The natural map $[\tilde{x}\tilde{y}\tilde{z}] \rightarrow [xyz]$ sends a point $\tilde{p} \in [\tilde{x}\tilde{y}] \cup [\tilde{y}\tilde{z}] \cup [\tilde{z}\tilde{x}]$ to the corresponding point $p \in [xy] \cup [yz] \cup [zx]$; for instance, if \tilde{p} lies on $[\tilde{y}\tilde{z}]$, then $p \in [yz]$ and $|\tilde{y} - \tilde{p}| = |y - p|$ (and therefore $|\tilde{z} - \tilde{p}| = |z - p|$).

2.17. Definition. *A triangle $[xyz]$ in a metric space is called thin (or fat) if the natural map $\tilde{\Delta}(xyz)_{\mathbb{E}^2} \rightarrow [xyz]$ is distance nonincreasing (or respectively distance nondecreasing).¹*

¹V: why do we need to define thin triangles in this book? I think they are only used in Exercise 2.19

2.18. Exercise. Show that any triangle in a CBB(0) space is fat.

2.19. Exercise. Let \mathcal{W} be \mathbb{R}^2 with the metric induced by a norm. Suppose that

(a) Any triangle in \mathcal{W} is thin.

(b) Any triangle in \mathcal{W} is fat.

Show that \mathcal{W} is isometric to the Euclidean plane \mathbb{E}^2 .

F Function comparison

Real-to-real functions. Choose $\lambda \in \mathbb{R}$. Let $s: \mathbb{I} \rightarrow \mathbb{R}$ be a locally Lipschitz function defined on an interval \mathbb{I} . The following statement are equivalent; if one (and therefore any) of them hold for s , then we say that s is λ -concave.

- ◊ We have inequality $s'' \leq \lambda$, where the second derivative s'' is understood in the sense of distributions.
- ◊ The function $t \mapsto s(t) - \lambda \cdot \frac{t^2}{2}$ is concave.
- ◊ The Jensen inequality

$$s(a \cdot t_0 + (1-a) \cdot t_1) \geq a \cdot s(t_0) + (1-a) \cdot s(t_1) + \frac{\lambda}{2} \cdot a \cdot (1-a) \cdot (t_1 - t_0)^2$$

holds for any $t_0, t_1 \in \mathbb{I}$ and $a \in [0, 1]$.

- ◊ for any $t_0 \in \mathbb{I}$ there is a quadratic polynomial $\ell = \frac{\lambda}{2} \cdot t^2 + a \cdot t + b$ that (locally) supports s at t_0 from above; that is, $\ell(t_0) = s(t_0)$ and $\ell(t) \geq s(t)$ for any t (in a neighborhood of t_0)

The equivalence of these definitions is assumed to be known. ²

We will also use that λ -concave functions are one-side differentiable.

Functions on metric space. A function on a metric space \mathcal{L} will usually mean a *locally Lipschitz real-valued function defined on an open subset of \mathcal{L}* . The domain of definition of a function f will be denoted by $\text{Dom } f$.

Let f be a function on a metric space \mathcal{L} . We say that f is λ -concave (briefly $f'' \leq \lambda$) if for any unit-speed geodesic $\gamma: \mathbb{I} \rightarrow \text{Dom } f$ the real-to-real function $t \mapsto f \circ \gamma(t)$ is λ -concave.

The following proposition is simple, but conceptual — it reformulates a global geometric condition into an infinitesimal condition on distance functions.

²V: I was asked by several people for references to this.

2.20. Proposition. *A geodesic space \mathcal{L} is CBB(0) if and only if $f'' \leq 1$ for any function f of the following type*

$$f: x \mapsto \frac{1}{2} \cdot |p - x|^2.$$

Proof. Choose a unit-speed geodesic γ in \mathcal{L} and two points $x = \gamma(t_0)$, $y = \gamma(t_1)$ for some $t_0 < t_1$. Consider the model triangle $[\tilde{p}\tilde{x}\tilde{y}] = \tilde{\Delta}(pxy)$. Let $\tilde{\gamma}: [t_0, t_1] \rightarrow \mathbb{E}^2$ be the unit-speed parametrization of $[\tilde{x}\tilde{y}]$ from \tilde{x} to \tilde{y} .

Set

$$\tilde{r}(t) := |\tilde{p} - \tilde{\gamma}(t)|, \quad r(t) := |p - \gamma(t)|.$$

Clearly, $\tilde{r}(t_0) = r(t_0)$ and $\tilde{r}(t_1) = r(t_1)$. Note that the point-on-side comparison (2.15c) is equivalent to

$$\textbf{1} \quad t_0 \leq t \leq t_1 \quad \implies \quad \tilde{r}(t) \leq r(t)$$

for any γ and $t_0 < t_1$.

Set

$$\tilde{h}(t) = \frac{1}{2} \cdot \tilde{r}^2(t) - \frac{1}{2} \cdot t^2, \quad h = \frac{1}{2} \cdot r^2(t) - \frac{1}{2} \cdot t^2.$$

Note that \tilde{h} is linear, $\tilde{h}(t_0) = h(t_0)$ and $\tilde{h}(t_1) = h(t_1)$. Observe that the Jensen inequality for the function h is equivalent to **1**. Hence the proposition follows. \square

G Comments

The CBB comparison is closely related to the so-called CAT comparison, which defines *upper* curvature bound in the sense of Alexandrov; this is the subject of our previous invitation [2].

In both comparisons we check certain condition on 6 distances between every pair of a 4-point subset. Michael Gromov [20, Section 1.19+] suggested to consider other conditions of that type for n -point subsets; see [18, 21, 33–38, 53, 62] for development of this idea.

All the discussed statements admit natural generalizations to CBB(κ) spaces. The proof are nearly the same, but the formulas are getting more complicated. It is common practice in Alexandrov geometry to write proofs in CBB(0) case and leave its generalization to CBB(κ) case as an exercise. Sometime these exercises are nontrivial.

For example, the function comparison of CBB(−1) states that $f'' \leq \leq f$ for any function of the type $f = \cosh \circ \text{dist}_p$. (The inequality used here will be defined in Section 4C.)

Similarly, the function comparison of $\text{CBB}(1)$ states that for any point p , we have $f'' \leq -f$ for the function $f = -\cos \circ \text{dist}_p$ defined in $B(p, \pi)$. The geometric meaning of these inequalities remains the same: *distance functions in $\text{CBB}(\kappa)$ are more concave than distance functions in $\mathbb{M}(\kappa)$.*

Lecture 3

Globalization

A Globalization

A metric space \mathcal{L} is locally CBB(0) if any point $p \in \mathcal{L}$ admits a neighborhood $U \ni p$ such that the CBB(0) comparison holds for any quadruple of points in U .

3.1. Globalization theorem. *Any locally CBB(0) compact geodesic space is CBB(0).*

Proof modulo the key lemma. Let \mathcal{L} be a locally CBB(0) compact geodesic space. Note that condition 2.14b holds in \mathcal{L} (the proof is the same as for CBB(0) spaces). It remains to prove that 2.14a holds in \mathcal{L} ; that is,

$$\bullet \quad \angle[x_y^p] \geq \tilde{\angle}(x_y^p).$$

First note that \bullet holds for hinges in a small neighborhood of any point; this can be proved the same way as 2.8 and 2.11, applying the local version of the CBB(0) comparison. Since \mathcal{L} is compact, there is $\varepsilon > 0$ such that \bullet holds if $|x - p| + |p - y| < \varepsilon$. Applying the key lemma several times we get that \bullet holds for any given hinge. \square

3.2. Key lemma. *Let \mathcal{L} be a locally CBB(0) geodesic space. Assume that the comparison*

$$\angle[x_q^p] \geq \tilde{\angle}(x_q^p)$$

holds for any hinge $[x_q^p]$ with $|x - y| + |x - q| < \frac{2}{3} \cdot \ell$. Then the comparison

$$\angle[x_q^p] \geq \tilde{\angle}(x_q^p)$$

holds for any hinge $[x_q^p]$ with $|x - p| + |x - q| < \ell$.

Let $[x_q^p]$ be a hinge in a metric space \mathcal{L} with defined angle measure. Denote by $\tilde{\gamma}[x_q^p]$ its model side; this is the opposite side in a flat triangle with the same angle and two adjacent sides as in $[x_q^p]$.

More precisely, consider the model hinge $[\tilde{x}_q^{\tilde{p}}]$ in \mathbb{E}^2 that is defined by

$$\begin{aligned}\angle[\tilde{x}_q^{\tilde{p}}]_{\mathbb{E}^2} &= \angle[x_q^p]_{\mathcal{L}}, \\ |\tilde{x} - \tilde{p}|_{\mathbb{E}^2} &= |x - p|_{\mathcal{L}}, \\ |\tilde{x} - \tilde{q}|_{\mathbb{E}^2} &= |x - q|_{\mathcal{L}};\end{aligned}$$

then

$$\tilde{\gamma}[x_q^p]_{\mathcal{L}} := |\tilde{p} - \tilde{q}|_{\mathbb{E}^2}.$$

Note that

$$\tilde{\gamma}[x_q^p] \geq |p - q| \iff \angle[x_q^p] \geq \tilde{\angle}(x_q^p).$$

We will use it in the following proof.

Proof. It is sufficient to prove the inequality

$$\textcircled{2} \quad \tilde{\gamma}[x_q^p] \geq |p - q|$$

for any hinge $[x_q^p]$ with $|x - p| + |x - q| < \ell$.

Consider a hinge $[x_q^p]$ such that

$$\frac{2}{3} \cdot \ell \leq |p - x| + |x - q| < \ell.$$

First, let us construct a new hinge $[x'_q^p]$ with

$$\textcircled{3} \quad |p - x| + |x - q| \geq |p - x'| + |x' - q|,$$

such that

$$\textcircled{4} \quad \tilde{\gamma}[x_q^p] \geq \tilde{\gamma}[x'_q^p].$$

Construction. Assume $|x - q| \geq |x - p|$; otherwise, switch the roles of p and q . Take $x' \in [xq]$ such that


$$\textcircled{5} \quad |p - x| + 3 \cdot |x - x'| = \frac{2}{3} \cdot \ell.$$

Choose a geodesic $[x'p]$ and consider the hinge $[x'_q^p]$ formed by $[x'p]$ and $[x'q] \subset [xq]$. The triangle inequality implies $\textcircled{3}$. Further, note that

$$|p - x| + |x - x'| < \frac{2}{3} \cdot \ell, \quad |p - x'| + |x' - x| < \frac{2}{3} \cdot \ell.$$

In particular,

$$\textcircled{6} \quad \angle[x_q^p] \geq \tilde{\angle}(x_q^p) \quad \text{and} \quad \angle[x'_q^p] \geq \tilde{\angle}(x'_q^p).$$



Now, let $[\tilde{x}\tilde{x}'\tilde{p}] = \tilde{\Delta}(xx'p)$. Take \tilde{q} on the extension of $[\tilde{x}\tilde{x}']$ beyond x' such that $|\tilde{x} - \tilde{q}| = |x - q|$ (and therefore $|\tilde{x}' - \tilde{q}| = |x' - q|$). By ❹,



$$\angle[x_q^p] = \angle[x_{x'}^p] \geq \tilde{\angle}(x_{x'}^p) \Rightarrow \tilde{\gamma}[x_q^p] \geq |\tilde{p} - \tilde{q}|.$$

Hence

$$\begin{aligned} \angle[\tilde{x}'\tilde{p}_{\tilde{q}}] &= \pi - \tilde{\angle}(x_{x'}^p) \geq \\ &\geq \pi - \angle[x_{x'}^p] = \\ &= \angle[x_q^p], \end{aligned}$$

and ❶ follows.

Let us continue the proof. Set $x_0 = x$. Let us apply inductively the above construction to get a sequence of hinges $[x_n^p]$ with $x_{n+1} = x'_n$. From ❶, we have that the sequence $s_n = \tilde{\gamma}[x_n^p]$ is nonincreasing.



The sequence might terminate at some n only if $|p - x_n| + |x_n - q| < \frac{2}{3} \cdot \ell$. In this case, by the assumptions of the lemma, $\tilde{\gamma}[x_n^p] \geq |p - q|$. Since the sequence s_n is nonincreasing, inequality ❷ follows.

Otherwise, the sequence $r_n = |p - x_n| + |x_n - q|$ is nonincreasing, and $r_n \geq \frac{2}{3} \cdot \ell$ for all n . Note that by construction, the distances $|x_n - x_{n+1}|$, $|x_n - p|$, and $|x_n - q|$ are bounded away from zero for all large n . Indeed, since on each step, we move x_n toward to the point p or q that is further away, the distances $|x_n - p|$ and $|x_n - q|$ become about the same. Namely, by ❺, we have that $|p - x_n| - |x_n - q| \leq \frac{2}{9} \cdot \ell$ for all large n . Since $|p - x_n| + |x_n - q| \geq \frac{2}{3} \cdot \ell$, we have $|x_n - p| \geq \frac{\ell}{100}$ and $|x_n - q| \geq \frac{\ell}{100}$. Further, since $r_n \geq \frac{2}{3} \cdot \ell$, ❺ implies that $|x_n - x_{n+1}| > \frac{\ell}{100}$.

Since the sequence r_n is nonincreasing, it converges. In particular, $r_n - r_{n+1} \rightarrow 0$ as $n \rightarrow \infty$. It follows that $\tilde{\angle}(x_n^{p_n}) \rightarrow \pi$, where $p_n = p$ if $x_{n+1} \in [x_n q]$, and otherwise $p_n = q$. Since $\angle[x_n^{p_n}] \geq \tilde{\angle}(x_n^{p_n})$, we have $\angle[x_n^{p_n}] \rightarrow \pi$ as $n \rightarrow \infty$.

It follows that

$$r_n - s_n = |p - x_n| + |x_n - q| - \tilde{\gamma}[x_n^p] \rightarrow 0.$$

Together with the triangle inequality

$$|p - x_n| + |x_n - q| \geq |p - q|$$

this yields

$$\lim_{n \rightarrow \infty} \tilde{\Upsilon}[x_n \frac{p}{q}] \geq |p - q|.$$

Finally, the monotonicity of the sequence $s_n = \tilde{\Upsilon}[x_n \frac{p}{q}]$ implies ❷. \square

B On general curvature bound

The globalization theorem can be generalized to $\text{CBB}(\kappa)$ spaces for any real κ . The case $\kappa \leq 0$ is proved in the same way, but the case $\kappa > 0$ requires minor modifications.

3.3. Exercise. Suppose $\kappa \leq K$. Show that

$$\tilde{\angle}(x \frac{y}{z})_{\mathbb{M}(\kappa)} \leq \tilde{\angle}(x \frac{y}{z})_{\mathbb{M}(K)}$$

if the right-hand side is defined.

Conclude that any $\text{CBB}(K)$ space is locally $\text{CBB}(\kappa)$.

The exercise and the globalization theorem (here we need a more general version 3.11) imply the following.

3.4. Claim. Suppose $K > \kappa$, then any geodesic $\text{CBB}(K)$ space is $\text{CBB}(\kappa)$.

Recall that $\text{CBB}(\kappa)$ stands for *curvature bounded below by κ* ; so, for geodesic spaces it makes sense. However, as you can see from the following exercise, it does not make much sense in general.

3.5. Exercise. Let \mathcal{X} be the set $\{p, x_1, x_2, x_3\}$ with the metric defined by

$$|p - x_i| = \pi, \quad |x_i - x_j| = 2 \cdot \pi$$

for all $i \neq j$. Show that \mathcal{X} is $\text{CBB}(1)$, but not $\text{CBB}(0)$.

3.6. Exercise. Let p and q be points in a $\text{CBB}(1)$ geodesic space \mathcal{L} . Suppose $|p - q| > \pi$. Denote by m the midpoint of $[pq]$. Show that for any hinge $[m \frac{x}{p}]$ we have either $\angle[m \frac{x}{p}] = 0$ or $\angle[m \frac{x}{p}] = \pi$. Conclude that \mathcal{L} is isometric to a line interval or a circle.

3.7. Exercise. Suppose \mathcal{L} is a complete geodesic $\text{CBB}(1)$ space and $\text{diam } \mathcal{L} \leq \pi$. Show that

$$|x - y| + |y - z| + |z - x| \leq 2 \cdot \pi$$

for any triple of points $x, y, z \in \mathcal{L}$.

C Remarks

The following question about 2.14a is a long-standing open problem (possibly dating back to Alexandrov); it was stated in [10, footnote in 4.1.5] but is much much older.

3.8. Open question. *Let \mathcal{L} be a complete geodesic space (you can also assume that \mathcal{L} is homeomorphic to \mathbb{S}^2 or \mathbb{R}^2) such that for any hinge $[x_y^p]$ in \mathcal{L} , the angle $\angle[x_y^p]$ is defined and*

$$\angle[x_y^p] \geq \tilde{\angle}(x_y^p).$$

Is it true that \mathcal{L} is CBB(0)?

The globalization theorem is also known as the *generalized Toponogov theorem*. Its two-dimensional case was proved by Paolo Pizzetti [55]; later it was reproved independently by Alexandr Alexandrov [7]. Victor Toponogov [61] proved it for Riemannian manifolds of all dimensions.

We took the proof from our book [3] and reduced generality. This proof is based on simplifications obtained by Conrad Plaut [56] and Dmitry Burago, Yuriy Burago, and Sergei Ivanov [10]. The same proof was rediscovered independently by Urs Lang and Viktor Schroeder [31]. Another proof was obtained by Katsuhiro Shiohama [59].

The compactness condition in our version of the theorem can be traded for completeness. The proof uses the following statement where $r(x)$ measures the size of a neighborhood of x where the comparison holds.

3.9. Exercise. *Let \mathcal{X} be a complete metric space. Suppose $r: \mathcal{X} \rightarrow \mathbb{R}$ is a positive continuous function. Show that for any $\varepsilon > 0$ there is a point $p \in \mathcal{X}$ such that*

$$r(x) > (1 - \varepsilon) \cdot r(p)$$

for any $x \in \overline{B}[p, \frac{1}{\varepsilon} \cdot r(p)]$.

Let us mention two more general versions of the globalization theorem. The first one was proved in the paper of Michael Gromov, Yuriy Burago, and Grigory Perelman [11]. The second one was proved by the second author [49]. Their assumptions overlap, but neither implies the other.

3.10. Theorem. *Any complete locally CBB(κ) length space is CBB(κ).*

3.11. Theorem. *Any locally CBB(κ) geodesic space is CBB(κ).*

Lecture 4

Calculus

A Space of directions

By 2.8, angle measure of a hinge in a $\text{CBB}(\kappa)$ space \mathcal{L} is defined. Given $p \in \mathcal{L}$, consider the set \mathfrak{S}_p of all nontrivial unit-speed geodesics starting at p . By 1.9, the triangle inequality holds for \angle on \mathfrak{S}_p , that is, (\mathfrak{S}_p, \angle) forms a semimetric space.

The metric space corresponding to (\mathfrak{S}_p, \angle) is called the space of geodesic directions at p , denoted by Σ'_p or $\Sigma'_p \mathcal{L}$. The elements of Σ'_p are called geodesic directions at p . Each geodesic direction is formed by an equivalence class of geodesics starting from p for the equivalence relation

$$[px] \sim [py] \iff \angle[p_y^x] = 0;$$

the direction of $[px]$ is denoted by $\uparrow_{[px]}$. By 2.9,

$$[px] \sim [py] \iff [px] \subset [py] \text{ or } [px] \supset [py].$$

The completion of Σ'_p is called the space of directions at p and is denoted by Σ_p or $\Sigma_p \mathcal{L}$. The elements of Σ_p are called directions at p .

4.1. Exercise. *Let \mathcal{L} be a complete geodesic $\text{CBB}(0)$ space. Assume that a sequence of geodesics $[px_n]$ converge to a geodesic $[px_\infty]$ in the sense of Hausdorff, and $x_\infty \neq p$.*

- (a) *Suppose Σ_p is compact. Show that $\uparrow_{[px_n]} \rightarrow \uparrow_{[px_\infty]}$ as $n \rightarrow \infty$.*
- (b) *Give an example showing that without compactness of Σ_p , part (a) does not hold.*

B Tangent space

The Euclidean cone $\mathcal{V} = \text{Cone } \mathcal{X}$ over a metric space \mathcal{X} is defined as the metric space whose underlying set consists of equivalence classes in $[0, \infty) \times \mathcal{X}$ with the equivalence relation “ \sim ” given by $(0, p) \sim (0, q)$ for any points $p, q \in \mathcal{X}$, and whose metric is given by the cosine rule

$$|(s, p) - (t, q)|_{\mathcal{V}} = \sqrt{s^2 + t^2 - 2 \cdot s \cdot t \cdot \cos \theta},$$

where $\theta = \min\{\pi, |p - q|_{\mathcal{X}}\}$.

Note that $\text{Cone } \mathbb{S}^n$ is isometric to \mathbb{E}^{n+1} — this is a leading example. Next, we need to generalize several notions from Euclidean space to Euclidean cones.

The point in \mathcal{V} that corresponds $(t, x) \in [0, \infty) \times \mathcal{X}$ will be denoted by $t \cdot x$. The point in \mathcal{V} formed by the equivalence class of $\{0\} \times \mathcal{X}$ is called the origin of the cone and is denoted by 0 or $0_{\mathcal{V}}$. For $v \in \mathcal{V}$ the distance $|0 - v|_{\mathcal{V}}$ is called the norm of v and is denoted by $|v|$ or $|v|_{\mathcal{V}}$. The scalar product $\langle v, w \rangle$ of $v = s \cdot p$ and $w = t \cdot q$ is defined by

$$\langle v, w \rangle := |v| \cdot |w| \cdot \cos \theta$$

where $\theta = \min\{\pi, |p - q|_{\mathcal{X}}\}$. The value θ is undefined if $v = 0$ or $w = 0$; in this cases we assume that $\langle v, w \rangle := 0$.

4.2. Exercise. Show that $\text{Cone } \mathcal{X}$ is geodesic if and only if \mathcal{X} is π -geodesic; that is any two points $x, y \in \mathcal{X}$ such that $|x - y|_{\mathcal{X}} < \pi$ can be joined by a geodesic in \mathcal{X} .

Tangent space. The Euclidean cone $\text{Cone } \Sigma_p$ over the space of directions Σ_p is called the tangent space at p and denoted by T_p or $T_p \mathcal{L}$ if we need to emphasise that p is a point of space \mathcal{L} . The elements of $T_p \mathcal{X}$ will be called tangent vectors at p (despite that T_p is only a cone — not a vector space). The space of directions Σ_p can be (and will be) identified with the unit sphere in T_p .

4.3. Proposition. Tangent spaces of $\text{CBB}(\kappa)$ space are $\text{CBB}(0)$.

Proof. Consider the tangent space $T_p = \text{Cone } \Sigma_p$ of a $\text{CBB}(\kappa)$ space \mathcal{L} at a point p . We need to show that the $\text{CBB}(0)$ comparison holds for a given quadruple $v_0, v_1, v_2, v_3 \in T_p$.

Recall that the space of geodesic directions Σ'_p is dense in Σ_p . It follows that the subcone $T'_p = \text{Cone } \Sigma'_p$ is dense in T_p . Therefore, it is sufficient to consider the case $v_0, v_1, v_2, v_3 \in T'_p$.

For each i , choose a geodesic γ_i from p in the direction of v_i ; assume γ_i has speed $|v_i|$ for each i . Since the angles are defined, we have

$$\textcircled{1} \quad |\gamma_i(\varepsilon) - \gamma_j(\varepsilon)|_{\mathcal{L}} = \varepsilon \cdot |v_i - v_j|_{T_p} + o(\varepsilon)$$

for $\varepsilon > 0$. The quadruple $\gamma_0(\varepsilon), \gamma_1(\varepsilon), \gamma_2(\varepsilon), \gamma_3(\varepsilon)$ meets the $\text{CBB}(\kappa)$ comparison. After rescaling all the distances by $\frac{1}{\varepsilon}$, it becomes the $\text{CBB}(\varepsilon^2 \cdot \kappa)$ comparison. Passing to the limit as $\varepsilon \rightarrow 0$ and applying **1**, we get the $\text{CBB}(0)$ comparison for v_0, v_1, v_2, v_3 . \square

C Semiconcave functions

Recall that λ -concave functions were defined in Section 2F; also recall that we assume that these functions are locally Lipschitz and defined on an open domain.

Let f be locally Lipschitz real-valued function defined in an open subset $\text{Dom } f$ of a $\text{CBB}(0)$ space \mathcal{L} . We will write $f'' \leq \varphi$ if for any point $x \in \text{Dom } f$ and any $\varepsilon > 0$ there is a neighborhood $U \ni x$ such that the restriction $f|_U$ is $(\varphi(x) + \varepsilon)$ -concave. Here we assume that φ is continuous and defined in $\text{Dom } f$.

If $f'' \leq \varphi$ for some continuous function φ , then f is called *semiconcave*.

4.4. Exercise. *Let f be a distance function on a geodesic $\text{CBB}(0)$ space \mathcal{L} ; that is, $f(x) \equiv |p - x|$ for some $p \in \mathcal{L}$. Show that $f'' \leq \frac{1}{f}$. In particular, f is semiconcave in $\mathcal{L} \setminus \{p\}$.*

D Differential

Let \mathcal{X} be a space with defined angles. Let f be a semiconcave function on \mathcal{X} and $p \in \text{Dom } f$. Choose a unit-speed geodesic γ that starts at p ; let $\xi \in \Sigma_p$ be its direction. Define

$$(\mathbf{d}_p f)(\xi) := (f \circ \gamma)^+(0),$$

here $(f \circ \gamma)^+$ denotes the right derivative of $(f \circ \gamma)$; it is defined since f is semiconcave.

By the following exercise, the value $(\mathbf{d}_p f)(\xi)$ is defined; that is, it does not depend on the choice of γ . Moreover, $\mathbf{d}_p f$ is a Lipschitz function on Σ'_p . It follows that the function $\mathbf{d}_p f: \Sigma'_p \rightarrow \mathbb{R}$ can be extended to a Lipschitz function $\mathbf{d}_p f: \Sigma_p \rightarrow \mathbb{R}$. Further, we can extend it to the tangent space by setting

$$(\mathbf{d}_p f)(r \cdot \xi) := r \cdot (\mathbf{d}_p f)(\xi)$$

for any $r \geq 0$ and $\xi \in \Sigma_p$. The obtained function $\mathbf{d}_p f: T_p \rightarrow \mathbb{R}$ is Lipschitz; it is called the differential of f at p .

4.5. Exercise. Let f be a semiconcave function on a geodesic $\text{CBB}(\kappa)$ space. Suppose γ_1 and γ_2 are unit-speed geodesics that start at $p \in \text{Dom } f$; denote by θ the angle between γ_1 and γ_2 at p . Show that

$$|(f \circ \gamma_1)^+(0) - (f \circ \gamma_2)^+(0)| \leq L \cdot \theta,$$

where L is the Lipschitz constant of f in a neighborhood of p .

4.6. Exercise. Let p and q be distinct points in a geodesic $\text{CBB}(0)$ space \mathcal{L} . Show that

(a) If ξ is the direction of a geodesic $[pq]$ at p , then

$$\mathbf{d}_p \text{dist}_q(v) \leq -\langle \xi, v \rangle$$

for any $v \in \mathbb{T}_p$.

(b) Suppose \mathcal{L} is proper. Let Ξ be the set of all direction of geodesics from q to p . Then

$$\mathbf{d}_p \text{dist}_q(v) = -\max_{\xi \in \Xi} \langle \xi, v \rangle$$

for any $v \in \mathbb{T}_p$.

E Gradient

4.7. Definition. Let f be a semiconcave function on a geodesic $\text{CBB}(\kappa)$ space. A tangent vector $g \in \mathbb{T}_p$ is called a gradient of f at p (briefly, $g = \nabla_p f$) if

- (a) $(\mathbf{d}_p f)(w) \leq \langle g, w \rangle$ for any $w \in \mathbb{T}_p$, and
- (b) $(\mathbf{d}_p f)(g) = \langle g, g \rangle$.

The following exercise provides a key property of gradients that will be important latter; see the first distance estimate (5.6).

4.8. Exercise. Let f be a λ -concave function on a geodesic $\text{CBB}(\kappa)$ space. Suppose that gradients $\nabla_x f$ and $\nabla_y f$ are defined. Show that

$$\langle \uparrow_{[xy]}, \nabla_x f \rangle + \langle \uparrow_{[yx]}, \nabla_y f \rangle + \lambda \cdot |x - y| \geq 0.$$

4.9. Proposition. Suppose that a semiconcave function f is defined in a neighborhood of a point p in a $\text{CBB}(\kappa)$ space. Then the gradient $\nabla_p f$ is uniquely defined.

Moreover, if $\mathbf{d}_p f \leq 0$, then we have $\nabla_p f = 0$; otherwise, $\nabla_p f = s \cdot \bar{\xi}$, where $s = \mathbf{d}_p f(\bar{\xi})$ and $\bar{\xi} \in \Sigma_p$ is the direction that maximize the value $\mathbf{d}_p f(\xi)$ for $\xi \in \Sigma_p$.

4.10. Key lemma. *Let f be a λ -concave function that is defined in a neighborhood of a point p in a geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Then for any $u, v \in T_p$, we have*

$$s \cdot \sqrt{|u|^2 + 2 \cdot \langle u, v \rangle + |v|^2} \geq (\mathbf{d}_p f)(u) + (\mathbf{d}_p f)(v),$$

where

$$s = \sup \{ (\mathbf{d}_p f)(\xi) : \xi \in \Sigma_p \}.$$

Note that if $T_p \stackrel{\text{iso}}{=} \mathbb{E}^m$ and $\mathbf{d}_p f$ is a concave function, then $2 \cdot (\mathbf{d}_p f)(\frac{u+v}{2}) \geq (\mathbf{d}_p f)(u) + (\mathbf{d}_p f)(v)$. The latter implies the statement since $|u + v| = \sqrt{|u|^2 + 2 \cdot \langle u, v \rangle + |v|^2}$. In general, T_p is not geodesic (and not even a length space), so concavity of $\mathbf{d}_p f$ does not make sense. The key lemma however says that in a certain sense $\mathbf{d}_p f$ behaves as a concave function.

It is instructive to solve the following exercise before reading the proof of the key lemma.

4.11. Exercise. *Let $[q^p_x]$ be a hinge in a $\text{CBB}(0)$ space and $y \in]qp[$. Suppose that γ is the unit speed parametrization of $[qx]$ from q to x . Show that*

$$|y - \gamma(t)| = |y - q| - t \cdot \cos(\angle[q^p_x]) + o(t).$$

Conclude that


$$(\mathbf{d}_q \text{dist}_y)(w) = -\langle \uparrow_{[qp]}, w \rangle$$

for any $w \in T_x$

Proof of 4.10. We will assume $\kappa = 0$; the general case requires only minor modifications. We can assume that $v \neq 0$, $w \neq 0$, and $\alpha = \angle(u, v) > 0$; otherwise, the statement is trivial.

Consider a model configuration of five points: $\tilde{p}, \tilde{u}, \tilde{v}, \tilde{q}, \tilde{w} \in \mathbb{E}^2$ such that

- ◇ $\angle[\tilde{p}\tilde{u}\tilde{v}] = \alpha$,
- ◇ $|\tilde{p} - \tilde{u}| = |u|$,
- ◇ $|\tilde{p} - \tilde{v}| = |v|$,
- ◇ \tilde{q} lies on an extension of $[\tilde{p}\tilde{v}]$ so that \tilde{v} is the midpoint of $[\tilde{p}\tilde{q}]$,
- ◇ \tilde{w} is the midpoint between \tilde{u} and \tilde{v} .



Note that

❶ $|\tilde{p} - \tilde{w}| = \frac{1}{2} \cdot \sqrt{|u|^2 + 2 \cdot \langle u, v \rangle + |v|^2}.$

Since the geodesic space of directions Σ'_p is dense in Σ_p , we can assume that there are geodesics in the directions of u and v . Choose such geodesics γ_u and γ_v and assume that they are parametrized with speed $|u|$ and $|v|$ respectively. For all small $t > 0$, consider points $u_t, v_t, q_t, w_t \in \mathcal{L}$ such that

- ◇ $v_t = \gamma_v(t)$, $q_t = \gamma_v(2 \cdot t)$
- ◇ $u_t = \gamma_u(t)$.
- ◇ w_t is the midpoint of $[u_t v_t]$.

Clearly

$$|p - u_t| = t \cdot |u|, \quad |p - v_t| = t \cdot |v|, \quad |p - q_t| = 2 \cdot t \cdot |v|.$$

Since $\angle(u, v)$ is defined, we have

$$|u_t - v_t| = t \cdot |\tilde{u} - \tilde{v}| + o(t), \quad |u_t - q_t| = t \cdot |\tilde{u} - \tilde{q}| + o(t).$$

From the point-on-side and hinge comparisons (2.15c+2.15d), we have

$$\tilde{\angle}(v_t \overset{p}{w_t}) \geq \tilde{\angle}(v_t \overset{p}{u_t}) \geq \angle[\tilde{v} \overset{\tilde{p}}{\tilde{u}}] + \frac{o(t)}{t}$$

and

$$\tilde{\angle}(v_t \overset{q_t}{w_t}) \geq \tilde{\angle}(v_t \overset{q_t}{u_t}) \geq \angle[\tilde{v} \overset{\tilde{q}}{\tilde{u}}] + \frac{o(t)}{t}.$$

Clearly, $\angle[\tilde{v} \overset{\tilde{p}}{\tilde{u}}] + \angle[\tilde{v} \overset{\tilde{q}}{\tilde{u}}] = \pi$. From the adjacent angle comparison (2.15b), $\tilde{\angle}(v_t \overset{p}{u_t}) + \tilde{\angle}(v_t \overset{q_t}{u_t}) \leq \pi$. Hence $\tilde{\angle}(v_t \overset{p}{w_t}) \rightarrow \angle[\tilde{v} \overset{\tilde{p}}{\tilde{w}}]$ as $t \rightarrow 0+$ and thus

$$|p - w_t| = t \cdot |\tilde{p} - \tilde{w}| + o(t).$$

Without loss of generality, we can assume that $f(p) = 0$. Since f is λ -concave, we have

$$\begin{aligned} 2 \cdot f(w_t) &\geq f(u_t) + f(v_t) + \frac{\lambda}{4} \cdot |u_t - v_t|^2 = \\ &= t \cdot [(\mathbf{d}_p f)(u) + (\mathbf{d}_p f)(v)] + o(t). \end{aligned}$$

Applying λ -concavity of f , we have

$$\begin{aligned} (\mathbf{d}_p f)(\uparrow_{[p w_t]}) &\geq \frac{f(w_t) - \frac{\lambda}{2} \cdot |p - w_t|^2}{|p - w_t|} \geq \\ &\geq \frac{t \cdot [(\mathbf{d}_p f)(u) + (\mathbf{d}_p f)(v)] + o(t)}{2 \cdot t \cdot |\tilde{p} - \tilde{w}| + o(t)}. \end{aligned}$$

By ❶, the key lemma follows. □

Proof of 4.9; uniqueness. If $g, g' \in T_p$ are two gradients of f , then

$$\langle g, g \rangle = (\mathbf{d}_p f)(g) \leq \langle g, g' \rangle, \quad \langle g', g' \rangle = (\mathbf{d}_p f)(g') \leq \langle g, g' \rangle.$$

Therefore,

$$|g - g'|^2 = \langle g, g \rangle - 2 \cdot \langle g, g' \rangle + \langle g', g' \rangle \leq 0.$$

It follows that $g = g'$.

Existence. Note first that if $\mathbf{d}_p f \leq 0$, then one can take $\nabla_p f = 0$.

Otherwise, if $s = \sup \{ (\mathbf{d}_p f)(\xi) : \xi \in \Sigma_p \} > 0$, it is sufficient to show that there is $\bar{\xi} \in \Sigma_p$ such that

$$\textcircled{2} \quad (\mathbf{d}_p f)(\bar{\xi}) = s.$$

Indeed, suppose $\bar{\xi}$ exists. Applying 4.10 for $u = \bar{\xi}$, $v = \varepsilon \cdot w$ with $\varepsilon \rightarrow 0+$, we get

$$(\mathbf{d}_p f)(w) \leq \langle w, s \cdot \bar{\xi} \rangle$$

for any $w \in T_p$; that is, $s \cdot \bar{\xi}$ is the gradient at p .

Take a sequence of directions $\xi_n \in \Sigma_p$, such that $(\mathbf{d}_p f)(\xi_n) \rightarrow s$. Applying 4.10 for $u = \xi_n$ and $v = \xi_m$, we get

$$s \geq \frac{(\mathbf{d}_p f)(\xi_n) + (\mathbf{d}_p f)(\xi_m)}{\sqrt{2 + 2 \cdot \cos \angle(\xi_n, \xi_m)}}.$$

Therefore $\angle(\xi_n, \xi_m) \rightarrow 0$ as $n, m \rightarrow \infty$; that is, the sequence ξ_n is Cauchy. Clearly, $\bar{\xi} = \lim_n \xi_n$ meets $\textcircled{2}$. \square

4.12. Exercise. Let f and g be locally Lipschitz semiconcave functions defined in a neighborhood of a point p in a CBB space. Show that

$$|\nabla_p f - \nabla_p g|_{T_p}^2 \leq s \cdot (|\nabla_p f| + |\nabla_p g|),$$

where

$$s = \sup \{ |(\mathbf{d}_p f)(\xi) - (\mathbf{d}_p g)(\xi)| : \xi \in \Sigma_p \}.$$

Conclude that if the sequence of restrictions $\mathbf{d}_p f_n|_{\Sigma_p}$ converges uniformly, then $\nabla_p f_n$ converges as $n \rightarrow \infty$. Here we assume that all functions f_1, f_2, \dots are semiconcave and locally Lipschitz.

4.13. Exercise. Let f be a locally Lipschitz λ -concave function on a complete geodesic CBB(κ) space \mathcal{L} .

(a) Suppose $s \geq 0$. Show that $|\nabla_x f| > s$ if and only if for some point y we have

$$f(y) - f(x) > s \cdot \ell + \lambda \cdot \frac{\ell^2}{2},$$

where $\ell = |x - y|$.

(b) Show that $x \mapsto |\nabla_x f|$ is lower semicontinuous; that is, if $x_n \rightarrow x_\infty$, then

$$|\nabla_{x_\infty} f| \leq \liminf_{n \rightarrow \infty} |\nabla_{x_n} f|.$$

Lecture 5

Gradient flow

A Velocity of curve

Let α be a curve in a geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . If for any choice of geodesics $[p\alpha(t_0 + \varepsilon)]$ the vectors

$$\frac{1}{\varepsilon} \cdot |p - \alpha(t_0 + \varepsilon)| \cdot \uparrow_{[p\alpha(t_0 + \varepsilon)]}$$

converge as $\varepsilon \rightarrow 0+$, then their limit in T_p is called the right derivative of α at t_0 ; it will be denoted by $\alpha^+(t_0)$. In addition, $\alpha^+(t_0) := 0$ if $\frac{1}{\varepsilon} \cdot |p - \alpha(t_0 + \varepsilon)| \rightarrow 0$ as $\varepsilon \rightarrow 0+$.

The tangent vector $v = |p - x| \cdot \uparrow_{[px]}$ can be called logarithm of x at p (briefly, $\log_p x$); it is a tangent vector at p of a geodesic path from p to x .

5.1. Claim. *Let α be a curve in a geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Suppose f a semiconcave Lipschitz function defined in a neighborhood of $p = \alpha(0)$, and $\alpha^+(0)$ is defined. Then*

$$(f \circ \alpha)^+(0) = (d_p f)(\alpha^+(0)).$$

Proof. Without loss of generality, we can assume that $f(p) = 0$. Suppose f and therefore $d_p f$ are L -Lipschitz.

Choose a constant-speed geodesic γ that starts from p , such that the distance $s = |\alpha^+(0) - \gamma^+(0)|_{T_p}$ is small. By the definition of differential,

$$(f \circ \gamma)^+(0) = d_p f(\gamma^+(0)).$$

By comparison and the definition of α^+ ,

$$|\alpha(\varepsilon) - \gamma(\varepsilon)|_{\mathcal{L}} \leq s \cdot \varepsilon + o(\varepsilon)$$

for $\varepsilon > 0$. Therefore,

$$|f \circ \alpha(\varepsilon) - f \circ \gamma(\varepsilon)| \leq L \cdot s \cdot \varepsilon + o(\varepsilon).$$

Suppose $(f \circ \alpha)^+(0)$ is defined. Then

$$|(f \circ \alpha)^+(0) - (f \circ \gamma)^+(0)| \leq L \cdot s.$$

Since $d_p f$ is L -Lipschitz, we also get

$$|d_p f(\alpha^+(0)) - d_p f(\gamma^+(0))| \leq L \cdot s.$$

It follows that the needed identity holds up to error $2 \cdot L \cdot s$. The statement follows since $s > 0$ can be chosen arbitrarily.

The same argument is applicable if in the place of $(f \circ \alpha)^+(0)$ we use any limit of $\frac{1}{\varepsilon_n} \cdot [f \circ \alpha(\varepsilon_n) - f(p)]$ for a sequence $\varepsilon_n \rightarrow 0+$. It proves that all such limits coincide; in particular, $(f \circ \alpha)^+(0)$ is defined and equals to $(d_p f)(\alpha^+(0))$. \square

B Gradient curves

5.2. Definition. Let $f: \mathcal{L} \rightarrow \mathbb{R}$ be a locally Lipschitz and semiconcave function on a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} .

A locally Lipschitz curve $\alpha: [t_{\min}, t_{\max}) \rightarrow \text{Dom } f$ will be called an f -gradient curve if

$$\alpha^+ = \nabla_{\alpha} f;$$

that is, for any $t \in [t_{\min}, t_{\max})$, $\alpha^+(t)$ is defined and $\alpha^+(t) = \nabla_{\alpha(t)} f$.

A complete proof of the following theorem takes about 5 pages [3]; it mimics the proof of the standard Picard theorem on the existence and uniqueness of solutions of ordinary differential equations. We omit the proof of existence; the uniqueness will be proved in the next section.

5.3. Picard theorem. Let $f: \mathcal{L} \rightarrow \mathbb{R}$ be a locally Lipschitz and λ -concave function on a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Then for any $p \in \text{Dom } f$, there are unique $t_{\max} \in (0, \infty]$ and f -gradient curve $\alpha: [0, t_{\max}) \rightarrow \mathcal{L}$ with $\alpha(0) = p$ such that any sequence $t_n \rightarrow t_{\max}-$, the sequence $\alpha(t_n)$ does not have a limit point in $\text{Dom } f$.

Note that the theorem says that the future of a gradient curve is determined by its present, but it says nothing about its past.

Here is an example showing that the past is not determined by the present. Consider the function $f: x \mapsto -|x|$ on the real line \mathbb{R} . The tangent space $T_x\mathbb{R}$ can be identified with \mathbb{R} . Note that

$$\nabla_x f = \begin{cases} 1 & \text{if } x < 0, \\ 0 & \text{if } x = 0, \\ -1 & \text{if } x > 0. \end{cases}$$


So, the f -gradient curves go to the origin with unit speed and then stand there forever. In particular, if α is an f -gradient curve that starts at x , then $\alpha(t) = 0$ for any $t \geq |x|$.

Here is a slightly more interesting example; it shows that gradient curves can merge even in the region where $|\nabla f| \neq 0$.

5.4. Example. Consider the function $f: (x, y) \mapsto -|x| - |y|$ on the (x, y) -plane. Note that f is concave; its gradient field is sketched on the figure.

Let α be an f -gradient curve that starts at (x, y) for $x > y > 0$. Then

$$\alpha(t) = \begin{cases} (x - t, y - t) & \text{for } 0 \leq t \leq x - y, \\ (x - t, 0) & \text{for } x - y \leq t \leq x, \\ (0, 0) & \text{for } x \leq t. \end{cases}$$



C Distance estimates

5.5. Observation. Let α be a gradient curve of a λ -concave function f defined on a complete geodesic CBB space. Choose a point p ; let $\ell(t) := \text{dist}_p \circ \alpha(t)$ and $q = \alpha(t_0)$. Then

$$\ell^+(t_0) \leq -\left(f(p) - f(q) - \frac{\lambda}{2} \cdot \ell^2(t_0)\right) / \ell(t_0)$$

Proof. Let γ be the unit-speed parametrization of $[qp]$ from q to p , so $q = \gamma(0)$. Then

$$\begin{aligned} \ell^+(t_0) &= (\mathbf{d}_q \text{dist}_p)(\nabla_q f) \leq \\ &\leq -\langle \uparrow_{[qp]}, \nabla_q f \rangle \leq \\ &\leq -\mathbf{d}_q f(\uparrow_{[qp]}) = \\ &= -(f \circ \gamma)^+(0) \leq \\ &\leq -\left(f(p) - f(q) - \frac{\lambda}{2} \cdot \ell^2(t_0)\right) / \ell(t_0) \end{aligned}$$

In the above calculations we consequently applied 5.1, 4.6, the definition of gradient, the definition of differential, and concavity of $t \mapsto f \circ \gamma(t) - \frac{\lambda}{2} \cdot t^2$. \square

Note that the following estimate implies uniqueness in the Picard theorem (5.3).

5.6. First distance estimate. *Let f be a λ -concave locally Lipschitz function on a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Then*

$$|\alpha(t) - \beta(t)| \leq e^{\lambda \cdot t} \cdot |\alpha(0) - \beta(0)|$$

for any $t \geq 0$ and any two f -gradient curves α and β .

Moreover, the statement holds for a locally Lipschitz λ -concave function defined in an open domain if there is a geodesic $[\alpha(t) \beta(t)]$ in $\text{Dom } f$ for any t .

Proof. Fix a choice of geodesic $[\alpha(t) \beta(t)]$ for each t . Let $\ell(t) = |\alpha(t) - \beta(t)|$. Note that

$$\ell^+(t) \leq -\langle \uparrow_{[\alpha(t)\beta(t)]}, \nabla_{\alpha(t)} f \rangle - \langle \uparrow_{[\beta(t)\alpha(t)]}, \nabla_{\beta(t)} f \rangle \leq \lambda \cdot \ell(t).$$

Here one has to apply 5.5 for distance to the midpoint m of $[\alpha(t) \beta(t)]$, and then apply the triangle inequality. Hence the result. \square

The following exercise describes a global geometric property of a gradient curve without direct reference to its function. It uses the notion of *self-contracting curves* introduced by Aris Daniilidis, Olivier Ley, and Stéphane Sabourau [14].

5.7. Exercise. *Let $f: \mathcal{L} \rightarrow \mathbb{R}$ be a locally Lipschitz and concave function on a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Then*

$$|\alpha(t_1) - \alpha(t_3)|_{\mathcal{L}} \geq |\alpha(t_2) - \alpha(t_3)|_{\mathcal{L}}.$$

for any f -gradient curve α and $t_1 \leq t_2 \leq t_3$.

5.8. Exercise. *Let f be a locally Lipschitz concave function defined on a $\text{CBB}(\kappa)$ space \mathcal{L} . Suppose $\hat{\alpha}: [0, \ell] \rightarrow \mathcal{L}$ is an arc-length reparametrization of an f -gradient curve. Show that $(f \circ \hat{\alpha})$ is concave.*

The following exercise implies that gradient curves for a uniformly converging sequence of λ -concave functions converge to the gradient curves of the limit function.

5.9. Exercise. *Let f and g be λ -concave locally Lipschitz functions on a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Suppose $\alpha, \beta: [0, t_{\max}) \rightarrow \mathcal{L}$*

are respectively f - and g -gradient curves. Assume $|f - g| < \varepsilon$; let $\ell: t \mapsto |\alpha(t) - \beta(t)|$. Show that

$$\ell^+ \leq \lambda \cdot \ell + \frac{2 \cdot \varepsilon}{\ell}.$$

Conclude that if $\alpha(0) = \beta(0)$ and $t_{\max} < \infty$, then

$$|\alpha(t) - \beta(t)| \leq c \cdot \sqrt{\varepsilon \cdot t}$$

for some constant $c = c(t_{\max}, \lambda)$.

D Gradient flow

Let \mathcal{L} be a complete geodesic CBB(κ) space and f be a locally Lipschitz semiconcave function defined on an open set of \mathcal{L} . If there is an f -gradient curve α such that $\alpha(0) = x$ and $\alpha(t) = y$, then we will write

$$\text{Flow}_f^t(x) = y.$$

The partially defined map Flow_f^t from \mathcal{L} to itself is called the f -gradient flow for time t . Note that

$$\text{Flow}_f^{t_1+t_2} = \text{Flow}_f^{t_1} \circ \text{Flow}_f^{t_2}.$$

In other words, one may think that gradient flow is an action of the semigroup $(\mathbb{R}_{\geq 0}, +)$ on the space.

From the first distance estimate 5.6, it follows that for any $t \geq 0$, the domain of definition of Flow_f^t is an open subset of \mathcal{L} . In some cases, it is globally defined. For example, if f is a λ -concave function defined on the whole space \mathcal{L} , then $\text{Flow}_f^t(x)$ is defined for all $x \in \mathcal{L}$ and $t \geq 0$; see [3, 16.19].

Now let us reformulate statements obtained earlier using this new terminology. From the first distance estimate, we have the following.

5.10. Proposition. *Let \mathcal{L} be a complete geodesic CBB(κ) space and $f: \mathcal{L} \rightarrow \mathbb{R}$ be a semiconcave function. Then the map $x \mapsto \text{Flow}_f^t(x)$ is locally Lipschitz.*

Moreover, if f is λ -concave, then Flow_f^t is $e^{\lambda \cdot t}$ -Lipschitz.

The next proposition follows from 5.9.

5.11. Proposition. *Let \mathcal{L} be complete geodesic CBB(κ) space. Suppose $f_n: \mathcal{L} \rightarrow \mathbb{R}$ is a sequence of λ -concave functions that converges to $f_\infty: \mathcal{L} \rightarrow \mathbb{R}$. Then for any $x \in \mathcal{L}$ and $t \geq 0$, we have*

$$\text{Flow}_{f_n}^t(x) \rightarrow \text{Flow}_{f_\infty}^t(x)$$

as $n \rightarrow \infty$.

E Gradient exponent

One of the technical difficulties in Alexandrov's geometry comes from nonextendability of geodesics. In particular, the exponential map, $\exp_p: T_p \rightarrow \mathcal{L}$, if defined the usual way, can be undefined in an arbitrary small neighborhood of the origin.

We construct its analog, the gradient exponential map

$$\text{gexp}_p: T_p \rightarrow \mathcal{L},$$

which practically solves this problem. It shares many properties with the ordinary exponential map, and better in certain respects, even in the Riemannian universe.

Let \mathcal{L} be Alexandrov's space and $p \in \mathcal{L}$, consider the function $f = \text{dist}_p^2/2$. Suppose $i_\lambda: \lambda \cdot \mathcal{L} \rightarrow \mathcal{L}$ sends a point in the rescaled copy $\lambda \cdot \mathcal{L}$ to the corresponding point in \mathcal{L} . Consider the one parameter family of maps

$$\Phi_f^t \circ i_{e^t}: e^t \cdot \mathcal{L} \rightarrow \mathcal{L}$$

where Φ_f^t denotes gradient flow. Note that $(e^t \cdot \mathcal{L}, p) \xrightarrow{\text{GH}} (T_p, o_p)$ as $t \rightarrow \infty$. Let us define the *gradient exponential map* as the limit

$$\text{gexp}_p = \lim_{t \rightarrow \infty} \Phi_f^t \circ i_{e^t}.$$

5.12. Proposition. *Let \mathcal{L} be a proper geodesic CBB(0) space. Then for any $p \in \mathcal{L}$ the gradient exponent $\text{gexp}_p: T_p \rightarrow \mathcal{L}$ is defined. Moreover, gexp_p is a short map and*

$$\text{gexp}_p(\gamma^+(0)) = \gamma(1)$$

for any geodesic path γ that starts at p .

The last statement in the proposition says that it is appropriate to use term *exponent* for gexp .

Proof. Note that $f'' \leq 1$. By the first distance estimate, we have that Φ_f^t is an e^t -Lipschitz. Therefore, the compositions $\Phi_f^t \circ i_{e^t}: e^t \cdot \mathcal{L} \rightarrow \mathcal{L}$ are short. Hence a partial limit $\text{gexp}_p: T_p \mathcal{L} \rightarrow \mathcal{L}$ exists, and it is a short map.

Clearly for any partial limit we have

$$\Phi_f^t \circ \text{gexp}_p(v) = \text{gexp}_p(e^t \cdot v).$$

Since Φ^t is e^t -Lipschitz, it follows that gexp_p is uniquely defined. \square

F Remarks

??? gradient exponent for $\kappa \neq 0$ and for nonproper.
 quasigeodesics

Lecture 6

Line splitting

A Busemann function

A half-line is a distance-preserving map from $\mathbb{R}_{\geq 0} = [0, \infty)$ to a metric space. In other words, a half-line is a geodesic defined on the real half-line $\mathbb{R}_{\geq 0}$. If $\gamma: [0, \infty) \rightarrow \mathcal{X}$ is a half-line, then the limit

$$\bullet \quad \text{bus}_\gamma(x) = \lim_{t \rightarrow \infty} |\gamma(t) - x| - t$$

is called the Busemann function of γ .

The Busemann function bus_γ mimics behavior of the distance function from the ideal point of γ .

6.1. Proposition. *For any half-line γ in a metric space \mathcal{X} , its Busemann function $\text{bus}_\gamma: \mathcal{X} \rightarrow \mathbb{R}$ is defined. Moreover, bus_γ is 1-Lipschitz and $\text{bus}_\gamma \circ \gamma(t) + t = 0$ for any t .*

Proof. By the triangle inequality, the function

$$t \mapsto |\gamma(t) - x| - t$$

is nonincreasing for any fixed x .

Since $t = |\gamma(0) - \gamma(t)|$, the triangle inequality implies that

$$|\gamma(t) - x| - t \geq -|\gamma(0) - x|.$$

Thus the limit in \bullet is defined, and it is 1-Lipschitz as a limit of 1-Lipschitz functions. The last statement follows since $|\gamma(t) - \gamma(t_0)| = t - t_0$ for all large t . \square

6.2. Exercise. *Any Busemann function on a geodesic $\text{CBB}(0)$ space is concave.*

B Splitting theorem

A line is a distance-preserving map from \mathbb{R} to a metric space. In other words, a line is a geodesic defined on the real line \mathbb{R} .

6.3. Exercise. *Let γ be a line in a metric space \mathcal{X} . Show that for any point x we have*

$$\text{bus}_+(x) + \text{bus}_-(x) \geq 0$$

where, bus_+ and bus_- , are the Busemann functions associated with half-lines $\gamma : [0, \infty) \rightarrow \mathcal{L}$ and $\gamma : (-\infty, 0] \rightarrow \mathcal{L}$ respectively.

Let \mathcal{X} be a metric space and $A, B \subset \mathcal{X}$. We will write

$$\mathcal{X} = A \oplus B$$

if there are projections $\text{proj}_A : \mathcal{X} \rightarrow A$ and $\text{proj}_B : \mathcal{X} \rightarrow B$ such that

$$|x - y|^2 = |\text{proj}_A(x) - \text{proj}_A(y)|^2 + |\text{proj}_B(x) - \text{proj}_B(y)|^2$$

for any two points $x, y \in \mathcal{X}$.

Note that if

$$\mathcal{X} = A \oplus B$$

then

- ◊ A intersects B at a single point,
- ◊ both sets A and B are convex sets in \mathcal{X} ; the latter means that any geodesic with the ends in A (or B) lies in A (or B).

6.4. Line splitting theorem. *Let \mathcal{L} be a complete geodesic CBB(0) space and γ be a line in \mathcal{L} . Then*

$$\mathcal{L} = \mathcal{L}' \oplus \gamma(\mathbb{R})$$

for some subset $\mathcal{L}' \subset \mathcal{L}$.

Before going into the proof, let us state a corollary of the theorem.

6.5. Corollary. *Let \mathcal{L} be a complete geodesic CBB(0) space. Then there is an isometric splitting*

$$\mathcal{L} = \mathcal{L}' \oplus H$$

where $H \subset \mathcal{L}$ is a subset isometric to a Hilbert space, and $\mathcal{L}' \subset \mathcal{L}$ is a convex subset that contains no line.

The following lemma is closely relevant to the first distance estimate (5.6); its proof goes along the same lines.

6.6. Lemma. *Suppose $f: \mathcal{L} \rightarrow \mathbb{R}$ be a concave 1-Lipschitz function on a geodesic $\text{CBB}(0)$ space \mathcal{L} . Consider two f -gradient curves α and β . Then for any $t, s \geq 0$ we have*

$$|\alpha(s) - \beta(t)|^2 \leq |p - q|^2 + 2 \cdot (f(p) - f(q)) \cdot (s - t) + (s - t)^2,$$

where $p = \alpha(0)$ and $q = \beta(0)$.

Proof. Since f is 1-Lipschitz, $|\nabla f| \leq 1$. Therefore

$$f \circ \beta(t) \leq f(q) + t$$

for any $t \geq 0$.

Set $\ell(t) = |p - \beta(t)|$. Applying 5.5, we get

$$\begin{aligned} (\ell^2)^+(t) &\leq 2 \cdot (f \circ \beta(t) - f(p)) \leq \\ &\leq 2 \cdot (f(q) + t - f(p)). \end{aligned}$$

Therefore

$$\ell^2(t) - \ell^2(0) \leq 2 \cdot (f(q) - f(p)) \cdot t + t^2.$$

It proves the needed inequality in case $s = 0$. Combining it with the first distance estimate (5.6), we get the result in case $s \leq t$. The case $s \geq t$ follows by switching the roles of s and t . \square

Proof of 6.4. Consider two Busemann functions, bus_+ and bus_- , associated with half-lines $\gamma: [0, \infty) \rightarrow \mathcal{L}$ and $\gamma: (-\infty, 0] \rightarrow \mathcal{L}$ respectively; that is,

$$\text{bus}_\pm(x) := \lim_{t \rightarrow \infty} |\gamma(\pm t) - x| - t.$$

According to 6.2, both functions bus_\pm are concave.

By 6.3, $\text{bus}_+(x) + \text{bus}_-(x) \geq 0$ for any $x \in \mathcal{L}$. On the other hand, by 2.20, $f(t) = \text{dist}_x^2 \circ \gamma(t)$ is 2-concave. In particular, $f(t) \leq t^2 + at + b$ for some constants $a, b \in \mathbb{R}$. Passing to the limit as $t \rightarrow \pm\infty$, we have $\text{bus}_+(x) + \text{bus}_-(x) \leq 0$. Hence

$$\text{bus}_+(x) + \text{bus}_-(x) = 0$$

for any $x \in \mathcal{L}$. In particular, the functions bus_\pm are affine; that is, they are convex and concave at the same time.

Note that for any x ,

$$\begin{aligned} |\nabla_x \text{bus}_\pm| &= \sup \{ \mathbf{d}_x \text{bus}_\pm(\xi) : \xi \in \Sigma_x \} = \\ &= \sup \{ -\mathbf{d}_x \text{bus}_\mp(\xi) : \xi \in \Sigma_x \} \equiv \\ &\equiv 1. \end{aligned}$$

Observe that α is a bus_\pm -gradient curve if and only if α is a geodesic such that $(\text{bus}_\pm \circ \alpha)^+ = 1$. Indeed, if α is a geodesic, then $(\text{bus}_\pm \circ \alpha)^+ \leq 1$ and the equality holds only if $\nabla_\alpha \text{bus}_\pm = \alpha^+$. Now suppose $\nabla_\alpha \text{bus}_\pm = \alpha^+$. Then $|\alpha^+| \leq 1$ and $(\text{bus}_\pm \circ \alpha)^+ = 1$; therefore

$$\begin{aligned} |t_0 - t_1| &\geq |\alpha(t_0) - \alpha(t_1)| \geq \\ &\geq |\text{bus}_\pm \circ \alpha(t_0) - \text{bus}_\pm \circ \alpha(t_1)| = \\ &= |t_0 - t_1|. \end{aligned}$$

It follows that for any $t > 0$, the bus_\pm -gradient flows commute; that is,

$$\text{Flow}_{\text{bus}_+}^t \circ \text{Flow}_{\text{bus}_-}^t = \text{id}_{\mathcal{L}}.$$

Setting

$$\text{Flow}^t = \begin{cases} \text{Flow}_{\text{bus}_+}^t & \text{if } t \geq 0 \\ \text{Flow}_{\text{bus}_-}^t & \text{if } t \leq 0 \end{cases}$$

defines an \mathbb{R} -action on \mathcal{L} .

Consider the level set $\mathcal{L}' = \text{bus}_+^{-1}(0) = \text{bus}_-^{-1}(0)$; it is a closed convex subset of \mathcal{L} , and therefore forms an Alexandrov space. Consider the map $h: \mathcal{L}' \times \mathbb{R} \rightarrow \mathcal{L}$ defined by $h: (x, t) \mapsto \text{Flow}^t(x)$. Note that h is onto. Applying Lemma 6.6 for $\text{Flow}_{\text{bus}_+}^t$ and $\text{Flow}_{\text{bus}_-}^t$ shows that h is short and non-contracting at the same time; that is, h is an isometry. \square

Recall that according our definition the real line \mathbb{R} is a CBB(1) space. However most of geodesic CBB(1) spaces have diameter at most π ; see 3.6.

6.7. Exercise. *Suppose \mathcal{X} is a complete geodesic space. Show that $\text{Cone } \mathcal{X}$ is CBB(0) if and only if $\mathcal{X} \stackrel{\text{iso}}{=} \text{Susp}^k \mathcal{Y}$ for some integer $k \geq 0$ and a geodesic CBB(1) space \mathcal{Y} such that $|x - y| < \pi$ for any $x, y \in \mathcal{Y}$.*

???
(For example, if $\mathcal{Y} = \emptyset$, then $\mathcal{X} \stackrel{\text{iso}}{=} \mathbb{S}^k$; if \mathcal{Y} is a one-point space, then \mathcal{X} is a hemisphere.)

C Comments

The splitting theorem has an interesting history that starts with Stefan Cohn-Vossen [13]. Our proof is based on the idea of Jeff Cheeger and Detlef Gromoll [12].

Lecture 7

Dimension

A Polar vectors

Here we give a corollary of 4.12. It will be used to prove basic properties of the tangent space.

7.1. Anti-sum lemma. *Let \mathcal{L} be a complete geodesic CBB space and $p \in \mathcal{L}$.*

Given two vectors $u, v \in T_p$, there is a unique vector $w \in T_p$ such that

$$\langle u, x \rangle + \langle v, x \rangle + \langle w, x \rangle \geq 0$$

for any $x \in T_p$, and

$$\langle u, w \rangle + \langle v, w \rangle + \langle w, w \rangle = 0.$$

7.2. Exercise. *Suppose $u, v, w \in T_p$ are as in 7.1. Show that*

$$|w|^2 \leq |u|^2 + |v|^2 + 2 \cdot \langle u, v \rangle.$$

If T_p were geodesic, then the lemma would follow from the existence of the gradient, applied to the function $T_p \rightarrow \mathbb{R}$ defined by $x \mapsto -(\langle u, x \rangle + \langle v, x \rangle)$ which is concave. However, the tangent space T_p might fail to be geodesic; see Halbeisen's example [3].

Applying the above lemma for $u = v$, we have the following statement.

7.3. Existence of polar vector. *Let \mathcal{L} be a complete geodesic CBB space and $p \in \mathcal{L}$. Given a vector $u \in T_p$, there is a unique vector $u^* \in T_p$ such that $\langle u^*, u^* \rangle + \langle u, u^* \rangle = 0$ and u^* is polar to u ; that is, $\langle u^*, x \rangle + \langle u, x \rangle \geq 0$ for any $x \in T_p$.*

In particular, for any vector $u \in T_p$ there is a polar vector $u^* \in T_p$ such that $|u^*| \leq |u|$.

7.4. Example. Let \mathcal{L} be the upper half plane in \mathbb{E}^2 ; that is, $\mathcal{L} = \{(x, y) \in \mathbb{E}^2 \mid y \geq 0\}$. It is a complete geodesic CBB(0) space. For $p = 0$, the tangent space T_p can be canonically identified with \mathcal{L} . If $y > 0$, then $u = (x, y) \in T_p$ has many polar vectors; it includes $u^* = (-x, 0)$ which is provided by 7.3, but the vector $w = (-x, y)$ is polar as well.

Proof of 7.1. By 4.11, we can choose two sequences of points a_n, b_n such that

$$\begin{aligned} \mathbf{d}_p \text{dist}_{a_n}(w) &= -\langle \uparrow_{[pa_n]}, w \rangle \\ \mathbf{d}_p \text{dist}_{b_n}(w) &= -\langle \uparrow_{[pb_n]}, w \rangle \end{aligned}$$

for any $w \in T_p$ and $\uparrow_{[pa_n]} \rightarrow u/|u|$, $\uparrow_{[pb_n]} \rightarrow v/|v|$ as $n \rightarrow \infty$

Consider a sequence of functions

$$f_n = |u| \cdot \text{dist}_{a_n} + |v| \cdot \text{dist}_{b_n}.$$

Note that

$$(\mathbf{d}_p f_n)(x) = -|u| \cdot \langle \uparrow_{[pa_n]}, x \rangle - |v| \cdot \langle \uparrow_{[pb_n]}, x \rangle.$$

Thus we have the following uniform convergence for $x \in \Sigma_p$:

$$(\mathbf{d}_p f_n)(x) \rightarrow -\langle u, x \rangle - \langle v, x \rangle$$

as $n \rightarrow \infty$, According to 4.12, the sequence $\nabla_p f_n$ converges. Let

$$w = \lim_{n \rightarrow \infty} \nabla_p f_n.$$

By the definition of gradient,

$$\begin{aligned} \langle w, w \rangle &= \lim_{n \rightarrow \infty} \langle \nabla_p f_n, \nabla_p f_n \rangle = & \langle w, x \rangle &= \lim_{n \rightarrow \infty} \langle \nabla_p f_n, x \rangle \geq \\ &= \lim_{n \rightarrow \infty} (\mathbf{d}_p f_n)(\nabla_p f_n) = & & \geq \lim_{n \rightarrow \infty} (\mathbf{d}_p f_n)(x) = \\ &= -\langle u, w \rangle - \langle v, w \rangle, & & = -\langle u, x \rangle - \langle v, x \rangle. \end{aligned}$$

□

B Linear subspace of tangent space

7.5. Definition. Let \mathcal{L} be a complete geodesic CBB(κ) space, $p \in \mathcal{L}$ and $u, v \in T_p$. We say that vectors u and v are opposite to each other, (briefly, $u + v = 0$) if $|u| = |v| = 0$ or $\angle(u, v) = \pi$ and $|u| = |v|$.
The subcone

$$\text{Lin}_p = \{ v \in T_p : \exists w \in T_p \text{ such that } w + v = 0 \}$$

will be called the linear subspace of T_p .

7.6. Proposition. Let \mathcal{L} be a complete geodesic CBB space and $p \in \mathcal{L}$. Given two vectors $u, v \in T_p$, the following statements are equivalent:

- (a) $u + v = 0$;
- (b) $\langle u, x \rangle + \langle v, x \rangle = 0$ for any $x \in T_p$;
- (c) $\langle u, \xi \rangle + \langle v, \xi \rangle = 0$ for any $\xi \in \Sigma_p$.

Proof. The equivalence $(b) \Leftrightarrow (c)$ is trivial.

The condition $u + v = 0$ is equivalent to

$$\langle u, u \rangle = -\langle u, v \rangle = \langle v, v \rangle;$$

thus $(b) \Rightarrow (a)$.

Recall that T_p is CBB(0). Note that the hinges $[0_x^u]$ and $[0_x^v]$ are adjacent. By 2.11, $\angle[0_x^u] + \angle[0_x^v] = 0$; hence $(a) \Rightarrow (b)$. \square

7.7. Exercise. Let \mathcal{L} be a complete geodesic CBB space and $p \in \mathcal{L}$. Then for any three vectors $u, v, w \in T_p$, if $u + v = 0$ and $u + w = 0$ then $v = w$.

Let $u \in \text{Lin}_p$; that is, $u + v = 0$ for some $v \in T_p$. Given $s < 0$, let

$$s \cdot u := (-s) \cdot v.$$

So we can multiply any vector in Lin_p by any real number (positive and negative). By 7.7, this multiplication is uniquely defined; by 7.6; we have identity

$$\langle -v, x \rangle = -\langle v, x \rangle;$$

later we will see that it extends to a linear structure on Lin_p .

7.8. Exercise. Suppose $u, v, w \in T_p$ are as in 7.1. Show that

$$\langle u, x \rangle + \langle v, x \rangle + \langle w, x \rangle = 0$$

for any $x \in \text{Lin}_p$.

7.9. Exercise. Let \mathcal{L} be a complete geodesic CBB space, $p \in \mathcal{L}$ and $u \in \text{T}_p$. Suppose $u^* \in \text{T}_p$ is provided by 7.3; that is,

$$\langle u^*, u^* \rangle + \langle u, u^* \rangle = 0 \quad \text{and} \quad \langle u^*, x \rangle + \langle u, x \rangle \geq 0$$

for any $x \in \text{T}_p$. Show that

$$u = -u^* \iff |u| = |u^*|.$$

7.10. Theorem. Let p be a point in a complete geodesic $\text{CBB}(\kappa)$. Then Lin_p is isometric to a Hilbert space.

Proof. Note that Lin_p is a closed subset of T_p ; in particular, it is complete.

If any two vectors in Lin_p can be connected by a geodesic in Lin_p , then the statement follows from the splitting theorem (6.4). By Menger's lemma (1.4), it is sufficient to show that for any two vectors $x, y \in \text{Lin}_p$ there is a midpoint $w \in \text{Lin}_p$.

Choose $w \in \text{T}_p$ to be the anti-sum of $u = -\frac{1}{2} \cdot x$ and $v = -\frac{1}{2} \cdot y$; see 7.1. By 7.2 and 7.8,

$$\begin{aligned} |w|^2 &\leq \frac{1}{4} \cdot |x|^2 + \frac{1}{4} \cdot |y|^2 + \frac{1}{2} \cdot \langle x, y \rangle, \\ \langle w, x \rangle &= \frac{1}{2} \cdot |x|^2 + \frac{1}{2} \cdot \langle x, y \rangle, \\ \langle w, y \rangle &= \frac{1}{2} \cdot |y|^2 + \frac{1}{2} \cdot \langle x, y \rangle, \end{aligned}$$

It follows that

$$\begin{aligned} |x - w|^2 &= |x|^2 + |w|^2 - 2 \cdot \langle w, x \rangle \leq \\ &\leq \frac{1}{4} \cdot |x|^2 + \frac{1}{4} \cdot |y|^2 - \frac{1}{2} \cdot \langle x, y \rangle = \\ &= \frac{1}{4} \cdot |x - y|^2. \end{aligned}$$

That is, $|x - w| \leq \frac{1}{2} \cdot |x - y|$, and similarly $|y - w| \leq \frac{1}{2} \cdot |x - y|$. Therefore w is a midpoint of x and y . In addition we get equality

$$|w|^2 = \frac{1}{4} \cdot |x|^2 + \frac{1}{4} \cdot |y|^2 + \frac{1}{2} \cdot \langle x, y \rangle.$$

It remains to show that $w \in \text{Lin}_p$. Let w^* be the polar vector provided by 7.3. Note that

$$|w^*| \leq |w|, \quad \langle w^*, x \rangle + \langle w, x \rangle = 0, \quad \langle w^*, y \rangle + \langle w, y \rangle = 0.$$

The same calculation as above shows that w^* is a midpoint of $-x$ and $-y$ and

$$|w^*|^2 = \frac{1}{4} \cdot |x|^2 + \frac{1}{4} \cdot |y|^2 + \frac{1}{2} \cdot \langle x, y \rangle = |w|^2.$$

By 7.9, $w = -w^*$; hence $w \in \text{Lin}_p$. \square

7.11. Exercise. Let p be a point in a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} and $f = \text{dist}_p$. Denote by S the subset of points $x \in \mathcal{L}$ such that $|\nabla_x f| = 1$.

(a) Show that S is a dense G -delta set.

(b) Show that

$$\nabla_x f + \uparrow_{[xp]} = 0$$

for any $x \in S$; in particular, $\uparrow_{[xp]} \in \text{Lin}_x$.

(c) Show that if $|\nabla_x f| = 1$, then $\mathbf{d}_x f(w) = \langle \nabla_x f, w \rangle$ for any $w \in \text{T}_x$.

Note that 7.11b implies the following.

7.12. Corollary. Given a countable set of points X in a complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} there is a G -delta dense set $S \subset \mathcal{L}$ such that $\uparrow_{[sx]} \in \text{Lin}_s$ for any $s \in S$ and $x \in X$.

C Linear dimension

Suppose \mathcal{L} is a complete geodesic $\text{CBB}(\kappa)$ space. Let us define its linear dimension $\text{LinDim } \mathcal{L}$ as the least upper bound on integers m such that the Euclidean space \mathbb{E}^m is isometric to a subspace of the tangent space $\text{T}_p \mathcal{L}$ at some point $p \in \mathcal{L}$. If not stated otherwise, dimension of a CBB space is its linear dimension.

7.13. $(n+1)$ -comparison. Let \mathcal{L} be a complete geodesic $\text{CBB}(0)$ space. Then for any finite set of points $p, x_1, \dots, x_n \in \mathcal{L}$, there is a model configuration $\tilde{p}, \tilde{x}_1, \dots, \tilde{x}_n \in \mathbb{E}^m$ such that

$$|\tilde{p} - \tilde{x}_i|_{\mathbb{E}^m} = |p - x_i|_{\mathcal{L}} \quad \text{and} \quad |\tilde{x}_i - \tilde{x}_j|_{\mathbb{E}^m} \geq |x_i - x_j|_{\mathcal{L}}$$

for any i and j . Moreover, we can assume that $m \leq \text{LinDim } \mathcal{L}$.

Proof. By 7.12, we can choose a point p' arbitrarily close to p so that $\text{Lin}_{p'} \ni \uparrow_{[p'x_i]}$ for any i . Let us identify \mathbb{E}^m with a subspace of $\text{Lin}_{p'}$ spanned by $\uparrow_{[p'x_1]}, \dots, \uparrow_{[p'x_n]}$. Note that $m \leq \text{LinDim } \mathcal{L}$.

Set $\tilde{p}' = 0 \in \mathbb{E}^m$ and $\tilde{x}_i = |p' - x_n| \cdot \uparrow_{[p'x_n]} \in \mathbb{E}^m$ for every i . Note that

$$|\tilde{p}' - \tilde{x}_i|_{\mathbb{E}^m} = |p' - x_i|_{\mathcal{L}}$$

for every i . Applying the comparison $\angle[p'_{x_i}] \geq \tilde{\angle}(p'_{x_j})$, we get

$$|\tilde{x}_i - \tilde{x}_j|_{\mathbb{E}^m} \geq |x_i - x_j|_{\mathcal{L}}$$

for any i and j . Passing to a limit configuration as $p' \rightarrow p$ we get the result. \square

7.14. Exercise. Let \mathcal{L} is a complete geodesic $\text{CBB}(0)$ space. Suppose $\text{LinDim } \mathcal{L} = m < \infty$. Show that $T_p \mathcal{L} \stackrel{\text{iso}}{=} \mathbb{E}^m$ for a G -delta dense set of points $p \in \mathcal{L}$.

7.15. Exercise. Show that a 1-dimensional complete geodesic $\text{CBB}(\kappa)$ space is homeomorphic to a 1-dimensional manifold, possibly with non-empty boundary.

7.16. Exercise. Let \mathcal{L} be a complete geodesic $\text{CBB}(0)$ space.

Show that $\text{LinDim } \mathcal{L} \geq m$ if and only if for some $m + 2$ points $p, x_0, \dots, x_m \in \mathcal{L}$ we have

$$\tilde{\angle}(p_{x_j}) > \frac{\pi}{2}$$

for any pair $i \neq j$.¹

D Space of directions

A metric space \mathcal{X} will be called ℓ -geodesic if any two points $x, y \in \mathcal{X}$ such that $|x - y| < \ell$ can be connected by a geodesic. Note that geodesic spaces are ∞ -geodesic.

7.17. Theorem. Let \mathcal{L} be a finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space. Then for any point $p \in \mathcal{L}$, its space of directions Σ_p is a compact π -geodesic $\text{CBB}(1)$ space.

7.18. Exercise. Prove the following in the assumptions of 7.17.

- (a) The tangent space T_p is a proper geodesic space.
- (b) $\dim \Sigma_p = \dim \mathcal{L} - 1$.
- (c) If $\dim \mathcal{L} > 1$, then Σ_p is geodesic.

Using 7.18b, one can prove results for all finite-dimensional complete geodesic $\text{CBB}(\kappa)$ spaces via induction on its dimension. Such proofs will be indicated below.

Proof. Note that 4.3 and ?? imply that Σ_p is $\text{CBB}(1)$.

¹If $m = \text{LinDim } \mathcal{L}$ then the map $q \mapsto (|x_1 - q|, \dots, |x_m - q|)$ induces a bi-Lipschitz embedding of a neighborhood of p to \mathbb{E}^m . (We mention it without proof, altho it is not hard to prove.)

Compactness. Choose $\varepsilon > 0$; suppose \mathcal{L} is m -dimensional. Assume can choose n directions $\xi_1, \dots, \xi_n \in \Sigma_p$ such that $\angle(\xi_i, \xi_j) > \varepsilon$ for any $i \neq j$. Without loss of generality, we may assume that each direction is geodesic; that is, there is a point $x_i \in \mathcal{L}$ such that $\xi_i = \uparrow_{[px_i]}$.

Choose $y_i \in [px_i]$ such that $|p - y_i| = r$ for each i and small fixed $r > 0$. Since r is small, we can assume that $\tilde{\angle}(p_{y_i}^{y_i}) > \varepsilon$ for any $i \neq j$. By 7.12, we can choose p' arbitrarily close to p such that $\uparrow_{[p'y_i]} \in \text{Lin } p'$ for any i . Since $|p' - p|$ is small, $\tilde{\angle}(p_{y_i}^{y_i}) > \varepsilon$ for any $i \neq j$. By comparison,

$$\angle[p_{y_i}^{y_i}] > \varepsilon.$$

Therefore $n \leq \text{pack}_\varepsilon \mathbb{S}^{m-1}$.

Since \mathbb{S}^{m-1} is compact, $\text{pack}_\varepsilon \mathbb{S}^{m-1} < \infty$. By the definition, the space of directions Σ_p is complete. Applying 8.16, we get that Σ_p is compact.

Geodesicness. Now we will show that if Σ_p is compact, then it is π -geodesic; we will not use the finiteness of dimension directly.

Choose two geodesic directions $\xi = \uparrow_{[px]}$ and $\zeta = \uparrow_{[py]}$; let

$$\alpha = \frac{1}{2} \cdot \angle[p_y^x] = \frac{1}{2} \cdot |\xi - \zeta|_{\Sigma_p}.$$

Suppose $\alpha < \pi$. Let us show that it is sufficient to construct an almost midpoint $\mu = \uparrow_{[pz]}$ of ξ and ζ in Σ_p ; that is, we need to show that for any $\varepsilon > 0$ there is a geodesic $[pz]$ such that

$$\angle[p_z^x] \leq \alpha + \varepsilon \quad \text{and} \quad \angle[p_z^y] \leq \alpha + \varepsilon.$$

Indeed, once it is done, the compactness of Σ_p can be used to get an actual midpoint for any two directions in Σ_p . After that Menger's lemma (1.4) will finish the proof.

Choose a sequence of small positive numbers $r_n \rightarrow 0$. Consider sequences $x_n \in [px]$ and $y_n \in [py]$ such that $|p - x_n| = |p - y_n| = r_n$. Let m_n be a midpoint of $[x_n y_n]$.

Since Σ_p is compact, we can pass to a sequence of r_n such that $\uparrow_{[pm_n]}$ converges; denote its limit by μ . Choose a geodesic $[pz]$ that runs at small angle from μ . Let us show that $\uparrow_{[pz]}$ is the needed almost midpoint.

Evidently, $\tilde{\angle}(p_{m_n}^{x_n}) = \tilde{\angle}(p_{m_n}^{y_n})$. By 2.5, we have

$$\tilde{\angle}(p_{m_n}^{x_n}) + \tilde{\angle}(p_{m_n}^{y_n}) \leq \tilde{\angle}(p_{y_n}^{x_n}).$$

Let $z_n \in [pz]$ be the point such that $|p - z_n| = |p - m_n|$. By construction, for all large n , we have $\angle[p_{m_n}^z] \approx 0$ with arbitrary small given error. By comparison, the value $\frac{|z_n - m_n|}{|p - z_n|}$ can be assumed to

be arbitrary small for all large n . Applying this observation and the definition of angle measure, we also have the following approximations

$$\begin{aligned}\tilde{\angle}(p_{y_n}^{x_n}) &\approx \angle[p_{y_n}^{x_n}], \\ \tilde{\angle}(p_{m_n}^{x_n}) &\approx \tilde{\angle}(p_{z_n}^{x_n}) \approx \angle[p_{z_n}^{x_n}], \\ \tilde{\angle}(p_{y_n}^{m_n}) &\approx \tilde{\angle}(p_{y_n}^{z_n}) \approx \angle[p_{y_n}^{z_n}],\end{aligned}$$

again, with arbitrary given error and all large n . It follows that $\uparrow_{[pz]}$ is an almost midpoint of $\uparrow_{[px]}$ and $\uparrow_{[py]}$, as required. \square

Notice that CBB comparison gives lower bounds on $\angle[p_z^x]$ and $\angle[p_z^y]$, but in the proof we needed upper bounds which were obtained by from the definition of angle measure and compactness of space of directions.

E Comments

Corollary 7.12 is the key to the proof that *all reasonable definitions of dimension give the same result on complete geodesic CBB spaces*. More precisely, we have the following theorem [3, 15.16].

7.19. Theorem. *For any complete length CBB(κ) space \mathcal{L} , we have*

$$\text{LinDim } \mathcal{L} = \text{TopDim } \mathcal{L} = \text{HausDim } \mathcal{L},$$

where TopDim and HausDim stands for Lebesgue covering dimension and Hausdorff dimension, respectively.

7.20. Open question. *Let p be a point in a geodesic CBB space \mathcal{L} . Suppose that $0 \neq v \in \text{Lin}_p$. Is it true that the tangent space T_p splits in the direction of v ?*

Halbeisen's example [3, 23] shows that compactness of space of directions is essential in the proof that space of directions is π -geodesic (see 7.17).

7.21. Open question. *Let \mathcal{L} be a proper length CBB(κ) space. Is it true that for any $p \in \mathcal{L}$, the tangent space T_p is a length space?*


Lecture 8

Limit spaces

A Hausdorff convergence

8.1. Definition. Let A_1, A_2, \dots be a sequence of closed sets in a metric space \mathcal{X} . We say that the sequence A_n converges to a closed set A_∞ in the sense of Hausdorff if, for any $x \in \mathcal{X}$, we have $\text{dist}_{A_n}(x) \rightarrow \text{dist}_{A_\infty}(x)$ as $n \rightarrow \infty$.

For example, suppose \mathcal{X} is the Euclidean plane and A_n is the circle with radius n and center at the point $(0, n)$; it converges to the x -axis.



Further, consider the sequence of one-point sets $B_n = \{(n, 0)\}$ in the Euclidean plane. It converges to the empty set; indeed, for any point x we have $\text{dist}_{B_n}(x) \rightarrow \infty$ as $n \rightarrow \infty$ and $\text{dist}_\emptyset(x) = \infty$ for any x .

The following exercise is an extension of the so-called Blaschke selection theorem to our version of Hausdorff convergence.

8.2. Exercise. Let \mathcal{X} be a proper metric space and A_1, A_2, \dots be a sequence of closed sets in \mathcal{X} . Show that the sequence A_1, A_2, \dots has a convergent subsequence in the sense of Hausdorff.

B Gromov–Hausdorff convergence

Let $\mathcal{X}_1, \mathcal{X}_2, \dots$ be a sequence of metric spaces. Suppose that there is a metric on the disjoint union


$$\mathbf{X} = \bigsqcup_{n \in \mathbb{N} \cup \{\infty\}} \mathcal{X}_n$$

that satisfies the following property:

8.3. Property. *The restriction of metric on each \mathcal{X}_n and \mathcal{X}_∞ coincides with its original metric, and $\mathcal{X}_n \rightarrow \mathcal{X}_\infty$ as subsets in \mathbf{X} in the sense of Hausdorff.*

In this case we say that the metric on \mathbf{X} defines a convergence $\mathcal{X}_n \rightarrow \mathcal{X}_\infty$ in the sense of Gromov–Hausdorff. The space \mathbf{X} makes it possible to talk about limits of sequences $x_n \in \mathcal{X}_n$ as $n \rightarrow \infty$, as well as weak limits of a sequence of Borel measures μ_n on \mathcal{X}_n and so on.

A limit space for this convergence is not uniquely defined. For example, if each space \mathcal{X}_n in the sequence is isometric to the half-line, then its limit might be isometric to the half-line or the whole line. The first convergence is evident and the second could be guessed from the diagram.



The following exercise states that if limit compact, then it is unique up to isometry.

8.4. Exercise. *Let $\mathcal{X}_1, \mathcal{X}_2, \dots$ be a sequence of metric spaces. Suppose \mathcal{X}_∞ and \mathcal{X}'_∞ are limit spaces for some Gromov–Hausdorff convergences of \mathcal{X}_n . Assume \mathcal{X}_∞ is compact, show that it is isometric to \mathcal{X}'_∞ .*

8.5. Exercise. *Let $\mathcal{X}_1, \mathcal{X}_2, \dots$ be a sequence of geodesic metric spaces. Suppose $\mathcal{X}_n \rightarrow \mathcal{X}_\infty$ is a convergence in the sense of Gromov–Hausdorff. Assume \mathcal{X}_∞ is proper, show that it is geodesic.*

Compact case. The Gromov–Hausdorff convergence defines a metrizable topology of the set of isometry classes of compact metric spaces; this space will be denoted by GH.

For two metric spaces \mathcal{X} and \mathcal{Y} , we write $\mathcal{X} \leq \mathcal{Y} + \varepsilon$ if there is a map $f: \mathcal{X} \rightarrow \mathcal{Y}$ such that

$$|x - x'|_{\mathcal{X}} \leq |f(x) - f(x')|_{\mathcal{Y}} + \varepsilon$$

for any $x, x' \in \mathcal{X}$.

8.6. Exercise. Let $\mathcal{X}_1, \mathcal{X}_2, \dots$ and \mathcal{X}_∞ are compact metric spaces. Show that there is a Gromov–Hausdorff convergence $\mathcal{X}_n \rightarrow \mathcal{X}_\infty$ if and only if for some sequence $\varepsilon_n \rightarrow 0$, we have

$$\mathcal{X}_\infty \leq \mathcal{X}_n + \varepsilon_n \quad \text{and} \quad \mathcal{X}_n \leq \mathcal{X}_\infty + \varepsilon_n.$$

Pointed convergence. Often the isometry class of the limit can be fixed by marking a point p_n in each space \mathcal{X}_n , it is called pointed Gromov–Hausdorff convergence — we say that (\mathcal{X}_n, p_n) converges to $(\mathcal{X}_\infty, p_\infty)$ if there is a metric on \mathbf{X} as in 8.3 such that $p_n \rightarrow p_\infty$. For example, the sequence $(\mathcal{X}_n, p_n) = (\mathbb{R}_+, 0)$ converges to $(\mathbb{R}_+, 0)$, while $(\mathcal{X}_n, p_n) = (\mathbb{R}_+, n)$ converges to $(\mathbb{R}, 0)$.

The pointed convergence works nicely for proper metric spaces; the following theorem is an analog of Gromov’s selection theorem for this convergence.

8.7. Theorem. Let $\mathcal{X}_1, \mathcal{X}_2, \dots$ be a sequence of proper metric spaces with marked points $p_n \in \mathcal{X}_n$. Assume that for any $R > 0$, the balls $\overline{B}[p_n, R]_{\mathcal{X}_n}$ are uniformly totally bounded; that is, for each R and $\varepsilon > 0$ there is N such that $\overline{B}[p_n, R]_{\mathcal{X}_n}$ can be covered by at most N ε -balls. Then there is a pointed Gromov–Hausdorff convergence $(\mathcal{X}_n, p_n) \rightarrow (\mathcal{X}_\infty, p_\infty)$ such that \mathcal{X}_∞ is proper.

Moreover, the space with marked point $(\mathcal{X}_\infty, p_\infty)$ is uniquely defined up to isometry that respects the marked point.

8.8. Exercise. Let p be a point in a finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} , and let $\lambda_n \rightarrow \infty$. Show that there is a pointed Gromov–Hausdorff convergence $(\lambda \cdot \mathcal{L}, p) \rightarrow (\mathbb{T}_p, 0)$ in the sense of Gromov–Hausdorff. Moreover, the convergence can be chosen so that

$$\lambda_n \cdot \gamma(t/\lambda_n) \rightarrow t \cdot \gamma^+(0)$$

for any geodesic γ that starts at p .

C Survival of curvature bounds

8.9. Theorem. For any $\kappa \in \mathbb{R}$, the $\text{CBB}(\kappa)$ spaces form a closed subset in GH.

Proof. Suppose $\mathcal{X}_n \rightarrow \mathcal{X}_\infty$ in the sense of Gromov–Hausdorff. Choose a quadruple of points $p_\infty, x_\infty, y_\infty, z_\infty \in \mathcal{X}_\infty$.

By the definition of Gromov–Hausdorff convergence, we can choose a quadruple $p_n, x_n, y_n, z_n \in \mathcal{X}_n$ such that each of 6 distances between pairs of p_n, x_n, y_n, z_n converge to the distance between the corresponding pairs of $p_\infty, x_\infty, y_\infty, z_\infty$.

By the assumption, $\text{CBB}(\kappa)$ comparison holds for each p_n, x_n, y_n, z_n . Passing to the limit, we get the $\text{CBB}(\kappa)$ comparison for $p_\infty, x_\infty, y_\infty, z_\infty$. \square

8.10. Exercise. Suppose that a sequence $\mathcal{L}_1, \mathcal{L}_2, \dots$ of compact geodesic $\text{CBB}(0)$ space converges to \mathcal{L}_∞ . Show that

$$\text{LinDim } \mathcal{L}_\infty \leq \lim_{n \rightarrow \infty} \text{LinDim } \mathcal{L}_n.$$

D Volume

Fix a positive integer m . The m -dimensional Hausdorff measure of a Borel set B in a metric space will be called its m -volume; it will be denoted by $\text{vol}_m B$. We assume that the Hausdorff measure is calibrated so that the unit cube in \mathbb{E}^m has unit volume.

This definition will be used mostly in m -dimensional complete geodesic $\text{CBB}(\kappa)$ spaces. In this case, we may write $\text{vol } B$ instead of $\text{vol}_m B$.

8.11. Bishop–Gromov inequality. Let \mathcal{L} be a complete geodesic $\text{CBB}(0)$ space. Suppose $\mathcal{L} = m < \infty$. Then

$$\text{vol } B(p, R) \leq \omega_m \cdot R^m,$$

where ω_m denotes the volume of the unit ball in \mathbb{E}^m . Moreover, the function

$$R \mapsto \frac{\text{vol } B(p, R)}{R^m}$$

is nonincreasing.

Proof. Given $x \in \mathcal{L}$ choose a geodesic path γ_x from p to x . Let $\log_p : \mathcal{L} \rightarrow T_p$ be defined by $\log_p : x \mapsto \gamma_x^+(0)$. By comparison, \log_p is distance-noncontracting. Note that \log_p maps $B(p, R)_\mathcal{L}$ to $B(0, R)_{T_p}$.

If $T_p \stackrel{\text{iso}}{=} \mathbb{E}^m$, then $\text{vol } B(0, R)_{T_p} = \omega_m \cdot R^m$, and the first statement follows. Otherwise, by 7.14, we can find a point p' arbitrarily close to

p such that $T_{p'} \stackrel{\text{iso}}{=} \mathbb{E}^m$. If $\varepsilon > |p - p'|$, then $B(p, R) \subset B(p', R + \varepsilon)$. Therefore,

$$\text{vol } B(p, R) \leq \omega_m \cdot (R + \varepsilon)^m$$

for any $\varepsilon > 0$. Hence the first statement follows in the general case.

For the second statement, choose $0 < R_1 < R_2$. Consider the map $w: \mathcal{L} \rightarrow \mathcal{L}$ defined by $w: x \mapsto \gamma_x(\frac{R_1}{R_2})$. By comparison,

$$|w(x) - w(y)| \geq \frac{R_1}{R_2} \cdot |x - y|.$$

It follows that

$$\text{vol } B \geq \left(\frac{R_1}{R_2}\right)^m \cdot \text{vol } A$$

if $B \supset w(A)$. Observe that

$$B(p, R_1) \supset w[B(p, R_2)]$$

— hence the result. \square

8.12. Exercise. *Let*

8.13. Exercise. *Given a positive integer m , show that any complete geodesic m -dimensional $\text{CBB}(0)$ space with bounded diameter is compact.*

E Gromov's selection theorem

8.14. Theorem. *The set of complete geodesic m -dimensional $\text{CBB}(\kappa)$ spaces with diameter $\leq D$ forms a compact subset in GH ; here we assume that m is finite.*

The theorem implies that one can select a converging sequence from any sequence of spaces from the described set.

Let X be a subset of a metric space \mathcal{W} . Recall that a set $Z \subset \mathcal{W}$ is called ε -net of X if for any point $x \in X$, there is a point $z \in Z$ such that $|x - z| < \varepsilon$.

We will use the following characterization of compact sets: *a closed subset X of a complete metric space is compact if and only if X admits a finite ε -net for any $\varepsilon > 0$.* The following statement is slightly more general.

8.15. Claim. *A closed subset X of a complete metric space is compact if and only if it admits a compact ε -net for any $\varepsilon > 0$.*

Proof. Let Z be a compact ε -net of X . Since Z is compact, it admits a finite ε -net F . Note that F is a $2 \cdot \varepsilon$ -net of X . Since $\varepsilon > 0$ is arbitrary, we get the result. \square

Let $\text{pack}_\varepsilon \mathcal{X}$ be the exact upper bound on the number of points $x_1, \dots, x_n \in \mathcal{X}$ such that $|x_i - x_j| \geq \varepsilon$ if $i \neq j$.

If $n = \text{pack}_\varepsilon \mathcal{X} < \infty$, then the collection of points x_1, \dots, x_n is called a maximal ε -packing.

8.16. Exercise. Show that any maximal ε -packing x_1, \dots, x_n is an ε -net. Conclude that a complete metric space \mathcal{X} is compact if and only if $\text{pack}_\varepsilon \mathcal{X} < \infty$ for any $\varepsilon > 0$.

Proof of 8.14. Denote by \mathbf{K} the set of isometry classes of complete geodesic CBB(0) spaces with dimension $\leq m$ and diameter $\leq D$. By 8.9, 8.5, and 7.16, \mathbf{K} is a closed subset of GH.

Choose a space $\mathcal{L} \in \mathbf{K}$; suppose $x_1, \dots, x_n \in \mathcal{L}$ is a collection of points such that $|x_i - x_j| \geq \varepsilon$ for all $i \neq j$. Note that the balls $B_i = B(x_i, \frac{\varepsilon}{2})$ do not overlap.

By Bishop–Gromov inequality, we get

$$\text{vol } B_i \geq \left(\frac{\varepsilon}{2 \cdot D}\right)^m \cdot \text{vol } \mathcal{L}$$

for any i and any small $\varepsilon > 0$. It follows that $n \leq \left(\frac{2 \cdot D}{\varepsilon}\right)^m$; that is,

$$\text{pack}_\varepsilon \mathcal{L} \leq N(\varepsilon) := \left(\frac{2 \cdot D}{\varepsilon}\right)^m$$

for all small $\varepsilon > 0$.

Choose a maximal ε -packing $x_1, \dots, x_n \in \mathcal{L}$. By 8.16, $\mathcal{F}_\varepsilon := \{x_1, \dots, x_n\}$ is an ε -net of \mathcal{L} . Observe that $|\mathcal{F}_\varepsilon - \mathcal{L}|_{\text{GH}} \leq \varepsilon$. Further, note that the set \mathbf{F}_ε of finite metric spaces with diameter $\leq D$ and at most $N(\varepsilon)$ points forms a compact subset in GH.

Summarizing, for any $\varepsilon > 0$ we can find a compact ε -net $\mathbf{F}_\varepsilon \subset \text{GH}$ of \mathbf{K} . It remains to apply ?? and 8.15. \square

8.17. Exercise. Let \mathcal{L} be complete geodesic CBB(0) spaces with dimension m and diameter $\leq D$. Suppose $\text{vol } \mathcal{L} \geq v_0 > 0$. Show that

$$\text{pack}_\varepsilon \mathcal{L} \geq \frac{c}{\varepsilon^m}$$

for some constant $c = c(m, D, v_0) > 0$.

Conclude that if \mathcal{L}_n is a sequence of complete geodesic m -dimensional CBB(0) spaces with diameter $\leq D$ and volume $\geq v_0$, then its Gromov–Hausdorff limit \mathcal{L}_∞ (if it is defined) has linear dimension m .¹

¹A stronger statement holds

$$\text{vol}_m \mathcal{L}_\infty = \lim_{n \rightarrow \infty} \text{vol}_m \mathcal{L}_n;$$

F Comments

Let us state a version of Bishop–Gromov inequality for curvature bound $\kappa = \pm 1$. Its proof is more technical. A weaker inequality can be obtained by applying the same argument as above, and it is sufficient to extend the rest of the result in the lecture.

8.18. Bishop–Gromov inequality. *Given a point p in a complete geodesic m -dimensional $\text{CBB}(\pm 1)$ space consider the function $v(R) = \text{vol}_m B(p, R)$; denote by $\tilde{v}(R)$ the volume of R ball in the corresponding m -dimensional model space (the unit m -dimensional sphere or the m -dimensional Lobachevsky space). Then*

$$v(R) \leq \tilde{v}(R)$$

and the function

$$R \mapsto \frac{v(R)}{\tilde{v}(R)}$$

is nonincreasing; in the $\text{CBB}(1)$ case, one has to assume that $R < \pi$.

The same inequality holds for complete m -dimensional Riemannian manifolds with Ricci curvature $\geq (m-1) \cdot \kappa$.

Gromov’s selection theorem is the main source of applications of CBB spaces. The homotopy-type finiteness theorem (9.5) in the next lecture illustrates this technique.

A version Gromov’s selection theorem (as well as Bishop–Gromov inequality) holds for m -dimensional Riemannian manifolds with a lower bound on Ricci curvature. It motivates the study of the so-called $\text{CD}(K, m)$ spaces; CD stands for curvature-dimension condition. This theory has serious applications in Alexandrov geometry; in particular, it provides a version of Liouville theorem about phase-space volume of geodesic flow in Alexandrov space [9].

in other words, if $\mathbf{K} \subset \text{GH}$ denotes the set of isometry classes of all compact geodesic $\text{CBB}(\kappa)$ spaces with dimension $\leq m$, then the function $\text{vol}_m: \mathbf{K} \rightarrow \mathbb{R}$ is continuous.

Lecture 9

Homotopy finiteness theorem

A Controlled concavity

While CBB spaces have plenty of semiconcave functions (for example dist_p^2), it is not at all easy to construct a strictly concave one. The following construction was found by Grigori Perelman [43, 46].

9.1. Theorem. *Let \mathcal{L} be a complete finite-dimensional geodesic CBB(κ) space. Then for any point $p \in \mathcal{L}$, there is a strictly concave function f defined in an open neighborhood of p .*

Moreover, given $0 \neq v \in T_p$, the differential, $\mathbf{d}_p f$, can be chosen arbitrarily close to $x \mapsto -\langle v, x \rangle$.

Sketch. Consider the real-to-real function

$$\varphi_{r,c}(x) = (x - r) - c(x - r)^2/r,$$

so we have

$$\varphi_{r,c}(r) = 0, \quad \varphi'_{r,c}(r) = 1 \quad \varphi''_{r,c}(r) = -2c/r.$$

Let γ be a unit-speed geodesic, fix a point q and let

$$\alpha(t) = \angle(\gamma^+(t), \uparrow_{[\gamma(t)q]}).$$

If $r > 0$ is sufficiently small and $|q - \gamma(t)|$ is sufficiently close to r , then direct calculations show that

$$(\varphi_{r,c} \circ \text{dist}_q \circ \gamma)''(t) \leq \frac{3 - c \cdot \cos^2[\alpha(t)]}{r}.$$

Now, assume $\{q_1, \dots, q_N\}$ is a finite set of points such that $|p - q_i| = r$ for any i . For a geodesic γ , set $\alpha_i(t) = \angle(\gamma^+(t), \uparrow_{[\gamma(t)q_i]})$. Assume we have a collection $\{q_i\}$ such that for any geodesic γ in $B(p, \varepsilon)$ we have $\max_i \{|\alpha_i(t) - \frac{\pi}{2}|\} \geq \varepsilon > 0$. Then taking in the above inequality $c > 3N/\cos^2 \varepsilon$, we get that the function

$$f = \sum_i \varphi_{r,c} \circ \text{dist}_{q_i}$$

is strictly concave in $B(p, \varepsilon')$ for some positive $\varepsilon' < \varepsilon$.

To construct the needed collection $\{q_i\}$, note that for small $r > 0$ one can choose $N \geq c/\delta^{(m-1)}$ points $\{q_i\}$ such that $|p - q_i| = r$ and $\angle(p, q_i) > \delta$ (here $c = c(\Sigma_p) > 0$). On the other hand, suppose γ runs from x to y . If $|\alpha_i(t) - \frac{\pi}{2}| < \varepsilon \ll \delta$, then applying the $(n+1)$ -comparison to $\gamma(t)$, x , y and all $\{q_i\}$ we get that $N \leq c(m)/\delta^{(m-2)}$. Therefore, for small $\delta > 0$ and yet smaller $\varepsilon > 0$, the set $\{q_i\}$ forms the needed collection.

If r is small, then points q_i can be chosen so that all directions $\uparrow_{[pq_i]}$ will be ε -close to a given direction ξ and therefore the second property follows. \square

Note that in 9.1 the function f can be chosen to have maximum value 0 at p , $f(p) = 0$ and with $\mathbf{d}_p f(x) \approx -|x|$. It can be constructed by taking the minimum of the functions in the theorem. Then the set $\Omega = \{x \in \mathcal{L} : f(x) \geq -\varepsilon\}$ forms an open convex neighborhood of p for any small $\varepsilon > 0$, so we get the following.

9.2. Corollary. *Any point p of a complete finite-dimensional geodesic CBB(0) space admits an arbitrary small convex neighborhood Ω and a strictly concave function f defined in a neighborhood of the closure $\bar{\Omega}$ such that p is the maximum point of f and $f|_{\partial\Omega} = 0$.*

B Liftings

Suppose that the Gromov–Haudorff distance $|\mathcal{L} - \mathcal{L}'|_{\text{GH}}$ is sufficiently small, so we may think that both spaces \mathcal{L} and \mathcal{L}' lie on small Haudorff distance in an ambient metric space \mathcal{W} . In particular, for any point $p \in \mathcal{L}$, we can choose a point $p' \in \mathcal{L}'$ such that $|p - p'|_{\mathcal{W}} < \varepsilon$ for small $\varepsilon > 0$; the point p' will be called an lifting (or ε -lifting) of p in \mathcal{L}' . We may choose a lifting $p' \in \mathcal{L}'$ for every point $p \in \mathcal{L}$, in this case the map $p \mapsto p'$ is called a (ε) -lifting map.


Note that the lifting is not uniquely defined. The lifting maps is not assumed to be continuous. When we talk about liftings, we assume that $\varepsilon > 0$, the inclusions $\mathcal{L}, \mathcal{L}' \hookrightarrow \mathcal{W}$, as well as \mathcal{W} are chosen.

Choose a compact geodesic m -dimensional $\text{CBB}(\kappa)$ space \mathcal{L} . Suppose \mathcal{L}' is another compact geodesic m -dimensional $\text{CBB}(\kappa)$ space such that $|\mathcal{L} - \mathcal{L}'|_{\text{GH}}$ is sufficiently small — smaller than some $\varepsilon = \varepsilon(\mathcal{L}) > 0$. Then the construction in \mathcal{L} from the previous section can be repeated in \mathcal{L}' for the liftings of all points and the same function φ . It produces a strictly concave function defined in a controlled neighborhood of the lifting p' of p .

The result of this and related constructions will be called liftings, say we can talk about a lifting from \mathcal{L} to \mathcal{L}' of a function provided by 9.1 (if the Gromov–Hausdorff distance $|\mathcal{L} - \mathcal{L}'|_{\text{GH}}$ is small, then these liftings are strictly concave) and a lifting of a convex neighborhood from 9.2. Here one cannot use 9.1 and 9.2 as black boxes — one has to understand the construction, but it is straightforward.

C Nerves

Let $\{\Omega_1, \dots, \Omega_k\}$ be a finite open cover of a compact metric space \mathcal{X} . Consider an abstract simplicial complex \mathcal{N} , with one vertex v_i for each set Ω_i such that a simplex with vertices v_{i_1}, \dots, v_{i_m} is included in \mathcal{N} if the intersection $\Omega_{i_1} \cap \dots \cap \Omega_{i_m}$ is nonempty. The obtained simplicial



complex \mathcal{N} is called the nerve of the covering $\{\Omega_i\}$. Evidently \mathcal{N} is a finite simplicial complex — it is a subcomplex of a simplex with the vertices $\{v_1, \dots, v_k\}$. Recall that Star_{v_i} denotes the union of all simplices in \mathcal{N} that shares vertex v_i .

The next statement follows from [25, 4G.3].

9.3. Nerve theorem. *Let $\{\Omega_1, \dots, \Omega_k\}$ be an open cover of a compact metric space \mathcal{X} and let \mathcal{N} be the corresponding nerve with vertices $\{v_1, \dots, v_k\}$. Suppose that every nonempty finite intersection $\Omega_{\alpha_1} \cap \dots \cap \Omega_{\alpha_k}$ is contractible. Then \mathcal{X} is homotopy equivalent to the nerve \mathcal{N} of the cover.*

Moreover homotopy equivalences $a: \mathcal{X} \rightarrow \mathcal{N}$ and $b: \mathcal{N} \rightarrow \mathcal{X}$ can be chosen so that if $x \in \Omega_i$, then $a(x) \in \text{Star}_{v_i}$, and if $y \in \mathcal{N}$ lies in the

simplex with vertices v_{i_1}, \dots, v_{i_m} , then $b(y) \in \Omega_{i_1} \cup \dots \cup \Omega_{i_m}$.

D Homotopy stability

9.4. Theorem. *Let $\mathcal{L}_1, \mathcal{L}_2, \dots, \mathcal{L}_\infty$ be a sequence of m -dimensional complete geodesic $\text{CBB}(\kappa)$ spaces and $m < \infty$. Suppose $\mathcal{L}_n \xrightarrow{\text{GH}} \mathcal{L}_\infty$ as $n \rightarrow \infty$. Then there is a \mathcal{L}_∞ is homotopically equivalent to \mathcal{L}_n for all large n .*

Moreover, given $\varepsilon > 0$ there are maps $h_n: \mathcal{L}_\infty \rightarrow \mathcal{L}_n$ that are homotopy equivalences and ε -liftings for all large n .

Applying this theorem with the Gromov's selection theorem (8.14) and Exercise 8.17, we get the following.

9.5. Theorem. *There are only finitely many homotopy types of m -dimensional $\text{CBB}(\kappa)$ spaces with diameter $\leq D$ and volume $\geq v_0$; here we assume that an integer m , and $v_0, D \in \mathbb{R}$ such that $v_0 > 0$ are given.*

Proof of 9.5 modulo 9.4. Assume the contrary, then we can choose a sequence of spaces $\mathcal{L}_1, \mathcal{L}_2, \dots$ that have different homotopy types and satisfy the assumptions of the theorem. By Gromov's compactness theorem, we can assume that \mathcal{L}_n converges to say \mathcal{L}_∞ in the sense of Gromov-Hausdorff.

By 8.17, $\text{LinDim } \mathcal{L}_\infty = m$. It remains to apply 9.4. □

Proof of 9.4. Since \mathcal{L}_∞ is compact, applying 9.2, we can find a finite open cover of \mathcal{L}_∞ by convex open sets $\Omega_1, \dots, \Omega_k$ such that for each Ω_i there is a strictly concave function f_i that is defined in a neighborhood of $\bar{\Omega}_i$ and such that $f_i|_{\partial\Omega_i} = 0$.

Subtracting from functions f_i some small value $\varepsilon > 0$, we can ensure that $\bigcap_{i \in S} \Omega_i \neq \emptyset$ if and only if $\bigcap_{i \in S} \bar{\Omega}_i \neq \emptyset$.

Suppose that $W = \bigcap_{i \in S} \Omega_i \neq \emptyset$. Then W is contractible. Indeed the function

$$f_S := \min_{i \in S} f_i$$

is strictly concave and it vanished on the boundary of W . The f_S -gradient flow $(t, x) \mapsto \text{Flow}_{f_S}^t(x)$ defines a homotopy $[0, \infty) \times W \rightarrow W$. Note that $\text{Flow}_{f_S}^t(x)$ converges to the (necessarily unique) maximum point of f_S as $t \rightarrow \infty$. Therefore, in the obtained homotoly we can parametrize $[0, \infty)$ by $[0, 1)$ and extend the homotopy by continuously to $[0, 1]$; this way we get that W is contractible. In other words, the cover $\{\Omega_1, \dots, \Omega_k\}$ meets the assumptions of the nerve theorem. Therefore \mathcal{L}_∞ is homotopy equivalent to the nerve \mathcal{N} of the cover.

The functions f_i and sets Ω_i can be lifted to \mathcal{L}_n keeping its properties for all large n . More precisely, there are liftings $f_{i,n}$ of all f_i to \mathcal{L}_n which are strictly concave for all large n and such that $\Omega_{i,n} = \{x \in \mathcal{L}_n : f_{i,n}(x) \geq 0\}$ is a compact convex set and $\Omega_{i,n} = \{x \in \mathcal{L}_n : f_{i,n}(x) > 0\}$ is an open convex set for each i .

Notice that $\{\Omega_{1,n}, \dots, \Omega_{k,n}\}$ is an open cover of \mathcal{L}_n for all large n . Indeed suppose we have $p_n \in \mathcal{L}_n \setminus (\Omega_{1,n} \cup \dots \cup \Omega_{k,n})$ for arbitrary large n . Since \mathcal{L}_∞ is compact, there is a limit point $p_\infty \in \mathcal{L}_\infty$ for a subsequence of p_n . But $p_\infty \in \Omega_i$ for some i and therefore $p_n \in \Omega_{i,n}$ for arbitrary large n — a contradiction.

In a similar fashion, we can show that if n is large, then any collection $\{\Omega_{i,n}\}_{i \in S}$ has a common point in \mathcal{L}_n if and only if $\{\Omega_i\}_{i \in S}$ has a common point in \mathcal{L}_∞ . Here we have to use that $\bigcap_{i \in S} \Omega_i \neq \emptyset$ if and only if $\bigcap_{i \in S} \bar{\Omega}_i \neq \emptyset$.

It follows that for any large n the following two covers the same nerve

- ◇ $\{\Omega_1, \dots, \Omega_k\}$ of \mathcal{L}_∞ and
- ◇ $\{\Omega_{1,n}, \dots, \Omega_{k,n}\}$ of \mathcal{L}_n .

Therefore, \mathcal{L}_n is homotopy equivalent to \mathcal{N} for all large n — a contradiction. \square

E Comments

Let us list some results that can be proved by applying Gromov's selection theorem in the same fashion as in the proof of homotopy-type finiteness theorem (9.5). The following theorem can be proved using this technique, altho Gromov's original proof [19] did not use Alexandrov geometry directly.

9.6. Betti-number theorem. *There is a constant $c = c(m, D, \kappa)$ such that*

$$\beta_0(M) + \beta_1(M) + \dots + \beta_m(M) \leq c$$

for any closed m -dimensional Riemannian manifold M with curvature $\geq \kappa$ and diameter $\leq D$. Here $\beta_i(M)$ denotes i^{th} Betti number of M .

The following result of the second author [52], and it uses the same technique.

9.7. Scalar curvature bound. *There is a constant $c = c(m, D, \kappa)$ such that*

$$\int_M \text{Sc} \leq c$$

for any closed m -dimensional Riemannian manifold M with curvature $\geq \kappa$ and diameter $\leq D$. Here Sc denotes the scalar curvature.

The following theorem is a more exact version of 9.4. It will play an important role in the following lecture.

9.8. Stability theorem. *Let $\mathcal{L}_1, \mathcal{L}_2, \dots, \mathcal{L}_\infty$ be a sequence of m -dimensional complete geodesic $\text{CBB}(\kappa)$ spaces and $m < \infty$. Suppose $\mathcal{L}_n \xrightarrow{\text{GH}} \mathcal{L}_\infty$ as $n \rightarrow \infty$. Then there is a \mathcal{L}_∞ is homotopically equivalent to \mathcal{L}_n for all large n .*

Moreover, given $\varepsilon > 0$ there are maps $h_n: \mathcal{L}_\infty \rightarrow \mathcal{L}_n$ that are homeomorphisms and ε -liftings for all large n .

This theorem was proved by Grigori Perelman [45]; the proof was rewritten with more details by the first author [28]. Around the same time, he made informal announcement that the homeomorphisms in the theorem can be made bi-Lipschitz with constants that depend on \mathcal{L}_∞ ; but the proof was not written and it save to consider it as a conjecture.

The last statement in the theorem implies the following finiteness result.

9.9. Homeomorphism-type finiteness. *There are only finitely many homeomorphism types of closed m -dimensional manifolds that admit a Riemannian metric with curvature $\geq \kappa$, and diameter $\leq D$.*

In fact, this theorem implies diffeomorphism-type finiteness in all dimensions except 4.

Lecture 10

Boundary

A Definition

Let us give an inductive definition the boundary for finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space.

Suppose \mathcal{L} is a 1-dimensional complete geodesic $\text{CBB}(\kappa)$ space. By Exercise 7.15, \mathcal{L} is homeomorphic to a 1-dimensional manifold (possibly with nonempty boundary). It allows us to define the boundary $\partial\mathcal{L} \subset \mathcal{L}$ as the boundary of a manifold.

Now assume that the notion of boundary is already defined in dimensions $1, \dots, m-1$. Suppose \mathcal{L} is m -dimensional complete geodesic $\text{CBB}(\kappa)$ space. We say that $p \in \mathcal{L}$ belongs to the boundary (briefly $p \in \partial\mathcal{L}$) if $\partial\Sigma_p \neq \emptyset$. By 7.17 and 7.18b, Σ_p is $(m-1)$ -dimensional complete geodesic $\text{CBB}(1)$ space; therefore its boundary is already defined.

10.1. Exercise. *Show that for closed convex set $K \subset \mathbb{E}^m$ with non-empty interior, the topological boundary of K as a subset of \mathbb{E}^m coincides with the boundary K described above.*

10.2. Exercise. *Let \mathcal{L} be a finite-dimensional complete geodesic $\text{CBB}(0)$ space. Suppose $\mathcal{L} \stackrel{\text{iso}}{=} \mathcal{L}_1 \times \mathcal{L}_2$. Show that*

$$\partial\mathcal{L} = (\partial\mathcal{L}_1 \times \mathcal{L}_2) \cup (\mathcal{L}_1 \times \partial\mathcal{L}_2).$$

B Conic neighborhoods

The following statement is close relative of Perelman's stability theorem 9.8, but its proof is slightly simpler [44]. We are going to use this

result without proof.

Recall that logarithm $\log_p x: \mathcal{L} \rightarrow T_p$ is defined on page 29.

10.3. Theorem. *Any point p in a finite-dimensional geodesic $\text{CBB}(\kappa)$ space \mathcal{L} and all sufficiently small $\varepsilon > 0$ there is a homeomorphism $h_\varepsilon: B(p, \varepsilon)_\mathcal{L} \rightarrow B(0, \varepsilon)_{T_p}$ such that $0 = h_\varepsilon(p)$.*

Moreover, we may assume that

$$\sup_{x \in B(p, \varepsilon)} \left\{ \frac{1}{\varepsilon} \cdot |\log_p x - h_\varepsilon(x)|_{T_p} \right\} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

This statement is often used together with *uniqueness of conic neighborhood* stated below.

Suppose that a neighborhood U of a point x in a metric space \mathcal{X} admits a homeomorphism to $\text{Cone } \Sigma$ such that x maps to the origin of the cone. In this case, we say that U has a conic neighborhood of x .

10.4. Uniqueness of conic neighborhood. *Any two conic neighborhoods of a given point in a metric space are pointed homeomorphic; that is, there is a homeomorphism between neighborhoods that maps the origin of the cone to the origin.*

10.5. Advanced exercise. *Prove 10.4 or read the original proof by Kyung Whan Kwun [30].*

10.6. Exercise. *Suppose $x \mapsto x'$ is a homeomorphism between finite-dimensional geodesic $\text{CBB}(\kappa)$ spaces \mathcal{L} and \mathcal{L}' . Show that*

- (a) $T_x \cong T_{x'}$,
- (b) $\text{Susp } \Sigma_x \cong \text{Susp } \Sigma_{x'}$.
- (c) *but in general $\Sigma_x \not\cong \Sigma_{x'}$.*

C Topology

The following theorem states that boundary is a topological property despite our definition used geometry.

10.7. Theorem. *Let \mathcal{L} and \mathcal{L}' be homeomorphic finite-dimensional geodesic $\text{CBB}(\kappa)$ spaces. Then $\dim \mathcal{L} = \dim \mathcal{L}'$ and*

$$\partial \mathcal{L} \neq \emptyset \iff \partial \mathcal{L}' \neq \emptyset$$

Before diving into the proof, it is instructive to solve the following exercise.

10.8. Exercise. Construct two plane subsets K_1 and K_2 such that $K_1 \not\cong K_2$, but $\mathbb{R} \times K_1 \cong \mathbb{R} \times K_2$.

Let \mathcal{L} be an m -dimensional geodesic $\text{CBB}(\kappa)$ space and $m < \infty$. Define rank of \mathcal{L} (briefly, $\text{rank } \mathcal{L}$) as the minimal value k such that $\mathcal{L} \stackrel{\text{iso}}{=} \mathbb{R}^{m-k} \times \mathcal{K}$, where \mathcal{K} is a k -dimensional geodesic $\text{CBB}(\kappa)$ space.

In the following proof we will apply induction on the rank of space.

Proof. The first statement follows from 7.19.

Suppose we have a counterexample, say $\partial\mathcal{L} \neq \emptyset$, but $\partial\mathcal{L}' = \emptyset$. Let $k := \text{rank } \mathcal{L}$ and $k' := \text{rank } \mathcal{L}'$. We can assume that the pair (k, k') is minimal in the lexicographic order; in particular, k is minimal. Let $x \mapsto x'$ be a homeomorphism from \mathcal{L} to \mathcal{L}' .

Choose $x \in \partial\mathcal{L}$. Since $\partial\mathcal{L}' = \emptyset$, we have $x' \notin \partial\mathcal{L}'$. Note that

$$\text{rank } T_x \leq k \quad \text{and} \quad \text{rank } T_{x'} \leq k',$$

By 10.6a, $T_x \cong T_{x'}$. Note that $\partial T_x \neq \emptyset$ and $\partial T_{x'} = \emptyset$. Therefore, we may assume that \mathcal{L} and \mathcal{L}' are Euclidean cones and the homeomorphism sends the origin to the origin. The remaining part of the proof is divided in three cases.

Case 1. Suppose $k > 1$. Let $\mathcal{L} \stackrel{\text{iso}}{=} \mathbb{R}^{m-k} \times \mathcal{C}$, where \mathcal{C} a k -dimensional geodesic $\text{CBB}(0)$ cone. Note that $\text{rank } T_y \leq \text{rank } \mathcal{L}$ for any $y \in \mathcal{L}$ and the equality holds only if y projects to the origin of \mathcal{C} .

Since $k > 1$ we can find $z \in \partial\mathcal{C}$ such that $z \neq 0$. Choose y that projects to z , so $\text{rank } T_y < \text{rank } \mathcal{L}$. By 10.6a, $T_y \cong T_{y'}$, $\partial T_y \neq \emptyset$ and $\partial T_{y'} = \emptyset$. The latter contradicts minimality of k .

Case 2. Suppose $k \leq 1$ and $k' > 1$. Since $\partial\mathcal{L} \neq \emptyset$, we get that $k = 1$ and $\mathcal{L} = \mathbb{R}^{m-1} \times \mathbb{R}_{\geq 0}$.

Let $\mathcal{L}' \stackrel{\text{iso}}{=} \mathbb{R}^{m-k'} \times \mathcal{C}'$, where \mathcal{C}' a k' -dimensional geodesic $\text{CBB}(0)$ cone. Since $\partial\mathcal{L} \cong \mathbb{R}^{m-1}$, the image of $\partial\mathcal{L}$ in \mathcal{L}' does not lie in $\mathbb{R}^{m-k'} \times \{0\}$. In other words, we can choose $y \in \partial\mathcal{L}$ such that its image $y' \in \mathcal{L}'$ has nonzero projection in \mathcal{C}' . Observe that $T_y \cong T_{y'}$,

$$\text{rank } T_y \leq k = 1, \quad \text{rank } T_{y'} < k', \quad \partial T_y = \emptyset, \quad \text{and} \quad \partial T_{y'} \neq \emptyset$$

— a contradiction.

Case 3. Suppose $k \leq 1$ and $k' \leq 1$. Since $\partial\mathcal{L} \neq \emptyset$, $k = 1$. By 7.15, $\mathcal{L} \cong \mathbb{R}_{\geq 0}$. Therefore, $\mathcal{L}' \cong \mathbb{R}$, and $\mathcal{L} \not\cong \mathcal{L}'$ — a contradiction. \square

10.9. Exercise. Let $x \mapsto x'$ be a homeomorphism $\Omega \rightarrow \Omega'$ between open subsets in finite-dimensional complete geodesic $\text{CBB}(\kappa)$ spaces \mathcal{L} and \mathcal{L}' . Show that $x \in \partial\mathcal{L}$ if and only if $x' \in \partial\mathcal{L}'$.

10.10. Exercise. Show that boundary of a finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space is a closed subset.

D Tangent space

Let X be a subset in a finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} . Choose $p \in \mathcal{L}$ and $\xi \in \Sigma_p$. Suppose ξ is a limit of directions $\uparrow_{[px_n]}$ for a sequence $x_1, x_2, \dots \in X$ that converges to p . Then we say that ξ is in the space of directions from p to X ; briefly $\xi \in \Sigma_p X$.

Further, the cone $\text{Cone}(\Sigma_p X)$ will be called tangent space to X at p ; it will be denoted by $T_p X$.

10.11. Theorem. *For any finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space \mathcal{L} , we have*

$$\partial(\Sigma_p \mathcal{L}) = \Sigma_p(\partial \mathcal{L}) \quad \text{and} \quad \partial(T_p \mathcal{L}) = T_p(\partial \mathcal{L}).$$

Proof. Choose a sequence $x_n \in \partial \mathcal{L}$ such that $x_n \rightarrow p$ and $\uparrow_{[px_n]} \rightarrow \xi$.

Let $\varepsilon_n = 2 \cdot |p - x_n|$, and let $h_{\varepsilon_n} : B(p, \varepsilon_n)_{\mathcal{L}} \rightarrow B(0, \varepsilon_n)_{T_p}$ be the homeomorphisms provided by 10.3; in particular, $\frac{2}{\varepsilon_n} \cdot h_{\varepsilon_n}(x_n) \rightarrow \xi$ as $n \rightarrow \infty$. By 10.9, $h_{\varepsilon_n}(x_n) \in \partial T_p$. By 10.10, $\xi \in \partial T_p$. Therefore,

$$\partial(\Sigma_p \mathcal{L}) \supset \Sigma_p(\partial \mathcal{L}) \quad \text{and} \quad \partial(T_p \mathcal{L}) \supset T_p(\partial \mathcal{L}).$$

Similarly, choose $\xi \in \partial \Sigma_p$. Let $h_{\varepsilon_n} : B(p, \varepsilon_n)_{\mathcal{L}} \rightarrow B(0, \varepsilon_n)_{T_p}$ be the homeomorphisms provided by 10.3 for a sequence $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. By 10.9, $x_n = h_{\varepsilon_n}^{-1}(\frac{\varepsilon_n}{2} \cdot \xi) \in \partial T_p$. By 10.3, $\uparrow_{[px_n]} \rightarrow \xi$. Hence

$$\partial(\Sigma_p \mathcal{L}) \subset \Sigma_p(\partial \mathcal{L}) \quad \text{and} \quad \partial(T_p \mathcal{L}) \subset T_p(\partial \mathcal{L}).$$

□

E Doubling

Let A be a closed subset in a metric space \mathcal{X} . The doubling \mathcal{W} of \mathcal{X} across A is two copies of \mathcal{X} glued along A ; more precisely, the underlying set of \mathcal{W} is the quotient $\mathcal{X} \times \{0, 1\} / \sim$, where $(a, 0) \sim (a, 1)$ for any $a \in A$ and it has minimal metric such that both maps $\mathcal{X} \rightarrow \mathcal{W}$ defined by $x \mapsto (x, 0)$ and $x \mapsto (x, 1)$ are distance-preserving.

The metric on \mathcal{W} can be also defined explicitly by

$$|(x, i) - (y, j)|_{\mathcal{W}} = \begin{cases} |x - y|_{\mathcal{X}} & \text{if } i = j. \\ \inf \{ |x - a|_{\mathcal{X}} + |y - a|_{\mathcal{X}} : a \in A \} & \text{if } i \neq j. \end{cases}$$

The last part of the following statement is the so-called doubling theorem.

10.12. Theorem. *Let \mathcal{L} be a finite-dimensional complete geodesic $\text{CBB}(\kappa)$ space with nonempty boundary. Suppose $f = \frac{1}{2} \cdot \text{dist}_p^2$ for some $p \in \mathcal{L}$. Then*

- (a) If $\dim \mathcal{L} \geq 2$, then $\text{dist}_{\partial \Sigma_x}(\xi) \leq \frac{\pi}{2}$ for any $x \in \partial \mathcal{L}$ and $\xi \in \Sigma_x$.
Moreover, if $\text{dist}_{\partial \Sigma_x}(\xi) = \frac{\pi}{2}$, then $\angle(\xi, \zeta) \leq \frac{\pi}{2}$ for any $\zeta \in \Sigma_x$.
- (b) $\nabla_x f \in \partial T_x$ for any $x \in \partial \mathcal{L}$.
- (c) If α is an f -gradient curve that starts at $x \in \partial \mathcal{L}$, then $\alpha(t) \in \partial \mathcal{L}$ for any t . Moreover, if $p \in \partial \mathcal{L}$, then $\text{gexp}_p(v) \in \partial \mathcal{L}$ for any $v \in \partial T_p$.
- (d) The doubling \mathcal{W} of \mathcal{L} across $\partial \mathcal{L}$ is a complete geodesic CBB(κ) space.

Proof. We will denote by $(a)_m, \dots, (d)_m$ the corresponding statement assuming $m = \dim \mathcal{L}$.

The proof goes by induction on m . Note that $(d)_1$ follows from 7.15 — this is the base of induction. The step of induction is a combination of several implications listed below.

$(d)_{m-1} \Rightarrow (a)_m$. If $m = 2$, then $\dim \Sigma_x = 1$; see 7.18b. By 7.15, Σ_x is isometric to a line segment $[0, \ell]$; we need to show that $\ell \leq \pi$.

Suppose $\ell > \pi$, then the tangent space $T_x = \text{Cone } \Sigma_x$ has several different lines thru the origin. Recall that T_x is a complete geodesic CBB(0) space; see 7.18. By 6.5, T_x is isometric to the Euclidean plane; the latter contradicts that Σ_x is a line segment.

Let $m = \dim \mathcal{L} > 2$, so $\dim \Sigma_x > 1$. Suppose $\text{dist}_{\partial \Sigma_x}(\xi) > \frac{\pi}{2}$ for some ξ . By $(d)_{m-1}$, the doubling Ξ of Σ_x is a geodesic CBB(κ) space. Denote by ξ_0 and ξ_1 the points in Ξ that correspond to ξ . Observe that $|\xi_0 - \xi_1|_{\Xi} > \pi$. The latter contradicts 3.6.

Finally, if $\text{dist}_{\partial \Sigma_x}(\xi) = \frac{\pi}{2}$, then $|\xi_0 - \xi_1|_{\Xi} = \pi$. Therefore $\text{Cone } \Xi$ contains a line in directions of ξ_0 and ξ_1 ; in other words, Ξ is a spherical suspension with poles ξ_0 and ξ_1 . In particular, every point of Ξ lies on distance at most $\frac{\pi}{2}$ from ξ_0 or ξ_1 . The natural projection $\Xi \rightarrow \Sigma_x$ does not increase distances and sends both ξ_0 and ξ_1 to ξ . Therefore, the second statement follows.

$(d)_{m-1} + (a)_{m-1} + (a)_m \Rightarrow (b)_m$. We can assume that $s = \nabla_x f \neq 0$. By 4.9, $\nabla_x f = s \cdot \bar{\xi}$, where $s = \mathbf{d}_x f(\bar{\xi})$ and $\bar{\xi} \in \Sigma_p$ is the direction that maximize $\mathbf{d}_x f(\bar{\xi})$.

Let $\zeta \in \partial \Sigma_x$ be a direction that minimize the angle $\angle(\bar{\xi}, \zeta)$. It is sufficient to show that $\zeta = \bar{\xi}$.

Assume $\zeta \neq \bar{\xi}$; let $\eta = \uparrow_{[\zeta \bar{\xi}]}$. By $(a)_m$, $\angle(\bar{\xi}, \zeta) \leq \frac{\pi}{2}$ and $(a)_{m-1}$ implies that

$$\textcircled{1} \quad \angle(\eta, \nu) \leq \frac{\pi}{2}$$

for any $\nu \in \Sigma_{\zeta} \Sigma_x$; if $m = 2$, then the last statement is evident.

Consider function $\varphi: \Sigma_x \rightarrow \mathbb{R}$ defined by $\varphi(\xi) := \mathbf{d}_x f(\xi)$. Applying 4.6a and $\textcircled{1}$, we get that $\mathbf{d}_{\xi} \varphi(\eta) \leq 0$. Since $\mathbf{d}_x f$ is convex, we have

that $\varphi'' + \varphi \leq 0$. If $\varphi(\zeta) \leq 0$, then it implies that $\varphi(\bar{\xi}) \leq 0$ — a contradiction. If $\varphi(\zeta) > 0$, then $\varphi(\bar{\xi}) < \varphi(\zeta)$ — a contradiction again.

$(b)_m \Rightarrow (c)_m$. Let α be an f -gradient curve and $\ell(t) = \text{dist}_{\partial L} \alpha(t)$. Note that ℓ is a Lipschitz function.

Choose t ; let $x = \alpha(t)$ and $y \in \partial \mathcal{L}$ be a closest point to x . By $(b)_m$, we have that $\nabla_y f \in \partial T_y$. Since the distance $|x - y|$ is minimal, we get $\langle \uparrow_{[yx]}, v \rangle \leq 0$ for any $v \in \partial T_y$. In particular,

$$\langle \uparrow_{[yx]}, \nabla_y f \rangle \leq 0$$

Applying Exercise 4.8 to x and y , we get

$$\ell'(t) \leq \ell(t)$$

if the left-hand side is defined. Since ℓ is Lipschitz, ℓ' is defined almost everywhere. Integrating the inequality, we get

$$\ell(t) \leq e^t \cdot \ell(0)$$


for any $t \geq 0$. In particular, if $\ell(0) = 0$, then $\ell(t) = 0$ for any $t \geq 0$. Since $\partial \mathcal{L}$ is closed (10.10), the statement follows.

$(c)_m + (d)_{m-1} \Rightarrow (d)_m$. We will consider the case $\kappa = 0$; other cases can be done the same way, but formulas getting more complicated.

Denote by \mathcal{L}_0 and \mathcal{L}_1 the two copies of \mathcal{L} in \mathcal{W} ; let us keep the notation $\partial \mathcal{L}$ for the common boundary of \mathcal{L}_0 and \mathcal{L}_1 . Next claim says that any geodesic in \mathcal{W} either lies in $\partial \mathcal{L}$ or it crosses $\partial \mathcal{L}$ at most once.

❷ *Let γ be a geodesic in \mathcal{W} . Then either γ has at most one interior point in $\partial \mathcal{L}$ or $\gamma \subset \partial \mathcal{L}$.*

Assume γ shares at least two points with $\partial \mathcal{L}$, say $x = \gamma(t_1)$ and $y = \gamma(t_2)$ and these are not endpoints of γ . Let us exchange $\gamma \cap \mathcal{L}_1$ to its reflection in \mathcal{L}_0 ; denote the obtained curve by $\hat{\gamma}$.



Note that any arc of $\hat{\gamma}$ with endpoint at x or y is a geodesic in \mathcal{L}_0 . Moreover, the arc of $\hat{\gamma}$ behind y lies in the image of map $t \mapsto \text{Flow}_{f_x}^t(y)$, where $f_x = \frac{1}{2} \cdot \text{dist}_x^2$. By (c), this arc lies in $\partial \mathcal{L}$.

Now choose a point z on this arc, so $z \in \partial \mathcal{L}$. Applying the same argument, we get that the arc of $\hat{\gamma}$ before y lies in $\partial \mathcal{L}$. Hence the claim follows. \triangle

Choose a point p in \mathcal{W} ; let $f := \frac{1}{2} \cdot \text{dist}_p^2$. Without loss of generality, we can assume that $p \in \mathcal{L}_0$. It is sufficient to show that $(f \circ \gamma)'' \leq 1$ for any t . If γ lies in $\partial \mathcal{L}$, then this inequality follows from the comparison in \mathcal{L}_0 .

If $\gamma(t) \in \mathcal{L}_0 \setminus \partial\mathcal{L}$, then $(f \circ \gamma)''(t) \leq 1$ follows from the comparison in \mathcal{L}_0 .

Assume $\gamma(t) \in \mathcal{L}_1 \setminus \partial\mathcal{L}$. Suppose $[p\gamma(t)]$ crosses $\partial\mathcal{L}$ at x . Let Σ_x be the space of directions of \mathcal{L} at x and let Ξ be its doubling. By $(d)_{m-1}$, Ξ is a geodesic CBB(1) space. Note that $\uparrow_{[x\gamma(t)]}$ and $\uparrow_{[xp]}$ lie in opposite sides of Ξ and $|\uparrow_{[x\gamma(t)]} - \uparrow_{[xp]}| \geq \pi$. Otherwise we could choose a direction $\xi \in \partial\Sigma$ such that

$$|\uparrow_{[x\gamma(t)]} - \xi| + |\xi - \uparrow_{[xp]}| < \pi.$$

Consider the radial curve $\alpha(t) = \text{gexp}_x(t \cdot \xi)$. By $(c)_m$, α lies in $\partial\mathcal{L}$. By 5.12

$$|p - \alpha(s)|_{\mathcal{L}_0} + |\gamma(t) - \alpha(s)|_{\mathcal{L}_1} < |p - \gamma(t)|_{\mathcal{W}}$$

for small values $s > 0$ — a contradiction.

Note that $\text{Cone } \Xi$ contains a line with directions $\uparrow_{[x\gamma(t)]}$ and $\uparrow_{[xp]}$. By splitting theorem $\text{Cone } \Xi$ split in these directions; in particular,

$$|\uparrow_{[x\gamma(t)]} - \xi| + |\xi - \uparrow_{[xp]}| = \pi.$$

for any $\xi \in \Xi$. It follows that for any $\xi \in \Xi$ there is $\xi' \in \partial\Sigma_x$ such that ξ and ξ' lie on a geodesic $[\uparrow_{[x\gamma(t)]} \uparrow_{[xp]}]_{\Xi}$.

Choose ξ' for $\xi = \uparrow_{[x\gamma(t+???)]}.$ Let $\alpha(s) := \text{gexp}_x(s \cdot \xi')$. Applying ??? we get that for suitably chosen parameter s we

the proof of inequality reminds the argument in part (a), but it is a bit more tricky. Finally if $\gamma(t) \in \partial\mathcal{L}$, then splitting argument shows that


$$(f \circ \gamma)^+(t) + (f \circ \gamma)^-(t) \leq 0.$$

These three statements imply that $(f \circ \gamma)''(t) \leq 1$ for any t .

Summarizing, we get that $(f \circ \gamma)'' \leq 1$ if γ lies entirely in \mathcal{L}_0 or \mathcal{L}_1 . If γ crosses $\partial\mathcal{L}$, then we know that it happens only once and at the crossing moment t_0 we have $f \circ \gamma^+(t_0) + f \circ \gamma^-(t_0) \leq 0$. All this implies that $(f \circ \gamma)'' \leq 1$. \square

10.13. Exercise. Let \mathcal{L} be a complete finite-dimensional geodesic CBB(1) space with nonempty boundary $\partial\mathcal{L}$. Show that $\partial\mathcal{L}$ is connected.

10.14. Exercise. Let \mathcal{L} be a complete finite-dimensional geodesic CBB(0) space with nonempty boundary $\partial\mathcal{L}$. Show that the distance



function to the boundary

$$\text{dist}_{\partial\mathcal{L}}: \mathcal{L} \rightarrow \mathbb{R}$$

is concave.

10.15. Exercise. Suppose γ is a geodesic in the length metric of boundary of a complete finite-dimensional geodesic CBB(0) space. Show that the function $t \mapsto \frac{1}{2} \cdot \text{dist}_p^2 \circ \gamma(t)$ is 1-concave for any point p .

F Remarks

Theorem 10.3 can be used to prove the following.

10.16. Topological stratification. Any m -dimensional Alexandrov space with $m < \infty$ can be subdivided into subsets topological manifolds S_0, \dots, S_m such that for every i we have $\dim S_i = i$ or $S_i = \emptyset$. Moreover,

- (a) closure of S_{m-1} is the boundary of the space, and
- (b) $S_{m-2} = \emptyset$.

Let us mention that this statement does not imply that (compact) finite-dimensional Alexandrov space is homeomorphic to CW-complex ???

Note that doubling of a finite-dimensional Alexandrov space across its boundary is an Alexandrov space without boundary. This observation can be used to deduce a statement about general finite-dimensional Alexandrov space to an Alexandrov space without boundary; so the following tools become available.

10.17. Fundamental-class lemma. Let \mathcal{L} be a compact m -dimensional Alexandrov space, and $m < \infty$. Then \mathcal{L} has a fundamental class in Alexander–Spanier cohomology with $\mathbb{Z}/2$ coefficients; that is,

$$H^n(\mathcal{L}, \mathbb{Z}/2) = \mathbb{Z}/2.$$

This lemma was proved by Karsen Grove and Peter Perersen [22]; It implies, for example, that gradient is an onto map for functions of the following type $f = \frac{1}{2} \text{dist}_p$ is surjective on finite-dimensional Alexandrov spaces. It is also used in the proof of a generalized domain invariance theorem stated below.

10.18. Domain invariance. Let \mathcal{L}_1 and \mathcal{L}_2 be two m -dimensional geodesic CBB(κ) spaces with empty boundary; m is finite. Suppose Ω_1

is an open subset in \mathcal{L}_1 and $f: \Omega_1 \rightarrow \mathcal{L}_2$ is an injective continuous map. Then $f(\Omega_1)$ is open in \mathcal{L}_2 .

Let f be a semiconcave function. A point $p \in \text{Dom } f$ is called critical point of f if $\mathbf{d}_p f \leq 0$; otherwise it is called regular.

The following statement plays technical role in the proof of stability theorem, but this is also a useful technical tool.

10.19. Morse lemma. *Let f be a semiconcave function on a finite-dimensional geodesic $\text{CBB}(\kappa)$ space without boundary. Suppose K is a compact set of regular points of f in its level set $f = a$. Then an open neighborhood Ω of K admits homeomorphism $x \mapsto (h(x), f(x))$ to a product space $\Lambda \times (a - \varepsilon, a + \varepsilon)$.*

Subsets that satisfy condition in 10.12c are called extremal. More precisely, a subset E in a complete geodesic CBB space is called extremal if for any $x \in E$ and f -gradient curve that starts in E remains in E ; here f is arbitrary function of the form $\frac{1}{2} \cdot \text{dist}_p^2$.

Extremal subsets were introduced by Grigori Perelman and the second author [43]. They play an important role in Alexandrov geometry and its applications; in particular they will pop up in the next lecture.

The following conjecture is one of the oldest questions in Alexandrov geometry that remains open.

10.20. Conjecture. *Let S be a component of the boundary of a complete geodesic finite-dimensional $\text{CBB}(\kappa)$ space. Then S equipped with the induced length metric is $\text{CBB}(\kappa)$.*

Lecture 11

Quotients

A Quotient space

Suppose that a group G acts isometrically on a metric space \mathcal{X} . Note that

$$|G \cdot x - G \cdot y|_{\mathcal{X}/G} := \inf \{ |x - g \cdot y|_{\mathcal{X}} : g \in G \}$$

defines a semimetric on the orbit space \mathcal{X}/G . Moreover, it is a genuine metric if the orbits of the action are closed.

11.1. Theorem. *Assume that group G acts isometrically on a proper CBB(0) space \mathcal{L} and has closed orbits. Then the quotient space \mathcal{L}/G is CBB(0).*

Proof. Denote by $\sigma: \mathcal{L} \rightarrow \mathcal{L}/G$ the quotient map.

Fix a quadruple of points $p, x_1, x_2, x_3 \in \mathcal{L}/G$. Choose an arbitrary $\hat{p} \in \mathcal{L}$ such that $\sigma(\hat{p}) = p$. Since \mathcal{L} is proper, we can choose the points $\hat{x}_i \in \mathcal{L}$ such that $\sigma(\hat{x}_i) = x_i$ and

$$|p - x_i|_{\mathcal{L}/G} = |\hat{p} - \hat{x}_i|_{\mathcal{L}}$$

for all i .

Note that

$$|x_i - x_j|_{\mathcal{L}/G} \leq |\hat{x}_i - \hat{x}_j|_{\mathcal{L}}$$

for all i and j . Therefore

$$\textcircled{1} \quad \tilde{\Delta}(p_{x_j}^{x_i}) \leq \tilde{\Delta}(\hat{p}_{\hat{x}_j}^{\hat{x}_i})$$

holds for all i and j .

By CBB(0) comparison in \mathcal{L} , we have

$$\tilde{\Delta}(\hat{p}_{\hat{x}_2}^{\hat{x}_1}) + \tilde{\Delta}(\hat{p}_{\hat{x}_3}^{\hat{x}_2}) + \tilde{\Delta}(\hat{p}_{\hat{x}_1}^{\hat{x}_3}) \leq 2 \cdot \pi.$$

Applying **1**, we get

$$\tilde{\mathcal{L}}(p_{x_2}^{x_1}) + \tilde{\mathcal{L}}(p_{x_3}^{x_2}) + \tilde{\mathcal{L}}(p_{x_1}^{x_3}) \leq 2 \cdot \pi;$$

that is, the CBB(0) comparison holds for any quadruple in \mathcal{L}/G . \square

11.2. Advanced exercise. *Let G be a compact Lie group with a bi-invariant Riemannian metric. Show that G is isometric to a quotient of the Hilbert space by an isometric group action.*

Conclude that $G \in \text{CBB}(0)$.

B Generalizations

A map $\sigma: \mathcal{X} \rightarrow \mathcal{Y}$ between the metric spaces \mathcal{X} and \mathcal{Y} is called a submetry if

$$\sigma(B(p, r)_{\mathcal{X}}) = B(\sigma(p), r)_{\mathcal{Y}}$$

for any $p \in \mathcal{X}$ and $r \geq 0$.

Suppose G and \mathcal{L} are as in 11.1. Observe that the quotient map $\sigma: \mathcal{L} \rightarrow \mathcal{L}/G$ is a submetry. The following two exercises show that this is not the only source of submetries.

11.3. Exercise. *Construct submetries*

(a) $\sigma_1: \mathbb{S}^2 \rightarrow [0, \pi]$

(b) $\sigma_2: \mathbb{S}^2 \rightarrow [0, \frac{\pi}{2}]$

(c) $\sigma_n: \mathbb{S}^2 \rightarrow [0, \frac{\pi}{n}]$ (for integer $n \geq 1$)

such that the fibers $\sigma_n^{-1}\{x\}$ are connected for any x .

11.4. Exercise. *Let $\sigma: \mathbb{E}^2 \rightarrow [0, \infty)$ be a submetry. Show that $K = \sigma^{-1}\{0\}$ is a closed convex set in \mathbb{E}^2 and $\sigma(x) = \text{dist}_K x$.*

The proof of 11.1 works for submetries. Therefore we get the following.

11.5. Generalization. *Let $\sigma: \mathcal{L} \rightarrow \mathcal{M}$ be a submetry. Suppose \mathcal{L} is a CBB(0) space, then so is \mathcal{M} .*

Theorems 11.1 and 11.5 admit straightforward generalizations to CBB(−1) spaces. In the CBB(1) case, the proof produces a slightly weaker statement — *any open $\frac{\pi}{2}$ -ball in the quotient of CBB(1) is CBB(1)*; in particular, the quotient space is *locally* CBB(1). If the space is geodesic, then the globalization theorem implies that it is globally CBB(1). The same holds for the targets of submetry from a CBB(1) space. In other words, we have the following two statements.

11.6. Theorem. *Let $\sigma: \mathcal{L} \rightarrow \mathcal{M}$ be a submetry. Suppose \mathcal{L} is CBB(−1) space, then so is \mathcal{M} .*

11.7. Theorem. *Let $\sigma: \mathcal{L} \rightarrow \mathcal{M}$ be a submetry. Suppose \mathcal{L} is a geodesic CBB(1) space, then so is \mathcal{M} .*

C Hopf's conjecture

Recall that Hopf's conjecture asks *does $\mathbb{S}^2 \times \mathbb{S}^2$ admit a Riemannian metric with positive sectional curvature?* The following partial result was obtained by Wu-Yi Hsiang and Bruce Kleiner [26].

11.8. Theorem. *There is no Riemannian metric on $\mathbb{S}^2 \times \mathbb{S}^2$ with sectional curvature ≥ 1 and a nontrivial isometric \mathbb{S}^1 -action.*

11.9. Key lemma. *Suppose $\mathbb{S}^1 \curvearrowright \mathbb{S}^3$ is an isometric action without fixed points and $\Sigma = \mathbb{S}^3/\mathbb{S}^1$ is its quotient space. Then there is a distance noncontracting map $\Sigma \rightarrow \frac{1}{2} \cdot \mathbb{S}^2$, where $\frac{1}{2} \cdot \mathbb{S}^2$ is the standard 2-sphere rescaled with a factor $\frac{1}{2}$.*

The proof of the lemma is done mostly by calculations; we present it in the form of guided exercise.

11.10. Exercise. *Suppose $\mathbb{S}^1 \curvearrowright \mathbb{S}^3$ is an isometric action without fixed points. Let us think that \mathbb{S}^3 is the unit sphere in \mathbb{R}^4 .*

- (a) *Show that one can identify \mathbb{R}^4 with \mathbb{C}^2 so that the action is given by matrix multiplication*

$$\begin{pmatrix} u^p & 0 \\ 0 & u^q \end{pmatrix},$$

where (p, q) is a pair of relatively prime positive integers and $u \in \mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$. In particular, our \mathbb{S}^1 is a subgroup in the torus that acts by matrix multiplication

$$\begin{pmatrix} v & 0 \\ 0 & w \end{pmatrix},$$

where $v, w \in \mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$.

Fix p and q as above. Let $\Sigma_{p,q} = \mathbb{S}^3/\mathbb{S}^1$ be the quotient space.

- (b) *Show that the $\Sigma_{p,q} = \mathbb{S}^3/\mathbb{S}^1$ is a topological sphere with \mathbb{S}^1 -symmetry. This symmetry has two fixed points, north pole and south pole, that correspond to the orbits of $(1, 0)$ and $(0, 1)$ in \mathbb{S}^3 . Denote by $S(r)$ the circle of radius r with the center at the north pole of $\Sigma_{p,q}$.*

- (c) *Show that the inverse image $T(r)$ of $S(r)$ in \mathbb{S}^3 is also an orbit of the torus action. Conclude that $a(r) = \pi^2 \cdot \sin r \cdot \cos r$, where $a(r)$ is the area $T(r)$.*

- (d) Let $b_{p,q}(r)$ be the length of the \mathbb{S}^1 -orbit in \mathbb{S}^3 that corresponds to a point on $S(r)$. Show that $b_{p,q} = \pi \cdot \sqrt{(p \cdot \sin r)^2 + (q \cdot \cos r)^2}$.
- (e) Let $c_{p,q}(r)$ be the length of $S(r)$. Show that $a(r) = c_{p,q}(r) \cdot b_{p,q}(r)$.
- (f) Show that $c_{p,q}(r) \leq c_{1,1}(r)$ for any pair (p, q) of relatively prime positive integers. Use it to show construct a distance noncontracting map $\Sigma_{p,q} \rightarrow \frac{1}{2} \cdot \mathbb{S}^2 \stackrel{\text{iso}}{=} \Sigma_{1,1}$.

Sketch of 11.8. Assume $\mathcal{M} = (\mathbb{S}^2 \times \mathbb{S}^2, g)$ is a counterexample. By the Toponogov theorem, \mathcal{M} is CBB(1). By 11.1, the quotient space $\mathcal{L} = \mathcal{M}/\mathbb{S}^1$ is CBB(1); evidently, \mathcal{L} is 3-dimensional.

Denote by $F \subset \mathcal{M}$ the fixed point set of the \mathbb{S}^1 -action. Each connected component of F is either an isolated point or a 2-dimensional geodesic submanifold in \mathcal{M} ; the latter has to have positive curvature and therefore it is either \mathbb{S}^2 or \mathbb{RP}^2 . Notice that

- ◊ each isolated point contributes 1 to the Euler characteristic of \mathcal{M} ,
- ◊ each sphere contributes 2 to the Euler characteristic of \mathcal{M} , and
- ◊ each projective plane contributes 1 to the Euler characteristic of \mathcal{M} .

Since $\chi(\mathcal{M}) = 4$, we are in one of the following three cases:

- ◊ F has exactly 4 isolated points,
- ◊ F has one 2-dimensional submanifold and at least 2 isolated points,
- ◊ F has at least two 2-dimensional submanifolds.

Each case is covered separately.

Case 1. Suppose F has exactly 4 isolated points x_1, x_2, x_3 , and x_4 . Denote by y_1, y_2, y_3 , and y_4 the corresponding points in \mathcal{L} . Note that $\Sigma_{y_i} \mathcal{L}$ is isometric to a quotient of \mathbb{S}^3 by an isometric \mathbb{S}^1 -action without fixed points.

By 11.10, each angle $\angle[y_i \begin{smallmatrix} y_j \\ y_k \end{smallmatrix}] \leq \frac{\pi}{2}$ for any three distinct points y_i, y_j, y_k . In particular, all four triangles $[y_1 y_2 y_3]$, $[y_1 y_2 y_4]$, $[y_1 y_3 y_4]$, and $[y_2 y_3 y_4]$ are nondegenerate. By the comparison, the sum of angles in each triangle is strictly greater than π .

Denote by σ the sum of all 12 angles in 4 triangles $[y_1 y_2 y_3]$, $[y_1 y_2 y_4]$, $[y_1 y_3 y_4]$, and $[y_2 y_3 y_4]$. From above,

$$\sigma > 4 \cdot \pi.$$

On the other hand, by 11.10 any triangle in $\Sigma_{y_1} \mathcal{L}$ has perimeter at most π . In particular,

$$\angle[y_1 \begin{smallmatrix} y_2 \\ y_3 \end{smallmatrix}] + \angle[y_1 \begin{smallmatrix} y_3 \\ y_4 \end{smallmatrix}] + \angle[y_1 \begin{smallmatrix} y_4 \\ y_2 \end{smallmatrix}] \leq \pi.$$

Apply the same argument in $\Sigma_{y_2}\mathcal{L}$, $\Sigma_{y_3}\mathcal{L}$, and $\Sigma_{y_4}\mathcal{L}$. Adding the results we get

$$\sigma \leq 4 \cdot \pi$$

— a contradiction.

Case 2. Let F contain one surface S . Note that the projection of S to \mathcal{L} forms its boundary $\partial\mathcal{L}$. Note that doubling $\hat{\mathcal{L}}$ of \mathcal{L} across its boundary has 4 singular points — each singular point of \mathcal{L} corresponds to two singular points of $\hat{\mathcal{L}}$.

By the doubling theorem, $\hat{\mathcal{L}}$ is a geodesic CBB(1) space. Therefore we arrive at a contradiction the same way as in the first case.

Case 3. Suppose F contains at least two surfaces. Then $\partial\mathcal{L}$ has at least two connected components; choose two of them A and B . Denote by γ a geodesic that minimizes the distance from A to B .

Let

$$\dots, \mathcal{L}_{-1}, \mathcal{L}_0, \mathcal{L}_1, \dots$$

be a two-sided infinite sequence of copies on $\partial\mathcal{L}$. Let us glue \mathcal{L}_i to \mathcal{L}_{i+1} along A if i is even and along B if i is odd.

By the doubling theorem, every point in the obtained space \mathcal{N} has a neighborhood that is isometric to a neighborhood of the corresponding point in \mathcal{L} or its doubling. By the globalization theorem, \mathcal{N} is CBB(1).

Note that the copies of γ in \mathcal{L}_i form a line in \mathcal{N} . By the splitting theorem, \mathcal{N} is isometric to a product $\mathcal{N}' \oplus \mathbb{R}$. Since $\dim \mathcal{N} > 1$, Exercise 7.15 implies that $\text{diam} \mathcal{N} \leq \pi$ — a contradiction. \square

D Erdős' problem rediscovered

A point p in a CBB(κ) space is called extremal if $\angle[p_y^x] \leq \frac{\pi}{2}$ for any hinge $[p_y^x]$ with the vertex at p .

11.11. Theorem. *Let \mathcal{L} be a compact m -dimensional CBB(0) space. Then it has at most 2^m extremal points.*

The proof is a translation of the proof of the following classical problem in discrete geometry to Alexandrov's language.

11.12. Erdős' problem. *Let F be a set of points in \mathbb{E}^m such that any triangle formed by three distinct points in F has no obtuse angles. Then $|F| \leq 2^m$. Moreover, if $|F| = 2^m$ then F consists of the vertexes of an m -dimensional rectangle.*

This problem was posed by Paul Erdős [17] and solved by Ludwig Danzer and Branko Grünbaum [15]. Grigori Perelman noticed that

after proper definitions, the same proof works in Alexandrov spaces [42]; so it proves 11.11.

Proof of 11.11. Let $\{p_1, \dots, p_N\}$ be extremal points in \mathcal{L} . For each p_i consider its open Voronoi domain V_i ; that is,

$$V_i = \{x \in \mathcal{L} : |p_i - x| < |p_j - x| \text{ for any } j \neq i\}.$$

Clearly $V_i \cap V_j = \emptyset$ if $i \neq j$.

Suppose $0 < \alpha \leq 1$. Given a point $x \in \mathcal{L}$, choose a geodesic $[p_i x]$ and denote by x_i the point on $[p_i x]$ such that $|p_i - x_i| = \alpha \cdot |p_i - x|$; let $\Phi_i: x \rightarrow x_i$ be the corresponding map. By the comparison,

$$|x_i - y_i| \geq \alpha \cdot |x - y|$$

for any x, y , and i . Therefore

$$\text{vol}(\Phi_i \mathcal{L}) \geq \alpha^m \cdot \text{vol } \mathcal{L}.$$

Suppose $\alpha < \frac{1}{2}$. Then $x_i \in V_i$ for any $x \in \mathcal{L}$. Indeed, assume $x_i \notin V_i$, then there is p_j such that $|p_i - x_i| \geq |p_j - x_i|$. Then from comparison, we have $\tilde{\angle}(p_j \frac{p_i}{x})_{\mathbb{E}^2} > \frac{\pi}{2}$; that is, p_j does not form a one-point extremal set. It follows that $\text{vol } V_i \geq \alpha^m \cdot \text{vol } \mathcal{L}$ for any $0 < \alpha < \frac{1}{2}$; hence

$$\text{vol } V_i \geq \frac{1}{2^m} \cdot \text{vol } \mathcal{L} \quad \text{and} \quad N \leq 2^m.$$

□

E Crystallographic actions

An isometric action $\Gamma \curvearrowright \mathbb{E}^m$ is called crystallographic if it is properly discontinuous (that is, for any compact set $K \subset \mathbb{E}^m$ and $x \in \mathbb{E}^m$ there are only finitely many $g \in \Gamma$ such that $g \cdot x \in K$) and cocompact (that is, the quotient space $\mathcal{L} = \mathbb{E}^m / \Gamma$ is compact).

Let F be a maximal finite subgroup of Γ ; that is, if H is a finite group H such that $F < H < \Gamma$, then $F = H$. Denote by $\#(\Gamma)$ the number of maximal finite subgroups of Γ up to conjugation.

11.13. Open question. *Let $\Gamma \curvearrowright \mathbb{E}^m$ be a crystallographic action. Is it true that $\#(\Gamma) \leq 2^m$?*

Note that any finite subgroup F of Γ fixes an affine subspace A_F in \mathbb{E}^m . If F is maximal, then A_F completely describes F . Indeed, since the action is properly discontinuous, the subgroup of Γ that fix A_F has to be finite. This subgroup must contain F , but since F is maximal, it must coincide with F .

Denote by $\#_k(\Gamma)$ the number of maximal finite subgroups $F < \Gamma$ (up to conjugation) such that $\dim A_F = k$.

Choose a finite subgroup $F < \Gamma$; consider a conjugate subgroup $F' = g \cdot F \cdot g^{-1}$. Note that $A_{F'} = g \cdot A_F$. In particular, the subspaces A_F and $A_{F'}$ have the same image in the quotient space $\mathcal{L} = \mathbb{E}^m / \Gamma$. Therefore, to count subgroups up to conjugation, we need to count the images of their fixed set. Therefore, by the lemma below, $\#_0(\Gamma)$ cannot exceed the number of extremal points in $\mathcal{L} = \mathbb{E}^m / \Gamma$. Combining this observation with 11.11, we get the following.

11.14. Proposition. *Let $\Gamma \curvearrowright \mathbb{E}^m$ be a crystallographic action. Then $\#_0(\Gamma) \leq 2^m$.*

11.15. Lemma. *Let $\Gamma \curvearrowright \mathbb{E}^m$ be a crystallographic action and F be a maximal finite subgroup of Γ that fixes an isolated point p . Then the image of p in the quotient space $\mathcal{L} = \mathbb{E}^m / \Gamma$ is an extremal point.*

Proof. Let q be the image of p . Suppose q is not extremal; that is, $\angle[q \begin{smallmatrix} y_1 \\ y_2 \end{smallmatrix}] > \frac{\pi}{2}$ for some hinge $[q \begin{smallmatrix} y_1 \\ y_2 \end{smallmatrix}]$ in \mathcal{L} .

Choose the inverse images $x_1, x_2 \in \mathbb{E}^m$ of $y_1, y_2 \in \mathcal{L}$ such that $|p - x_i|_{\mathbb{E}^m} = |q - y_i|_{\mathcal{L}}$. Note that $\angle[p \begin{smallmatrix} x_1 \\ x_2 \end{smallmatrix}] \geq \angle[q \begin{smallmatrix} y_1 \\ y_2 \end{smallmatrix}] > \frac{\pi}{2}$. Moreover, since p is fixed by F , we have

$$\textcircled{1} \quad \angle[p \begin{smallmatrix} x_1 \\ g \cdot x_2 \end{smallmatrix}] > \frac{\pi}{2}$$

for any $g \in F$.

Denote by z the barycenter of the orbit $G \cdot x_2$. Note that z is a fixed point of F . By $\textcircled{1}$, $z \neq p$; so F must fix the line pz . But p is an isolated fixed point of F — a contradiction. \square

11.16. Exercise. *Let $\Gamma \curvearrowright \mathbb{E}^m$ be a crystallographic action. Show that*

- (a) $\#_{m-1}(\Gamma) \leq 2$, and
- (b) if $\#_{m-1}(\Gamma) = 1$, then $\#_0(\Gamma) \leq 2^{m-1}$.

Construct crystallographic actions with equalities in (a) and (b).

F Remarks

It is expected that *no geodesic CBB(1) space with a nontrivial isometric \mathbb{S}^1 -action can be homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^2$* ; so 11.8 holds for general CBB(1) space. The proof of 11.8 would work if we had the following generalization of 11.9; see [24].

11.17. Open question. *Let Σ be a geodesic CBB(1) space homeomorphic to \mathbb{S}^3 . Suppose \mathbb{S}^1 acts on Σ isometrically and without fixed*

points. Is it true that any triangle in Σ/\mathbb{S}^1 has perimeter at most π ? and, what about the existence of distance-noncontracting map $\Sigma/\mathbb{S}^1 \rightarrow \frac{1}{2} \cdot \mathbb{S}^2$?

11.18. Advanced exercise. Suppose that closed local geodesic γ is the fixed point of an isometric \mathbb{S}^1 action on a geodesic CBB(1) space that is homeomorphic to \mathbb{S}^3 . Show that $\text{length } \gamma \leq 2 \cdot \pi$.

It seems that the same question for a \mathbb{Z}_2 -action is open.

Compact geodesic m -dimensional CBB(0) spaces with the maximal number of extremal points include m -dimensional rectangles and the quotients of flat tori by reflections across a point. (This action has 2^m isolated fixed points; each corresponds to an extremal point in the quotient space $\mathcal{L} = \mathbb{T}^m/\mathbb{Z}_2$.) Nina Lebedeva has proved [32] that every m -dimensional geodesic CBB(0) space with 2^m extremal points is a quotient of Euclidean space by a crystallographic action.

Counting maximal finite subgroups in a crystallographic group Γ is equivalent to counting the so-called primitive extremal subsets in the quotient space $\mathcal{L} = \mathbb{E}^m/\Gamma$. So, 11.14 would follow from the next conjecture.

11.19. Conjecture. Any m -dimensional compact geodesic CBB(0) space has at most 2^m primitive extremal subset.

A closed subset E in a CBB(κ) space is called extremal if $\angle[p_y^x] \leq \leq \frac{\pi}{2}$ for any $x \notin E$ and $p \in E$ such that $|x-p|$ takes minimal value. An extremal subset is called primitive if it contains no proper extremal subsets. For example, the whole space and empty set are also extremal in any space. Also every vertex, edge, or face (as well as their union) of the cube is an extremal subset of the cube. Vertices of the cube are the only its primitive extremal subsets.

Lecture 12

Surface of convex body

Let us define a convex body as a compact convex subset in \mathbb{E}^3 with nonempty interior.

Suppose B is a convex body. Then the surface of B is defined as its boundary ∂B equipped with the induced length metric.

12.1. Exercise. *Show that surface of a convex body is homeomorphic to \mathbb{S}^2 .*

In this lecture, we will prove that *surface of a convex body is CBB(0)*.

A Surface of convex polyhedra

Convex polyhedron is a convex body with finite number of extremal points that called its vertices.

Observe that a surface, say Σ , of a convex polyhedron P admits a triangulation such that each triangle is isometric to a plane triangle. In other words, Σ is a polyhedral surfaces; that is, it is a 2-dimensional manifold with length metric that admits a triangulation such that each triangle is isometric to a solid plane triangle. A triangulation of polyhedral surface will be assumed to satisfy this condition.

The total angle around a vertex p in Σ is defined as the sum of angles at p of all triangles in the triangulation that contain p .

Note that if a point p is not a vertex of P , then

- ◇ p lies in the interior of a face of P , and its neighborhood in Σ is a piece of plane, or
- ◇ p lies on an edge, and its neighborhood is two half-planes glued along the boundary.

In both cases, a neighborhood of p in Σ (with the induced length metric) is isometric to an open domain of the plane.

12.2. Claim. *Let Σ be the surface of a convex polyhedron P . Then, the total angle around a vertex in Σ cannot exceed $2\cdot\pi$.*

In the proof, we will need the triangle inequality for angles (or the spherical triangle inequality). A proof of this statement is given in the classical geometry textbook by Andrei Kiselyov [29, § 47], it also follows from 1.9. (In fact our proof of 1.9 is a straightforward generalization of the argument in [29, § 47].)

12.3. Spherical triangle inequality. *Let w_1, w_2, w_3 be unit vectors in \mathbb{E}^3 . Denote by $\theta_{i,j}$ the angle between the vectors v_i and v_j . Show that*

$$\theta_{1,3} \leq \theta_{1,2} + \theta_{2,3}$$

and in case of equality, the vectors w_1, w_2, w_3 lie in a plane.

Proof of 12.2. Consider the intersection of P with a small sphere centered at p ; it is a convex spherical polygon, say F . Applying rescaling we may assume that the sphere has unit radius. Then we need to show that the perimeter of F does not exceed $2\cdot\pi$.

Note that F lies in a hemisphere, say H . Moreover, there is a decreasing sequence

$$H = H_0 \supset H_1 \supset \cdots \supset H_n = F,$$

such that H_{i+1} is obtained from H_i by cutting along a chord.

By 12.3, we have

$$2\cdot\pi = \text{perim } H = \text{perim } H_0 \geq \text{perim } H_1 \geq \cdots \geq \text{perim } H_n = \text{perim } F$$

— hence the result. \square

A vertex of a triangulation of a polyhedral surface is called essential if the total angle around it is not $2\cdot\pi$.

12.4. Exercise. *Let v be a point on the surface Σ of a convex polyhedron P . Show that v is a vertex of P if and only if v is an essential vertex of Σ .*

12.5. Exercise. *Show that geodesics on a surface of convex polyhedron do not pass thru its essential vertices.*

B Curvature

Let p be a vertex of a polyhedron. If θ_p is the total angle around p , then the value $2\pi - \theta_p$ is called the curvature of the polyhedral surface at p ; if p is not a vertex, then its curvature is defined to be zero.

12.6. Exercise. Assume that the surface of a nondegenerate tetrahedron T has curvature π at each of its vertices. Show that

- (a) all faces of T are congruent;
- (b) the line passing thru midpoints of opposite edges of T intersects these edges at right angles.

Claim 12.2 says that *surface of a convex polyhedron has nondegenerate curvature*. However this definition works only for polyhedral surfaces. Now we show that it agrees with the CBB(0) definition.

12.7. Proposition. A polyhedral surface with nonnegative curvature at each vertex is CBB(0).

Proof. Denote the surface by Σ . By 2.20, it is sufficient to check that $\text{dist}_p^2 \circ \gamma$ is 1-concave for any geodesic γ and a point p in Σ .

We can assume that p is not a vertex; the vertex case can be done by approximation. Further, by 12.5, we may assume that γ does not contain vertices.

Given a point $x = \gamma(t_0)$, choose a geodesic $[px]$. Again, by 12.5, $[px]$ does not contain vertices. Therefore a small neighborhood of $U \supset [px]$ can be unfolded on a plane; that is, there is an injective length-preserving map $z \mapsto \tilde{z}$ of U into the Euclidean plane. Note that this way we map part of γ in U to a line segment. Let

$$\tilde{f}(t) := \frac{1}{2} \cdot \text{dist}_p^2 \circ \tilde{\gamma}(t).$$

Note that $\tilde{f}(t_0) \geq f(t_0)$. Further, since the unfolding $z \mapsto \tilde{z}$ preserves lengths of curves, we get $\tilde{f}(t) \geq f(t)$ if t is sufficiently close to t_0 . That is, \tilde{f} is a local upper support of f at t_0 . Evidently, $\tilde{f}'' \equiv 1$; therefore $f'' \leq 1$. It remains to apply 2.20. \square

12.8. Exercise. Prove the converse to the proposition; that is, show that if a polyhedral surface is CBB(0), then it has nonnegative curvature at each vertex.

C Surface of convex body

12.9. Advanced exercise. Let $K_\infty, K_1, K_2, \dots$ be convex bodies in \mathbb{E}^3 . Denote by S_n the surface of K_n with induced length metric. Suppose $K_n \rightarrow K_\infty$ in the sense of Hausdorff. Show that $S_n \rightarrow S_\infty$ in the sense of Gromov–Hausdorff.

12.10. Proposition. The surface of a convex body is CBB(0).

Note that any convex body is a Hausdorff limit of a sequence of convex polyhedra. Therefore, the proposition follows from 12.7, 12.9, and 8.9.

D Comments

Note that 12.1 and 12.7 imply that surface of convex body is a sphere with nonnegative curvature in the sense of Alexandrov. The celebrated theorem of Alexandrov states that the converse also holds if we allow degeneration of convex bodies to plane figures; the surface of a plane figure is defined as its doubling across the boundary. In other words, any geodesic CBB(0) metric on the sphere is isometric to a surface of convex body. Moreover this convex body is unique up to congruence. The last result is due to Alexei Pogorelov [58].

Originally, Alexandrov proved the statement for polyhedral metrics on the sphere; this proof is sketched in the appendix. Then he used 12.9 to extend the result to arbitrary geodesic CBB(0) metric on the sphere.

12.11. Advanced exercise. Let Σ be the surface of a nondegenerate convex body $K \subset \mathbb{E}^3$; we assume that Σ is equipped with its induced length metric.

- (a) Show that any geodesic γ in Σ is one-side differentiable as a curve in \mathbb{E}^3
- (b) Let γ_1 and γ_2 be geodesic paths in Σ that start at one point $p = \gamma_1(0) = \gamma_2(0)$. Suppose $x_1 = \gamma_1(1)$, $x_2 = \gamma_2(1)$, $y_1 = p + \gamma_1^+(0)$, and $y_2 = p + \gamma_2^+(0)$. Show that

$$|x_1 - x_2|_\Sigma \leq |y_1 - y_2|_W,$$

where W is the complement to the interior of K .

Appendix A

Alexandrov's embedding theorem

BY NINA LEBEDEVA AND ANTON PETRUNIN

A Introduction

Intrinsic distance between two points on the surface of a convex polyhedron is defined as the length of a shortest curve on the surface between these points.

Recall that the sum of angles at the tip of a convex polyhedral angle is less than $2\cdot\pi$; this statement can be found in a school textbook [29, § 48].

It is easy to see that the surface of a convex polyhedron is homeomorphic to the sphere. Therefore the statements above imply that the surface of a convex polyhedron equipped with its intrinsic metric is an example of a *polyhedral metric on the sphere with the sum of angles around each vertex at most $2\cdot\pi$* ; a metric is called polyhedral if the sphere admits a triangulation such that every triangle is congruent to a plane triangle.

Alexandrov's theorem states that the converse holds if one includes in the consideration *twice covered polygons*. In other words, we assume that a polyhedron can degenerate to a plane polygon; in this case, its surface is defined as two copies of the polygon glued along their boundary.

Further, we assume that a polyhedron can degenerate to a plane polygon.

A.1. Alexandrov's theorem.

- I. A polyhedral metric on the sphere is isometric to the surface of a convex polyhedron if and only if the sum of angles around each of its vertex is not greater than 2π .
- II. Moreover, a convex polyhedron is defined up to congruence by the intrinsic metric on its surface.


A. D. Alexandrov has many remarkable theorems, but in our opinion, this theorem is the most remarkable. At the same time, its proof is elementary; it could be explained to anyone familiar with basic topology.

This theorem has many applications. In particular, it is used in the proof of its generalization [6] that gives a complete description of intrinsic metrics on the sphere that are isometric to convex surfaces in the Euclidean space. The latter statement is fundamental in a branch of modern mathematics — the so-called *Alexandrov geometry*.

The first part is central; it is called the *existence theorem*. The second part is called the *uniqueness theorem*; it is a slight variation of Cauchy's theorem about polyhedrons. (There is another uniqueness theorem of Alexandrov that generalizes Minkowski's theorem about polyhedrons.)

According to the theorem, a convex polyhedron is completely defined by the intrinsic metric of its surface. In particular, knowing the metric we could find the position of the edges. However, in practice, it is not easy to do. For example, the surface glued from a rectangle as shown on the diagram defines a tetrahedron. Some of the glued lines appear inside facets of the tetrahedron and some edges (dashed lines) do not follow the sides of the rectangle.

The theorem was proved by A. D. Alexandrov in 1941 [5]; we will present a sketch of his proof. A complete proof is nicely written by A. D. Alexandrov in his book [4]. Yet another proof was found by Yu. A. Volkov in his thesis [63]; it uses a deformation of three-dimensional polyhedral space.



B Space of polyhedrons and metrics

Space of polyhedrons. Let us denote by Φ the space of all convex polyhedrons in the Euclidean space, including polyhedrons that degenerate to a plane polygon. Polyhedra in Φ will be considered up to a motion of the space, and the whole space Φ will be considered

with the natural topology (an intuitive meaning of closeness of two polyhedrons should be sufficient).

Further, denote by Φ_n the polyhedrons in Φ with exactly n vertices. Since any polyhedron has at least 3 vertices, the space Φ admits a subdivision into a countable number of subsets Φ_3, Φ_4, \dots

Space of polyhedral metrics. The space of polyhedral metrics on the sphere with the sum of angles around each point at most $2\cdot\pi$ will be denoted by Ψ . The metrics in Ψ will be considered up to an isometry, and the whole space Ψ will be equipped with the natural topology (again, an intuitive meaning of closeness of two metrics is sufficient).

A point on the sphere with the sum of angles strictly less than $2\cdot\pi$ will be called an essential vertex. The subset of Ψ of all metrics with exactly n essential vertices will be denoted by Ψ_n . It is easy to see that any metric in Ψ has at least 3 essential vertices. Therefore Ψ is subdivided into countably many subsets Ψ_3, Ψ_4, \dots

From a polyhedron to its surface. Recall that the surface of a convex polyhedron is a sphere with a polyhedral metric such that the sum of angles around each point is at most $2\cdot\pi$. Therefore passing from a polyhedron to its surface defines a map

$$\iota: \Phi \rightarrow \Psi.$$

Note that the number of vertices of a polyhedron is equal to the number of essential vertices of its surface. In other words, $\iota(\Phi_n) \subset \Psi_n$ for any $n \geq 3$.

C About the proof

Using the notation introduced in the previous section, we can give the following more exact formulation of Alexandrov's theorem:

A.2. Reformulation. *For any integer $n \geq 3$, the map ι is a bijection from Φ_n to Ψ_n .*

We sketch the original proof of A. D. Alexandrov. It is based on the construction of a one-parameter family of polyhedrons that starts at arbitrary polyhedron and ends at a polyhedron with its surface isometric to the given one. This type of argument is called the continuity method; it is often used in the theory of differential equations.

The two parts of the first formulation will be proved separately.

Part II. Let us show that the map $\iota: \Phi_n \rightarrow \Psi_n$ is injective; in other words, a convex polyhedron is defined by the intrinsic metric on its surface up to a motion of the space.

The last statement is analogous to the Cauchy theorem about polyhedrons, and the proof goes along the same lines.

The Cauchy theorem states that facets of a polyhedron together with the gluing rule completely describe a convex polyhedron; its proof is given in many classical popular texts [1, 16, 60].

Part I. Let us prove that $\iota: \Phi_n \rightarrow \Psi_n$ is surjective. This part of the proof is subdivided into the following lemmas:

A.3. Lemma. *For any integer $n \geq 3$, the space Ψ_n is connected.*

The proof of this lemma is not complicated, but it requires ingenuity; it can be done by the direct construction of a one-parameter family of metrics in Ψ_n that connects two given metrics. Such a family can be obtained by a sequential application of the following construction and its inverse.

Let M be a sphere with metric from Ψ_n . Suppose v and w are essential vertices in M . Let us cut M along a shortest line from v to w . Note that the shortest line cannot pass thru an essential vertex of M . Further, note that there is a three-parameter family of patches that can be used to patch the cut so that the obtained metric remains in Ψ_n ; in particular, the obtained metric has exactly n essential vertices (after the patching, the vertices v and w may become inessential).

A.4. Lemma. *The map $\iota: \Phi_n \rightarrow \Psi_n$ is open, that is, it maps any open set in Φ_n to an open set in Ψ_n .*

In particular, for any $n \geq 3$, the image $\iota(\Phi_n)$ is open in Ψ_n .

This statement is very close to the so-called invariance of domain theorem; the latter states that a continuous injective map between manifolds of the same dimension is open.

According to part II, ι is injective. The proof of the invariance of domain theorem can be adapted to our case since both spaces Φ_n and Ψ_n are $(3 \cdot n - 6)$ -dimensional and both look like manifolds, altho, formally speaking, they are *not* manifolds. In a more technical language, Φ_n and Ψ_n have the natural structure of $(3 \cdot n - 6)$ -dimensional orbifolds, and the map ι respects the orbifold structure.

We will only show that both spaces Φ_n and Ψ_n are $(3 \cdot n - 6)$ -dimensional.

Choose a polyhedron P in Φ_n . Note that P is uniquely determined by the $3 \cdot n$ coordinates of its n vertices. We can assume that the first vertex is the origin, the second has two vanishing coordinates and the

third has one vanishing coordinate; therefore, all polyhedrons in Φ_n that lie sufficiently close to P can be described by $3 \cdot n - 6$ parameters. If P has no symmetries then this description can be made one-to-one; in this case, a neighborhood of P in Φ_n is a $(3 \cdot n - 6)$ -dimensional manifold. If P has a nontrivial symmetry group, then this description is not one-to-one but it does not have an impact on the dimension of Φ_n .

The case of polyhedral metrics is analogous. We need to construct a subdivision of the sphere into plane triangles using only essential vertices. By Euler's formula, there are exactly $3 \cdot n - 6$ edges in this subdivision. Note that the lengths of edges completely describe the metric, and slight changes of these lengths produce a metric with the same property.

A.5. Lemma. *The map $\iota: \Phi_n \rightarrow \Psi_n$ is closed; that is, the image of a closed set in Φ_n is closed in Ψ_n .*

In particular, for any $n \geq 3$, the set $\iota(\Phi_n)$ is closed in Ψ_n .

Choose a closed set Z in Φ_n . Denote by \bar{Z} the closure of Z in Φ ; note that $Z = \Phi_n \cap \bar{Z}$. Assume $P_1, P_2, \dots \in Z$ is a sequence of polyhedrons that converges to a polyhedron $P_\infty \in \bar{Z}$. Note that $\iota(P_n)$ converges to $\iota(P_\infty)$ in Ψ . In particular, $\iota(\bar{Z})$ is closed in Ψ .

Since $\iota(\Phi_n) \subset \Psi_n$ for any $n \geq 3$, we have $\iota(Z) = \iota(\bar{Z}) \cap \Psi_n$; that is, $\iota(Z)$ is closed in Ψ_n .

Summarizing, $\iota(\Phi_n)$ is a nonempty closed and open set in Ψ_n , and Ψ_n is connected for any $n \geq 3$. Therefore, $\iota(\Phi_n) = \Psi_n$; that is, $\iota: \Phi_n \rightarrow \Psi_n$ is surjective. \square

Acknowledgments. We want to thank Stephanie Alexander, Yuri Burago, and Jules Tsukahara for help. The authors were partially supported by RFBR grant 20-01-00070 and NSF grant DMS-2005279.

Semisolutions

1.3. Given a pair of points p and q , choose a sequence of paths γ_n from p to q such that

$$\text{length } \gamma_n \rightarrow |p - q| \quad \text{as } n \rightarrow \infty;$$

it exists since we are in a length space. Note that we can assume that each γ_n is parametrized proportional to the arc length; in particular, γ_n are equicontinuous. Show that paths γ_n lie in a closed ball, say $\overline{B}[p, r]$ for some $r < \infty$. Since the space is proper, $\overline{B}[p, r]$ is compact. By Arzelà–Ascoli theorem, we can pass to a converging subsequence of γ_n . Show that its limit is a geodesic path from p to q .

1.2. Choose a Cauchy sequence x_n in $(\mathcal{X}, \|* - *\|)$; it is sufficient to show that a subsequence of x_n converges.

Observe that the sequence x_n is Cauchy in $(\mathcal{X}, |* - *|)$; denote its limit by x_∞ .

Passing to a subsequence, we can assume that $\|x_n - x_{n+1}\| < \frac{1}{2^n}$. It follows that there is a 1-Lipschitz path γ in $(\mathcal{X}, \|* - *\|)$ such that $x_n = \gamma(\frac{1}{2^n})$ for each n and $x_\infty = \gamma(0)$. Therefore,

$$\|x_\infty - x_n\| \leq \text{length } \gamma|_{[0, \frac{1}{2^n}]} \leq \frac{1}{2^n}.$$

In particular, x_n converges to x_∞ in $(\mathcal{X}, \|* - *\|)$.

Source: [27, Corollary]; see also [47, Lemma 2.3].

1.1. Choose a sequence of positive numbers $\varepsilon_n \rightarrow 0$ and a finite ε_n -net N_n of K for each n . We can assume that $\varepsilon_0 > \text{diam } K$, and N_0 is a one-point set. If $|x - y| < \varepsilon_k$ for some $x \in N_{k+1}$ and $y \in N_k$, then connect them by a curve of length at most ε_k .

Let K' be the union of all these curves and K . Show that K' is compact and path-connected.

Source: This problem is due to Eugene Bilokopytov [8].

1.5. Choose a sequence $\varepsilon_n > 0$ that converges to zero very fast, say such that $\sum_n 10^n \cdot \varepsilon_n$ is small. Follow the argument in the proof of Menger's lemma, taking ε_n -midpoints at the n^{th} stage.

1.6. Let us write the Riemannian metric on $\mathbb{M}(\kappa)$ in the polar coordinates (θ, r) ; it has form $\begin{pmatrix} h^2 & 0 \\ 0 & 1 \end{pmatrix}$, where $h = h(\kappa, r)$. Calculate $h(\kappa, r)$. Show that for fixed r , the function $r \mapsto h(\kappa, r)$ is nonincreasing in the domain of definition. Suppose $\kappa < K$, consider the map $\mathbb{M}(\kappa) \rightarrow \mathbb{M}(K)$ that sends a point to the point with the same polar coordinates. Show that this map is short in the domain of definition. Use it to prove the statement in the exercise.

1.7. Show and use that $\tilde{\angle}(p_y^x)_{\mathbb{S}^2} - \tilde{\angle}(p_y^x)_{\mathbb{E}^2} = O(|p - x|^2 + |p - y|^2)$ and $\tilde{\angle}(p_y^x)_{\mathbb{E}^2} - \tilde{\angle}(p_y^x)_{\mathbb{H}^2} = O(|p - x|^2 + |p - y|^2)$.

1.8. Consider a hinge in the plane \mathbb{R}^2 with a metric defined by norm, say by ℓ^∞ -norm.

1.10. Assume $\angle[p_z^x] + \angle[p_z^y] < \pi$. By 1.9, $\angle[p_y^x] < \pi$. Therefore, $\tilde{\angle}(p_{\bar{y}}^{\bar{x}}) < \pi$ for some $\bar{x} \in [px]$ and $\bar{y} \in [py]$. Hence

$$|p - \bar{x}| + |\bar{y} - p| < |\bar{x} - \bar{y}|$$

— a contradiction.

1.11. Denote by α the arc-length parametrization of $[qp]$ from q to p . Choose $\varepsilon > 0$. Observe that

$$|\gamma(t) - \alpha(\frac{1}{\varepsilon} \cdot t)|^2 \leq t^2 \cdot (1 - \frac{2}{\varepsilon} \cdot \cos \varphi + \frac{1}{\varepsilon^2}) + o(t^2),$$

where $\varphi = \angle[q_x^p]$. By the triangle inequality

$$|p - \gamma(t)| \leq |\gamma(t) - \alpha(\frac{1}{\varepsilon} \cdot t)| + |q - p| - \frac{1}{\varepsilon} \cdot t.$$

Conclude that

$$|p - \gamma(t)| \leq |q - p| - t \cdot \cos \varphi + \delta(\varepsilon) \cdot t + o(t),$$

where $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. The statement follows since $\varepsilon > 0$ is arbitrary.

2.2. The definition of CBB spaces (2.1) reduces our question to the following. *Any spherical triangle has perimeter at most $2 \cdot \pi$.* Choose a spherical triangle $[xyz]$. Let x' be the antipode of x ; that is $x' = -x$. The spherical triangle inequality implies that

$$|x - z|_{\mathbb{S}^2} \leq |y - x'|_{\mathbb{S}^2} + |x' - z|_{\mathbb{S}^2}.$$

Observe that

$$|x - y|_{\mathbb{S}^2} + |y - x'|_{\mathbb{S}^2} = \pi, \quad \text{and} \quad |x - z|_{\mathbb{S}^2} + |z - x'|_{\mathbb{S}^2} = \pi.$$

Hence

$$|x - y|_{\mathbb{S}^2} + |x - z|_{\mathbb{S}^2} + |y - z|_{\mathbb{S}^2} \leq 2 \cdot \pi.$$

2.3. For the only-if part consider the following two cases.

If $\tilde{\angle}(p_{x_1}^{x_2}) + \tilde{\angle}(p_{x_3}^{x_2}) \geq \pi$, then choose two model triangles $[qy_1y_2] = \tilde{\Delta}(px_1x_2)$ and $[qy_2y_3] = \tilde{\Delta}(px_2x_y)$ that lie on the opposite sides of $[qy_2]$. By comparison, $|y_1 - y_3| \geq |x_1 - x_3|$. Therefore the obtained configuration meets all the conditions.

If $\tilde{\angle}(p_{x_1}^{x_2}) + \tilde{\angle}(p_{x_3}^{x_2}) \geq \pi$, then choose two model triangles $[qy_1y_2] = \tilde{\Delta}(px_1x_2)$ and take y_3 on the extension of $[y_1q]$ behind q such that $|q - y_3| = |p - x_3|$. Then $\angle[q_{y_3}^{y_2}] \geq \tilde{\angle}(p_{x_3}^{x_2})$, therefore $|y_2 - y_3| \geq |x_2 - x_3|$. Further, $|y_2 - y_3| = |x_2 - p| + |p - x_3| \geq |x_2 - x_3|$, and again, the obtained configuration meets all the conditions.

To prove the if part, choose a configuration q, y_1, y_2, y_3 that meets all the conditions and maximize the sum

$$|y_1 - y_2| + |y_2 - y_3| + |y_3 - y_1|.$$

Show that that q lies in the solid triangle $y_1y_2y_3$; in particular

$$\angle[q_{y_2}^{y_1}] + \angle[q_{y_3}^{y_2}] + \angle[q_{y_1}^{y_3}] = 2 \cdot \pi.$$

Moreover, $|q - y_i| = |p - x_i|$ for each i . Applying that increasing the opposite side in a plane triangle increases the corresponding angle, we get

$$\tilde{\angle}(p_{x_2}^{x_1}) + \tilde{\angle}(p_{x_3}^{x_2}) + \tilde{\angle}(p_{x_1}^{x_3}) \leq 2 \cdot \pi.$$

2.5. Consider model triangles $[\tilde{p}\tilde{x}\tilde{z}] = \tilde{\Delta}(pxz)$ and $[\tilde{p}\tilde{y}\tilde{z}] = \tilde{\Delta}(pyz)$ that share side $[\tilde{p}\tilde{z}]$ and lie on its opposite sides. Note that

$$\begin{aligned} |\tilde{x} - \tilde{y}|_{\mathbb{E}^2} &\geq |\tilde{x} - \tilde{y}|_{\mathbb{E}^2} + |\tilde{x} - \tilde{y}|_{\mathbb{E}^2} = \\ &= |x - z|_{\mathcal{X}} + |z - y|_{\mathcal{X}} = \\ &= |x - y|_{\mathcal{X}}, \end{aligned}$$

where \mathcal{X} is our metric space. It remains to apply the monotonicity of angle in a triangle with respect to its opposite side.

2.7. Apply 2.6.

2.9. Without loss of generality, we can assume that $|p - x| \leq |p - y|$. Choose $\bar{x} \in [px]$; let $\bar{y} \in [py]$ be such that $|p - \bar{x}| = |p - \bar{y}|$. Apply 2.6 to show that $\bar{x} = \bar{y}$. Conclude that $[px] \subset [py]$.

2.10. Assume that there are two distinct geodesics from z to x . Then we can choose distinct points p and q on these geodesics such that $|z - p| = |z - q|$. Observe that $\tilde{\angle}(z \frac{p}{q}) > 0$. By triangle inequality, we get

$$|x - p| + |p - y| \leq |x - p| + |p - z| + |z - y| = |x - z| + |z - y|$$

Observe that $\tilde{\angle}(z \frac{x}{y}) = \pi$. Therefore $\angle[z \frac{x}{y}] = \pi$ for any geodesic $[zx]$.

2.11. By 1.10, we have

$$\angle[p \frac{x}{z}] + \angle[p \frac{y}{z}] \geq \pi.$$

Since $z \in]xy[$ we have

$$\tilde{\angle}(z \frac{\bar{x}}{\bar{y}}) = \pi$$

for any $\bar{x} \in [xz[$ and $\bar{y} \in]zy]$. By comparison, we have that

$$\tilde{\angle}(z \frac{\bar{x}}{\bar{p}}) + \tilde{\angle}(z \frac{\bar{p}}{\bar{y}}) \leq \pi$$

for any $\bar{p} \in]zp]$. Passing to the limit as $|z - \bar{x}| \rightarrow 0$, $|z - \bar{y}| \rightarrow 0$, and $|z - \bar{p}| \rightarrow 0$, we get the statement.

2.12. Without loss of generality, we can assume that x , v , w , and y appear on $[xy]$ in this order. By 2.6,

$$\tilde{\angle}(x \frac{y}{p}) \geq \tilde{\angle}(x \frac{w}{p}) \geq \tilde{\angle}(x \frac{v}{p}).$$

Hence, \Rightarrow follows.

By Alexandrov's lemma,

$$\begin{aligned} \tilde{\angle}(x \frac{y}{p}) = \tilde{\angle}(x \frac{v}{p}) &\iff \tilde{\angle}(y \frac{x}{p}) = \tilde{\angle}(y \frac{v}{p}), \\ \tilde{\angle}(x \frac{y}{p}) = \tilde{\angle}(x \frac{w}{p}) &\iff \tilde{\angle}(y \frac{x}{p}) = \tilde{\angle}(y \frac{w}{p}). \end{aligned}$$

Whence, \Leftarrow follows.

2.13. Suppose $\angle[x_\infty \frac{y_\infty}{z_\infty}] > \alpha$. Then we can choose $\bar{y}_\infty \in]x_\infty y_\infty]$ and $\bar{z}_\infty \in]x_\infty z_\infty]$ such that $\tilde{\angle}(x_\infty \frac{\bar{y}_\infty}{\bar{z}_\infty}) > \alpha$. Now choose $\bar{y}_n \in]x_n y_n]$ and $\bar{y}_n \in]x_n z_n]$ such that $\bar{y}_n \rightarrow \bar{y}_\infty$ and $\bar{z}_n \rightarrow \bar{z}_\infty$. Observe that

$$\varliminf_{n \rightarrow \infty} \angle[x_n \frac{y_n}{z_n}] \geq \varliminf_{n \rightarrow \infty} \tilde{\angle}(x_n \frac{\bar{y}_n}{\bar{z}_n}) \geq \alpha,$$

hence the result.

2.16. The Urysohn space provides an example; see for example [54, Lecture 2].

2.18. The natural map will be denoted by $\tilde{p} \mapsto p$. Apply 2.8 twice to show that $|\tilde{p} - \tilde{q}| \leq |p - q|$ for any $\tilde{p} \in [\tilde{x}\tilde{y}]$ and $\tilde{q} \in [\tilde{x}\tilde{z}]$. Conclude that $[xyz]$ is fat.

2.19; (a). Choose a triangle $[0vw]$. Note that $m = \frac{1}{2}(v + w)$ is the midpoint of $[vw]$.

Since $[0vw]$ is thin, we get

$$2 \cdot \left| \frac{1}{2}(v + w) \right|^2 + 2 \cdot \left| \frac{1}{2}(v - w) \right|^2 \leq |v|^2 + |w|^2.$$

Note this inequality implies the opposite one; it follows if we rewrite it via $x = \frac{1}{2}(v + w)$ and $y = \frac{1}{2}(v - w)$. Hence we have

$$2 \cdot \left| \frac{1}{2}(v + w) \right|^2 + 2 \cdot \left| \frac{1}{2}(v - w) \right|^2 = |v|^2 + |w|^2$$

for any v, w . That is the norm is quadratic and the statement follows.

(b). Apply the same argument changing the signs of inequalities.

3.3.

3.5. Note that \mathcal{X} has no defined spherical model angles; therefore it is CBB(1).

However, \mathcal{X} is not CBB(0) since

$$\tilde{\angle}(p_{x_2}^{x_1})_{\mathbb{E}^2} = \tilde{\angle}(p_{x_3}^{x_2})_{\mathbb{E}^2} = \tilde{\angle}(p_{x_3}^{x_1})_{\mathbb{E}^2} = \pi.$$

3.6. Suppose $\angle[m_p^x] \neq 0$ and $\angle[m_p^x] \neq \pi$, or equivalently $\angle[m_q^x] \neq 0$.

We can assume that $|p - q|$ only slightly exceeds π , so $|p - m| < \pi$ and $|q - m| < \pi$. We can also assume that $|x - m| < \pi$. Use the comparison to show that

$$|p - x| + |q - x| < |p - q|,$$

and arrive at a contradiction with the triangle inequality.

Extend $[pq]$ to a maximal local geodesic γ . It might be a closed or a line segment. Argue as above to show that any point lies on γ and make a conclusion.

3.7. Arguing by contradiction, suppose

$$\textbf{①} \quad |p - q| + |q - r| + |r - p| > 2 \cdot \pi$$

for $p, q, r \in \mathcal{L}$. Rescaling the space slightly, we can assume that $\text{diam } \mathcal{L} < \pi$, but the inequality **①** still holds. By 3.4, after rescaling \mathcal{L} is still CBB(1).

Take $z_0 \in [qr]$ on maximal distance from p . Consider the following model configuration: two geodesics $[\tilde{p}\tilde{z}_0]$, $[\tilde{q}\tilde{r}]$ in \mathbb{S}^2 such that

$$\begin{aligned} |\tilde{p} - \tilde{z}_0| &= |p - z_0|, & |\tilde{q} - \tilde{r}| &= |q - r|, \\ |\tilde{z}_0 - \tilde{q}| &= |z_0 - q|, & |\tilde{z}_0 - \tilde{r}| &= |z_0 - q|, \end{aligned}$$

and

$$\angle[\tilde{z}_0 \tilde{q} \tilde{p}] = \angle[\tilde{z}_0 \tilde{r} \tilde{p}] = \frac{\pi}{2}.$$

Let $\tilde{z} \in [\tilde{q}\tilde{r}]$, and let $z \in [qr]$ be the corresponding point. By comparison, $|p - z| \leq |\tilde{p} - \tilde{z}|$ for points z near z_0 . Moreover, this inequality holds as far as

$$|\tilde{p} - \tilde{z}_0| + |\tilde{z}_0 - \tilde{z}| + |\tilde{p} - \tilde{z}| < 2 \cdot \pi.$$

But this inequality holds for all \tilde{z} since $|\tilde{p} - \tilde{z}_0| < \pi$, $|\tilde{z}_0 - \tilde{q}| < \pi$, and $|\tilde{z}_0 - \tilde{r}| < \pi$. Hence we get $|p - q| \leq |\tilde{p} - \tilde{q}|$ and $|p - r| \leq |\tilde{p} - \tilde{r}|$. The latter contradicts **1**.

3.9. Suppose such point does not exist; that is, for any $p \in \mathcal{X}$ there is a point p' such that $r(p') \leq (1 - \varepsilon) \cdot r(p)$ and $|p - p'| < \frac{1}{\varepsilon} \cdot r(p)$. Construct a sequence of points p_0, p_1, \dots such that $p_n = p'_{n-1}$ for any n . Show that this sequence is Cauchy; denote its limit by p_∞ . Arrive at a contradiction by showing that $r(p_\infty) \leq 0$.

4.1; (a). Suppose $\uparrow_{[px_n]} \not\rightarrow \uparrow_{[px_\infty]}$. Since Σ_p is compact we may pass to a converging subsequence of $\uparrow_{[px_n]}$; denote by ξ its limit. We may assume that $\angle(\uparrow_{[px_\infty]}, \xi) > 0$.

Denote by γ_n and γ_∞ the arc-length parametrization of $[px_n]$ and $[px_\infty]$ from p . Choose a geodesic α that starts from p and goes in a direction sufficiently close to ξ . By comparison we can choose α so that

$$|\alpha(t) - \gamma_n(t)| < \varepsilon \cdot t$$

for all large n and all sufficiently small t . Moreover, we can assume that

$$|\alpha(t) - \gamma_\infty(t)| > a \cdot t$$

for some fixed $a > 0$ and all small t . These two inequalities imply that

$$|\gamma_n(t) - \gamma_\infty(t)| > \frac{a}{2} \cdot t$$

for all small t and all large n . On the other hand, by assumption, $|\gamma_n(t) - \gamma_\infty(t)| \rightarrow 0$ as $n \rightarrow \infty$ — a contradiction.

(b) ???

4.2. Note that any point of Cone \mathcal{X} can be connected to the origin by a geodesic. Given a nonzero element $v \in \text{Cone } \mathcal{X}$, denote by v' its projection in \mathcal{X} .

Suppose \mathcal{X} is π -geodesic. Choose two nonzero elements $v, w \in \text{Cone } \mathcal{X}$; let $\alpha = \angle(v, w) = |v' - w'|_{\mathcal{X}}$. If $\alpha \geq \pi$, then the product of geodesics $[v0] \cup [0w]$ forms a geodesic $[vw]$. If $\alpha < \pi$, there is a geodesic $\gamma: [0, \alpha] \rightarrow \mathcal{X}$ from v' to w' . Consider hinge $[\tilde{o} \tilde{v}_{\tilde{w}}]$ in the plane such that $\angle[\tilde{o} \tilde{v}_{\tilde{w}}] = \alpha$, $|\tilde{o} - \tilde{v}| = |v|$, and $|\tilde{o} - \tilde{w}| = |w|$. Let $t \mapsto (\varphi(t), r(t))$ be geodesic $[\tilde{v}\tilde{w}]$ written in the polar coordinates with origin \tilde{o} , so that $\varphi(0) = 0$. Show that $t \mapsto r(t) \cdot \gamma \circ \varphi(t)$ is a geodesic from v to w ; here we identify \mathcal{X} with the unit sphere in Cone \mathcal{X} .

To prove the converse, try to revert the steps in the argument above.

4.4. From 2.20, this inequality follows in the sense of distributions, and hence in any other sense.

4.5. Since angles are defined, it follows that

$$|\gamma_1(t) - \gamma_2(t)| \leq \theta \cdot t$$

for all small $t > 0$. Since f is L -Lipschitz, we get

$$|f(\gamma_1(t)) - f(\gamma_2(t))| \leq L \cdot \theta \cdot t,$$

hence the statement.

4.6; (a) Note that we can assume that there is a geodesic in the direction of v , and apply 1.11.

(b). By (a) we have an \leq inequality. Suppose this inequality is strict for some v . We can assume that $|v| = 1$ and there is a geodesic, say γ in the direction of v . Suppose ... Let $\alpha = ???$.

The function $f = \text{dist}_q \circ \gamma$ is Lipschitz; By Rademacher's theorem it is differentiable almost everywhere; moreover,

$$f(t) - f(0) = \int_0^t f'(t) \cdot dt.$$

Suppose $f'(t)$ is defined. Use (a) to show that $f'(t) = -\cos \alpha(t)$, where $\alpha(t)$ is the angle between γ and a geodesic from $\gamma(t)$ to q . Note that we can choose a sequence $t_n \rightarrow 0$ such that

$$\lim_{n \rightarrow \infty} \alpha(t_n) \leq \alpha.$$

Consider a sequence of geodesics $[p\gamma(t_n)]$. Since the space is proper, we can pass to its convergent subsequence. Its limit is a geodesic from p

to q , denote it by $[pq]$. Observe that $[pq]$ makes angle at most α with γ — a contradiction.

4.8. Let $\gamma: [0, \ell] \rightarrow \mathcal{L}$ be the geodesic $[xy]$ parametrized from x to y , and let $\varphi = f \circ \gamma$. Observe that

$$\varphi'(0) = \mathbf{d}_x f(\uparrow_{[xy]}) \leq \langle \uparrow_{[xy]}, \nabla_x f \rangle.$$

The same way we get $-\varphi'(\ell) \leq \langle \uparrow_{[yx]}, \nabla_y f \rangle$. Since f is λ -concave, we have

$$\begin{aligned} f(y) &\leq f(x) + \varphi'(0) \cdot \ell + \frac{\lambda}{2} \cdot \ell^2, \\ f(x) &\leq f(y) - \varphi'(\ell) \cdot \ell + \frac{\lambda}{2} \cdot \ell^2. \end{aligned}$$

Hence the statement follows.

4.11.

4.12.

4.13.

5.7. Note that $f \circ \alpha$ is a nondecreasing function. Apply 4.6a and the definition of gradient to show that

$$-\mathbf{d}_{\alpha(t)} \text{dist}_{\alpha(t_3)}(\nabla_{\alpha(t)} f) \geq \langle \nabla_{\alpha(t)}, \uparrow_{[\alpha(t)\alpha(t_3)]} \rangle \geq \mathbf{d}_{\alpha(t)}(\uparrow_{[\alpha(t)\alpha(t_3)]}) \geq 0$$

for any $t < t_3$. Conclude that the function $t \mapsto \text{dist}_{\alpha(t_3)} \circ \alpha(t)$ is noncreasing for $t \leq t_3$.

5.8.

5.9.

6.2. Apply 4.4.

6.3. By the triangle inequality,

$$|\gamma(-t) - x| + |\gamma(t) - x| - 2 \cdot t \geq 0$$

for any $t \geq 0$. Passing to the limit as $t \rightarrow \infty$, we get the result.

6.7.

7.2. Observe that

$$\begin{aligned} \langle u, u \rangle + \langle v, u \rangle + \langle w, u \rangle &\geq 0, \\ \langle u, v \rangle + \langle v, v \rangle + \langle w, v \rangle &\geq 0, \\ \langle u, w \rangle + \langle v, w \rangle + \langle w, w \rangle &= 0. \end{aligned}$$

Add the first two inequalities and subtract the last identity.

7.7.

7.8. Show and use that

$$\langle u, x \rangle + \langle v, x \rangle + \langle w, x \rangle \geq 0$$

and

$$\langle u, -x \rangle + \langle v, -x \rangle + \langle w, -x \rangle \geq 0.$$

7.9. Part \Rightarrow is evident. To prove part \Leftarrow , observe that

$$\langle u^*, u^* \rangle = -\langle u, u^* \rangle \leq \langle u, u \rangle$$

and since $|u| = |u^*|$, we have equality.

7.11; (a). Let $S_n \subset \mathcal{L}$ be defined by inequality $|\nabla_x f| > 1 - \frac{1}{n}$. Apply 4.13a to show that S_n is open. Choose a point $q \neq p$, observe that $|\nabla_x f| = 1$ for any point $x \in]pq[$. Conclude that S_n is dense in \mathcal{L} . Observe and use that $S = \bigcap_n S_n$.

(b)+(c). Apply 7.9.

7.14. Apply 7.11b to show that for any finite set of points p_1, \dots, p_n there is a G-delta dense set of points x such that $\text{Lin}_x \ni \uparrow_{[xp_i]}$ for every i . ???

7.15.

7.16.

7.18.

8.5.

8.8.

8.10.

8.13.

8.16.

8.17.

10.1.

10.2.

10.5.

10.6. (a) (b) (c)

10.8.

10.9.

10.10.

10.13.

10.14.

10.15.

11.2.

11.3.

11.4.

11.10; (a). Our \mathbb{S}^1 is a commutative subgroup of $\mathrm{SO}(3)$. Therefore it is a subgroup of a maximal torus in $\mathrm{SO}(3)$. Show that the described torus action is induced by a maximal torus in $\mathrm{SO}(3)$. Use that maximal tori in $\mathrm{SO}(3)$ are conjugate.

(b). Cut \mathbb{S}^3 into two solid tori the Clifford torus $\frac{1}{\sqrt{2}} \cdot \mathbb{S}^1 \times \mathbb{S}^1$. Observe that the quotient of each solid torus is a disc; conclude that $\Sigma_{p,q}$ is a sphere. The torus action on \mathbb{S}^3 induce the needed \mathbb{S}^1 -action on $\Sigma_{p,q}$.

(c)+(d)+(e). Straightforward calculations.

(f). Consider the map $\Sigma_{p,q} \rightarrow \Sigma_{1,1}$ that sends poles to poles, preserve the distance to the poles and respects the \mathbb{S}^1 action.

11.16; (a). Suppose $\#_{m-1}(\Gamma) \geq 3$; that is $\mathcal{L} = \mathbb{E}^m/\Gamma$ has at least 3 boundary components. Follow Case 3 in the proof 11.8 to glue a train-space from copies of \mathcal{L} using two of these components. Show that the obtained space splits and arrive at a contradiction.

(Alternatively, apply a similar construction to all components of the boundary. Show that obtained space has exponential volume growth; that is, there is $a > 1$ such that $\mathrm{vol} B(p, r) > a^r$ for all large r . Arrive at a contradiction with Bishop–Gromov inequality.)

(b). Apply the doubling theorem as in Case 2 in the proof 11.8.

12.1. We can assume that the origin lies in the interior of the convex body. Consider the central projection from the surface, say Σ to the sphere \mathbb{S}^2 centered at the origin. Show that this projection $\Sigma \rightarrow \mathbb{S}^2$ is a homeomorphism.

12.4. Follow the argument in 12.2. Show that the inequality is strict if and only if F has opposite points.

12.5. Suppose a geodesic γ pass thru a vertex v . Denote by α and β the angles that γ cuts at v . Since v is essential, $\alpha + \beta < 2\pi$. Therefore $\alpha < \pi$ or $\beta < \pi$. Arrive at a contradiction by showing that γ is not length-minimizing.

12.6; (a). Cut the surface of T along three edges coming from one vertex v_1 and unfold the obtained surface on the plane. Show that this way we get a triangle, the three vertices correspond to v_1 and the midpoints of sides correspond to the remaining three vertices. Make a conclusion.

(b). Suppose that $0, v_1, v_2, v_3 \in \mathbb{R}^3$ are the vertices of T . From (a), we have that

$$|v_1| = |v_2 - v_3|, \quad |v_2| = |v_3 - v_1|, \quad |v_3| = |v_1 - v_2|.$$

Use it to show that $\langle v_1, v_2 + v_3 - v_1 \rangle = 0$. Make a conclusion.

12.8. We need to show that if a polyhedral surface is CBB(0), then the total angle θ of any of its vertex p it at most 2π .

Assume that $\theta > 2\pi$, let $\varphi = \max\{\pi, \frac{1}{3}\theta\}$. Note that we can choose three points x_1, x_2 , and x_3 close to p such that $\angle[p_{x_j}^{x_i}] = \varphi$ for $i \neq j$. Since the points x_i are close to p , we have $\angle[p_{x_j}^{x_i}] = \tilde{\angle}(p_{x_j}^{x_i})$. The latter contradicts CBB(0) comparison.

12.9. We will use that the closest-point projection from the Euclidean space to a convex body is short; that is, distance-nonexpanding [48, 13.3].

Assume K_∞ is nondegenerate. Without loss of generality, we may assume that

$$\overline{B}(0, r) \subset K_\infty \subset \overline{B}(0, 1)$$

for some $r > 0$. Note that there is a sequence $\varepsilon_n \rightarrow 0$ such that

$$K_n \subset (1 + \varepsilon_n) \cdot K_\infty \quad \text{and} \quad K_\infty \subset (1 + \varepsilon_n) \cdot K_n$$

for each large n .

Given $x \in K_n$, denote by $g_n(x)$ the closest-point projection of $(1 + \varepsilon_n) \cdot x$ to K_∞ . Similarly, given $x \in K_\infty$, denote by $h_n(x)$ the closest point projection of $(1 + \varepsilon_n) \cdot x$ to K_n . Note that

$$|g_n(x) - g_n(y)| \leq (1 + \varepsilon_n) \cdot |x - y|$$

and

$$|h_n(x) - h_n(y)| \leq (1 + \varepsilon_n) \cdot |x - y|.$$

Denote by Σ_∞ and Σ_n the surface of K_∞ and K_n respectively. The above inequalities imply

$$|g_n(x) - g_n(y)|_{\Sigma_\infty} \leq (1 + \varepsilon_n) \cdot |x - y|_{\Sigma_n}$$

for any $x, y \in \Sigma_n$, and

$$|h_n(x) - h_n(y)|_{\Sigma_n} \leq (1 + \varepsilon_n) \cdot |x - y|_{\Sigma_\infty}.$$

for any $x, y \in \Sigma_\infty$.

Note that the maps g_n and h_n are onto. Apply 8.6 to finish the proof.

Alternatively, since the closest-point projection cannot increase the length of curve, we also get

$$\begin{aligned} |x - h_n \circ g_n(x)|_{\Sigma_\infty} &\leq 10 \cdot \varepsilon_n \\ |y - g_n \circ h_n(y)|_{\Sigma_n} &\leq 10 \cdot \varepsilon_n. \end{aligned}$$

for all large n . Therefore, g_n is a δ_n -isometry $\Sigma_n \rightarrow \Sigma_\infty$ for a sequence $\delta_n \rightarrow 0$.

Comments. More generally, if a sequence of geodesic CBB(κ) spaces $\mathcal{L}_1, \mathcal{L}_2, \dots$ converges to \mathcal{L}_∞ and $\dim \mathcal{L}_n = \dim \mathcal{L}_\infty < \infty$ for any n , then $\partial \mathcal{L}_n$ equipped with induced length metric converge to $\partial \mathcal{L}_\infty$. This statement is a partial case of the theorem about extremal subsets proved by the second author [51, 1.2].

12.11; (a). By 10.15, the function $f_p: t \mapsto \text{dist}_p \circ \gamma(t)$ is semiconcave for any $p \in K$. In particular, one-sided derivatives $f_p^+(t)$ are defined for every t .

Given $x = \gamma(t)$, choose three points $p_1, p_2, p_3 \in K$ in general position; that is, the four points x, p_1, p_2, p_3 do not lie in one plane. Observe that distance functions dist_{p_i} give smooth coordinates in a neighborhood of x . From above the functions f_{p_i} have one-sided derivatives at t . Since the coordinates are smooth we get that $\gamma^+(t)$ is defined as well.

(b). If the plane py_1y_2 supports K , then $\angle[p_{y_2}^{y_1}]_{\mathbb{E}^3} = \angle[p_{x_2}^{x_1}]_\Sigma$. In this case, the statement follows from 12.10.

Now suppose that the line segment $[y_1y_2]_{\mathbb{E}^3}$ intersects K . Choose a geodesic $[y_1y_2]_W$; note that it contains a point of K , say z . Now consider one parameter family of points $y_i(t) := \gamma(t) + \gamma^+(t) \cdot (1-t) \cdot |p - x_i|_\Sigma$. Note that this family is not continuous.

Show that for any point $p \in K$ the function $t \mapsto |p - \gamma_i(t)|_{\mathbb{E}^3}$ is nonincreasing. Conclude that the function $t \mapsto |p - \gamma_i(t)|_W$ is nonincreasing for any $p \in \Sigma$. Therefore,

$$\begin{aligned} |y_1 - y_2|_W &= |y_1(0) - y_2(0)|_W = \\ &= |y_1(0) - z|_W + |y_2(0) - z|_W \geq \\ &\geq |y_1(1) - z|_W + |y_2(1) - z|_W \geq \\ &\geq |x_1 - x_2|_\Sigma. \end{aligned}$$

The last inequality follows since the closest point projection $W \rightarrow \Sigma$ is short.

It remains to consider the case when the plane py_1y_2 does not support K , and $[y_1y_2]_{\mathbb{E}^3}$ does not intersect K . In this case the plane py_1y_2 intersects K along a convex figure F that lies in the solid triangle py_1y_2 and contains its vertex p .

Choose points $y'_1 \in [py_1]_{\mathbb{E}^3}$ and $y'_2 \in [py_2]_{\mathbb{E}^3}$ such that $[y'_1y'_2]$ touches F . Denote by $x'_1 \in [px_1]_\Sigma$ and $x'_2 \in [px_2]_\Sigma$ the corresponding points; that is, $|p - y'_1|_{\mathbb{E}^3} = |p - x'_1|_\Sigma$ and $|p - y'_2|_{\mathbb{E}^3} = |p - x'_2|_\Sigma$. From above, we have that $|y'_1 - y'_2|_{\mathbb{E}^3} \geq |x'_1 - x'_2|_\Sigma$; in other words,

$$\tilde{\Delta}(p_{y'_2}^{y'_1}) \geq \tilde{\Delta}(p_{x'_2}^{x'_1});$$

here we think that $[py'_1y'_2]$ is a triangle in \mathbb{E}^3 , but $[px'_1x'_2]$ is a triangle in Σ . Note that

$$\tilde{\Delta}(p_{y'_2}^{y'_1}) = \tilde{\Delta}(p_{y_2}^{y_1}) \quad \text{and} \quad \tilde{\Delta}(p_{x_2}^{x_1}) \leq \tilde{\Delta}(p_{x'_2}^{x'_1});$$

the second inequality follows from 2.7. Hence the remaining case follows.

Comments. Part (a) was originally proved by Joseph Liberman [39]; the proof of 10.15 is a generalized version of the so-called Liberman lemma — the main tools in studying geodesics on convex surfaces.

Part (b) is a result of Anatolii Milka [41, Theorem 2].

Index

- $A \oplus B$, 38
- $[**]$, 3
- CBB comparison, 7
- \mathbb{I} , 3
- $\uparrow_{[pq]}$, 21
- ε -net, 53
- $[*^*]$, 5
- λ -concave function, 12
- $\tilde{\Delta}$, 4
- ∇ , 24
- rank \mathcal{L} (rank), 65
- $\tilde{\gamma}[x_q^p]$, 16
- $\tilde{\mathcal{Z}}$
- $\tilde{\mathcal{Z}}(*^*)$, 5
- $[***]$, 4
- $v = \log_p x$ (logarithm), 29
- adjacent hinges, 6
- affine function, 39
- Alexandrov's lemma, 8
- almost midpoint, 4, 47
- Busemann function, 37
- closed ball, 1
- comparison
 - adjacent angle comparison, 11
 - hinge comparison, 11
 - point-on-side comparison, 11
- cone, 22
- conic neighborhood, 64
- continuity method, 85
- convex body, 79
- convex polyhedron, 79
- critical point, 70
- crystallographic action, 76
- curvature, 81
- differential, 23
- differential of a function, 23
- direction, 21
- doubling, 66
- doubling theorem, 66
- essential vertex, 80, 85
- extremal point, 75
- extremal set, 78
- extremal subset, 70
- fat triangle, 11
- geodesic, 3
- geodesic direction, 21
- geodesic path, 3
- geodesic space, 3
 - π -geodesic, 22
- gradient, 24
- gradient curve, 30
- gradient exponential map, 34
- gradient flow, 33
- Gromov–Hausdorff limit, 50
- half-line, 37
- Hausdorff dimension, 48
- Hausdorff limit, 49
- hinge, 5
- hinge comparison, 11
- hyperbolic model triangle, 4
- induced length metric, 2

- invariance of domain, 86
- Jensen inequality, 12
- Lebesgue covering dimension, 48
- length metric, 2
- length space, 2
- lifting, 58, 59
- line, 38
- linear dimension, 45
- linear subspace, 43
- locally CBB(0), 15
- logarithm, 29
- maximal packing, 54
- model angle, 5
- model side, 16
- model triangle, 4
- natural map, 11
- nerve, 59
- norm, 22
- open ball, 1
- opposite vectors, 43
- orbifold, 86
- origin, 22
- pointed convergence, 51
- pointed homeomorphic, 64
- polar vectors, 41
- polyhedral space, 83
- polyhedral surfaces, 79
- primitive extremal set, 78
- proper space, 1
- properly discontinuous, 76
- rank, 65
- regular point, 70
- right derivative, 23, 29
- scalar product, 22
- semiconcave function, 23
- short map, 99
- space of directions, 21
 - of a subset, 66
 - space of geodesic directions, 21
 - spherical model triangles, 4
 - submetry, 72
 - surface, 79
- tangent space, 22
 - of a subset, 66
- tangent vector, 22
- thin triangle, 11
- triangle, 4
- triangulation, 79
- uniformly totally bounded sets, 51
- vertex, 79
- volume, 52
- Voronoi domain, 76

Bibliography

- [1] M. Aigner and G. Ziegler. *Proofs from the Book*. Any edition.
- [2] S. Alexander, V. Kapovitch, and A. Petrunin. *An invitation to Alexandrov geometry: CAT(0) spaces*. SpringerBriefs in Mathematics. 2019.
- [3] S. Alexander, V. Kapovitch, and A. Petrunin. *Alexandrov geometry—foundations*. Vol. 236. Graduate Studies in Mathematics. 2024.
- [4] А. Д. Александров. *Выпуклые многогранники*. 1950. [English translation: Alexandrov, A. D., *Convex polyhedra*, 2005].
- [5] А. Д. Александров. «Существование выпуклого многогранника и выпуклой поверхности с заданной метрикой». *Матем. сб.* 11(53).1-2 (1942), 15–65.
- [6] А. Д. Александров. *Внутренняя геометрия выпуклых поверхностей*. 1948.
- [7] A. D. Alexandrow. “Über eine Verallgemeinerung der Riemannschen Geometrie”. *Schr. Forschunsinst. Math.* 1 (1957), 33–84.
- [8] E. Bilokopytov. *Is it possible to connect every compact set?* MathOverflow. eprint: <https://mathoverflow.net/q/359390>.
- [9] E. Bruè, A. Mondino, and D. Semola. “The metric measure boundary of spaces with Ricci curvature bounded below”. *Geom. Funct. Anal.* 33.3 (2023), 593–636.
- [10] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*. Vol. 33. Graduate Studies in Mathematics. 2001. [Русский перевод: Бурого Д. Ю., Бурого Ю. Д., Иванов С. В. «Курс метрической геометрии», 2004.]
- [11] Г. Я. Перельман Ю. Д. Бурого М. Л. Громов. «Пространства А. Д. Александрова с ограниченными снизу кривизнами». *УМН* 47.2(284) (1992), 3–51. [English translation: Yu. D. Burago, M. L. Gromov, G. Ya. Perel'man, “A. D. Alexandrov spaces with curvature bounded below”, Russian Math. Surveys, 47:2 (1992), 1–58].
- [12] J. Cheeger and D. Gromoll. “The splitting theorem for manifolds of nonnegative Ricci curvature”. *J. Differential Geometry* 6 (1971/72), 119–128.
- [13] S. Cohn-Vossen. „Totalkrümmung und geodätische Linien auf einfach zusammenhängenden offenen vollständigen Flächenstücken“. *Mat. Sb.* 1(43).2 (1936).
- [14] A. Daniilidis, O. Ley, and S. Sabourau. “Asymptotic behaviour of self-contracted planar curves and gradient orbits of convex functions”. *J. Math. Pures Appl. (9)* 94.2 (2010), 183–199.

- [15] L. Danzer und B. Grünbaum. „Über zwei Probleme bezüglich konvexer Körper von P. Erdős und von V. L. Klee“. *Math. Z.* 79 (1962), 95–99.
- [16] Н. П. Долбилин. *Жемчужины теории многогранников*. 2000.
- [17] P. Erdős. “Some unsolved problems”. *Michigan Math. J.* 4 (1957), 291–300.
- [18] A. Eskenazis, M. Mendel, and A. Naor. “Nonpositive curvature is not coarsely universal”. *Invent. Math.* 217.3 (2019), 833–886.
- [19] M. Gromov. “Curvature, diameter and Betti numbers”. *Commentarii Mathematici Helvetici* 56 (1981), 179–195.
- [20] M. Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Vol. 152. 1999.
- [21] M. Gromov. “CAT(κ)-spaces: construction and concentration”. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 280 (2001), 100–140, 299–300.
- [22] K. Grove and P. Petersen. “A radius sphere theorem”. *Inventiones mathematicae* 112.1 (1993), 577–583.
- [23] S. Halbeisen. “On tangent cones of Alexandrov spaces with curvature bounded below”. *Manuscripta Math.* 103.2 (2000), 169–182.
- [24] J. Harvey and C. Searle. “Positively curved Riemannian orbifolds and Alexandrov spaces with circle symmetry in dimension 4”. *Doc. Math.* 26 (2021), 1889–1927.
- [25] A. Hatcher. *Algebraic topology*. 2002. [Русский перевод: Хатчер А. «Алгебраическая топология», 2011.]
- [26] W.-Y. Hsiang and B. Kleiner. “On the topology of positively curved 4-manifolds with symmetry”. *J. Differential Geom.* 29.3 (1989), 615–621.
- [27] T. Hu and W. A. Kirk. “Local contractions in metric spaces”. *Proc. Amer. Math. Soc.* 68.1 (1978), 121–124.
- [28] V. Kapovitch. “Perelman’s stability theorem”. *Surveys in differential geometry. Vol. XI*. Vol. 11. Surv. Differ. Geom. Int. Press, Somerville, MA, 2007, 103–136.
- [29] A. P. Kiselev. *Kiselev’s Geometry: Stereometry*. 2008.
- [30] K. W. Kwun. “Uniqueness of the open cone neighborhood”. *Proc. Amer. Math. Soc.* 15.3 (1964), 476–479.
- [31] U. Lang and V. Schroeder. “On Toponogov’s comparison theorem for Alexandrov spaces”. *Enseign. Math.* 59.3-4 (2013), 325–336.
- [32] N. Lebedeva. “Alexandrov spaces with maximal number of extremal points”. *Geom. Topol.* 19.3 (2015), 1493–1521.
- [33] N. Lebedeva, A. Petrunin, and V. Zolotov. “Bipolar comparison”. *Geom. Funct. Anal.* 29.1 (2019), 258–282.
- [34] N. Lebedeva and A. Petrunin. “5-point CAT(0) spaces after Tetsu Toyoda”. *Anal. Geom. Metr. Spaces* 9.1 (2021), 160–166.
- [35] N. Lebedeva and A. Petrunin. “Graph comparison meets Alexandrov”. *Sib. Math. J.* 64.3 (2023). Translation of *Sibirsk. Mat. Zh.* **64** (2023), no. 3, 579–584., 624–628.
- [36] N. Lebedeva and A. Petrunin. “Five-point Toponogov theorem”. *Int. Math. Res. Not. IMRN* 5 (2024), 3601–3624.

- [37] N. Lebedeva and A. Petrunin. “Trees meet octahedron comparison”. *Journal of Topology and Analysis* (July 2023), 1–5.
- [38] N. Lebedeva. “On open flat sets in spaces with bipolar comparison”. *Geom. Dedicata* 203 (2019), 347–351.
- [39] И. М. Либрман. «Геодезические линии на выпуклых поверхностях». 32.2 (1941), 310–312.
- [40] K. Menger. „Untersuchungen über allgemeine Metrik“. *Math. Ann.* 100.1 (1928), 75–163.
- [41] А. Д. Милка. «Геодезические и кратчайшие линии на выпуклых гиперповерхностях. I». *Украин. геом. сб.* 25 (1982), 95–110.
- [42] G. Perelman. “Spaces with curvature bounded below”. *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*. 1995, 517–525.
- [43] G. Perelman and A. Petrunin. “Extremal subsets in Aleksandrov spaces and the generalized Liberman theorem”. *Algebra i Analiz* 5.1 (1993), 242–256.
- [44] G. Ya. Perelman. “Elements of Morse theory on Aleksandrov spaces”. *St. Petersburg Math. J.* 5.1 (1994), 205–213.
- [45] G. Perelman. *Alexandrov spaces with curvatures bounded from below II*. 1991.
- [46] Г. Я. Перельман. «Начала теории Морса на пространствах Александрова». *Алгебра и анализ* 5.1 (1993), 232–241.
- [47] A. Petrunin and S. Stadler. “Metric-minimizing surfaces revisited”. *Geom. Topol.* 23.6 (2019), 3111–3139.
- [48] A. Petrunin and S. Zamora Barrera. *What is differential geometry: curves and surfaces*. 2021. arXiv: 2012.11814 [math.HO].
- [49] A. Petrunin. “A globalization for non-complete but geodesic spaces”. *Math. Ann.* 366.1-2 (2016), 387–393.
- [50] A. Petrunin. “Semiconcave functions in Alexandrov’s geometry”. *Surveys in differential geometry. Vol. XI*. Vol. 11. Surv. Differ. Geom. 2007, 137–201.
- [51] A. Petrunin. “Applications of quasigeodesics and gradient curves”. *Comparison geometry*. 1997, 203–219.
- [52] А. М. Петрунин. “Верхняя оценка на интеграл кривизны”. *Алгебра и анализ* 20.2 (2008), 134–148. [Translation: Petrunin, A. M., An upper bound for the curvature integral, *St. Petersburg Math. J.* 20(2009), no.2, 255–265.]
- [53] A. Petrunin. “In search of a five-point Aleksandrov type condition”. *Algebra i Analiz* 29.1 (2017), 296–298.
- [54] A. Petrunin. *Pure metric geometry*. SpringerBriefs in Mathematics. 2023.
- [55] P. Pizzetti. “Paragone fra due triangoli a lati uguali”. *Atti della Reale Accademia dei Lincei, Rendiconti (5). Classe di Scienze Fisiche, Matematiche e Naturali* 16.1 (1907), 6–11.
- [56] C. Plaut. “Spaces of Wald curvature bounded below”. *J. Geom. Anal.* 6.1 (1996), 113–134.
- [57] C. Plaut. “Metric spaces of curvature $\geq k$ ”. *Handbook of geometric topology*. 2002, 819–898.
- [58] А. В. Погорелов. *Однозначная определённость общих выпуклых поверхностей*. Монографии института математики, вып. II. 1952.

- [59] K. Shiohama. *An introduction to the geometry of Alexandrov spaces*. Vol. 8. Lecture Notes Series. 1993.
- [60] D. Fuchs and S. Tabachnikov. *Mathematical omnibus*. Thirty lectures on classic mathematics. 2007. [Русский перевод: Табачников, С. Л., Фукс, Д. Б. «Математический дивертисмент» 2016].
- [61] В. А. Топоногов. «Римановы пространства кривизны, ограниченной снизу». *Успехи математических наук* 14.1 (85) (1959), 87–130.
- [62] T. Toyoda. “An intrinsic characterization of five points in a CAT(0) space”. *Anal. Geom. Metr. Spaces* 8.1 (2020), 114–165.
- [63] Ю. А. Волков. «Существование многогранника с данной разверткой». *Зап. научн. сем. ПОМИ* (2018), 50–78.